

# Jolts in the Jade factory: A route for subduction fluids and their implications for mantle wedge seismicity

Samuel Angiboust, Jesus Muñoz-Montecinos, Aitor Cambeses, Tom

Raimondo, Damien Deldicque, Antonio Garcia-Casco

## ▶ To cite this version:

Samuel Angiboust, Jesus Muñoz-Montecinos, Aitor Cambeses, Tom Raimondo, Damien Deldicque, et al.. Jolts in the Jade factory: A route for subduction fluids and their implications for mantle wedge seismicity. Earth-Science Reviews, 2021, 220, 10.1016/j.earscirev.2021.103720. insu-03590019

## HAL Id: insu-03590019 https://insu.hal.science/insu-03590019

Submitted on 2 Aug 2023  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1	Jolts in the Jade Factory:
2	a Route for Subduction Fluids and
3	their Implications for Mantle Wedge Seismicity
4	
5	Samuel Angiboust <sup>1</sup>
6	Jesus Muñoz-Montecinos <sup>1,2</sup>
7	Aitor Cambeses <sup>1,2</sup>
8	Tom Raimondo <sup>3</sup>
9	Damien Deldicque <sup>4</sup>
10	Antonio Garcia-Casco <sup>2,5</sup>
11	
12 13	<sup>1</sup> Université de Paris, Institut de Physique du Globe de Paris, F-75005 Paris, France (* corresponding author: angiboust@ipgp.fr)
14 15	<sup>2</sup> Department of Mineralogy and Petrology, Faculty of Sciences, University of Granada, Campus Fuentenueva s/n, 18002 Granada, Spain
16	<sup>3</sup> UniSA STEM, University of South Australia, GPO Box 2471, Adelaide SA 5001, Australia
17	<sup>4</sup> Laboratoire de Géologie, Ecole Normale Supérieure PSL, F-75005 Paris, France
18 19	<sup>5</sup> Instituto Andaluz de Ciencias de la Tierra, CSIC-Universidad de Granada, 18100 Armilla, Granada, Spain.
20	
21	
22	Keywords
23	Subduction, Fluids, Jadeitite, Earthquakes, Mantle wedge, Foliated cataclasites

### 24 Abstract

25 An increasing number of seismological studies report transient seismicity clusters in the mantle wedge 26 several kilometers above the subduction interface. Their physical significance with respect to subduction 27 zone seismo-tectonics remains poorly understood. Jadeitites are known to form and/or be associated 28 with mantle wedge serpentinites in the c. 30-70 km depth range, and thus may yield information on 29 deformation mechanisms in this region of deep subduction environments. We herein document and 30 compare brittle-viscous features recorded in jadeitites from Polar Urals (Russia), Kashin state (Myanmar) 31 and Motagua fault region (Guatemala) - some of the most important jadeitite occurrences worldwide. In 32 the Polar Urals we identified ultramafic-hosted pristine jadeitite-bearing veins c. 1 km above a Devonian 33 paleo-subduction interface, interpreted as metasomatized former felsic dyke networks crosscutting the 34 mantle wedge peridotites. Here, both jadeitites and associated amphibole-rich dark granofels display 35 widespread brittle-ductile deformation fabrics such as shear bands, foliated cataclasites and breccias, 36 cemented through dissolution-precipitation processes by omphacite and sodic amphiboles, a mineral 37 assemblage typical of high-pressure-low-temperature subduction zone conditions. Electron probe and 38 laser ablation ICP-MS mapping indicate that these brittle-viscous networks display a substantial 39 metasomatic imprint highlighted in the dark granofels by variations in major and trace elements. 40 Switches between viscous and brittle deformation patterns are attested by crystallographic-preferred 41 orientations of jadeite in some of the shear zones that crosscut the host jadeitites. Strikingly similar 42 mineral assemblages and deformation patterns were observed in the Kashin and Motagua samples. 43 Observed deformation features in these localities can be classified into three categories (tectonic 44 breccias, foliated cataclasites and hydraulic breccias), which may occasionally form in sequence and 45 exhibit mutually overprinting textures. Some of the foliated cataclasites contain fine-grained and foliated 46 "shard-like" features forming a radial omphacite-jadeite spherulitic texture, interpreted as former 47 pseudotachylyte that evokes a paleo-seismic origin. We interpret these healed fault networks as

recording external fluid influx within fracture zones that repeatedly ruptured along former "dyke" networks. These high permeability drains likely (i) contribute to the transfer of highly pressurized plateinterface metamorphic fluids into the mantle wedge; and (ii) trigger seismic instabilities recorded in the basal part of active mantle wedge sections. These findings provide new insights into the current understanding of the rheology (e.g., serpentinization ratio) and stress state in the mantle wedge, with implications for subduction interface seismogenesis.

#### 54 **1. Introduction**

55 Understanding processes rooted in the mantle wedge region of subduction zones is of critical 56 importance because its structure strongly controls the rheology of the subduction interface at the 57 downdip end of the seismogenic zone (Hyndman & Peacock, 2003; Dessa et al., 2009; Agard et al., 2018). 58 Fluids released by metamorphic reactions in the downgoing plate are thought to control the degree of 59 hydration (serpentinization) of the fore-arc mantle wedge (e.g., Hacker et al., 2003; Deschamps et al., 60 2010; Bostock, 2013). The bulk serpentinization ratio of the mantle wedge, estimated based on the 61 perturbation of seismic wave velocities, is generally considered low (i.e., <20 vol.%; Abers et al., 2017) 62 except for specific subduction environments where it may reach up to 60 vol.% (e.g., Central Japan: 63 Hyndman & Peacock, 2003; Mariana margin: Hussong, 1981). The serpentinized mantle wedge has long 64 been considered as mostly aseismic due to the presence of weak minerals such as antigorite or talc that 65 are known to substantially reduce rock strength (e.g., Hilairet et al., 2007). Yet, insights gained from 66 detailed studies on the Sumatra 2004 (M<sub>w</sub> = 9.1) and Maule 2010 (M<sub>w</sub> = 8.8) mega-earthquakes have 67 shown that some of these exceptional rupture events may nucleate or propagate at the base of the "cold 68 nose" region (e.g., Dessa et al., 2009; Wang et al., 2020). This paradox raises important questions 69 regarding the rheology of the partly serpentinized mantle above the subduction interface. The report of 70 slow earthquakes near the seismogenic downdip end of several subduction megathrusts also changed 71 our vision regarding stress distribution along the deep interface (e.g., Fu & Freymuller, 2013; Frank et al.,

72 2015; Audet & Kim, 2016), with potential implications for mega-earthquake prediction (Obara & Kato, 73 2016; Bouchon et al., 2018). Whereas it is now accepted that fluids dramatically impact the mechanical 74 stability of the deep serpentinized interface, very little *in situ* information is known about fluid-rock 75 interaction processes or the feedback between upward transported fluids and seismicity (e.g., Angiboust 76 et al., 2014; Locatelli et al., 2018).

77 Over the last decade, an increasing number of high-resolution seismological studies have 78 identified the presence of seismicity nests in the partly serpentinized mantle wedge (e.g., Halpaap et al., 79 2019 and references therein). These events are generally interpreted as mechanically related to the 80 influx of fluids or melts between 30 and 70 km depth (e.g., Davey & Ristau, 2011). A majority of the 81 clusters identified in literature concentrates in the first 15 km above the plate interface (e.g., Greece: 82 Halpaap et al., 2019; Japan: Nakajima & Uchida, 2018; New Zealand: Davey & Ristau, 2011; Central Chile: 83 Wang et al., 2020; Colombia: Chang et al., 2019) along steeply-dipping, planar features commonly seen as "vent-like" structures, apparently channelizing plate-interface fluids towards the inner wedge. 84 85 Normal, strike-slip and thrust focal mechanisms are reported for a large majority of these supra-slab 86 earthquakes, with magnitudes generally spanning a range between 2 and 5 (e.g., Halpaap et al., 2019). 87 The physics of the rupture and the nature of the material where these earthquakes are nested remain 88 unknown. Moreover, the source and the composition of the fluids passing through these networks is also a matter of discussion since the precise location of hydrous mineral breakdown reactions strongly 89 90 depends on the subduction thermal structure (e.g., Hermann et al., 2006; Syracuse et al., 2010).

Natural mantle wedge samples, despite their scarcity worldwide, represent a unique opportunity
to shed light on these deep-seated processes (e.g., Kepezhinskas et al., 1995; Horn et al., 2020).
Jadeitites are known to represent fossilized fluid pathways from the base of the hydrated mantle wedge
(Harlow & Sorensen, 2005; Harlow et al., 2015). From this perspective, they may provide insights on the
physical nature of supra-slab seismic events. However, primary structures from jadeitite-bearing

localities have been almost systematically overprinted during long-term subduction, extensive 96 97 serpentinization and exhumation (e.g., Central America: Flores et al., 2015; Kawamoto et al., 2018; 98 Myanmar: Shi et al., 2009a; Japan: Morishita et al., 2007). In the Polar Urals (Russia), the Pus'yerka locality exhibits a relatively undisturbed contact between a jadeitite "dyke" network and its ultramafic 99 100 host (e.g., Meng et al., 2011; Angiboust et al., 2021), thus providing an opportunity to identify 101 deformation processes rooted in these jadeitite bodies. Through a combined petrological, 102 microstructural and geochemical investigation of Polar Urals samples, we provide new evidence for 103 brittle-ductile switches in jadeitites (and associated amphibole-phlogopite granofels) microstructures. 104 These structures are compared with those from samples of loose jadeitite boulders from Myanmar and 105 Guatemala, settings where pristine structures are only exceptionally exposed (e.g., Sorensen et al., 106 2010). We then evaluate their potential meaning in terms of fluid pathways and the genesis of seismic 107 instabilities in the basal region of active mantle wedges.

108

#### 2. Geological setting

## 109

#### 2.1 Pus'yerka jadeitite deposit (Polar Urals, Russia)

110 The Polar Urals belt formed during closure of the Uralian ocean by subduction and by the 111 eastward burial of the European continental margin under an oceanic volcanic arc (e.g., Savelieva et al., 112 2002). In the Polar Urals, the Main Ural Thrust (MUT; Fig.1a) corresponds to a major crustal-scale shear zone with peridotites that were thrust over eclogitized continental crust (Marun-Keu complex; e.g., 113 114 Udovkina, 1971; Dobretsov & Sobolev, 1984; Glodny et al., 2003). Rare ophiolitic, blueschist-facies 115 mélange exposures are restricted to the base of the MUT (e.g., Kazak et al., 1976). Locally, the MUT has 116 been subject to moderate reactivation as a detachment fault during exhumation (e.g., Sychev & Kulikova, 117 2012). In the Polar Urals,



Figure 1. a. Simplified geological map of the Polar Urals locating the three main mantle wedge sections 119 exposed in this region and the study area at the base of the Syum-Keu massif. The inset localizes the Polar 120 121 Urals in northern Russia. Green dots correspond to jadeitite-bearing localities (modified after Angiboust et 122 al., 2021). b. Geological map showing the structures above the Main Ural Thrust, interpreted as an ancient subduction interface. The studied jadeitite "dyke" crops out as a boudinaged sliver wrapped by a network 123 124 of sheared serpentinites, approximately 1 km above the Main Ural Thrust. c. Cross-section highlighting the 125 geometry of the structures identified in the field (modified after Angiboust et al., 2021). d. Pressure-126 Temperature-time sketch summarizing the long-term evolution of the Pus'yerka jadeitite dyke structure (after Angiboust et al., 2021). Jd<sub>50</sub>Di<sub>50</sub> reaction line is after Maruyama & Liou (1988). Reaction (1): 127 Analcime = Albite + Nepheline. e. Schematic view of the white jadeitite "dyke" as observed in situ in the 128 129 Polar Urals with dismembered patches of dark granofels and dark amphibole-phlogopite bearing 130 blackwalls forming at the contact with the host serpentinized ultramafic rocks. Mineral abbreviations after 131 Whitney & Evans (2010).

132

133 three peridotite massifs (the Rai-Iz, Syum-Keu and Voikar massifs) were locally transformed into 134 antigorite-schists along the MUT hangingwall (Fig.1b). These large mantle exposures, mostly harzburgitic 135 and Iherzolitic in composition (Savelieva & Suslov, 2014; Shmelev et al., 2011), exhibit large chromite 136 deposits and are crosscut by numerous subvertical dunitic channels interpreted as the melt extraction 137 pathways (Batanova et al., 2011). The MUT hanging wall displays (i) a serpentinization gradient towards 138 the underlying unit; (ii) the presence of jadeite (NaAlSi $_2O_6$ ) veins in the serpentinites from the hanging 139 wall; and (iii) a high pressure-low temperature (HP-LT) metamorphic imprint in the footwall units (e.g., 140 Dobretsov & Ponomareva, 1968; Glodny et al., 2003; Batanova et al., 2011; Shmelev, 2011; Meng et al., 141 2016). The Rai-Iz, Syum-Keu and Voikar peridotite massifs in the Polar Urals can thus be viewed as good 142 analogues of a supra-subduction setting, enabling the understanding of deep-seated processes below an 143 inferred Paleozoic island arc (e.g., Batanova et al., 2011; Savelieva et al., 2002, 2016; Angiboust et al., 144 2021), in a region of the deep subduction interface that is not commonly exhumed (Guillot et al., 2009; 145 Agard et al., 2018).

The studied Pus'yerka exposure is located along a jadeitite-bearing serpentinized shear zone that is several hundred meters thick and 5 km long, striking N-S and dipping E approximately 1 km above the Main Ural Thrust (MUT; **Fig.1b**). In a recent study, Angiboust et al. (2021) have interpreted the MUT hanging wall as a rare natural case study highlighting the structure along the base of a subduction mantle wedge. This major jadeitite deposit, discovered and mined in the 1980s (Kuznetsov et al., 1986; Fishman, 2006), represents a unique locality to investigate the structural contacts between the jadeite bodies and its host (e.g., Meng et al., 2011; Angiboust et al., 2021).

The basal serpentinites as well as the jadeitite-bearing network exhibit a regional foliation parallel with the MUT (**Fig.1c**). Field, geochemical and density measurements indicate serpentinization ratios in the range of 35-65 vol.% (in agreement with the 45-65 vol.% estimates from Makeyev, 1992), with up to 90-100% approaching the MUT and within the jadeite-bearing networks (**Fig.1b**; Angiboust et al., 2021 and references therein). The serpentinite schists that host the jadeitite boudins are composed of antigorite with minor amounts of brucite, magnetite and phlogopite (Makeyev, 1992) as well as chlorite, tremolite and magnesite. Within the serpentinite shear zone, tens of whitish jadeitite lenses are observed, elongated *en echelon* and stretched parallel to the main foliation dip direction (**Fig.1b**). The thickness of these lenses ranges between several tens of centimeters to several meters in the few places where the lenses were observed *in situ* (e.g., Kuznetsov et al., 1986; Meng et al., 2011).

163 Ion probe U-Pb dating of zircon crystals from the main jadeitite body yielded U-Pb ages of  $404 \pm 7$ 164 Ma (Meng et al., 2011) and 409 ± 3.3 Ma (Konovalov & Sergeev, 2015). These Devonian ages were 165 interpreted by Meng et al. (2011) as dating intra-oceanic subduction initiation of the Uralian ocean 166 realm. Angiboust et al. (2021) proposed that these jadeitite bodies derive from the metasomatic replacement of a former trondhjemitic dyke that crystallized from slab-derived melts within an 167 168 ultramafic mantle wedge setting, in a subduction initiation context (i.e., at a temperature regime much 169 higher than expected in a long-lived subduction context; e.g., Soret et al., 2016 and references therein). 170 Ar-Ar plateau ages and multi-mineral Rb-Sr dating yield ages for phlogopite and amphibole-bearing 171 domains ranging from 410–395 Ma, interpreted by Angiboust et al. (2021), in line with the pioneering study of Dobretsov & Ponomareva (1968), as marking the re-equilibration of dyke material during secular 172 cooling from supra-solidus to HP-LT conditions (i.e., from  $T > 700^{\circ}C$  to  $T < 500^{\circ}C$  for a pressure on the 173 174 order of 1.5 GPa). The studied samples were collected in the main jadeitite "dyke" (Fig.1b,e).

175

## 2.2 Kashin state Jade Mines Belt (northern Myanmar)

One of the world's largest jadeitite-bearing suture zones is exposed in the Kashin state of Myanmar in the famous Jade Mines Belt (e.g., Shi et al., 2012; Nyunt et al., 2017), where loose jadeitite fragments are found in conglomerates, river beds or exceptionally embedded within strongly-weathered 179 antigorite schists (e.g., Harlow et al., 2015 and references therein; Ridd et al., 2019). The original 180 jadeitite-bearing structures (likely ancient felsic dykes; e.g., Bleeck, 1908) were formed within a 181 serpentinized mantle wedge from a subduction zone of debated Late Jurassic to Late Cretaceous age (Goffé et al., 2002; Shi et al., 2009a, 2012; Yui et al., 2012; Harlow et al., 2015). These metasomatized 182 183 dykes were extensively affected by exhumation and subsequent strike-slip deformation related to the Sagaing transform fault system (Harlow et al., 2015; Searle et al., 2017; Ridd et al., 2019). Protracted 184 185 deformation led to the formation of a serpentinite mélange in which jadeitite "dykes" and blocks were 186 disrupted and disseminated in the antigorite schist matrix, together with other lenses of seafloor origin 187 such as graphite schists, glaucophane schists, garnet amphibolites and garnet micaschists (Nyunt et al., 2017). 188

189 Although the original thickness of the jadeitite "dykes" is challenging to evaluate due to poor 190 exposure conditions and late deformation, some studies mention typical thicknesses on the order of 191 several meters (Shi et al., 2012, Harlow et al., 2015 and references therein), in line with the structures 192 observed in situ in the Polar Urals (Fig.1e). Texturally secondary chlorite schists ("blackwalls") as well as 193 albitite bands are reported at the contact between the dyke structure and the host (Bleeck, 1908; 194 Chhibber, 1934). Na-amphibole-rich bands (mostly eckermannite and glaucophane; see Oberti et al., 195 2015 for further details on mineralogy) are also found either embedded within the jadeitite "dyke" or 196 lining the contact with the ultramafic host (Bleeck, 1908; Nyunt, 2009). Most pressure-temperature 197 estimates for jadeitite formation in the Jade Mine Belt region span a wide range from 1.0-1.5 GPa and 198 300–500°C (Mével & Kiénast, 1986; Goffé et al., 2002; Shi et al., 2003). The herein studied samples, 199 provided by a local miner, were found as boulders in a conglomerate near the Lonkin township (near 200 Hpakan).

201

2.3 Motagua fault zone (Guatemala)

202 The Guatemala suture zone is an E-W-trending major plate boundary zone that separates the 203 Caribbean and North American plates. This strike-slip suture, which puts in contact the Maya block to the 204 north with the Chortis block to the south (e.g., Ortega-Gutierrez et al., 2007 and references therein), 205 contains many mafic and ultramafic blocks as well as eclogite-, blueschist- and garnet amphibolite-facies 206 crustal and sedimentary fragments interpreted as remnants from a Cretaceous meta-ophiolite (e.g., 207 Brueckner et al., 2009). The Motagua fault zone separates two distinct terranes likely exhumed during 208 two distinct collisional events: the North Motagua and the South Motagua mélanges (e.g., Beccaluva et 209 al., 1995; Gendron et al., 2002; Harlow et al., 2004; Harlow, 2011). Different metamorphic ages on 210 jadeitites and eclogites were reported for these two terranes, ranging from c. 100–60 Ma for the North Motagua and c. 160–110 Ma for the South Motagua mélanges (see Flores et al., 2013 and references 211 212 therein). The highest-pressure rocks from these two mélanges are also slightly different, with 500–650 °C 213 and 1.5–2.3 GPa for the North Motagua mélange (Harlow et al., 2008; Tsujimori et al., 2004) and 470– 214 520 °C and 2.0–2.7 GPa (Tsujimori et al., 2006; Endo et al., 2012) for the South Motagua mélange.

215 Jadeitites occur as meter-sized blocks within the serpentinite mélanges but are most commonly 216 found as pebbles within streams or as loose blocks in slope debris. As for the Jade Mines Belt in 217 Myanmar, pristine tectonic relationships are extremely difficult to document. The mineralogy of jadeitites is quite varied in terms of minerals and mineral abundances, and with contrasting assemblages 218 219 in the north and south Motagua mélanges (Harlow et al., 2011; Flores et al., 2013). The herein studied 220 samples were collected as loose boulders in streams from the southern Motagua mélange (Rio El Tambor 221 area) and contain essentially jadeite plus minor omphacite and lawsonite, in agreement with the 222 mineralogy of jadeitites from this mélange (Harlow et al., 2011).

#### 223 3. Analytical methods

#### 224 3.1 Electron probe microanalysis

225 Mineral compositions were quantified via electron probe microanalysis (EPMA) using a Cameca 226 SXFive operated in the CAMPARIS analytical facility at Paris University. Standard analytical conditions (15 227 keV, 10 nA, beam diameter 5  $\mu$ m) and a set of synthetic and natural crystals for calibration standards 228 were used: Fe<sub>2</sub>O<sub>3</sub> (Fe), MnTiO<sub>3</sub> (Mn, Ti), diopside (Mg, Si), CaF<sub>2</sub> (F), orthoclase (AI, K), anorthite (Ca) and 229 albite (Na). X-ray maps were acquired on the same instrument using analytical conditions of 15 keV, 250 230 nA, a dwell time of 60 milliseconds and a step size of 2 µm. Some of the X-ray images were processed 231 and quantified with DWImager software (Torres-Roldán and Garcia-Casco, unpublished; see Garcia-232 Casco, 2007). A scanning electron microscope (SEM) Zeiss EVO MA10 at the Institut de Physique du 233 Globe de Paris using internal calibration standards was used for microscopic investigations, energy dispersive X-ray spectral (EDS) mapping and surface composition characterizations. Mineral 234 abbreviations are from Whitney & Evans (2010). Clinopyroxene and clinoamphibole compositions, 235 236 including estimation of Fe<sup>3+</sup>, are calculated according to the schemes of Morimoto (1989) and 237 Hawthorne & Oberti (2007), respectively. Classification of these minerals also follows the same authors.

#### 238 3.2 Laser Ablation Inductively Coupled Plasma Mass Spectrometry

239 Laser Ablation Inductively Coupled Plasma Mass Spectrometry (LA-ICP-MS) trace element maps 240 were acquired using a Resonetics M-50-LR 193 nm excimer laser coupled to an Agilent 7700x 241 Quadrupole ICP-MS housed at Adelaide Microscopy, University of Adelaide. Instrument conditions and 242 mapping protocols similar to that employed in this study are outlined in Raimondo et al. (2017). Pre-243 ablation of each raster scan was completed to minimize the effect of redeposition (19 µm, 75% overlap), 244 followed by 15 s washout and 10 s of background measurement. A beam diameter of 19  $\mu$ m, line spacing 245 of 19 µm and repetition rate of 10 Hz were employed for sample PU2, resulting in an energy density of 246 3.5 J/cm<sup>2</sup> at the target. Standards were analyzed in duplicate every 2 h during the mapping session, including reference glasses NIST 610 (Pearce et al., 1997; Jochum et al., 2011a) and GSD-1D (Jochum et 247 248 al., 2011b). A beam diameter of 51  $\mu$ m was used for all standard analyses, and included 5 pre-ablation

shots (51 µm, 75% overlap) followed by 20 s washout, 30 s background measurement and 40 s ablation
time. Data acquisition was performed in time-resolved analysis mode as a single continuous experiment.
Each analysis comprised a suite of 38 elements, and dwell times were as follows: 0.01 s (Li), 0.002 s (Na,
Mg, Al, Si, K, Ca, Mn, Fe, Ni), 0.005 s (Sc, Ti, V, Cr, Nb, Ba, Hf, Th, U), and 0.008 s (Sr, Y, Zr, Ta, Pb, REEs).
The total sweep time was 0.297 s. Post-acquisition processing was performed using the software lolite
(Woodhead et al. 2007; Hellstrom et al. 2008; Paton et al. 2011), with data reduction and image
processing procedures following those outlined by Raimondo et al. (2017) and Hyppolito et al. (2018).

#### 256 3.3 Electron Back-Scattered Diffraction and cathodoluminescence mapping

257 Electron Back Scattered Diffraction mapping (EBSD) has been performed at the Laboratoire de 258 Géologie of the Ecole Normale Supérieure of Paris using a ZEISS SIGMA Field Emission Gun Scanning 259 Electron Microscope equipped with an EDX (Energy Dispersive X-ray Spectroscopy; X-MAX) and an EBSD 260 detector (Nordlys Nano, Oxford Instruments). An acceleration voltage of 15 keV, a beam current of 5 nA, 261 an aperture of 120 µm, an inclination of 70°, an acquisition rate of 100 Hz, a working distance of 14 mm 262 and a mapping step size of 2.8 µm were the analytical parameters chosen for the mapping. Data 263 acquisition, post-processing treatment and statistical analysis were performed using Aztec, Channel 5 264 and MTEX software (Bachmann et al., 2010; Bachmann et al., 2011). For noise reduction, every single-265 pixel isolated data point was removed and followed by denoising MTEX procedures. Cathodoluminescence (CL) mosaic images were acquired using the Cathodyne (NEWTEC) device 266 267 equipped with a motorized stage, a 12kV and 120  $\mu$ A plasma, and 2 seconds of image acquisition time.

268

## 4. Field constraints on Polar Urals jadeitite body

In the Pus'yerka locality of the Polar Urals, the ore jadeitite body is mostly formed by a white jadeitite core that represents more than 90% of the "dyke" volume (see schematic dyke structure in **Fig.1e**). Locally, remnants of felsic lithologies (comprising an albitite groundmass with paragonite flakes 272 surrounded by a jadeite-bearing corona) were observed, suggesting that the white jadeitite formed by 273 replacement of a leucocratic dyke (see the model in Angiboust et al., 2021; see also Dobretsov & 274 Ponomareva, 1968 and Kuznetsov et al., 1986). The whitish jadeitite-rich domains comprise weakly to 275 strongly foliated mafic blocks that host millimeter-sized intricate amphibole and phlogopite crystals 276 (sample PU2). These domains, referred hereafter to as dark granofels (sample UR11b), were interpreted 277 by Angiboust et al. (2021) to be produced by the influx of alkali-rich fluids in a warm mantle wedge 278 environment, before the emplacement of the leucocratic dyke. Such amphibole-rich blocks occupying a 279 similar structural position were reported in the Jade Mine Belt area (Myanmar) by Bleek (1908; see also 280 Harlow et al., 2015).

281 The white jadeitite is transected by centimeter to decimeter-long cracks filled by phlogopite as well 282 as dark-blueish amphibole-rich domains and emerald-green Cr-rich clinopyroxene (Fig.2a, b). While the 283 bulk of the white jadeitite mass looks at a first sight microstructurally homogeneous, detailed 284 observations highlight numerous locations where structures are brecciated, ranging from crackle- to 285 mosaic-type breccias (e.g., Woodcock & Mort, 2008). Chaotic breccias, where substantial disruption of 286 the original structure occurred, are also observed in jadeitites and associated dark granofels (Fig.2b). 287 Lastly, centimeter- to decimeter-thick, strongly sheared phlogopite-rich metasomatic rinds are observed 288 along the margins of the jadeitite body at the contact with the host serpentinite (Fig.1e; see also Kuznetsov et al., 1986 and Angiboust et al., 2021). Further south, directly below the Rai-Iz peridotite 289 290 massif, occurs a tectonic mélange zone ("Nephrite brook", Kazak, 1978; Fig.1a) that contains blocks of 291 nephrite and rare jadeitite within a schistose serpentinite matrix. Sample UR25 represents one of these 292 jadeitite pods, considered to be derived from a former "dyke" structure that has been fully disrupted by 293 late subduction zone tectonic deformation.



Figure 2. a. Field photo of a brecciated white jadeitite block, showing a discrete crackle breccia texture
transected by a hydrofracture filled with green Cr- rich jadeite. b. Field photo highlighting the structure of
a mosaic breccia, with disrupted white jadeitite fragments cemented by a Cr-rich jadeitic clinopyroxene.

The nomenclature used in Table 1 is defined based on the following deformation criteria: Type (I) 298 299 corresponds to breccias formed by centimeter- to tens of centimeter-sized clasts exhibiting a substantial 300 shearing component, fracturing and size reduction through indentation processes. Such breccias can typically reflect a damage zone deformation pattern. Type (II) brecciated materials are defined as a 301 302 highly localized fault zone with evidence of pervasive shearing, grain comminution and flow banding, 303 with pulverized wall clasts floating in a fine-grained matrix. Such microstructures are generally known in the vicinity of fault cores and are hereafter termed "foliated cataclasites". Type (III) corresponds to 304 305 hydraulic breccias with millimeter to centimeter-sized clasts (which may have undergone rotation) cemented by clinopyroxene or amphibole. Space-filling material can be either dendritic, oscillatory, or 306 307 strained.

308

## **5.** Structure of Polar Urals jadeitites

The bulk of the white jadeitite matrix is formed by idiomorphic, oscillatory and intricate 100–500 μm long jadeite crystals (Meng et al., 2011). Their outer rims are commonly lined by Ca-rich, omphacitic

311 compositions as well as interstitial phlogopitic micas (Angiboust et al., 2021; Fig.S1; Table 1 and Table 312 **S1**). Despite an apparently homogeneous macroscopic texture, detailed petrographic investigations 313 reveal that jadeite crystals exhibit widespread fracturing, dissolution and replacement textures 314 (Fig.3a,b). Vein systems that range in color from white (jadeite composition) to green (Cr-rich jadeite or 315 Cr-rich omphacite; Fig.2a) are ubiquitously found crosscutting the host jadeitite matrix (see also Franz et 316 al, 2014). These veins are filled by idiomorphic to fibrous clinopyroxene crystals with locally oscillatory 317 and/or dendritic textures (Kuznetsov et al., 1986; sample UR03b in Angiboust et al., 2021: Type III). 318 Clinopyroxene crystals from brecciated samples exhibit clear evidence for pressure-solution (as shown by 319 indentation and truncation textures), as well as solution-precipitation with fracture healing and 320 overgrowth by a new (generally more omphacitic) clinopyroxene composition (Fig.3a,b; Fig. S2).



322 Figure 3. Backscattered electron (BSE) images for Polar Urals samples. a. White jadeitite specimen (PU2) 323 showing highly disrupted clasts with evidence for fracturing, dissolution, indentation and re-precipitation. 324 Several clinopyroxene generations can be identified with increasing diopside component towards clast 325 rim. b. Example of a cryptic jadeitite breccia showing extensive fracturing, comminution and dissolution of 326 dark white jadeitite clasts (locally exhibiting oscillatory zoning pattern). Location of sample UR25 given in 327 figure 1a. c. Dark, amphibole-rich granofels that is heavily brecciated, with similar dissolution-328 reprecipitation features as well as extensive healed microfractures (sample PU2). d. Fractures associated 329 with the growth of several amphibole compositions such as richterite (Rct) and eckermannite (Eck) 330 overgrowing magnesio-katophorite (Mkt). Note that omphacite grows lately within a crack in textural 331 equilibrium with Rct and Eck (sample UR11b).

332

333 The studied white jadeitite sample UR03d exhibits coarse-grained, idiomorphic and sector-zoned jadeite 334 crystals in its matrix (Fig.4a). These crystals, generally extremely rich in jadeitic molecule (>90 mol.%), 335 are hereafter referred to as  $Jd_1$ . The jadeite growth structures document how early structures have been 336 transected by a localized fine-grained shear zone of oriented jadeite crystals with interstitial omphacite-337 rich grains. Orientation maps (Fig.4b) show that [001] axes are dominant along the fine-grained shear 338 zone, whereas [100] and [010] axes become important approaching the margins of the shear band. 339 Intracrystalline misorientation EBSD maps, depicting misorientation angles between each data point and 340 the mean orientation of the parent crystal (Fig. S3), show that grains at the shear zone margins are 341 highly misoriented. In addition, the finer grains within the shear band are consistently devoid of 342 intragranular misorientation except for strongly misoriented larger grains.

We interpret that the strongly misoriented clasts may represent fragmented remnants affected by brittle localized shear followed by dynamic recrystallization, resulting in the ubiquitous lack of misorientation in the surrounding finer-grained material. This pattern is highlighted by microscopic observations and strong crystallographic-preferred orientation (CPO), with [001] axes maxima but also forming a weak girdle along the stretching lineation and the foliation plane, respectively, whereas [010] and [100] display 348 maxima subperpendicular to it (normal to and within the foliation plane, respectively; Fig.4c). This fabric, 349 similar SL-type tectonites, is reported in many previous omphacite (a mineral rheologically similar to 350 jadeite and diopside) fabrics in eclogites (e.g., Philippot & Van Roermund, 1992; Godard & Van 351 Roermund, 1995; Keppler et al., 2016), compatible with near-plain strain dislocation creep deformation 352 mechanisms (e.g., Zhang et al., 2006; see also the review paper from Keppler, 2018). In addition, 353 truncation of oscillatory zoning in jadeite crystals from the shear band denotes the contribution of 354 solution-precipitation mechanisms. Thus, the fabrics herein observed indicate that a large part of the 355 material involved in the shear zone (in particular the dark-shaded, fine-grained jadeite rims visible in 356 Figure 4a) grew syn-kinematically, most likely via crystal-plastic deformation processes coupled with 357 dissolution-precipitation creep.



358

**Figure 4. a.** EDS X-ray map (counts of Ca, brighter shades indicating greater elemental concentrations) showing the structure of a shear band that transects a white jadeitite sample (PU3) from the Pus'yerka deposit. The coarse-grained oscillatory white  $Jd_1$  crystals are deformed within the shear band into oriented and truncated aggregates of jadeite crystals with local omphacite overgrowths. **b.** EBSD orientation map colored according to the inverse pole figure key (IPF) of jadeite (bottom right), showing a shape preferred orientation to the X direction of the strain ellipsoid (i.e., stretching orientation). View (XZ plane) corresponding to the Y axis of the finite strain ellipsoid. c. Pole diagrams of jadeite from the shear
band (black square in panel b) represented in an upper hemisphere equal-area projection for [100], [010]
and [001] crystallographic axes. Contours are multiples of uniform density distributions. The bold dot on
the L axis represents the stretching lineation direction (X direction of the strain ellipsoid), and the black
line represents the foliation plane.

370

#### 6. Texture and mineral chemistry of Polar Urals dark granofels

371 The dark granofels found within and along the white jadeitite "dykes" from the Pus'yerka locality exhibits 372 striking evidence for ductile and brittle shearing. Coarse idiomorphic calcic to sodic-calcic amphibole 373 cores of edenite (NaCa<sub>2</sub>Mg<sub>5</sub>Si<sub>7</sub>AlO<sub>22</sub>(OH)<sub>2</sub>) to Mg-katophorite (Na(CaNa)Mg<sub>4</sub>AlSi<sub>7</sub>AlO<sub>22</sub>(OH)<sub>2</sub>; Amp<sub>1</sub> 374 brighter Ca-rich cores in BSE imaging mode) form a dense network of dark, sealed fractures as well as 375 indentation and dissolution-precipitation features (Fig.3c,d). Various mutually overgrowing generations 376 of amphiboles ranging from Mg-katophorite ( $Amp_2$  on Fig.3d) to sodic amphiboles such as eckermannite 377 (NaNa<sub>2</sub>(Mg<sub>4</sub>Al)Si<sub>8</sub>O<sub>22</sub>(OH)<sub>2</sub>; Amp<sub>3</sub>) or richterite (Na(CaNa)(Mg,Fe)<sub>5</sub>Si<sub>8</sub>O<sub>22</sub>(OH)<sub>2</sub>) fill the breccia inter-clast 378 space in apparent textural equilibrium with omphacite (see also Angiboust et al., 2021). In sample PU2, a 379 dark granofels layer is observed (adjacent to a micro-fractured white jadeitite domain), containing 380 oriented sodic-calcic amphibole-phlogopite (± omphacite ± clinochlore) crystals (Fig.5a). Pervasive grain 381 size reduction occurred through micro-brecciation (Fig.5b,c,d) followed by further comminution that 382 ultimately led to the formation of anastomosing foliated cataclasite networks (Fig.5e,f,g; Type II). During 383 (or after) grain size reduction, the large primary amphibole porphyroclasts of edenitic to magnesio-384 katophoritic compositions (with irregular Cr enrichments; Fig.5c,g) were re-equilibrated along their rims 385 and fractures with very fine-grained pulverized domains with richteritic to eckermannitic compositions 386 (Fig.5c,f). It thus appears that fracturing and milling occurred after (or during) the formation of Mgkatophorite around the edenitic rims. LA-ICP-MS trace element mapping shows that the Mg-katophorite 387 388 clasts are enriched in Cr, Nd, Zr and Y, whereas the fine richterite-eckermannite intergrowths are 389 relatively enriched in Li and Ni (see Fig. S4).



Figure 5. a. EDS X-ray map (counts of Na) showing the internal structure of a shear band crosscutting a
 dark granofels (sample PU2, Polar Urals). Primary edenite compositions (large clast) are gradually
 transformed along their rims into Mg-katophorite to richteritic compositions. Remnants of the original

394 edenite clasts are found floating in the Mg-rich domain, associated with oriented phlogopite (Phl) as well 395 as rare omphacite and clinochlore crystals. b and c. EPMA X-ray maps showing a close-up on the foliated, 396 matrix-rich, clast-supported chaotic breccia. d. BSE image showing dissolution features of the primary 397 edenitic amphibole. e. BSE image showing a region of the shear band exhibiting a typical foliated 398 cataclasite texture with Mg-katophorite clasts wrapped in a richterite+eckermannite fine-grained matrix. **f** 399 and g. EPMA X-ray maps showing the internal structure of the foliated cataclasite as well as the presence 400 of pre-fracturing Cr sector zoning in Mg-katophorite crystals. Some trace element maps for this sample 401 are provided in Fig. S4. Texturally late Na- and Ca-rich zeolites grow as patches and along fractures parallel 402 with the main foliation.

403 An example of Type I mosaic breccia can be seen in Fig. 6a (sample UR11b), where clasts of Mg-404 katophoritic composition were also heavily affected by multiple fracturing events. Remnants from the 405 most pristine amphiboles lie along the edenite-Mg-katophorite transition (see Table S1 and Fig. S4 for 406 chemical properties). Omphacitic clinopyroxene is very common as inclusions within clast cores as well 407 as along sealed fractures (Fig.6d). Several generations of Mg-katophorite (with the younger exhibiting 408 increasing Na(B) and decreasing Ca contents; Fig.6c) can be identified within and around the clasts. 409 Texturally late eckermannite crystals (together with nyboïte: NaNa<sub>2</sub>(Mg<sub>3</sub>Al<sub>2</sub>)Si<sub>7</sub>AlO<sub>22</sub>(OH)<sub>2</sub>) are found filling the clast's cracks as well as replacing quadrangular inclusions in the cores of the fragments (Fig.6b; 410 411 see Angiboust et al., 2021 for details on mineral chemistry). The inter-clast space is filled with fine-412 grained Mg-katophorite fragments that coexist with zoned clinopyroxene (ranging from jadeite to omphacite in composition; Fig.6d) as well as micrometer- to tens of micrometer-sized chromian spinel 413 414 crystals (surrounded by Cr-clinopyroxene; Fig.6b; Fig. S2). Electron probe mapping demonstrates that 415 the inter-clast domain is relatively enriched in Al, Na, Ca and Cr with respect to the Mg-katophorite clasts 416 (Fig. S5).

LA-ICP-MS trace element mapping of the brecciated region from the dark granofels (UR11b) shows Mg-katophorite cores exhibiting Cr oscillations equivalent to sample PU2 (**Fig.5g**), revealing cryptic fractures healed with amphiboles substantially enriched in Zr, Hf, Y, Cr, Li and Ba (**Fig.6e,f,g; Fig. S5**).

420	Similar enrichments are distinguished within clinopyroxene-rich fracture-fill material (Fig. S5). Patchy
421	enrichment in Li, Ba, Sr, Y and Ce (among other elements) is also visible in the amphibole-clinopyroxene
422	inter-clast domain (see Fig. S5 for further maps as well as normalized trace-element spider diagrams).

423

424

425

426 Figure 6. a. EDS X-ray map (counts of Al, brighter shades indicating greater elemental concentrations) 427 depicting the chaotic breccia texture with the various healed fracture networks, as well as the location of 428 both LA-ICP-MS and electron probe maps (sample UR11b, Polar Urals). b. EPMA-based masked phase map 429 showing the various Mg-katophorite generations (green), eckermannite (blue) and nyboïte (violet). 430 Omphacite is in light yellow shades, jadeite in deep orange and chromite (Chr) in red. Note how the 431 various fracture generations are mutually overprinting. c. Amphibole masked map showing the Na(B) content. Increasing Na(B) is commonly viewed as reflecting a decreasing P/T ratio (closer to HP-LT 432 433 conditions). d. Masked map of Na content in clinopyroxene showing two distinct compositions in the 434 breccia-filling material. e, f and g. LA-ICP-MS masked trace element maps (amphiboles only) showing Y, Zr 435 and Cr concentrations (in ppm). The full analytical dataset is given in Fig. S5. Black arrows indicate Zr and Y 436 enrichments in the healed fractures and in the inter-clast matrix.



#### 439 **7.** Deformation and mineral chemistry of Motagua (Guatemala) jadeitite breccias

440 Macroscopic evidence for brecciation of the Motagua region jadeitites is widespread in the studied 441 samples (see also Harlow, 2011 and references therein). Three types of deformation patterns were 442 observed: (i) deep green fractures crosscutting white jadeitites (Type I, Fig.7a,b); (ii) finely comminuted 443 foliated breccia and cataclastic patterns (Type II, Fig.7e); and (iii) hydraulic-like breccias where 444 millimeter- to centimeter-sized angular clasts are infiltrated by a fibrous, oriented jadeitite (bluish) 445 cement (Type III, Fig.7c,d; see below). Whereas the clasts forming the breccias are systematically whitish 446 to light-greenish along their rims, the fracture- and breccia-filling material is always darker, ranging from 447 dark green (Fig.7a,b) to deep blue (Fig.7c,d,e). The inter-clast domain filling these breccias is always 448 enriched in Ca, Mg and Fe and depleted in Na and Al, as a consequence of omphacite growth. The 449 replacement of the original jadeite (Jd1, of near pure jadeite composition) by a secondary clinopyroxene 450 ( $Jd_2$ , commonly slightly poorer in jadeite molecule ( $80 < Jd_2 < 90 \text{ mol.\%}$ ) and Omp) is pervasive, affecting 451 the entire breccia and not only the clast margins (Fig.8a,b). Fracture-filling clinopyroxene exhibits 452 feather-like omphacitic crystals (Fig. S2) as well as a texturally strained appearance (Fig. 8a). The Mg EDS 453 map in Fig.8a shows relict white jadeitite clast cores (also visible on the hand specimen image in Fig.7e) 454 affected by pervasive re-equilibration (dissolution-precipitation) by Ca-rich jadeite and omphacite 455 compositions along crystal joints, fractures, grain boundaries and micro-faults.

## Tectonized jadeitites from Motagua fault zone area (Guatemala)



Tectonized jadeitites from Hpakan area (Kashin region, Myanmar)





469 Strongly foliated white jadeitite matrix in between two foliated dark granofels bands (Type II). Note the
470 large amount of blue amphibole clasts wrapped in the main clear foliation. The white boudin corresponds
471 to a remnant of the pre-deformation pure jadeitite (*Jd*<sub>1</sub>).

472 In Type II fault rocks, a centimeter-thick layering with variable shades of blue-grey (Fig.7c) is 473 observed transecting the original white jadeitite (Fig.9a). Pale, clast-like patches (of pure jadeite 474 composition) are macroscopically visible both in the host jadeitite as well as within the deep blue layers 475 (see **Table S1** for representative mineral compositions). Microscopically, the bands are composed of very 476 fine-grained (5–15  $\mu$ m on average diameter) jadeitite fragments cemented by omphacite-rich 477 compositions in interstitial positions (Fig.9b,c,d,g). Interestingly, this apparently pulverized domain also 478 hosts "shard-like" features ranging in size from several tens to hundreds of microns (Fig. 9e,f,g). These 479 shards (locally showing ptygmatic folds and contortion) define a weak lamination, resembling the 480 "fiamme" textures known in ignimbrites and other pyroclastic flow deposits. Many of these shards 481 display a fibro-radial intergrowth of jadeite and omphacite fibers that evoke the textures reported in 482 spherulites formed after the devitrification of a former glass (Fig.9g). SEM-based surface estimates of 483 these shards yield compositions ranging between those measured for jadeite and omphacite crystals 484 (Fig.9c,h). The shard-bearing fault zone domain shown in Fig.9b is transected by a dark, very fine-grained 485 omphacitite band that seems connected with the network of omphacite-bearing hydrofractures that 486 transect the rock volume. Remnants from foliated fragments of the host (namely the white jadeitite  $Jd_1$ but also the foliated shard-bearing domain) are observed, strongly dissolved, within this omphacitite 487 488 band. A metasomatic reaction front also appears to have formed at the contact between these two 489 domains (Fig.9b).



490

Figure 8. a. EDS X-ray map (counts of Mg, brighter shades indicating greater elemental concentrations) depicting a Type III hydraulic breccia texture with fibrous clinopyroxene fills (see Fig. S2 for further images; Motagua region, Guatemala). Note how the  $Jd_1$  clasts are corroded and replaced by a secondary jadeite (richer in omphacitic component) along grain boundaries, micro-fractures as well as inside the clasts. b. BSE image of a brecciated jadeitite with fibrous eckermannite (sodic amphibole) infills (Myanmar). c. BSE image showing dissolved primary  $Jd_1$  generation (nearly  $Jd_{100}$ ) replaced by a secondary  $Jd_2$  composition (slightly more enriched in omphacite).



Figure 9. a. Hand specimen image of a Motagua jadeitite showing deformation features of the pristinewhite jadeitite on the right, the foliated clast-bearing domain in the center (Type II texture) and a

502 secondary omphacitite band on the left. b. EDS X-ray map (counts of Mg) depicting a Type II matrix 503 hosting white jadeitite clasts (in black), foliated "shards" as well as late omphacite-bearing hydrofracture 504 networks. c. Clinopyroxene composition in the Jd-Di-Aeg triangle (after Morimoto, 1989; MTG03). Solvus 505 domains for clinopyroxene are derived from Garcia-Casco et al. (2009). d. BSE image showing the typical 506 appearance of a Type II foliated cataclasite matrix where shards and partly re-equilibrated white jadeitite 507 clasts coexist. Note how the shards are randomly folded but yet define a mild foliation. e and f. Quantified 508 EPMA X-ray maps of TiO<sub>2</sub> and CaO contents (in wt.%) showing elemental distributions in a clast and shard-509 bearing, Type II foliated cataclastic matrix. g. High-magnification BSE image showing a shard that exhibits a 510 fibro-radial internal structure that evokes devitrification spherulites. Note how the finely-comminuted 511 clasts forming the pulverized matrix were overgrown by a new (locally facetted) omphacitic clinopyroxene 512 composition during fault consolidation and fluid-enhanced sealing. h. Clinopyroxene composition in the 513 Jd-Di-Aeg triangle (sample MTG12). Note how shards are systematically enriched in omphacitic content.

514

#### 8. Deformation and mineral chemistry of Hpakan (Myanmar) jadeitite breccias

515 Further evidence for brecciation of jadeitites can be obtained observing the hand specimens 516 from Myanmar shown in Figs 7f, 7g and 7h. These samples exhibit the most complex sequence of 517 mutually crosscutting brittle events (Table 1). In Fig.7f, white (and slightly rounded) jadeitite clasts are 518 rimmed by omphacite (light green domains) and wrapped by dark blue amphibole (eckermannite)-519 clinopyroxene intergrowths (**Fig.8b**). The pristine jadeitic pyroxene composition  $(Id_1)$  is only preserved as 520 islands in the middle of a brighter (in BSE) more omphacitic pyroxene ( $Jd_2$  and Omp; Fig.8c). In sample 521 KAS10, dissolved cores exhibiting exsolution features can also be observed, with two pyroxenes forming 522 at the expense of a former one (Fig. S2). Jd<sub>1</sub> seems to have been replaced along grain boundaries by a 523 fluid that triggered pervasive re-equilibration (mixed Type I and Type III deformation). In other samples, 524 the same meso-scale structure exhibits a weak, omphacite-rich foliation wrapping remnants of the white jadeitite and dismembered amphibole-rich fragments (Fig.7g). This texture suggests that viscous flow 525 526 may occasionally overprint previously formed brittle (Type I) brecciated zones.

527 Sample KAS07 displays a whitish, foliated band separating two dark granofels domains (**Fig.7h**). The 528 leucocratic band comprises small, flattened dark amphibole fragments oriented parallel with the main 529 foliation (Fig.10a), as well as green trails characteristic of Cr-bearing omphacitic clinopyroxenes. A 530 cathodoluminescence image (Fig.10b) of the whitish domain reveals the presence of three distinct 531 jadeite-forming events in a Type II pattern deformation zone. The first is a green-colored aggregate (Jd<sub>1</sub>; 532 white jadeite:  $Jd_{91}Quad_5Aeg_4$  on average) wrapped within a pink-colored foliated band (mostly  $Jd_2$ ) that 533 hosts numerous angular, green-shaded fragments clearly derived from the green  $Jd_1$  aggregate on the 534 left of the image (Fig.10b). These green clasts range in size from several tens to hundreds of microns, and 535 display widespread evidence of fracturing, dissolution and re-precipitation/overgrowth by a brighter, 536 slightly more omphacitic composition (see for example **Fig.10c**; Jd<sub>84</sub>Quad<sub>6</sub>Aeg<sub>10</sub> on average). This fabric 537 evokes a deformation by micro-fracturing, with substantial comminution accompanied with shearing and flow banding of the pink cataclastic domain. 538

539 In the same sample, an eckermannite-filled extensional vein (Type III) is observed crosscutting at low 540 angles the foliated cataclastic domain (Fig. S2). A late pure jadeite formation event ( $Jd_3$ , in purple on 541 **Fig.10b**) also occurs as foliation parallel bands, as patchy domains in the green-colored  $Jd_1$  augen on the 542 left of the image, as well as filling late cracks that crosscut all previous features. Sheared amphibole 543 fragments (mostly eckermannite) appear dark in the cathodoluminescence image in the main foliated 544 area (Fig.10b). BSE imaging and composition analysis reveal that the amphibole-rich band from the left 545 part of this thin section (Fig.10a) locally contains Mg-katophorite cores rimmed by eckermannite or 546 nyboïte compositions (see Table S1 for representative mineral compositions). This brecciated amphibole-547 rich domain (likely a former Type I) has been cemented by a clinopyroxene-Na-amphibole mixture and 548 foliated parallel to the adjacent foliated cataclasite domain (Fig.7h, 10b). A striking feature of this dark 549 domain is the systematic decrease of Ca content in amphibole from the cores (Mg-katophorite; 7 wt.% 550 CaO) to the foliated matrix, where amphiboles are extremely enriched in Na<sub>2</sub>O (>11 wt.%) and depleted 551 in CaO (<1.5 wt.%). Similar observations were made in sample KAS06 where cores are edenitic (up to 10

- 552 wt.% CaO), whereas crystals lining the foliation correspond to eckermannite, also closely associated with
- 553 strained jadeite crystals (see **Fig. S2** for EDS mapping and further details on mineral chemistry).



555 Figure 10. a. EDS X-ray map (counts of Ca) showing the internal structure of a foliated and brecciated dark 556 granofels layer with Mg-katophorite cores (locally containing up to 8 wt.% CaO) wrapped within a jadeite-557 eckermannite-bearing foliation. b. Cathodoluminescence image of the whitish, foliated part of sample 558 KAS07. Note the white jadeitite ( $Jd_1$ ; CL-colored in green) fragments now dispersed in the Type II foliated 559 cataclasite matrix ( $Jd_2$ ; CL-colored in pink) and the black trails parallel with the foliation that are mostly 560 made of sodic amphibole (e.g., eckermannite). Late Cr-rich jadeite ( $Jd_3$ ; CL-colored in purple) is also 561 observed parallel with the main foliation as well as along cracks. c. EPMA X-ray map showing FeO 562 distribution in the whitish foliated cataclastic domain. Note the random (Fe-poor)  $Jd_1$  clast size distribution 563 as well as the apparent absence of internal deformation (such as plasticity) of  $Jd_1$  clasts.

564 The most compelling evidence pointing to repeated brittle deformation events is exhibited in 565 Fig.11 (sample KAS04). Three chemically distinct domains can be distinguished: (i) the white jadeitite 566 clasts (likely a former Type I structure); (ii) the blue amphibole fibrous veins that "inject" between 567 breccia clasts (e.g., hydrofracture-filling material); and (iii) a very fine-grained, greenish Type II foliated 568 omphacitite domain at the top of the sample (Fig.11a, Fig.12f). Locally, the contact between the blue 569 amphibole domains and the white clasts exhibits structures that evoke Type III hydrofracturing processes 570 (with local embayments pointing to interface-coupled dissolution-precipitation mechanisms; Fig.11b; e.g., Putnis & Austrheim, 2013). A large surface of the same sample (KAS04) has been mapped using EDS 571 572 (Fig.11c) in order to image millimeter-scale relationships between these three domains. The largest 573 rounded, Al-rich white jadeite clast (colored in red) is transected along its base by a sharp micro-fault 574 plane, along which the green, very fine-grained material ("foliated cataclasite") is found. The green 575 omphacitite domain is formed by tens of micrometer-sized crystals (Fig.11d) that commonly contain dark 576 clasts (in BSE mode) with a composition similar to the large white jadeitite clasts (Fig.12f). 577 Cathodoluminescence imaging reveals that the clasts in the foliated cataclasites have a similar purple CL 578 color as the large white jadeitite clasts, confirming their genetic link (Fig.12a,b,c). These clasts are 579 rimmed by two distinct clinopyroxene compositions (Jd2 and Omp) with increasing omphacitic content 580 (Fig.11e, 12f), the interfaces between





582 Figure 11. a. Polished hand specimen of a brecciated jadeitite from Myanmar (KAS04), where white 583 jadeitite clasts are transected by a green, strongly foliated cataclastic domain (Type II). The sample is lately 584 affected by a stage of pervasive, fibrous sodic amphibole (eckermannite) growth, infiltrating along cracks. 585 **b**. Higher magnification image showing how the white jadeitite has been hydrofractured and filled by sodic 586 amphibole, leaving embayments that may be interpreted as resulting from the dissolution by a reactive 587 fluid. Note also the distribution of sodic amphibole fibers infiltrating the Type III breccia. c. EDS X-ray map 588 (counts of Al) showing the great complexity of the internal structure in sample KAS04, where several 589 brittle deformation and fluid-rock interaction events can be identified, including deformation features 590 along ancient fault planes with injecting cataclastic material as well as damage zone formation. d. BSE 591 image of the foliated cataclasite domain where white  $Jd_1$  clasts are now dissolved and embedded within a 592  $Jd_2$ -omphacite matrix. **e**. BSE close-up image of the faulted area in panel **c** showing the foliated cataclasite 593 domain, the micro-fault plane as well as the partly re-equilibrated damage zone. The contact between the 594 large white jadeitite clast has been re-activated and filled by a clinochlore-eckermannite-celsian 595 paragenesis.

596 which appear to be dissolved (see black arrows on **Fig.11d**).

597 Elongated patches with omphacitic composition are aligned with the main foliation, and invade the fault 598 as well as its damage zone (Fig.12a,b). The foliated nature of this fine-grained jadeite-omphacite rich 599 domain is evidenced using EBSD mapping (Fig.12d). This image shows a shape-preferred orientation, 600 with elongated jadeitic clinopyroxene grains (aspect ratios up to 6:1) defining a lineation subparallel to 601 the strong crystallographic-preferred orientation well visible in [001] (Fig.12e). Similarly to Polar Urals 602 jadeitites, the [010] crystallographic axes are highly concentrated normal to the foliation plane but [100] are aligned within the foliation plane perpendicular to [010]. This defines a fabric akin to LS-type 603 604 tectonites, confirming the likely presence of crystal-plastic deformation mechanisms associated with a 605 near-plain strain geometry, as also supported by experimental investigations and numerical simulations 606 (Ulrich & Mainprice, 2005; Zhang et al., 2006). In addition, intragranular misorientation maps show that 607 several highly misoriented grains are surrounded by clusters of finer-grained crystals devoid of internal

608 deformation. The former likely represent remnant fragments, and the latter dynamically recrystallized

609 grains along the foliation (**Fig.12d**, **Fig. S3**).



610

**Figure 12. a, b, c.** Cathodoluminescence images showing the relationships between the different textural domains of sample KAS04. Note how the  $Jd_1$  fragments (pink colors) are dragged within the foliated cataclasite domain. **d.** EBSD orientation map colored according to the inverse pole figure key (IPF) of jadeite (bottom right), showing a shape preferred orientation of jadeite to the X direction of the strain ellipsoid (i.e., stretching orientation). View (XZ plane) corresponding to the Y axis of the finite strain ellipsoid. **e.** Pole diagrams of jadeite crystals smaller than 100 µm in diameter (larger crystals are considered as clasts, as indicated in the figure and more detailed in **Fig. S3**), represented in an upper

hemisphere equal-area projection for [100], [010] and [001] crystallographic axes. Contours are multiples
of uniform density distributions. The bold dot on the L axis represents the stretching lineation direction (X
direction of the strain ellipsoid) and the black line the foliation plane. f. Projection in a CaO-Al<sub>2</sub>O<sub>3</sub>(FeO+MgO)-Na<sub>2</sub>O system of the different groups of mineral compositions, as well as local surface
estimates of bulk major element composition (WR: whole-rock). Circles are single point measurements
and squares are SEM-based surface compositional estimates (WR).

624 A late faulting event that led to the precipitation of blue amphibole as well as clinochlore along 625 micro-fault planes affects the white jadeitite domain and the foliated cataclasites (Fig.11c,e). Celsian 626 crystals (a Ba-rich feldspar) are commonly found texturally associated with these apparently late 627 clinochlore-bearing fault zones (black arrows on Fig.11c). To summarize, textural relationships visible in 628 sample KAS04 suggest the following deformation sequence: (i) tectonic brecciation of the white jadeitite; 629 (ii) cataclasis of some of the white jadeitite and foliation of this cataclased domain; (iii) re-brecciation 630 (hydraulic) of the volume and precipitation of the Na-amphibole-rich domain between brecciated and 631 cataclased domains; and (iv) crystallization of clinochlore and celsian along late discrete fault zones 632 oblique to the previous fault structures (Fig.11c).

#### 633 9. Discussion

## 634 9.1 Worldwide jadeitites record mixed deformation regimes

## 635 Our comparative and analytical investigation reveals that most of the studied samples from Russia, 636 Guatemala and Myanmar display widespread and similar markers of fracturing that have received little 637 attention in previous studies (e.g., Dobretsov & Ponomareva, 1968; Shi et al., 2009b). Brittle deformation 638 markers span a broad range, from Type I breccias typical of damage zones (e.g., Figs.2, 6, 10a, 13a), to 639 Type II variably foliated cataclasites that are more diagnostic of strongly localized fault systems (e.g., 640 Rowe et al., 2011; Angiboust et al., 2015; Oncken et al., in revision; Figs.5, 9, 13b), to Type III hydraulic 641 breccias thought to develop at high fluid pressure conditions as extensional veining (e.g. Woodcock et 642 al., 2007). A common feature of all studied materials are arrays of healed mineral fractures commonly

643 coupled with grain size reduction, crystal indentation and dissolution-precipitation processes. This 644 pattern, sometimes hardly visible using optical microscopy, requires the use of specific analytical 645 approaches such as cathodoluminescence imaging or electronic microscopy to be documented (e.g., 646 Figs.3, 12; Shi et al., 2003; Sorensen et al., 2006; Takahashi et al., 2017). Pure extensional fracturing (e.g., 647 hydrofracturing; Type III; Fig.13c) has been identified in several samples, with fragments of the host that 648 appear to have been snatched into clinopyroxene or amphibole-rich veins (Fig.10b). When several brittle 649 events can be distinguished, Type III events always occur late in the sequence (Table 1). In all localities, it 650 is always the original white jadeitite material  $(Jd_1)$  that undergoes shearing-related brecciation and subsequent cementation by a darker clinopyroxene ( $Jd_2$  or Omp; Fig.7e) and/or by sodic amphiboles 651 652 (e.g., eckermannite; Figs.7f, 11a).

653 Evidence for viscous deformation has been observed in a shear band transecting the coarse-654 grained Polar Urals white jadeitite body (Fig.4a), as well as within Myanmar jadeitites (Shi et al., 2009b). 655 This observation reveals that the white jadeitite body, once crystalized, underwent differential stresses 656 high enough to generate crystal plasticity (most likely via dislocation creep) yielding a CPO. Following the 657 argument developed in Angiboust et al. (2021), it is hypothesized that this shearing occurred at c. 650 °C, 658 i.e., above the antigorite stability field (Fig.14d), because no evidence for Cr enrichment was found in this shear band (even though other processes may also contribute to Cr availability in the system; e.g., 659 Huang et al., 2019). Assuming a strain rate of 10<sup>-14</sup> s<sup>-1</sup> (a median value commonly inferred for ductile 660 661 shear zones; e.g., Pfiffner & Ramsay, 1982) and using the experimentally-based dislocation creep flow 662 laws from Orzol et al. (2006) and Zhang et al. (2006), it is possible to estimate the differential stresses 663 required for creating the shear band of sample UR03 to be between 125 and 40 MPa, respectively. Note 664 that these values likely represent an upper bound since other competing deformation mechanisms such 665 as pressure solution creep would likely be activated at lower stress values (e.g. Godard & Van Roermund, 666 1995). The implications of these values will be discussed hereafter.



668 Figure 13. a, b and c. Sketches showing the key features of the three main categories of tectonized 669 jadeitites identified in this comparative study. d. Sketch depicting the white jadeitite structure (derived 670 from observations from the Pus'yerka body) at the time of brecciation, indicating the various elements 671 identified in the field and their position with respect to the main fault and host ultramafics. e. Sketch 672 highlighting the structure of jadeitite blocks as exposed in HP-LT serpentinite mélanges, where they are 673 commonly associated with blueschist and/or eclogite-facies crustal rocks. We posit that the faulting 674 frequently observed within jadeitite blocks has been acquired earlier, when the jadeitite was a dyke-like 675 structure hosted by weakly serpentinized mantle wedge peridotites.

676

The CPO observed in KAS04 sample and other Type II samples likely results from a combination of dislocation creep and solution-precipitation processes in a dominant simple shear regime but also with some degree of pure shear, as suggested by the observed LS-type fabric but also a weak girdle of [001], respectively (**Figs.4c,12e**). These deformation mechanisms have been already documented in eclogite-facies clinopyroxene by Godard & Van Roermund (1995; see also Ulrich and Mainprice, 2005) (e.g., **Fig.12b**). It is clear from the fragmented jadeitite clast distribution (along with the

683 cathodoluminescence color of the various crystals; Fig.12b) that cataclastic flow (during brittle 684 deformation) operated through localized faulting before CPO formation (see also Oncken et al., in 685 revision; Fig.10c). In other words, we infer that the observed foliated cataclasite structure is the 686 protracted result of consecutive brittle flow followed by slower, viscous creep. This sequence of events 687 may be the record of one or several seismic cycles where fast co-seismic slip is immediately followed by 688 a slower after-slip event (e.g., Sibson, 1986; Rowe et al., 2011). Indeed, many of the reported fabrics 689 resemble the structures documented in fault cores at shallower/colder environments where multiple 690 episodes of particle fluidization are reported on the principal slip zone (Figs.5f, 10c; e.g., Sibson, 1986; 691 Snoke et al., 2014; Cox & Munroe, 2016; Muñoz-Montecinos et al., 2021a). Interestingly, the main brecciation/cataclasis event in sample KAS04 is followed by an episode of hydraulic brecciation where 692 693 Na-amphibole precipitates Type III; Fig. S2). Similar deformation patterns have been observed in dark 694 granofels that also comprise breccias (Fig.6), foliated cataclasites (Fig.5) and multiple fracturing events 695 (Fig.3c; Fig. S5; Table 1).

696 Exceptionally, the finding of elongated spherulitic shards in a fine-grained sheared domain 697 (Figs.7c,9d) – possibly formed by the devitrification of former glass fragments – may be indicative of 698 former pseudotachylytes (i.e., a glassy injection domain formed by melting during fault slip at seismic 699 strain rates; e.g., Austrheim & Boundy, 1994; Austrheim & Andersen, 2004). The Jd1 clasts found floating 700 in the comminuted matrix display extensive fracturing, grain-size reduction and stress-driven corrosion 701 that characterize pseudotachylyte-bearing systems (e.g., Kirkpatrick & Rowe, 2013). Glassy shards similar 702 to those shown in Fig.9g are mostly reported in ignimbrite deposits (e.g., Bull & McPhie, 2007) and in 703 rocks from ancient impact craters (e.g., Dressler & Reimold, 2001). Although, to the best of our 704 knowledge, such shard-like textures have not been described in ancient fault zones, the discovery of 705 unreported textures must be expected because the formation environment of these slip events strongly 706 differs from other paleo-earthquake reports made in the downgoing lithosphere (see below).

707 9.2 Jadeitite formation and pressure-temperature conditions of brittle deformation

708 In most serpentinite mélange settings, the original structure at the time of jadeitite formation has 709 been obliterated or disrupted by subduction zone tectonics (e.g., Harlow et al., 2015). Whereas in several 710 cases jadeitites appear to be directly precipitated from hydrothermal fluids ("P-type" jadeitite, according 711 the classification from Tsujimori & Harlow, 2012; e.g., Garcia-Casco et al., 2009; Harlow et al., 2011), 712 some localities preserve textural features typical of a replacement process ("R-type" jadeitite according 713 to the same classification), including igneous zircon crystals with older crystallization ages (Yui et al., 714 2012; Hertwig et al., 2016) or the presence of incompletely "jadeitized" trondhjemitic remnants in the 715 dyke volume (Angiboust et al., 2021). Indeed, early observations from Dobretsov & Ponomareva (1968) 716 and Kuznetsov et al. (1986) in Russia, as well as Bleek (1908) and Chhibber (1934) in Myanmar, reveal 717 that original jadeitite may form as dyke-like structures. It is plausible that these dykes were jadeitized 718 during long-term cooling within the high-pressure field, coeval with replacement of a former felsic 719 intrusive body formed earlier in the mantle wedge above the subduction interface, and possibly shortly 720 after subduction initiation in a warm thermal regime (see the recent model by Angiboust et al., 2021; see 721 also Meng et al., 2011 and Fig.14). Such secular cooling has also been documented above the 722 metamorphic soles from New Caledonia (e.g. Soret et al., 2016) and Oman ophiolites (e.g. Prigent et al., 723 2018).

A number of recent studies (e.g., Garcia-Casco et al., 2009; Schertl et al., 2012; Angiboust et al., 2021) have suggested that jadeitites may commonly form at temperatures far above 500 °C, perhaps up to 700 °C. Garcia-Casco et al. (2009) and Cárdenas-Párraga et al. (2012) have also suggested that exsolution features, as observed in sample KAS10 (**Fig. S2**), could evidence the cooling of a clinopyroxene of intermediate jadeite-omphacite composition below the solvus at 500 °C. This implies that white jadeitites including the Myanmar samples formed at temperatures higher than generally expected. Consistent with this interpretation, Harlow & Sorensen (2005) suggested that the whitish color of the

731 relatively high temperature jadeitite may be due to Cr stability in the host ultramafic chromite and 732 clinopyroxene crystals. In other words, it is likely that Cr-poor white jadeitite bodies formed at T 733 conditions above the antigorite stability field, i.e., before serpentinization of the host ultramafic (Fig.14), 734 even though some Cr may also be hosted in high temperature chloritite blackwall (not documented in 735 the studied localities; see Cardenas-Parraga et al., 2012). Following this logic, it can be concluded that 736 the systematic observation of darker-shaded clinopyroxene precipitates/overgrowths within breccias or 737 cataclasites in many jadeitite localities indicate that fracturing and fluid-rock interaction events occurred 738 during long-term cooling at T<600 °C (e.g., Garcia-Casco et al., 2009; Schertl et al., 2012; this study).

739 Whereas breccias and cataclasites are classically reported for rather shallow - and hence colder -740 environments (<15 km depth, T<300 °C; Sibson, 1986), a number of field examples have recently shown 741 that such features can form at depths greater than 30 km in the downgoing subducted crust (e.g., 742 Angiboust et al., 2012; 2015; Hertgen et al., 2017; Muñoz-Montecinos et al., 2021a; Oncken et al., in 743 revision). Minerals filling the studied breccias and growing within these fault zones encompass (i) 744 omphacite (and locally lawsonite) in Guatemalan samples; (ii) omphacite, jadeite, phlogopite and Na-rich 745 amphiboles in the Polar Urals samples; and (iii) Na-amphiboles, jadeitic and omphacitic clinopyroxenes in 746 Myanmar samples. These parageneses point to HP-LT conditions ranging between 1–2 GPa and 300–500 747 °C as documented in the reviews of Tsujimori & Harlow (2012) and Harlow et al. (2015; Fig.14d). Angiboust et al. (2021) have demonstrated that the phlogopite-bearing fracturing event seen 748 749 crosscutting Polar Urals jadeitites occurred at least 15 Ma after the HT jadeite-forming event (see also 750 Meng et al., 2011), thus providing an independent confirmation of the secondary nature of the 751 brecciation process. We conclude that in the studied set of samples - and perhaps as a rule in fossil and 752 active subduction systems – brittle deformation of jadeitites is a deep process occurring at or near peak-753 burial depths in a cooling environment (as suggested in Fig.14), rather than as a consequence of late 754 exhumation-related deformation. In order to undergo brittle deformation, the rock must be in-situ and

- surrounded by mechanically strong material, rather than as a tectonic block floating in a serpentinite-
- 756 matrix mélange where stress is dissipated in the weak matrix.

#### 757



758

Figure 14. a, b and c. Sketches depicting the long-term tectonic evolution of jadeitite "dykes" along the 759 760 base of the mantle wedge, from their formation (by replacement of a previous dyke structure or from 761 direct precipitation from a fluid) to their fracturing, dismembering and wrapping as blocks in a chaotic 762 serpentinite mélange. Previous geochronological data (Angiboust et al., 2021) confirms that several tens 763 of millions of years of subduction activity and mantle wedge hydration are required to achieve a structure 764 as shown in panel (c). Atg-: antigorite dehydration reaction. d. Pressure-Temperature diagram indicating 765 the conditions of formation of the main white jadeitite bodies documented worldwide, and the conditions 766 of the green jadeitite tectonic overprint (and associated sodic amphibole generation). After, and only after 767 brecciation (yellow stars), the dyke fragments can be exhumed along the plate interface, provided that the hangingwall serpentinization rate is sufficiently high to enable return-flow of these dense blocks in a 768 769 buoyant, serpentinized matrix (e.g., Gerya et al., 2002; Guillot et al., 2009). P-T conditions for jadeitites

770 are sourced from the compilations by Tsujimori & Harlow (2012) and Harlow et al. (2015). Also shown are 771 the slab-top temperature estimates as calculated by Antriasian et al. (2019, Hikurangi), Wang et al. (2020, 772 central Chile), Halpaap et al. (2019, W Greece) and Peacock (2003, NE and SW Japan). References used for 773 the depth of mantle wedge earthquakes (MW EQ) are provided in text. e. Location of one of the MW 774 clusters identified by Halpaap et al. (2019) in W. Greece and interpreted as marking the trace of plate-775 interface fluids migrating along a vent structure. The formation of this EQ nest (corresponding to the 776 yellow stars from panel d) likely implies that the serpentinization ratio of the subduction interface hanging 777 wall is very low, thus pointing to a structure analogous to the sketch drawn in panel (b).

778

## 9.3 Origin of brecciated dark granofels and relationships with fluid-rock interaction events

779 A striking similarity characterizes the textural relationships between the dark granofels and 780 jadeitites from the Polar Urals and Myanmar. In both localities, they occur as thin layers or pods of 781 "amphibolites" of several tens of centimeters (Chhibber, 1934; Angiboust et al., 2021; Fig.1e), generally 782 striking parallel to the main "dyke" foliation and deformed coeval with the associated jadeitites. Their 783 amphibole compositional patterns are also nearly identical, with relicts of Ca-rich amphiboles in the 784 granofels mineral cores (mostly edenite and Mg-katophorite), and Ca-poor, Na-rich amphiboles along 785 their rims and along highly metasomatized domains (Shi et al., 2003; Fig. S2). Field observations 786 (Chhibber, 1934; Angiboust et al., 2021) and petrological investigations suggest that two distinct dark 787 granofels occurrences may exist: (i) as disaggregated fragments (the "schistose amphibolite inclusions" 788 from Chhibber, 1934; see also Harlow et al., 2015) inherited from a pre-jadeitization event; and (ii) as 789 amphibole-rich blackwalls formed at the contact between the jadeitite body and the host ultramafic (Shi 790 et al., 2003). The latter should in theory have formed lately at T<350 °C (Shi et al., 2003) and only exhibit 791 Na-amphiboles (since no Ca chemical potential gradient exists between ultramafics and jadeitites), thus 792 hampering the formation of calcic and sodic-calcic amphiboles such as edenite or Mg-katophorite in the 793 reaction blackwall. The former should instead contain these sodic-calcic amphibole species (e.g., 794 Figs.5a,6b,10a) as remnants of an early HT event, associated with the infiltration of alkali-rich fluids 795 and/or melts (likely trondhjemitic in composition) within ultramafics in a subduction initiation setting.

This process is known to occur in anomalously high subduction thermal regimes (see also the discussion in Angiboust et al., 2021 and similar structures and rock assemblages in Lazaro et al., 2011 and Soret et al., 2016). Brecciation, followed by boudinage, late tectonic disaggregation and block-in-matrix deformation, most likely blurred the primary contacts, hampering an accurate reconstruction of the geometry of the original jadeitite-dark granofels boundaries (e.g., **Fig.10a**).

801

## 9.4 Fluid ingress and metasomatic imprint

802 A common feature reported in jadeitites worldwide is the enrichment in Ca, Fe and Mg of jadeitic 803 clinopyroxenes due to the infiltration of Ca-Fe-Mg-rich fluids associated with dissolution-precipitation 804 processes during late fluid-rock interaction events (e.g., Sorensen et al., 2006; Garcia-Casco et al., 2009; 805 Harlow et al., 2011; Cardenas-Parraga et al., 2012; Angiboust et al., 2021; this study). The formation of 806 dissolution features (Fig.3c), re-equilibration of clast rims (Fig.8a) and the metasomatic overprint left in 807 the rock record (Fig.9b) are witnesses to the apparently corrosive nature of the infiltrating fluids, which 808 were at thermodynamic disequilibrium with respect to the infiltrated material. The absence of 809 carbonates in most jadeitite occurrences suggests that the fluids were CO<sub>2</sub>-poor, most likely because the 810 CO<sub>2</sub> precipitated as carbonates before reaching the jadeitite-forming location. Substantial Cr 811 enrichments are also visible in some of the sheared domains and between breccia fragments (as shown 812 for instance by the growth of chromian spinel around Mg-katophorite clasts in sample UR11b; Fig.6b; see 813 also the green overgrowths around white jadeitite clast in Fig.7f).

As discussed earlier, the release of Cr may be related to serpentinization of the host peridotite during fluid-rock interaction. It is important to note that (i) jadeite/omphacite is found growing within brecciated white jadeitites (**Figs.3b**, **8a and 11c**) and amphibole-rich dark granofels (**Fig.6d**); and (ii) amphibole growth is observed in brecciated dark granofels (**Fig.5b**) and between white jadeitite breccia fragments (**Figs.7f,11a,11b**). From these observations, it appears that the incoming fluid composition 819 was not buffered by the wall-rock composition and that brittle events contributed to a chemical and 820 mechanical homogenization of the original dyke-forming lithologies. Furthermore, the widespread 821 formation of texturally late micas (e.g., phlogopite in the Polar Urals: Angiboust et al., 2021, see also Fig. 822 **S1**; phengite or phlogopite in Guatemala: Harlow et al., 2011; Flores et al., 2013; white mica in W. Japan: 823 Shigeno et al., 2012) associated with enrichments in LILEs and other fluid mobile elements (Figs. S3 and 824 S4) confirm that fluid sources were variable in time, switching from a mafic oceanic crust signature to a 825 more hybridized composition with transient highs in sedimentary input (e.g., Sorensen et al., 2006, 2010; 826 Morishita et al., 2007; Meng et al., 2011, 2016; Harlow et al., 2015; Chen et al., 2018; Cardenas-Parraga 827 et al., 2012). This temporal variability of incoming fluid chemistry from a more mafic to a more sedimentary signature could likely be explained by variations in the amount of sedimentary material 828 829 undergoing devolatilization reactions in the underlying subduction channel (e.g., Bebout, 2007; 830 Scambelluri et al., 2019). This chronological sequence confirms that the same fluid pathways (namely the 831 jadeitite "dykes") were used over several millions of years as major drains for highly pressurized plate-832 interface fluids, on their way to the partly hydrated mantle wedge (e.g. Doglioni et al., 2009; Spandler & 833 Pirard, 2013).

The distribution of jadeitite "dykes" as seen in the Polar Urals (Fig.1b; see also Angiboust et al., 834 835 2021) demonstrates that fluid pathways are rather discrete in the overlying plate (i.e., jadeitite does not 836 precipitate randomly everywhere in the fore-arc mantle). It also indicates that less mechanical energy is 837 required to re-fracture a pre-existing physical discontinuity (e.g., a dyke-like structure) than creating new 838 channels for draining plate-interface fluids. This implicitly requires that breccia sealing must have been 839 faster than rupture recurrence to explain the mutual crosscutting relationships documented between 840 the various events identified in our study (see also Woodcock et al., 2007). It seems clear that nearly 841 lithostatic pore fluid pressure was sustained throughout the entire jadeitite drain activity as 842 demonstrated by the ubiquitous evidence for fluid-rock interaction in the three types of brittle features

reported here (**Table 1**; see also Kuznetsov et al., 1986 and Angiboust et al., 2021). Yet, transient overpressures likely triggered the formation of some of the (explosive) breccia features (Type III) within already-tectonized Type I and Type II domains (**Figs.11c,13a,b**). These findings are consistent with the episodic opening of vein systems and filling of voids by omphacitic compositions documented by Garcia-Casco et al. (2009) and Cardenas-Parraga et al. (2012) in the Sierra del Convento serpentinite mélange (E. Cuba), and also in other subduction HP-LT mélanges environments (e.g., Muñoz-Montecinos et al., 2021b).

## 850

### 9.5 Insights on slip properties in a fluid-saturated mantle fault zone

851 Brecciated rocks have long been recognized as potential markers of seismic deformation (e.g., 852 Sibson, 1986; Angiboust et al., 2012; Melosh et al., 2014). Evidence of fault-zone rocks with diagnostic 853 elements such as damage zones and fault cores (Fig.13a, b) highlights for the first time that jadeitites 854 (and associated dark amphibole-rich granofels) host abundant brittle deformation events, most likely 855 along fast-slipping seismogenic fault planes. Estimating slip or strain rates along paleo-fault surfaces (or 856 shear zones) is a challenging task, subject to great uncertainties. While Oncken et al. (in revision) have 857 recently demonstrated that foliated cataclasites can form in slow slip environments for strain rates in the range  $10^{-3}$  to  $10^{-5}$  s<sup>-1</sup>, the lack of well-preserved pseudotachylytes in the studied samples hampers a 858 859 direct identification of seismic slip rates (i.e., on the order of m/s). However, the discovery of a foliated 860 fault zone with abundant "shards" exhibiting structures analogous to devitrified spherulites (Fig.9g) 861 opens the possibility for constraining former slip events at strain rates approaching seismic slip 862 velocities, fast enough to generate co-seismic temperatures as high as 1500 °C (e.g., Sibson et al., 2006; 863 Menant et al., 2018) and trigger frictional melting. Note that chemical modification of the system is 864 required to explain the enrichment in Ca, Fe and Mg of the shard-bearing fault zone as shown in Fig.9b, c. As earlier stated by Magloughlin (1992) and Swanson (1992), cataclasites and pseudotachylytes may 865 866 be closely intricated in fault zone rocks, both exhibiting evidence for open-system modification of the

867 pristine host composition. Thus, the structures herein observed may have formed through (i) 868 metasomatic fault-zone alteration associated with fluid influx within a finely-crushed cataclastic domain 869 (likely a structure as shown in Fig.13b), followed by (ii) fast-slip along narrow zones where local melting 870 of the previously-formed cataclasites occurred (thus explaining the enrichment in omphacitic component 871 of "shards" and devitrified spherulites, Fig.9d-h). Subsequent shear deformation contributed to the 872 apparent banding systematically observed in Type II fault rocks (e.g., Fig.9b). It is therefore proposed 873 that the reported features can be explained by fluctuations in slip rate velocities in a fluid-saturated fault 874 zone.

#### 875

## 9.6 Implications for mantle wedge seismicity and plate-interface rheology

876 Observations made on the Myanmar and Guatemalan loose samples are challenging to link to 877 their respective formation context because pristine field relationships were obliterated during block-in-878 matrix dismembering as well as sedimentary transport into conglomerate deposits. A few exceptional 879 observations in Myanmar document the presence of jadeitite-bearing "dykes" that are several hundred 880 meters long and meters to tens of meters thick (e.g., Bleek, 1908), but most of these bodies have been 881 mined. The Polar Urals Pus'yerka locality (and to a lesser extent other jadeitite localities in Siberia; 882 Dobretsov & Ponomareva, 1968) has the potential to yield in situ information on the dimensions of 883 ruptured bodies as well as refining the chronology of the processes at stake. Kuznetsov et al. (1986) report that jadeitite "dykes" in the Pus'yerka locality (also mined in the 1990s and mostly exhausted) 884 885 were locally up to several tens of meters thick, pinching out to meter-thick bands at the northern and 886 southern terminations of the several kilometer-long dyke structure (Fig.1b). These observations provide 887 critical insights into the minimum size for mantle wedge earthquakes. Assuming a planar geometry for a single event that ruptured the Pus'yerka jadeitite body (before the dyke structure was boudinaged), a 888 squared 3x3 km rupture, a 10 cm co-seismic displacement and a shear modulus of 84 GPa for jadeitite 889 890 (Hao et al., 2020), a magnitude of  $M_w$  = 4.5 can be calculated (Sibson, 1989). This magnitude estimates

891 ranges between 3 and 5 when changing fault dimensions and displacements within bounds compatible892 with field and structural observations.

893 Is this range of magnitudes comparable with the events reported in the forearc lithospheric 894 mantle clusters in active subduction zones? Davey & Ristau (2011) report a subvertical, 10 km-large 895 cluster of earthquakes (average maximum magnitude of 4.5 for normal, inverse and strike-slip events) 896 above the subduction interface at 40–50 km depth in the northern Hikurangi margin. These authors note 897 their spatial relationship with the inferred 700 °C isotherm, where they expect full antigorite breakdown 898 and relate these earthquakes to dehydration embrittlement processes. Released upwelling fluids 899 infiltrate the mantle wedge, possibly explaining volcanism in the above back-arc basin. Nakajima & 900 Uchida (2018) have identified a nest of supra-slab micro-seismicity (M<sub>w</sub>= 1–3) located 10-15 km above 901 the Philippine plate in central Japan, between 25–35 km depth with focal mechanisms corresponding to 902 normal and strike-slip events. Interestingly, seismic activity markedly increased after the 2011 Tohoku 903 megathrust rupture, leading Nakajima & Uchida (2018) to consider these events as marking the trace of 904 highly pressurized fluids, cyclically drained above the plate interface along a "highly fractured, pre-905 existing mature pathway". According to this model, several months are required for these fluid pulses to 906 travel the few kilometers that separate these nests from the megathrust area (see also White et al., 907 2016). Similar structural conclusions were drawn by Yu and Zhao (2020) who focused on 35–55 km depth 908 microseismicity events in the NE Japan forearc mantle using P-wave velocity perturbations. Halpaap et 909 al. (2019) have demonstrated that several seismicity nests ranging in magnitude from 2–4 occur under 910 Western Greece in a weakly serpentinized mantle. These events (mostly extensional) occur between 45-911 55 km depth along a diffuse structure that dips at 45° to the subduction interface plane (see hypocenter 912 distribution plotted in Fig.14e). Halpaap et al. (2019) conclude that plate-interface fluids transiently 913 migrate along fault-bounded vents, diminishing the pore pressure along the interface, thus decreasing 914 plate-interface seismicity. This non-exhaustive list of studies confirms that fluids are unequivocally 915 required to explain the seismotectonic features documented in the remote mantle wedge. It also
916 appears that these nests are transient in the context of subduction zone evolution and spatially localized
917 within well-defined structures.

918 The formation of mantle wedge earthquakes has critical implications for plate-interface rheology and 919 more specifically may provide precious constraints on the volumetric abundance of serpentinite in the 920 forearc mantle. A vibrant debate exists on the abundance and distribution of serpentinite in the mantle 921 wedge based on the interpretation of seismic wave velocity perturbations and the effect of serpentinite 922 anisotropy (e.g., Hyndman & Peacock, 2003; Liu et al., 2020; Luo & Wang, 2021). It is likely that 923 serpentinite distribution is highly uneven in the mantle wedge, with the highest serpentinization rates 924 directly above the interface and fairly low rates (likely less than 15 vol.%) within the innermost forearc 925 mantle (e.g., Reynard, 2013; Abers et al., 2017; Halpaap et al., 2019). Stress accumulation in the mantle 926 must be inhibited by serpentinite formation, which is known to flow at very low stresses (e.g., Escartin et 927 al., 2001; Reynard, 2013 and references therein). The stress level required for enabling plastic 928 deformation of jadeitites (40–125 MPa according to our calculations) is several orders of magnitude 929 greater than the stresses required for the flow of serpentinites (approximately 1 MPa at 600 °C using the 930 calibration from Hilairet et al., 2007). In order to transfer plate-interface stresses into the overlying 931 mantle and trigger plastic deformation of jadeitites (Fig.4), the host peridotite must have been relatively dry (i.e., with serpentinization rates far lower than 10 vol.%.) as inferred for many active subduction 932 933 settings (e.g., Abers et al., 2017; Halpaap et al., 2019). The triggering of multiple cataclastic events as 934 documented by our observations (Fig.9b) could be explained by transient increases in strain rate (during 935 slow slip events or earthquakes), by an increase of pore fluid pressures, or by a combination of both. The 936 latter option is supported by the discovery of fabrics typically observed in faulted jadeitites (e.g., Figs.9, 937 12), as well as major and trace element chemical arguments that point to mass transfer and opening of 938 the system to external fluids (e.g., Fig.6), most likely sourced in the subduction interface region.

939 From these considerations, we can conclude that (i) the seismic events recorded above the subduction 940 interface in several active subduction zones likely correspond to the brittle deformation of jadeitite 941 bodies (Fig.14d,e), which according to field observations are themselves derived from the metasomatic 942 replacement of former felsic dyke bodies or direct precipitation from fluids (see discussion in Harlow et 943 al., 2015 and Angiboust et al., 2021); and (ii) brittle jadeitite deformation may have occurred in the fossil 944 subduction record and in present-day settings in a relatively dry peridotitic mantle, thus providing 945 valuable insights into the serpentinization degree of the interface hangingwall and the stress distribution 946 at the base of the megathrust region (Halpaap et al., 2019; Luo & Wang, 2021). This record of the deep 947 conduits draining plate-interface fluids into the overlying mantle wedge sheds light on the structure of the "cold nose" above the subduction interface, with implications for volatile budget estimates, the 948 949 rheology of the plate interface itself (including the various types of seismicity), and the interpretation of 950 Vp/Vs and Poisson's ratios from active subduction settings worldwide.

#### 951 10. Conclusions

952 A vein network from the Polar Urals composed of jadeitite and amphibole-phlogopite granofels is hosted 953 in mildly serpentinized peridotites. This structure formed by metasomatic replacement of a former 954 magmatic dyke above the subduction interface at c. 50 km depth, in a young and dry mantle wedge 955 environment. During subsequent cooling of the subduction zone, this jadeitized "dyke" experienced 956 mixed brittle-viscous deformation regimes associated with several serpentinite- and oceanic crust-957 derived fluids. Extensive fracturing, well-visible in the amphibole-rich granofels and more cryptic in the 958 jadeitite bodies, led to the local formation of breccias and foliated cataclasites. The associated 959 metasomatic imprint confirms the importance of these fault zones as major drains for overpressurized 960 plate-interface metamorphic fluids. Very similar features, including omphacite- and amphibole-rich 961 breccias together with foliated cataclasites, are also observed in loose jadeitite boulders from Myanmar 962 and Guatemala serpentinite mélanges. We propose that supra-slab seismicity nests recorded in present-963 day subduction margins (e.g., N Hikurangi, NE Japan, W Greece) may reflect the repeated fluid-assisted breakage of planar bodies formed by jadeitite "dykes". The local presence of these rupture events 964 965 implies that the serpentinization rate was locally low (probably <10-15 vol.%) along the base of the

966 mantle wedge during "dyke" faulting activity. These results shed light on the importance of physical
967 discontinuities such as jadeitite bodies for draining plate-interface fluids into the mantle wedge, and
968 provide key constraints on the stress distribution in deep subduction margins.

## 969 Acknowledgements

970 Johannes Glodny (GFZ Potsdam) is acknowledged for insightful discussions that initiated this project. 971 François Gendron (MNHN) is also warmly acknowledged for sharing his knowledge about the Motagua 972 region and information on archeological material. Ksenia Kulikova is thanked for sharing her knowledge 973 on the regional geology of the Urals. J.M-M express his gratitude to Ralf Hielscher, Rüdiger Kilian and Luiz 974 Morales for assistance with the MTEX software. Michael Popov, Stephan Borensztajn, Arthur Delorme, 975 Julia Cozzolino and the Camparis team are acknowledged for technical assistance. Michael Bostock and 976 Gaston Godard are acknowledged for insightful discussions. The careful and efficient editorial handling 977 by Carlo Doglioni is much appreciated. Comments from Bill Leeman and two anonymous reviewers 978 helped to greatly increase the clarity of this manuscript. D.D. was funded by the European Research 979 Council grant REALISM (2016-grant 681346) to Alexandre Schubnel. This study was funded with an IDEX-980 USPC research chair grant (#16C538) and was partly supported by the IdEx Université de Paris ANR-18-981 IDEX-0001. This is IPGP contribution #40xx.

## 982 References

- Abers, G. A., Van Keken, P. E., & Hacker, B. R. (2017). The cold and relatively dry nature of mantle
  forearcs in subduction zones. *Nature Geoscience*, *10*(5), 333-337.
- Agard, P., Plunder, A., Angiboust, S., Bonnet, G., & Ruh, J. (2018). The subduction plate interface: rock
  record and mechanical coupling (from long to short timescales). *Lithos*, *320*, 537-566.
- Angiboust, S., Agard, P., Yamato, P., & Raimbourg, H. (2012). Eclogite breccias in a subducted ophiolite: A
   record of intermediate-depth earthquakes?. *Geology*, *40*(8), 707-710.
- Angiboust, S., Pettke, T., De Hoog, J. C., Caron, B., & Oncken, O. (2014). Channelized fluid flow and
  eclogite-facies metasomatism along the subduction shear zone. *Journal of petrology*, *55*(5), 883916.
- Angiboust, S., Kirsch, J., Oncken, O., Glodny, J., Monié, P., & Rybacki, E. (2015). Probing the transition
   between seismically coupled and decoupled segments along an ancient subduction interface.
   *Geochemistry, Geophysics, Geosystems, 16*(6), 1905-1922.

- Angiboust, S., Glodny, J., Cambeses, A., Raimondo, T., Monié, P., Popov, M., & Garcia-Casco, A. (2021).
  Drainage of subduction interface fluids into the forearc mantle evidenced by a pristine jadeitite
  network (Polar Urals). *Journal of Metamorphic Geology*. *39*(4), 473-500.
- Antriasian, A., Harris, R. N., Tréhu, A. M., Henrys, S. A., Phrampus, B. J., Lauer, R., ... & Barker, D. (2019).
  Thermal regime of the northern Hikurangi margin, New Zealand. *Geophysical Journal International*,
  216(2), 1177-1190.
- Audet, P., & Kim, Y. (2016). Teleseismic constraints on the geological environment of deep episodic slow
   earthquakes in subduction zone forearcs: A review. *Tectonophysics*, 670, 1-15.
- 1003 Austrheim, H., & Boundy, T. M. (1994). Pseudotachylytes generated during seismic faulting and 1004 eclogitization of the deep crust. *Science*, 265(5168), 82-83.
- Austrheim, H., & Andersen, T. B. (2004). Pseudotachylytes from Corsica: fossil earthquakes from a
   subduction complex. *Terra nova*, *16*(4), 193-197.
- Bachmann, F., Hielscher, R., & Schaeben, H. (2010). Texture analysis with MTEX–free and open source
   software toolbox. In Solid State Phenomena (Vol. 160, pp. 63-68). *Trans Tech Publications Ltd*.
- Bachmann, F., Hielscher, R., & Schaeben, H. (2011). Grain detection from 2d and 3d EBSD data—
   Specification of the MTEX algorithm. *Ultramicroscopy*, 111(12), 1720-1733.
- Batanova, V. G., Belousov, I. A., Savelieva, G. N., & Sobolev, A. V. (2011). Consequences of channelized
  and diffuse melt transport in supra-subduction zone mantle: evidence from the Voykar ophiolite
  (Polar Urals). *J. Petrol.* 52(12), 2483-2521.
- Bebout, G. E. (2007). Metamorphic chemical geodynamics of subduction zones. *Earth and Planetary Science Letters*, 260(3-4), 373-393.
- Beccaluva, L., Bellia, S., Coltorti, M., Dengo, G., Giunta, G., Mendez, J., Romero, J., Rotolo, S. and Siena,
   F., 1995. The northwestern border of the Caribbeanplate in Guatemala: New geological and
   petrological data on the Motagua ophiolitic belt. Ofioliti, 20, 1-15.
- Bleek, A.W.G (1908). Jadeite in the Kachin Hills, Upper Burma. *Records of the Geological Survey India*,
  36(4), 254-285.
- 1021 Bostock, M. G. (2013). The Moho in subduction zones. *Tectonophysics*, 609, 547-557.
- Bouchon, M., Marsan, D., Jara, J., Socquet, A., Campillo, M., & Perfettini, H. (2018). Suspected deep
   interaction and triggering between giant earthquakes in the Chilean subduction zone. *Geophysical Research Letters*, 45(11), 5454-5460.
- 1025 Brueckner, H. K., Lallemant, H. G. A., Sisson, V. B., Harlow, G. E., Hemming, S. R., Martens, U., ... &
- 1026 Sorensen, S. S. (2009). Metamorphic reworking of a high pressure–low temperature mélange along

- the Motagua fault, Guatemala: a record of Neocomian and Maastrichtian transpressional tectonics.
   *Earth and Planetary Science Letters*, 284(1-2), 228-235.
- Bull, K. F., & McPhie, J. (2007). Fiamme textures in volcanic successions: Flaming issues of definition and
   interpretation. *Journal of Volcanology and Geothermal Research*, *164*(4), 205-216.
- 1031 Cárdenas-Párraga, J., García-Casco, A., Harlow, G. E., Blanco-Quintero, I. F., Agramonte, Y. R., & Kröner,
- 1032A. (2012). Hydrothermal origin and age of jadeitites from Sierra del Convento Mélange (Eastern1033Cuba). European Journal of Mineralogy, 24(2), 313-331.
- 1034 Chang, Y., Warren, L. M., Zhu, L., & Prieto, G. A. (2019). Earthquake focal mechanisms and stress field for
   1035 the intermediate-depth Cauca cluster, Colombia. *Journal of Geophysical Research: Solid Earth*,
   1036 124(1), 822-836.
- Chen, Y., Huang, F., Shi, G. H., Wu, F. Y., Chen, X., Jin, Q. Z., ... & Nyunt, T. T. (2018). Magnesium isotope
   composition of subduction zone fluids as constrained by jadeitites from Myanmar. *Journal of Geophysical Research: Solid Earth*, *123*(9), 7566-7585.
- 1040 Chhiber HL. 1934. The Mineral Resources of Burma. London: MacMillan
- 1041 Cox, S. F., & Munroe, S. M. (2016). Breccia formation by particle fluidization in fault zones: implications
   1042 for transitory, rupture-controlled fluid flow regimes in hydrothermal systems. *American Journal of* 1043 *Science*, *316*(3), 241-278.
- Davey, F. J., & Ristau, J. (2011). Fore-arc mantle wedge seismicity under northeast New Zealand.
   *Tectonophysics*, *509*(3-4), 272-279.
- 1046 Deschamps, F., Guillot, S., Godard, M., Chauvel, C., Andreani, M., & Hattori, K. (2010). In situ 1047 characterization of serpentinites from forearc mantle wedges: timing of serpentinization and 1048 behavior of fluid-mobile elements in subduction zones. *Chemical Geology*, *269*(3-4), 262-277.
- Dessa, J. X., Klingelhoefer, F., Graindorge, D., André, C., Permana, H., Gutscher, M. A., ... & SUMATRA OBS Scientific Team. (2009). Megathrust earthquakes can nucleate in the forearc mantle: Evidence
   from the 2004 Sumatra event. *Geology*, *37*(7), 659-662.
- 1052 Dobretsov, N. L., & N. V. Sobolev (1984). Glaucophane schists and eclogites in the folded systems of 1053 northern Asia. *Ofioliti*, 9. 401-424.
- Dobretsov, N. L., & Ponomareva, L. G. (1968). Comparative characteristics of jadeite and associated rocks
   from Polar Ural and Prebalkhash region. *International Geology Review*, 10(3), 247-279.
- Doglioni, C., Tonarini, S., & Innocenti, F. (2009). Mantle wedge asymmetries and geochemical signatures
   along W-and E–NE-directed subduction zones. *Lithos*, *113*(1-2), 179-189.

- 1058 Dressler, B. O., & Reimold, W. U. (2001). Terrestrial impact melt rocks and glasses. *Earth-Science* 1059 *Reviews*, *56*(1-4), 205-284.
- Endo, S., Wallis, S. R., Tsuboi, M., Torres De León, R., & Solari, L. A. (2012). Metamorphic evolution of
   lawsonite eclogites from the southern Motagua fault zone, Guatemala: insights from phase
   equilibria and Raman spectroscopy. *Journal of Metamorphic Geology*, *30*(2), 143-164.
- Escartin, J., Hirth, G., & Evans, B. (2001). Strength of slightly serpentinized peridotites: Implications for
   the tectonics of oceanic lithosphere. *Geology*, *29*(11), 1023-1026.
- 1065 Fishman, A.M. (2006): Gems in the north Ural and Timan. *Geoprint*, Syktyvkar, 88 p. (in Russian).
- Flores, K. E., Skora, S., Martin, C., Harlow, G. E., Rodríguez, D., & Baumgartner, P. O. (2015). Metamorphic
   history of riebeckite-and aegirine-augite-bearing high-pressure–low-temperature blocks within the
   Siuna Serpentinite Mélange, northeastern Nicaragua. *International Geology Review*, *57*(5-8), 943 977.
- Flores, K. E., Martens, U. C., Harlow, G. E., Brueckner, H. K., & Pearson, N. J. (2013). Jadeitite formed
   during subduction: In situ zircon geochronology constraints from two different tectonic events
   within the Guatemala Suture Zone. *Earth and Planetary Science Letters*, *371*, 67-81.
- Frank, W. B., Shapiro, N. M., Husker, A. L., Kostoglodov, V., Bhat, H. S., & Campillo, M. (2015). Along-fault
   pore-pressure evolution during a slow-slip event in Guerrero, Mexico. *Earth and Planetary Science Letters*, 413, 135-143.
- Franz, L., Sun, T. T., Hänni, H. A., de Capitani, C., Thanasuthipitak, T., Atichat, W., ... & Capitani, C. D.
  (2014). A comparative study of jadeite, omphacite and kosmochlor jades from Myanmar, and
  suggestions for a practical nomenclature. *J. Gemmol*, 34, 210-229.
- Fu, Y., & Freymueller, J. T. (2013). Repeated large slow slip events at the southcentral Alaska subduction
   zone. *Earth and Planetary Science Letters*, *375*, 303-311.
- García-Casco, A. (2007). Magmatic paragonite in trondhjemites from the Sierra del Convento mélange,
   Cuba. *American Mineralogist*, 92(7), 1232-1237.
- García-Casco, A., Vega, A. R., Párraga, J. C., Iturralde-Vinent, M. A., Lázaro, C., Quintero, I. B., ... & Torres Roldán, R. L. (2009). A new jadeitite jade locality (Sierra del Convento, Cuba): first report and some
   petrological and archeological implications. Contributions to Mineralogy and Petrology, 158(1), 1.
- Gendron, F., Smith, D. C., & Gendron-Badou, A. (2002). Discovery of jadeite-jade in Guatemala confirmed
   by non-destructive Raman microscopy. *Journal of Archaeological Science*, 29(8), 837-851.
- Gerya, T. V., Stöckhert, B., & Perchuk, A. L. (2002). Exhumation of high-pressure metamorphic rocks in a
   subduction channel: A numerical simulation. *Tectonics*, 21(6), 6-1.

- Glodny, J., Austrheim, H., Molina, J. F., Rusin, A. I., & Seward, D. (2003). Rb/Sr record of fluid-rock
   interaction in eclogites: The Marun-Keu complex, Polar Urals, Russia. *Geochimica et Cosmochimica Acta*, 67(22), 4353-4371.
- Godard, G., & van Roermund, H. L. (1995). Deformation-induced clinopyroxene fabrics from eclogites.
   *Journal of Structural Geology*, 17(10), 1425-1443.
- Goffé, B., C. Rangin, and H. Maluski (2002), Jade and associated rocks from the jade Mines area,
   Northern Myanmar as record of a polyphased high pressure metamorphism. Himalaya–Karakoram–
   Tibet Workshop meeting (abstract) *J. Asian Earth Sci.*,20,16–17.
- Guillot, S., Hattori, K., Agard, P., Schwartz, S., & Vidal, O. (2009). Exhumation processes in oceanic and
   continental subduction contexts: a review. *Subduction zone geodynamics*, 175-205.
- Hacker, B. R., Abers, G. A., & Peacock, S. M. (2003). Subduction factory 1. Theoretical mineralogy,
  densities, seismic wave speeds, and H2O contents. *Journal of Geophysical Research: Solid Earth*,
  102 108(B1).
- Halpaap, F., Rondenay, S., Perrin, A., Goes, S., Ottemöller, L., Austrheim, H., ... & Eeken, T. (2019).
  Earthquakes track subduction fluids from slab source to mantle wedge sink. *Science Advances*, 5(4),
  eaav7369.
- Hao, M., Pierotti, C. E., Tkachev, S., Prakapenka, V., & Zhang, J. S. (2019). The single-crystal elastic
  properties of the jadeite-diopside solid solution and their implications for the compositiondependent seismic properties of eclogite, *American Mineralogist*, 104(7), 1016-1021.
- Harlow, G. E., Hemming, S. R., Lallemant, H. G. A., Sisson, V. B., & Sorensen, S. S. (2004). Two high pressure–low-temperature serpentinite-matrix mélange belts, Motagua fault zone, Guatemala: a
   record of Aptian and Maastrichtian collisions. *Geology*, *32*(1), 17-20.
- Harlow, G. E., & Sorensen, S. S. (2005). Jade (nephrite and jadeitite) and serpentinite: metasomatic
  connections. *International Geology Review*, 47(2), 113-146.
- Harlow, G. E., Sisson, V. B., Tsujimori, T., Sorensen, S. S., & Brueckner, H. K. (2008). P–T conditions of
   eclogite/garnet–amphibolite from serpentinite mélanges along the Motagua fault zone, Guatemala.
- 1116 Abs. Ann. Meet. Geol. Soc. Amer.: http://acs. confex. com/crops/2008am/webprogram/Paper50090.
  1117 html.
- Harlow, G. E., Sisson, V. B., & Sorensen, S. S. (2011). Jadeitite from Guatemala: new observations and
  distinctions among multiple occurrences. *Geologica Acta*, 9(3-4), 0363-387.
- Harlow, G. E., Tsujimori, T., & Sorensen, S. S. (2015). Jadeitites and plate tectonics. *Annual Review of Earth and Planetary Sciences*, *43*, 105-138.

- Harlow, G. E., Flores, K. E., & Marschall, H. R. (2016). Fluid-mediated mass transfer from a
  paleosubduction channel to its mantle wedge: evidence from jadeitite and related rocks from the
  Guatemala Suture Zone. *Lithos*, *258*, 15-36.
- Hawthorne, F. C., & Oberti, R. (2007). Classification of the amphiboles. *Reviews in Mineralogy and Geochemistry*, 67(1), 55-88.
- Hellstrom, J., C. Paton, J. Woodhead, and J. Hergt (2008), Iolite: software for spatially resolved LA-(quad
  and MC) ICPMS analysis, *Mineralogical Association of Canada short course series*, 40, 343-348.
- Hermann, J., Spandler, C., Hack, A., & Korsakov, A. V. (2006). Aqueous fluids and hydrous melts in high pressure and ultra-high pressure rocks: implications for element transfer in subduction zones.
   *Lithos, 92*(3-4), 399-417.
- Hertgen, S., Yamato, P., Morales, L. F., & Angiboust, S. (2017). Evidence for brittle deformation events at
  eclogite-facies PT conditions (example of the Mt. Emilius klippe, Western Alps). *Tectonophysics*, *706*,
  1134 1-13.
- Hertwig, A., McClelland, W. C., Kitajima, K., Schertl, H. P., Maresch, W. V., Stanek, K., ... & Sergeev, S. A.
  (2016). Inherited igneous zircons in jadeitite predate high-pressure metamorphism and jadeitite
  formation in the Jagua Clara serpentinite mélange of the Rio San Juan Complex (Dominican
  Republic). *Contributions to Mineralogy and Petrology*, 171(5), 48.
- Hilairet, N., Reynard, B., Wang, Y., Daniel, I., Merkel, S., Nishiyama, N., & Petitgirard, S. (2007). Highpressure creep of serpentine, interseismic deformation, and initiation of subduction. *Science*, *318*(5858), 1910-1913.
- Horn, C., Bouilhol, P., & Skemer, P. (2020). Serpentinization, deformation, and seismic anisotropy in the
  subduction mantle wedge. *Geochemistry, Geophysics, Geosystems, 21*(4), e2020GC008950.
- Huang, J., Hao, J., Huang, F., & Sverjensky, D. (2019). Mobility of chromium in high temperature crustal
  and upper mantle fluids. *Geochemical Perspectives Letters*, 12, 1-6.
- Hussong, D. M. (1981). Tectonic processes and the history of the Mariana arc: a synthesis of the results
  of Deep Sea Drilling Project Leg 60. *Initial reports of the deep sea drilling project*, 60, 909-929.
- Hyndman, R. D., & Peacock, S. M. (2003). Serpentinization of the forearc mantle. *Earth and Planetary Science Letters*, 212(3-4), 417-432.
- 1150 Hyppolito, T., Cambeses, A., Angiboust, S., Raimondo, T., García-Casco, A., & Juliani, C. (2018).
- 1151 Rehydration of eclogites and garnet-replacement processes during exhumation in the amphibolite
- facies. *Geological Society, London, Special Publications*, 478, SP478.473. doi:10.1144/sp478.3.

- Jochum, K. P., et al. (2011a), Determination of Reference Values for NIST SRM 610–617 Glasses Following
   ISO Guidelines, *Geostandards and Geoanalytical Research*, 35(4), 397-429.
- 1155 Jochum, K.P., Wilson, S.A., Abouchami, W., Amini, M., Chmeleff, J., Eisenhauer, A., Hegner, E., Iaccheri,

1156 L.M., Kieffer, B., Krause, J., McDonough, W.F., Mertz-Kraus, R., Raczek, I., Rudnick, R.L., Scholz, D.,

- 1157 Steinhoefel, G., Stoll, B., Stracke, A., Tonarini, S., Weis, D., Weis, U. & Woodhead, J.D. (2011b). GSD-
- 1158 1G and MPI-DING Reference Glasses for In Situ and Bulk Isotopic Determination. *Geostandards and*
- 1159 *Geoanalytical Research*, 35, 193-226, doi: 10.1111/j.1751-908X.2010.00114.x.
- Kawamoto, T., Hertwig, A., Schertl, H. P., & Maresch, W. V. (2018). Fluid inclusions in jadeitite and
  jadeite-rich rock from serpentinite mélanges in northern Hispaniola: Trapped ambient fluids in a
  cold subduction channel. *Lithos*, 308, 227-241.
- Kazak, A.P., Dobretsov, N.L., Moldavantsev, J.E. (1976). Glaucophane schists, jadeitites, vesuvianites and
   nephrites of the Rai-Iz ultramafic massif. *Geology and Geophysics* (in Russian), 2, 60-66.
- Kepezhinskas, P. K., Defant, M. J., & Drummond, M. S. (1995). Na-metasomatism in the island-arc mantle
  by slab melt—peridotite interaction: evidence from mantle xenoliths in the North Kamchatka Arc. *Journal of Petrology*, 36(6), 1505-1527.
- Keppler, R., Stipp, M., Behrmann, J. H., Ullemeyer, K., & Heidelbach, F. (2016). Deformation inside a
   paleosubduction channel–Insights from microstructures and crystallographic preferred orientations
   of eclogites and metasediments from the Tauern Window, Austria. *Journal of Structural Geology*, 82,
- 1171 60-79.
- 1172 Keppler, R. (2018). Crystallographic preferred orientations in eclogites–A review. *Journal of Structural* 1173 *Geology*, 115, 284-296.
- Kirkpatrick, J. D., & Rowe, C. D. (2013). Disappearing ink: How pseudotachylytes are lost from the rock
   record. *Journal of Structural Geology*, *52*, 183-198.
- 1176 Konovalov A.L. & Sergeev S.S. (2015). About the age of zircons from jadeitite ultrabasite array of Syomke
  1177 in the zone of the Main Uralian fault (the Polar Urals). *Regionalnaya geologiya i metallogeniya*. 64,
- 1178 41–47. (In Russian).
- Kuznetsov, Y. V., Kuznetsova, N. A., & Tsyutskiy, S. S. (1986). Features of jadeite mineralization in the
   Pus'yerka deposit (Polar Urals). *International Geology Review*, 28(9), 1096-1101.
- 1181 Lázaro, C., Blanco-Quintero, I. F., Marchesi, C., Bosch, D., Rojas-Agramonte, Y., & García-Casco, A. (2011).
- 1182 The imprint of subduction fluids on subducted MORB-derived melts (Sierra del Convento Mélange,
- 1183 Cuba). *Lithos*, 126(3-4), 341-354.

- Liu, W., Zhang, J., Cao, Y., & Jin, Z. (2020). Geneses of two contrasting antigorite crystal preferred orientations and their implications for seismic anisotropy in the forearc mantle. *Journal of Geophysical Research: Solid Earth*, *125*(9), e2020JB019354.
- Locatelli, M., Verlaguet, A., Agard, P., Federico, L., & Angiboust, S. (2018). Intermediate-depth
  brecciation along the subduction plate interface (Monviso eclogite, W. Alps). *Lithos*, *320*, 378-402.
- Luo, H., & Wang, K. (2021). Postseismic geodetic signature of cold forearc mantle in subduction zones.
   *Nature Geoscience*, 1-6.
- Magloughlin, J. F. (1992). Microstructural and chemical changes associated with cataclasis and frictional
   melting at shallow crustal levels: the cataclasite-pseudotachylyte connection. *Tectonophysics*,
   204(3-4), 243-260.
- Makeyev, A.B. (1992) Mineralogy of alpine-type ultramafics in the Ural. Nauka, St. Petersburg. 195 p. (in
  Russian).
- Maruyama, S., & Liou, J. G. (1988). Petrology of Franciscan metabasites along the jadeite-glaucophane
   type facies series, Cazadero, California. *Journal of Petrology*, 29(1), 1-37.
- Melosh, B. L., Rowe, C. D., Smit, L., Groenewald, C., Lambert, C. W., & Macey, P. (2014). Snap, Crackle,
  Pop: Dilational fault breccias record seismic slip below the brittle–plastic transition. *Earth and Planetary Science Letters*, 403, 432-445.
- Menant, A., Angiboust, S., Monié, P., Oncken, O., & Guigner, J. M. (2018). Brittle deformation during
   Alpine basal accretion and the origin of seismicity nests above the subduction interface. *Earth and Planetary Science Letters*, 487, 84-93.
- Meng, F., Yang, H. J., Makeyev, A. B., Ren, Y., Kulikova, K. V., & Bryanchaninova, N. I. (2016). Jadeitite in
   the Syum-Keu ultramafic complex from Polar Urals, Russia: insights into fluid activity in subduction
   zones. *European Journal of Mineralogy*, 28(6), 1079-1097.
- Meng, F., Makeyev, A. B., & Yang, J. (2011). Zircon U–Pb dating of jadeitite from the Syum-Keu ultramafic
   complex, Polar Urals, Russia: constraints for subduction initiation. *Journal of Asian Earth Sciences*,
   42(4), 596-606.
- 1210 Mével, C., & Kienast, J. R. (1986). Jadeite-kosmochlor solid solution and chromian sodic amphiboles in 1211 jadeitites and associated rocks from Tawmaw (Burma). *Bulletin de Minéralogie*, *109*(6), 617-633.
- 1212 Morimoto, N. (1989). Nomenclature of pyroxenes. *Mineralogical Journal*, 14(5), 198-221.

- 1213 Morishita, T., Arai, S., & Ishida, Y. (2007). Trace element compositions of jadeite (+ omphacite) in 1214 jadeitites from the Itoigawa-Ohmi district, Japan: Implications for fluid processes in subduction 1215 zones. *Island Arc*, 16(1), 40-56.
- Muñoz-Montecinos, J., Angiboust, S., Garcia-Casco, A., Glodny, J., & Bebout, G. (2021b). Episodic
   hydrofracturing and large-scale flushing along deep subduction interfaces: Implications for fluid
   transfer and carbon recycling (Zagros Orogen, southeastern Iran). *Chemical Geology*, 571, 120173.
- Muñoz-Montecinos J, Angiboust S, Garcia-Casco A, (2021a) Blueschist-facies paleo-earthquakes in a
   serpentinite channel enlighten seismogenesis in Mariana-type subduction margins, Accepted (*Earth and planetary science letters*).
- Nakajima, J., & Uchida, N. (2018). Repeated drainage from megathrusts during episodic slow slip. *Nature Geoscience*, *11*(5), 351-356.
- Nyunt, T. T., Massonne, H. J., & Sun, T. T. (2017). Jadeitite and other high-pressure metamorphic rocks
   from the Jade Mines Belt, Tawmaw area, Kachin State, northern Myanmar. *Geological Society, London, Memoirs, 48*(1), 295-315.
- Nyunt, T. T. (2009). Petrological and geochemical contribution to the origin of jadeitite and associated
   rocks of the Tawmaw Area, Kachin State, Myanmar. Stuttgart PhD thesis.
- 1229 Obara, K., & Kato, A. (2016). Connecting slow earthquakes to huge earthquakes. *Science*, 353(6296), 2531230 257.
- Oberti, R., Boiocchi, M., Hawthorne, F. C., Ball, N. A., & Harlow, G. E. (2015). Eckermannite revised: The
   new holotype from the Jade Mine Tract, Myanmar—crystal structure, mineral data, and hints on the
   reasons for the rarity of eckermannite. *American Mineralogist*, 100(4), 909-914.
- 1234 Oncken O, Angiboust S, Dresen G, Slow slip in subduction zones: Reconciling deformation fabrics with 1235 instrumental observations and laboratory results, In revision.
- Ortega-Gutiérrez, F., Solari, L. A., Ortega-Obregon, C., Elias-Herrera, M., Martens, U., Moran-Ical, S., ... &
   Schaaf, P. (2007). The Maya-Chortís boundary: a tectonostratigraphic approach. *International*
- 1238 *Geology Review, 49*(11), 996-1024.
- Orzol, J., Stöckhert, B., Trepmann, C. A., & Rummel, F. (2006). Experimental deformation of synthetic wet
   jadeite aggregates. *Journal of Geophysical Research: Solid Earth*, 111(B6).
- Paton, C., J. Hellstrom, B. Paul, J. Woodhead, and J. Hergt (2011), Iolite: Freeware for the visualisation
  and processing of mass spectrometric data, *Journal of Analytical Atomic Spectrometry*, 26(12), 25082518.

Peacock, S. M. (2003). Thermal structure and metamorphic evolution of subducting slabs. *Geophysical Monograph-American Geophysical Union*, *138*, 7-22.

Pearce, N. J. G., W. T. Perkins, J. A. Westgate, M. P. Gorton, S. E. Jackson, C. R. Neal, and S. P. Chenery
(1997), A compilation of new and published major and trace element data for NIST SRM 610 and
NIST SRM 612 glass reference materials, Geostandards Newsletter - *Journal of Geostandards and Geoanalysis*, 21(1), 115-144.

- Pfiffner, O. A., & Ramsay, J. G. (1982). Constraints on geological strain rates: arguments from finite strain
   states of naturally deformed rocks. *Journal of Geophysical Research: Solid Earth*, 87(B1), 311-321.
- Philippot, P., & van Roermund, H. L. (1992). Deformation processes in eclogitic rocks: evidence for the
   rheological delamination of the oceanic crust in deeper levels of subduction zones. *Journal of structural geology*, 14(8-9), 1059-1077.
- Prigent, C., Guillot, S., Agard, P., Lemarchand, D., Soret, M., & Ulrich, M. (2018). Transfer of subduction
  fluids into the deforming mantle wedge during nascent subduction: Evidence from trace elements
  and boron isotopes (Semail ophiolite, Oman). *Earth and Planetary Science Letters*, 484, 213-228.
- Putnis, A., & Austrheim, H. (2013). Mechanisms of metasomatism and metamorphism on the local
   mineral scale: The role of dissolution-reprecipitation during mineral re-equilibration. In
   *Metasomatism and the chemical transformation of rock* (pp. 141-170). Springer, Berlin, Heidelberg.
- 1261 Raimondo, T., Payne, J., Wade, B., Lanari, P., Clark, C., & Hand, M. (2017). Trace element mapping by LA-
- 1262 ICP-MS: assessing geochemical mobility in garnet. *Contributions to mineralogy and petrology*, 1263 172(4), 17.
- 1264 Reynard, B. (2013). Serpentine in active subduction zones. *Lithos, 178,* 171-185.
- Ridd, M. F., Crow, M. J., & Morley, C. K. (2019). The role of strike-slip faulting in the history of the
  Hukawng Block and the Jade Mines Uplift, Myanmar. *Proceedings of the Geologists' Association*,
  1267 130(2), 126-141.
- Rowe, C. D., Meneghini, F., & Moore, J. C. (2011). Textural record of the seismic cycle: Strain-rate
  variation in an ancient subduction thrust. *Geological Society, London, Special Publications, 359*(1),
  77-95.
- Savelieva, G. N., Sharaskin, A. Y., Saveliev, A. A., et al. (2002). Ophiolites and zoned mafic–ultramafic
   massifs of the Urals: a comparative analysis and some tectonic implications. *Mountain building in the Uralides*, 135-153.

- Savelieva, G. N., & Suslov, P. V. (2014). Structure and composition of mantle peridotites at the boundary
  with crustal complexes of ophiolites in the Syumkeu massif, Polar Urals. *Geotectonics*, 48(5), 347358.
- Savelieva, G. N., Batanova, V. G., & Sobolev, A. V. (2016). Pyroxene–Cr-spinel exsolution in mantle
   Iherzolites of the Syum-Keu ophiolite massif (Arctic Urals). *Russian Geology and Geophysics*, *57*(10),
   1419-1436.
- Scambelluri, M., Cannaò, E., & Gilio, M. (2019). The water and fluid-mobile element cycles during
   serpentinite subduction. A review. *European Journal of Mineralogy*, 31(3), 405-428.
- Schertl, H. P., Maresch, W. V., Stanek, K. P., Hertwig, A., Krebs, M., Baese, R., & Sergeev, S. S. (2012). New
  occurrences of jadeitite, jadeite quartzite and jadeite-lawsonite quartzite in the Dominican Republic,
  Hispaniola: petrological and geochronological overview. *European Journal of Mineralogy*, 24(2), 199216.
- Searle, M. P., Noble, S. R., Cottle, J. M., Waters, D. J., Mitchell, A. H. G., Hlaing, T., & Horstwood, M. S. A.
  (2007). Tectonic evolution of the Mogok metamorphic belt, Burma (Myanmar) constrained by U-ThPb dating of metamorphic and magmatic rocks. *Tectonics*, *26*(3).
- Shi, G. H., Cui, W. Y., Tropper, P., Wang, C. Q., Shu, G. M., & Yu, H. (2003). The petrology of a complex
  sodic and sodic–calcic amphibole association and its implications for the metasomatic processes in
  the jadeitite area in northwestern Myanmar, formerly Burma. *Contributions to Mineralogy and Petrology*, *145*(3), 355-376.
- Shi, G. H., Jiang, N., Liu, Y., Wang, X., Zhang, Z. Y., & Xu, Y. J. (2009a). Zircon Hf isotope signature of the
  depleted mantle in the Myanmar jadeitite: implications for Mesozoic intra-oceanic subduction
  between the Eastern Indian Plate and the Burmese Platelet. *Lithos*, *112*(3-4), 342-350.
- Shi, G., Wang, X., Chu, B., & Cui, W. (2009b). Jadeite jade from Myanmar: its texture and gemmological
  implications. *The journal of gemmology and proceedings of the Gemmological Association of Great Britain*, 31(5), 185.
- Shi, G., Harlow, G. E., Wang, J., Wang, J., Ng, E., Wang, X., ... & Cui, W. (2012). Mineralogy of jadeitite and
  related rocks from Myanmar: a review with new data. *European Journal of Mineralogy*, 24(2), 345370.
- Shigeno, M., Mori, Y., Shimada, K., & Nishiyama, T. (2012). Jadeitites with metasomatic zoning from the
   Nishisonogi metamorphic rocks, western Japan: fluid–tectonic block interaction during exhumation.
   *European Journal of Mineralogy*, 24(2), 289-311.

- Shmelev, V. R. (2011). Mantle ultrabasites of ophiolite complexes in the Polar Urals: petrogenesis and
   geodynamic environments. *Petrology*, *19*(6), 618-640.
- Sibson, R. H. (1986). Brecciation processes in fault zones: inferences from earthquake rupturing. *Pure and Applied Geophysics*, *124*(1), 159-175.
- Sibson, R. H. (1989). Earthquake faulting as a structural process. *Journal of structural geology*, *11*(1-2), 11310 14.
- Sibson, R. H., Toy, V. G., Abercrombie, R., & McGarr, A. (2006). The habitat of fault-generated
   pseudotachylyte: Presence vs. absence of friction-melt. *GEOPHYSICAL MONOGRAPH-AMERICAN GEOPHYSICAL UNION*, *170*, 153.
- 1314 Snoke, A. W., Tullis, J., & Todd, V. R. (Eds.). (2014). *Fault-related rocks: a photographic atlas* (Vol. 410).
  1315 Princeton University Press.
- Sorensen, S., Harlow, G. E., & Rumble, D. (2006). The origin of jadeitite-forming subduction-zone fluids:
   CL-guided SIMS oxygen-isotope and trace-element evidence. *American Mineralogist*, *91*(7), 979-996.
- 1318 Sorensen, S. S., Sisson, V. B., Harlow, G. E., & Avé Lallemant, H. G. (2010). Element residence and
- transport during subduction-zone metasomatism: evidence from a jadeitite-serpentinite contact,
  Guatemala. *International Geology Review*, *52*(9), 899-940.
- Soret, M., Agard, P., Dubacq, B., Vitale-Brovarone, A., Monie, P., Chauvet, A., ... & Villemant, B. (2016).
  Strain localization and fluid infiltration in the mantle wedge during subduction initiation: Evidence
  from the base of the New Caledonia ophiolite. *Lithos*, 244, 1-19.
- Spandler, C., & Pirard, C. (2013). Element recycling from subducting slabs to arc crust: A review. *Lithos*,
  170, 208-223.
- Swanson, M. T. (1992). Fault structure, wear mechanisms and rupture processes in pseudotachylyte
   generation. *Tectonophysics*, 204(3-4), 223-242.
- Sychev, S. N., & Kulikova, K. V. (2012). Structural evolution of the Main Ural Fault Zone in the western
   framework of the Voikar-Synya ophiolite massif. *Geotectonics*, *46*(6), 427-434.
- Syracuse, E. M., van Keken, P. E., & Abers, G. A. (2010). The global range of subduction zone thermal
   models. *Physics of the Earth and Planetary Interiors*, *183*(1-2), 73-90.
- 1332 Takahashi, N., Tsujimori, T., Kayama, M., & Nishido, H. (2017). Cathodoluminescence petrography of P-
- type jadeitites from the New Idria serpentinite body, California. *Journal of Mineralogical and Petrological Sciences*, *112*(5), 291-299.

- Torres-Roldan, R. L., Garcia-Casco, A., & Garcia-Sanchez, P. A. (2000). CSpace: An integrated workplace
   for the graphical and algebraic analysis of phase assemblages on 32-bit Wintel platforms. *Computers* & *Geosciences*, *26*(7), 779-793.
- Tsujimori, T., & Harlow, G. E. (2012). Petrogenetic relationships between jadeitite and associated high pressure and low-temperature metamorphic rocks in worldwide jadeitite localities: a review.
   *European Journal of Mineralogy*, 24(2), 371-390.
- Tsujimori, T., Liou, J. G., & Coleman, R. G. (2004, November). Comparison of two contrasting eclogites
   from the Motagua fault zone, Guatemala: southern lawsonite eclogite versus northern zoisite
   eclogite. In *Geological Society of America Abstracts with Programs* (Vol. 36, No. 5, p. 136).
- 1344 Uchida, N., Kirby, S. H., Okada, T., Hino, R., & Hasegawa, A. (2010). Supraslab earthquake clusters above
- 1345 the subduction plate boundary offshore Sanriku, northeastern Japan: Seismogenesis in a graveyard
- 1346 of detached seamounts?. *Journal of Geophysical Research: Solid Earth*, 115(B9).
- 1347 Udovkina, N.G. (1971). Eclogites of Polar Urals Nauka, Moscow (in Russian).
- Ulrich, S., & Mainprice, D. (2005). Does cation ordering in omphacite influence development of lattice preferred orientation?. *Journal of Structural Geology*, 27(3), 419-431.
- 1350 Wang, K., Huang, T., Tilmann, F., Peacock, S. M., & Lange, D. (2020). Role of Serpentinized Mantle Wedge
- in Affecting Megathrust Seismogenic Behavior in the Area of the 2010 M= 8.8 Maule Earthquake.
   Geophysical Research Letters, 47(22), e2020GL090482.
- White, L., Rawlinson N., Lister G., Tanner D., Macpherson C., and Morgan J. (2016). Pulses of earthquake
  activity in the mantle wedge track the route of slab fluid ascent. EGU General Assembly Conference
  Abstracts, 18. 2016.
- Whitney, D. L., & Evans, B. W. (2010). Abbreviations for names of rock-forming minerals. *American mineralogist*, *95*(1), 185-187.
- Woodcock, N. H., Dickson, J. A. D., & Tarasewicz, J. P. T. (2007). Transient permeability and reseal
   hardening in fault zones: evidence from dilation breccia textures. *Geological Society, London, Special Publications, 270*(1), 43-53.
- Woodcock, N. H., & Mort, K. (2008). Classification of fault breccias and related fault rocks. *Geological Magazine*, *145*(3), 435-440.

1363 Woodhead, J. D., J. Hellstrom, J. M. Hergt, A. Greig, and R. Maas (2007). Isotopic and Elemental Imaging

of Geological Materials by Laser Ablation Inductively Coupled Plasma-Mass Spectrometry,
 *Geostandards and Geoanalytical Research*, 31(4), 331-343.

- Yu, Z., & Zhao, D. (2020). Seismic evidence for water transportation in the forearc off Northern Japan. *Journal of Geophysical Research: Solid Earth*, 125(4), e2019JB018600.
- Yui, T. F., Fukoyama, M., Iizuka, Y., Wu, C. M., Wu, T. W., Liou, J. G., & Grove, M. (2013). Is Myanmar
  jadeitite of Jurassic age? A result from incompletely recrystallized inherited zircon. *Lithos*, *160*, 268282.
- Yui, T. F., Maki, K., Wang, K. L., Lan, C. Y., Usuki, T., Iizuka, Y., ... & Grove, M. (2012). Hf isotope and REE
  compositions of zircon from jadeitite (Tone, Japan and north of the Motagua fault, Guatemala):
  implications on jadeitite genesis and possible protoliths. *European Journal of Mineralogy*, 24(2),
  263-275.
- Zhang, J., Green II, H. W., & Bozhilov, K. N. (2006). Rheology of omphacite at high temperature and
   pressure and significance of its lattice preferred orientations. *Earth and Planetary Science Letters*,
   246(3-4), 432-443.
- 1378
- 1379

**Table 1.** Summary of selected samples for this comparative study, including their paragenesis and the various fracturing patterns identified therein (see text for details on the criteria used for this classification). Numbers in the last three columns refer to the chronological sequence of fracturing events identified in each sample.

					Frac	Fracturing pattern	
	Sample	Region	Pre-fracturing assemblage	Paragenesis associated w/ brittle def.	type I	type II	type III
Polar Urals	PU2	Pus'yerka	white Jd (Jd <sub>1</sub> ), Ed	Omp, Mg-ktp, Rct, Eck, Phl, Clc		x (1)	x (2)
	PU5	Pus'yerka	white Jd (Jd <sub>1</sub> )	Omp, Phl, Clc (±Cal)		х	
	UR11b	Pus'yerka	Ed or Mg-ktp with Omp inc.	Jd <sub>2</sub> , Omp, Mg-ktp, Eck/Nyb, Ttn, Chr	х		
	UR03b,03d	Pus'yerka	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Omp			x
	UR25	Nephrite brook	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Omp, Ttn	х		
Guatemala	MTG00	Motagua region	white Jd	Omp (not analyzed)	х		
	MTG01	South Motagua	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Omp, Lws, Ttn	х		
	MTG02	South Motagua	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Omp, Ttn			х
	MTG03	South Motagua	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Omp, Ttn (±REE-rich Ep)		х	
	MTG12	South Motagua	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Omp, Ttn		х	
Myanmar	KAS01	Hpakan (Kashin)	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Omp, Eck	x (1)		x (2)
	KAS04	Hpakan (Kashin)	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Omp, Eck, Clc (±Cls)	x (1)	x (2)	x (3)
	KAS06	Hpakan (Kashin)	Ed (+Omp?)	Jd <sub>2</sub> , Omp, Rct, Eck	х		
	KAS07	Hpakan (Kashin)	white Jd (Jd1) and Mg-ktp	Jd <sub>2</sub> , Omp, Eck, Chr		x (1)	x (2)
	KAS10	Hpakan (Kashin)	white Jd (Jd <sub>1</sub> )	Jd <sub>2</sub> , Eck		?	

Table 1



H<sub>2</sub>O, Ca,Fe,Mg (+LILEs, ±Cr, ±Ti ±REEs)

Present-day occurrence in serpentinite mélanges