



**HAL**  
open science

## X-ray haloes and star formation in early-type galaxies

Andrea Negri, Silvia Pellegrini, Luca Ciotti

► **To cite this version:**

Andrea Negri, Silvia Pellegrini, Luca Ciotti. X-ray haloes and star formation in early-type galaxies. Monthly Notices of the Royal Astronomical Society, 2015, 451, pp.1212-1228. 10.1093/mnras/stv968 . insu-03644883

**HAL Id: insu-03644883**

**<https://insu.hal.science/insu-03644883v1>**

Submitted on 20 Apr 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# X-ray haloes and star formation in early-type galaxies

Andrea Negri,<sup>1,2★</sup> Silvia Pellegrini<sup>1</sup> and Luca Ciotti<sup>1</sup>

<sup>1</sup>*Department of Physics and Astronomy, University of Bologna, viale Bertoni Pichat 6/2, I-40127 Bologna, Italy*

<sup>2</sup>*Institut d'Astrophysique de Paris, 98bis Boulevard Arago, F-75014 Paris, France*

Accepted 2015 April 28. Received 2015 April 24; in original form 2015 January 28

## ABSTRACT

High-resolution 2D hydrodynamical simulations describing the evolution of the hot interstellar medium (ISM) in axisymmetric two-component models of early-type galaxies well reproduced the observed trends of the X-ray luminosity ( $L_X$ ) and temperature ( $T_X$ ) with galaxy shape and rotation, however they also revealed the formation of an exceedingly massive cooled gas disc in rotating systems. In a follow-up of this study, here we investigate the effects of star formation in the disc, including the consequent injection of mass, momentum and energy in the pre-existing ISM. It is found that subsequent generations of stars originate one after the other in the equatorial region; the mean age of the new stars is  $>5$  Gyr, and the adopted recipe for star formation can reproduce the empirical Kennicutt–Schmidt relation. The results of the previous investigation without star formation, concerning  $L_X$  and  $T_X$  of the hot gas, and their trends with galactic shape and rotation, are confirmed. At the same time, the consumption of most of the cold gas disc into new stars leads to more realistic final systems, whose cold gas mass and star formation rate agree well with those observed in the local Universe. In particular, our models could explain the observation of kinematically aligned gas in massive, fast-rotating early-type galaxies.

**Key words:** ISM: evolution – galaxies: elliptical and lenticular, cD – galaxies: ISM – galaxies: kinematics and dynamics – X-rays: galaxies – X-rays: ISM.

## 1 INTRODUCTION

Early-type galaxies (ETGs) are embedded in hot gaseous haloes produced mainly by stellar winds, and heated to X-ray temperatures by Type Ia supernovae (SNIa) explosions and by the thermalization of stellar motions (Fabbiano 1989; Ciotti et al. 1991; David, Forman & Jones 1991; O’Sullivan, Forbes & Ponman 2001). The thermalization is due to the interaction between the stellar and SNIa ejecta and the pre-existing hot interstellar medium (ISM; e.g. Mathews 1989; Parriott & Bregman 2008). Recent high-resolution 2D hydrodynamical simulations of hot gas flows (Negri, Ciotti & Pellegrini 2014a; Negri et al. 2014b, hereafter N14) showed that the presence of ordered rotation in the stellar component can alter significantly the ISM evolution with respect to that shown by fully velocity dispersion supported systems of same total mass and mass distribution. First, it is found that the rotation field of the ISM in rotating galaxies is very similar to that of the stars, with a consequent negligible heating contribution from thermalization of the ordered motions. Secondly, conservation of angular momentum in the ISM of rotating galaxies results in the formation of a centrifugally supported cold equatorial disc, with the consequent reduction of both the X-ray luminosity  $L_X$  and temperature  $T_X$  of the hot ISM. These

results compared well with observations, which show a dependence of  $L_X$  and  $T_X$  on the galactic shape and internal dynamics:  $L_X$  is observed to be high only in round and slowly rotating galaxies, and is limited to lower values for flatter, fast-rotating ones (Eskridge, Fabbiano & Kim 1995; Pellegrini, Held & Ciotti 1997; Sarzi et al. 2010, 2013; Li et al. 2011; Pellegrini 2012);  $T_X$  of slowly rotating systems is consistent just with the thermalization of the stellar random kinetic energy, estimated from  $\sigma_e$  (the stellar velocity dispersion averaged within one effective radius  $R_e$ ), while fast-rotating systems show  $T_X$  values below 0.4 keV, and not scaling with  $\sigma_e$  (Sarzi et al. 2013, see also Pellegrini 2011, Posacki, Pellegrini & Ciotti 2013).

A major outcome of the previous simulations is that cold material can be accumulated in considerable amounts, during the lifetime of rotating galaxies. The cold gas typically settles in the equatorial plane, where it forms an extended disc (of 0.5–10 kpc radius), that can be as massive as  $\simeq 10^{10} M_\odot$ , in the largest galaxies. This result brings in a few important questions: Are these cold discs observed? Do they become, as seems natural, a site for star formation (hereafter SF)? Is SF observed? What is the impact of SF in the disc on  $L_X$  and  $T_X$ ? In recent years, evidence has been accumulating that ETGs host significant quantities of cold gas, in the form of atomic and molecular hydrogen (Morganti et al. 2006; Combes, Young & Bureau 2007; di Serego Alighieri et al. 2007; Grossi et al. 2009; Young et al. 2011); approximately 50 per cent of massive ETGs (of

\*E-mail: [andrea.negri@unibo.it](mailto:andrea.negri@unibo.it)

stellar mass  $M_* \gtrsim 10^{10} M_\odot$ ) contain  $10^7$ – $10^9 M_\odot$  of H I and/or H<sub>2</sub> (Cappellari et al. 2011; Davis et al. 2011, 2013; Serra et al. 2012, 2014; Young et al. 2014). A large, systematic investigation of the ATLAS<sup>3D</sup> sample of ETGs with the Westerbork Synthesis Radio Telescope found that  $\simeq 40$  per cent of galaxies outside Virgo, and  $\simeq 10$  per cent of galaxies inside it, are detected in H I, with  $M_{\text{HI}} \gtrsim 10^7 M_\odot$ ; the majority (2/3) of the detections consists of settled configurations, where the cold gas is in discs or rings. Small discs (size of a few kpc), confined within the stellar body, share the same kinematics of the stars; large discs (up to  $5 \times 10^9 M_\odot$ ) extend to tens of kpc, and in half of the cases are kinematically misaligned with the stars (Serra et al. 2012). In particular, fast-rotating Virgo galaxies have kinematically aligned gas, and the most massive ( $M_* \gtrsim 8 \times 10^{10} M_\odot$ ) fast-rotating ETGs always have kinematically aligned gas, independent of environment; this alignment leads to hypothesize that the gas has been internally generated (Davis et al. 2011, 2013). The H I discs/rings around slowly rotating ETGs, instead, are usually not fully settled, which suggests an external origin (in mergers, or in accretion from satellite galaxies, or from the intergalactic medium); an external origin was considered likely also for those ETGs showing stellar/gas kinematic misalignment. Interestingly, while H I is more ubiquitous, molecular gas is detected only in fast rotators across the entire ATLAS<sup>3D</sup> sample (Davis et al. 2011, 2013; Young et al. 2011, 2014).

Besides giving indication about its origin through its morphology and kinematics, the observed cold gas seems also to provide material for SF. Low level SF activity is present in approximately one-third of ETGs (Yi et al. 2005; Suh et al. 2010; Ko et al. 2014); Sarzi et al. (2006) showed ongoing SF signatures in the optical spectra of at least  $\sim 10$  per cent of nearby ETGs. In the ATLAS<sup>3D</sup> sample, galaxies with H I within  $\sim 1 R_e$  exhibit ongoing SF in  $\sim 70$  per cent of the cases,  $\sim 5$  times more frequently than galaxies without H I (Serra et al. 2012). Interestingly, as for molecular gas, some degree of SF and young stellar populations are detected only in fast rotators, in the ATLAS<sup>3D</sup> sample (Kuntschner et al. 2010; Sarzi et al. 2013). Integral-field spectroscopy showed that ETGs host frequently a rotating stellar component younger and more metal rich than the bulge (Krajnović et al. 2008). The presence of this component, and the occurrence of SF, imply an important role for the cold gas during the evolution of ETGs (Khochfar et al. 2011; Cappellari et al. 2013; Naab et al. 2014).

Finally, the cold gas content of ETGs is also an important prediction of  $\Lambda$  cold dark matter hydrodynamical simulations of galaxy formation, that include cold gas evolution (e.g. Martig et al. 2013), and accretion via various processes during the secular evolution of galaxies (Oser et al. 2010, see also Lagos et al. 2014, Dubois et al. 2013).

In conclusion, the cold gas has become a tool to gain insight into recent (and less recent) galaxy evolution. In order to correctly interpret the variety of observational results, and to use them properly as constraints for different scenarios for the origin of the structure and SF history of ETGs, it is crucial to establish what is the relative importance of the various gas production processes (internal and external to galaxies), gas depletion ones (AGN and SF-driven outflows, environmental stripping), and gas consumption ones (SF). Cold gas is usually thought to come from accretion from the surrounding medium or satellites, as well as from gas-rich mergers; in the cases of the giant central-dominant galaxies in groups or clusters it can also come from cooling of hot gas (Edge et al. 2010; McDonald et al. 2011). The numerical investigation of N14 showed an additional *internal* contribution of cold gas, coming from the evolution of the passive stellar population, that can be substantial,

or even too large with respect to observed values. A natural sequel of the N14 work should address the questions of whether this gas can possibly lead to SF, when SF takes place, and what is the fate of the cold discs, whether they are consumed or they are continuously replenished by cooling hot gas. In this paper, we add SF to the simulations of N14, and we explore the ISM evolution including the removal of cold gas, and the injection of mass, momentum and energy appropriate for the newly (and continuously) forming stellar population. In this way we aim at establishing whether (i) the N14 results for the general trends of the hot gas properties with galaxy shape and stellar kinematics still hold; (ii) the formation of stars can reduce the amount of cold gas in the simulations, thus bringing it more in agreement with observed values; (iii) a significant channel for SF, previously neglected, should be taken into consideration for rotating systems, and whether this can account for the low-level SF activity currently seen to be ongoing.

This paper is organized as follows. In Section 2, we describe the main ingredients of the simulations, such as the galaxy models and the input physics. In Section 3, we present and discuss the results of the simulations. In Section 4, we summarize the conclusions.

## 2 THE SIMULATIONS

N14 performed a large set of 2D hydrodynamical simulations with the ZEUS MP2 code to fully explore the large parameter space of realistic (axisymmetric) galaxy models, characterized by different stellar mass, intrinsic flattening, distribution of dark matter, and internal kinematics. The galaxy flattening was either fully supported by ordered rotation, originating the set of models that are isotropic rotators, or by tangential anisotropy, originating the set of fully velocity dispersion supported models. These two extreme configurations were built adopting the Satoh decomposition, respectively, with Satoh parameter  $k = 1$  and 0. The galaxy models were tailored to reproduce the observed properties and scaling laws of ETGs (see also Posacki et al. 2013). In this work, we perform hydrodynamical simulations for a representative subset of rotating models already investigated by N14 (Section 2.1), including SF in the simulations (as described in Section 2.2 below), but keeping the code (numerical set-up, grid properties) in all equal to that used by N14 (see N14 for details). Also, a less extreme value of  $k = 0.1$  is explored. A logarithmically spaced numerical mesh ( $R, z$ ) of  $960 \times 480$  gridpoints is employed, with a resolution of 90 pc in the first 10 kpc from the centre.

### 2.1 The galaxy models

N14 built axisymmetric two-component galaxy models where the stellar component has two different intrinsic flattening, corresponding to the E4 and E7 shapes, while the dark matter halo is spherical. The luminous matter is described by the deprojection (Mellier & Mathez 1987) of the de Vaucouleurs (1948) law, generalized for ellipsoidal axisymmetric distributions; the dark matter profile is the (Navarro, Frenk & White 1997) one, with the dark mass  $M_h$  amounting at  $\simeq 20$  times the total stellar mass  $M_*$ . For any fixed galaxy mass and shape (E4 or E7), N14 built two models: the first one, called ‘FO-built’, when seen face-on has the same  $R_e$  of the spherical E0 counterpart, thus its stellar mass distribution becomes more and more concentrated than in the E0 model, as it gets flatter; the second one, called ‘EO-built’, when seen edge-on has the same circularized  $R_e$  of the E0 counterpart, which makes its stellar mass distribution to expand with increasing flattening.

**Table 1.** Main properties of the galaxy models.

Name	$L_B$ ( $10^{11} L_\odot$ )	$R_e$ (kpc)	$M_*$ ( $10^{11} M_\odot$ )	$M_h$ ( $10^{11} M_\odot$ )	$\sigma_{e8}$ ( $\text{km s}^{-1}$ )	$f_{DM}$	$c$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
EO4 <sup>200</sup> <sub>IS</sub>	0.26	4.09	1.25	25	166	0.63	37
EO7 <sup>200</sup> <sub>IS</sub>	0.26	4.09	1.25	25	124	0.66	37
FO4 <sup>200</sup> <sub>IS</sub>	0.26	4.09	1.25	25	178	0.59	37
FO7 <sup>200</sup> <sub>IS</sub>	0.26	4.09	1.25	25	150	0.57	37
EO4 <sup>250</sup> <sub>IS</sub>	0.62	7.04	3.35	67	207	0.62	28
EO7 <sup>250</sup> <sub>IS</sub>	0.62	7.04	3.35	67	154	0.66	28
EO4 <sup>300</sup> <sub>IS</sub>	1.32	11.79	7.80	160	248	0.64	22
EO7 <sup>300</sup> <sub>IS</sub>	1.32	11.79	7.80	160	185	0.68	22
FO4 <sup>300</sup> <sub>IS</sub>	1.32	11.79	7.80	160	266	0.60	22
FO7 <sup>300</sup> <sub>IS</sub>	1.32	11.79	7.80	160	224	0.59	22

*Notes.* (1) Model name: ‘EO’ or ‘FO’ indicate the procedure (EO-building or FO-building) applied to the spherical (E0) model to obtain the shape indicated by the number (E4 or E7 shapes); the superscript indicates  $\sigma_{e8}$  of the corresponding E0 model; the subscript ‘IS’ indicates the kinematical configuration of the isotropic rotator (for example, FO4<sup>200</sup><sub>IS</sub> is an E4 isotropic rotator, FO-built from a spherical model with  $\sigma_{e8} = 200 \text{ km s}^{-1}$ ). (2) Luminosities in the  $B$  band. (3) Effective radius (for an FO view for FO-built models, and an EO view for EO-built models). (4) Total stellar mass of the original (old) stellar population. The first four models are LM models, the next two are intermediate mass models, and the last four are HM models. (5) Total dark matter mass. (6) Luminosity-weighted average of the stellar velocity dispersion within a circular aperture of radius  $R_{e/8}$ ; for non-spherical models,  $\sigma_{e8}$  is the edge-on viewed value. (7) Dark matter fraction enclosed within a sphere of radius  $R_e$ . (8) Concentration parameter of the dark matter profile.

The properties listed above, for the EO4<sup>250</sup> and EO7<sup>250</sup> models with  $k = 0.1$ , are not reported, since they are equal to those of the EO4<sup>250</sup><sub>IS</sub> and EO7<sup>250</sup><sub>IS</sub> ones, except for  $\sigma_{e8}$ , which is, respectively,  $223 \text{ km s}^{-1}$  for the EO4<sup>250</sup> <sub>$k=0.1$</sub>  model, and  $184 \text{ km s}^{-1}$  for the EO7<sup>250</sup> <sub>$k=0.1$</sub>  model.

In this work, we re-simulate a few flat (E4 and E7) rotating models (isotropic rotators, with  $k = 1$ ) of N14 including SF in them. In order to explore the effects of SF at the high and low ends of the galaxy mass range explored by N14, we choose four flat models with luminosity-weighted stellar velocity dispersion within  $R_{e/8}$  of  $\sigma_{e8} = 300 \text{ km s}^{-1}$  for the parent spherical model (and we call these high-mass models, ‘HM’ models), and four with  $\sigma_{e8} = 200 \text{ km s}^{-1}$  for the E0 counterpart (low-mass models, ‘LM’ models). In N14, the first set was found to host inflows, with the creation of a massive, centrifugally supported cold gaseous disc, while the second set was found in a global wind, or close to the transition to it. In a global wind, the gas has very low density and positive velocity (directed outwards) through most of the galaxy (e.g. Mathews & Baker 1971).<sup>1</sup> In addition, we re-simulated two intermediate-mass models (again E4 and E7), with  $\sigma_{e8} = 250 \text{ km s}^{-1}$  for the E0 counterpart. For these two models, we also built a moderately rotating stellar kinematical configuration ( $k = 0.1$ ), not explored by N14. The main structural properties of the ten re-simulated models, identical to those presented in N14, are listed in Table 1; the two new models with  $k = 0.1$  differ from the intermediate-mass models only in the  $\sigma_{e8}$  value, which is given in the notes to the table.

<sup>1</sup> In fact the phase of the gas flow (that can range from a wind to an inflow) is basically determined by the different relative importance of SN heating in galaxies of different mass, as already thoroughly discussed in Ciotti et al. 1991.

## 2.2 Star formation in the code

We employed two different schemes for SF, a passive (pSF) one and an active (aSF) one. In the pSF, only cold gas removal from the numerical mesh is allowed; in the aSF, we also consider the injection of mass, momentum and energy from the newly forming stellar population, for simplicity limiting in this work to the evolution of stars more massive than  $8 M_\odot$  (ending with SNII explosions); note however that for reasonable stellar initial mass functions (IMF) these massive stars are major contributors to the total mass return rate. We describe below the main inputs and sinks of mass and energy for the gas flow, due to both the original stellar population (of total mass  $M_*$ ) and the newly forming stars, and the corresponding equations of hydrodynamics solved by the code.

### 2.2.1 The mass injection and sink terms

The mass inputs are stellar winds and SNIa’s ejecta produced during the passive evolution of the original stellar population of the galaxy (at a rate per unit volume, respectively, of  $\dot{\rho}_*$  and  $\dot{\rho}_{\text{Ia}}$ , for which we adopt the standard recipes coming from the stellar evolution theory; e.g. N14), and the Type II supernovae ejecta produced by the newly born stellar population (at a rate of  $\dot{\rho}_{\text{II}}$ , calculated as detailed below).

The mass sink is due to SF subtracting gas from the grid, at an adopted rate per unit volume of:

$$\dot{\rho}_{\text{SF}} = \frac{\eta_{\text{SF}} \rho}{t_{\text{SF}}}, \quad t_{\text{SF}} = \max(t_{\text{cool}}, t_{\text{dyn}}), \quad (1)$$

where  $\rho$  is the gas density, and  $\eta_{\text{SF}}$  is the SF efficiency, for which we adopt two values of  $\eta_{\text{SF}} = 0.01$  and  $0.1$ . The rate  $\dot{\rho}_{\text{SF}}$  depends on the maximum between the cooling time-scale  $t_{\text{cool}} \equiv E/\mathcal{L}$  (where  $E$  is the ISM internal energy density, and  $\mathcal{L}$  is the ISM bolometric luminosity per unit volume), and the dynamical time-scale  $t_{\text{dyn}} \equiv \sqrt{3\pi/32G\rho}$  (see also Ciotti & Ostriker 2007).

When SF takes place in a computational cell, the ISM is removed and an equal mass in stars  $\Delta M_*$  appears at the same place. These new stars are assumed to form with a Salpeter IMF, thus for a given  $\Delta M_*$ , the number of stars having a mass greater than  $8 M_{\odot}$ , that will explode as SNII, is  $N_{\text{II}} \simeq 7 \times 10^{-3} \Delta M_*(M_{\odot})$ . These new stars in turn inject mass into the ISM. By integration of the mass difference between the mass of the progenitor and that of the remnant, for the Salpeter IMF, one finds that the mass injected by SNII's is  $\simeq 0.2\Delta M_*$ . The final SNII mass source term  $\dot{\rho}_{\text{II}}$  comes from considering that a given SF episode generates SNII's that inject mass at a rate exponentially declining on a time-scale  $\tau_{\text{II}} = 2 \times 10^7$  yr, and that at a certain time during the evolution of an SF episode, another episode may take place, forming younger SNII's that in turn eject mass into the ISM. The whole process can be formalized as

$$\frac{d\dot{\rho}_{\text{II}}}{dt} = -\frac{\dot{\rho}_{\text{II}}}{\tau_{\text{II}}} + \frac{0.2\dot{\rho}_{\text{SF}}}{\tau_{\text{II}}}, \quad (2)$$

(Ciotti & Ostriker 2007). The equation above is very useful in numerical works, as it allows us to compute the mass return of the new stars formed at each timestep without storing the whole SF history at each gridpoint, but only the current value and the value of the mass return at the previous timestep. For more complex recipes, built on the scheme above, see Calura, Ciotti & Nipoti (2014).

### 2.2.2 The energy injection and sink terms

Energy is injected into the ISM by the thermalization of the kinetic energy of SNIa and SNII explosions, and by the thermalization of random and streaming stellar motions for the stellar winds. The rate of SNIa's explosions is the same entering  $\dot{\rho}_{\text{Ia}}$  defined in Section 2.2.1. The energy input rate per unit volume from the original (old) stellar population is then:

$$\dot{E}_{\text{old}} = \dot{E}_{\text{Ia}} + \frac{\dot{\rho}_{\text{Ia}} + \dot{\rho}_*}{2} \left[ \left\| v_{\varphi} \hat{e}_{\varphi} - \mathbf{u} \right\|^2 + \text{Tr}(\boldsymbol{\sigma}^2) \right], \quad (3)$$

where  $\dot{E}_{\text{Ia}} = \dot{\rho}_{\text{Ia}} \vartheta_{\text{SNIa}} E_{\text{SN}} / 1.4 M_{\odot}$ ,  $\vartheta_{\text{SNIa}}$  is the thermalization efficiency (for which we adopt the value of 0.85, see Thornton et al. 1998; Tang & Wang 2005),  $E_{\text{SN}} = 10^{51}$  erg, and  $1.4 M_{\odot}$  is the mass that is ejected by one SNIa event;  $v_{\varphi}$  is the stellar streaming velocity field,  $\mathbf{u}$  is the velocity of the ambient gas,  $\boldsymbol{\sigma}^2$  is the stellar velocity dispersion tensor of the stars.

The energy input rate  $\dot{E}_{\text{new}}$  from the newly born stellar population is derived as follows. A mass of newly formed stars  $\Delta M_*$  injects energy through SNII with an efficiency

$$\varepsilon_{\text{II}} = \frac{N_{\text{II}} E_{\text{SN}}}{\Delta M_* c^2} \simeq 3.9 \times 10^{-6}. \quad (4)$$

Consistently with equation (2), the SNII energy injection rate  $\dot{E}_{\text{II}}$  due to the thermalization of the ejecta is given by

$$\frac{d\dot{E}_{\text{II}}}{dt} = -\frac{\dot{E}_{\text{II}}}{\tau_{\text{II}}} + \frac{\varepsilon_{\text{II}} c^2 \dot{\rho}_{\text{SF}} \vartheta_{\text{SNII}}}{\tau_{\text{II}}}, \quad (5)$$

where we take the fiducial value of  $\vartheta_{\text{SNII}} = \vartheta_{\text{SNIa}}/5 = 0.17$  to account for the lower thermalization efficiency of SNII exploding in a cold and dense medium. We assume that the new stars inherit the kinematical configuration of the original stellar component in the

place where they are born (i.e. the velocity dispersion and rotation of the new stars are the same as those of the original stellar distribution in the same place). This allows us to treat the heating terms described in equation (7) below by using the same properties of the old stars. Note however that this choice is not unreasonable, because the bulk of SF takes place in the cold and rotationally supported disc, whose rotational properties are very similar to those of the stars. Thus, the total heating rate per unit volume due to Type II SNe is

$$\dot{E}_{\text{new}} = \dot{E}_{\text{II}} + \frac{\dot{\rho}_{\text{II}}}{2} \left[ \left\| v_{\varphi} \hat{e}_{\varphi} - \mathbf{u} \right\|^2 + \text{Tr}(\boldsymbol{\sigma}^2) \right]. \quad (6)$$

The total energy injection due to the old and new stellar populations is then  $\dot{E} = \dot{E}_{\text{old}} + \dot{E}_{\text{new}}$ , that is:

$$\dot{E} = \dot{E}_{\text{Ia}} + \dot{E}_{\text{II}} + \frac{\dot{\rho}_{\text{Ia}} + \dot{\rho}_* + \dot{\rho}_{\text{II}}}{2} \left[ \left\| v_{\varphi} \hat{e}_{\varphi} - \mathbf{u} \right\|^2 + \text{Tr}(\boldsymbol{\sigma}^2) \right]. \quad (7)$$

Finally, an energy and momentum sink associated with SF are present; they are, respectively, written as

$$\dot{E}_{\text{SF}} = \frac{\eta_{\text{SF}} E}{t_{\text{SF}}}, \quad \dot{\mathbf{m}}_{\text{SF}} = \frac{\eta_{\text{SF}} \mathbf{m}}{t_{\text{SF}}}, \quad (8)$$

where  $E$  and  $\mathbf{m}$  are the internal energy and momentum density of the ISM.

### 2.2.3 The hydrodynamical equations

The hydrodynamical equations are the same as in N14, with the addition of all the source and sink terms related with the SF process described above:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = \dot{\rho}_{\text{Ia}} + \dot{\rho}_* + \dot{\rho}_{\text{II}} - \dot{\rho}_{\text{SF}},$$

$$\rho \frac{\partial \mathbf{u}}{\partial t} + \rho (\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla p - \rho \nabla \Phi_{\text{tot}} + (\dot{\rho}_{\text{Ia}} + \dot{\rho}_* + \dot{\rho}_{\text{II}})(v_{\varphi} \hat{e}_{\varphi} - \mathbf{u}),$$

$$\frac{\partial E}{\partial t} + \nabla \cdot (E \mathbf{u}) = -p \nabla \cdot \mathbf{u} - \mathcal{L} + \dot{E} - \dot{E}_{\text{SF}},$$

where  $\rho$ ,  $\mathbf{u}$ ,  $E$ ,  $p$ ,  $\Phi_{\text{tot}}$ , and  $\mathcal{L}$  are, respectively, the ISM mass density, velocity, internal energy density, pressure, total gravitational potential, and bolometric luminosity per unit volume. The gas is assumed to be an ideal monoatomic fully ionized plasma, so that  $p = (\gamma - 1)E$ , where  $\gamma = 5/3$  is the adiabatic index. The chemical composition is fixed to solar ( $\mu \simeq 0.62$ ), and the gas self-gravity is neglected.

In the pSF scheme, the source terms  $\dot{\rho}_{\text{II}}$  and  $\dot{E}_{\text{II}}$  are zero, thus they either do not enter the hydrodynamical equations above and they do not contribute to  $\dot{E}$ . In the more realistic aSF,  $\dot{\rho}_{\text{II}}$  and  $\dot{E}_{\text{II}}$  are non-zero. Most galaxy models are simulated with pSF and aSF; for each of the two schemes, we adopted two values of the SF efficiency in equations (1) and (8),  $\eta_{\text{SF}} = 10^{-1}$  and  $\eta_{\text{SF}} = 10^{-2}$  (see Section 3 below).

We tested that the code conserves the total mass and energy; the total mass at any time  $t$  is  $M_{\text{gas}} + M_{\text{esc}} = M_{\text{inj}} + M_{\text{inj}}^{\text{II}} - M_*^{\text{new}}$ , where  $M_{\text{gas}}$  is the gas mass within the simulation box,  $M_{\text{esc}}$  is the cumulative escaped mass out of the simulation box, until that time;  $M_{\text{inj}}$ ,  $M_{\text{inj}}^{\text{II}}$ , and  $M_*^{\text{new}}$  are the cumulative masses, respectively, injected by the passively evolving stellar population, injected by SNII events due to the newly born stellar population, and of the new stars. The values of these masses at the end of the simulations are given in Tables 2–5. Finally, the radiative cooling is implemented by adopting a modified version of the cooling law of Sazonov et al. (2005),

**Table 2.** Relevant quantities at  $t = 13$  Gyr for the rotating models ( $k = 1$ ) with  $\sigma_{\text{as}} = 300 \text{ km s}^{-1}$ .

Name	SF	$\eta_{\text{SF}}$	$M_{\text{inj}}$ ( $10^9 M_{\odot}$ )	$M_{\text{esc}}$ ( $10^9 M_{\odot}$ )	$M_{\text{gas}}$ ( $10^9 M_{\odot}$ )	$M_{\text{not}}$ ( $10^9 M_{\odot}$ )	$M_{\text{c}}$ ( $10^9 M_{\odot}$ )	$M_{\text{new}}^*$ ( $10^9 M_{\odot}$ )	$M_{\text{inj}}^{\text{II}}$ ( $10^9 M_{\odot}$ )	$(t)_{\text{new}}^*$ (Gyr)	$M_{\text{new}}^{\text{ew}} / (t)_{\text{new}}^*$ ( $M_{\odot} \text{ yr}^{-1}$ )	SFR ( $M_{\odot} \text{ yr}^{-1}$ )	$L_{\text{X}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$L_{\text{SNIIa}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$L_{\text{SNII}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$T_{\text{X}}$ (keV)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)
EO4 <sub>IS</sub> <sup>300</sup>	×	–	69.38	14.10	56.04	6.52	49.8	–	–	–	–	–	1.34	47.7	–	0.68
EO7 <sub>IS</sub> <sup>300</sup>	×	–	65.46	15.87	50.24	6.13	44.5	–	–	–	–	–	1.04	45.0	–	0.56
FO4 <sub>IS</sub> <sup>300</sup>	×	–	71.81	11.93	60.92	6.11	54.7	–	–	–	–	–	1.35	49.4	–	0.68
FO7 <sub>IS</sub> <sup>300</sup>	×	–	71.93	11.14	61.82	5.07	56.7	–	–	–	–	–	0.90	49.5	–	0.59
EO4 <sub>IS</sub> <sup>300</sup>	P	0.01	70.53	13.85	14.87	6.86	7.93	41.81	–	5.02	8.33	2.24	1.47	48.5	–	0.69
EO7 <sub>IS</sub> <sup>300</sup>	P	0.01	66.56	15.68	17.02	6.50	10.4	33.86	–	5.34	6.34	2.10	1.07	45.8	–	0.58
FO4 <sub>IS</sub> <sup>300</sup>	P	0.01	72.94	11.75	14.15	6.23	7.85	47.04	–	4.90	9.6	2.47	1.38	50.2	–	0.68
FO7 <sub>IS</sub> <sup>300</sup>	P	0.01	73.01	10.83	16.13	5.07	10.9	46.06	–	4.98	9.25	2.54	0.89	50.2	–	0.55
EO4 <sub>IS</sub> <sup>300</sup>	P	0.1	70.53	13.89	8.23	6.90	1.24	48.41	–	4.36	11.10	1.99	1.54	48.5	–	0.68
EO7 <sub>IS</sub> <sup>300</sup>	P	0.1	66.56	15.77	7.82	6.45	1.31	42.96	–	4.42	9.72	1.66	1.04	45.8	–	0.59
FO4 <sub>IS</sub> <sup>300</sup>	P	0.1	72.94	11.71	7.73	6.40	1.22	53.50	–	4.25	12.59	2.12	1.50	50.2	–	0.68
FO7 <sub>IS</sub> <sup>300</sup>	P	0.1	73.01	10.87	6.86	5.30	1.47	55.29	–	4.19	13.2	2.03	0.92	50.2	–	0.58
EO4 <sub>IS</sub> <sup>300</sup>	A	0.01	70.53	13.83	15.98	6.95	8.97	50.89	10.17	5.12	9.94	3.08	1.56	48.5	11.5	0.69
EO7 <sub>IS</sub> <sup>300</sup>	A	0.01	66.56	15.71	18.53	6.22	12.3	40.38	8.07	5.47	7.38	2.67	0.99	45.8	10.2	0.58
FO4 <sub>IS</sub> <sup>300</sup>	A	0.01	72.94	11.79	15.06	6.27	8.73	57.60	11.51	4.94	11.66	3.21	1.44	50.2	11.9	0.68
FO7 <sub>IS</sub> <sup>300</sup>	A	0.01	73.01	10.86	17.79	5.36	12.4	55.45	11.08	5.07	10.94	3.09	0.94	50.2	11.8	0.57
EO4 <sub>IS</sub> <sup>300</sup>	A	0.1	70.53	13.95	8.55	7.05	1.42	60.03	12.00	4.43	13.55	2.51	1.65	48.5	9.83	0.69
EO7 <sub>IS</sub> <sup>300</sup>	A	0.1	66.56	15.89	8.23	6.58	1.58	53.03	10.60	4.50	11.78	2.07	1.13	45.8	7.76	0.58
FO4 <sub>IS</sub> <sup>300</sup>	A	0.1	72.94	11.89	8.04	6.61	1.36	66.25	13.24	4.29	15.44	2.65	1.65	50.2	10.1	0.70
FO7 <sub>IS</sub> <sup>300</sup>	A	0.1	73.01	11.04	7.40	5.72	1.59	68.21	13.63	4.23	16.13	2.49	1.09	50.2	9.42	0.57

*Notes.* (1) Model name. (2) SF scheme adopted ( $\times = \text{noSF}$ ,  $\text{P} = \text{pSF}$ ,  $\text{A} = \text{aSF}$ ). (3) SF efficiency. (4)–(5) Cumulative mass injected by the evolution of the original stellar population, and mass escaped from the numerical grid, respectively. Differences in  $M_{\text{inj}}$  for models of same  $L_{\text{B}}$  are accounted for by different sampling of the stellar density profile over the numerical grid. (6) Total ISM mass retained within the galaxy. (7)–(8) ISM mass with  $T > 10^6 \text{ K}$ , and  $T < 2 \times 10^4 \text{ K}$ , respectively. (9) Mass of the new stars. (10) Cumulative mass ejected by type II SNe. (11)–(13) Mean formation time of the new stars, mean SFR, and SFR at the end of the simulations. (14) ISM X-ray luminosity in the 0.3–8 keV band. (15)–(16): SNIa and SNII kinetic energy input per unit time (Section 2.2.2). (17) ISM X-ray emission weighted temperature in the 0.3–8 keV band.

**Table 3.** Relevant quantities at  $t = 13$  Gyr for the rotating models ( $k = 1$ ) with  $\sigma_{\text{es}} = 200 \text{ km s}^{-1}$ .

Name	SF	$\eta_{\text{SF}}$	$M_{\text{inj}}$ ( $10^9 M_{\odot}$ )	$M_{\text{esc}}$ ( $10^9 M_{\odot}$ )	$M_{\text{gas}}$ ( $10^9 M_{\odot}$ )	$M_{\text{hot}}$ ( $10^9 M_{\odot}$ )	$M_{\text{c}}$ ( $10^9 M_{\odot}$ )	$M_{\text{new}}^{\text{new}}$ ( $10^9 M_{\odot}$ )	$M_{\text{inj}}^{\text{II}}$ ( $10^9 M_{\odot}$ )	$(t)_{\text{new}}^{\text{new}}$ (Gyr)	$M_{\text{new}}^{\text{new}} / (t)_{\text{new}}^{\text{new}}$ ( $M_{\odot} \text{ yr}^{-1}$ )	SFR ( $M_{\odot} \text{ yr}^{-1}$ )	$L_{\text{X}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$L_{\text{SNIIa}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$L_{\text{SNII}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$T_{\text{X}}$ (keV)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)
EO4 <sub>IS</sub> <sup>200</sup>	×	–	11.92	9.76	2.39	0.15	2.24	–	–	–	–	–	0.0007	10.2	–	0.50
EO7 <sub>IS</sub> <sup>200</sup>	×	–	11.93	10.34	1.84	0.24	1.57	–	–	–	–	–	0.002	10.2	–	0.49
FO4 <sub>IS</sub> <sup>200</sup>	×	–	12.00	3.32	8.80	1.73	6.97	–	–	–	–	–	0.449	10.3	–	0.37
FO7 <sub>IS</sub> <sup>200</sup>	×	–	11.96	3.41	8.69	1.36	7.16	–	–	–	–	–	0.248	10.2	–	0.32
EO4 <sub>IS</sub> <sup>200</sup>	P	0.01	12.07	4.32	3.28	1.63	1.57	4.47	–	6.61	0.68	0.45	0.110	10.3	–	0.39
EO7 <sub>IS</sub> <sup>200</sup>	P	0.01	12.09	10.43	0.34	0.27	3.99E-2	1.32	–	3.84	0.34	0.02	0.003	10.3	–	0.50
FO4 <sub>IS</sub> <sup>200</sup>	P	0.01	12.14	6.65	0.68	0.43	2.19E-1	4.80	–	4.20	1.14	0.08	0.006	10.4	–	0.40
FO7 <sub>IS</sub> <sup>200</sup>	P	0.01	12.08	2.98	3.12	1.57	1.40	5.99	–	5.39	1.11	0.37	0.163	10.3	–	0.36
EO4 <sub>IS</sub> <sup>200</sup>	P	0.1	12.07	4.65	2.00	1.77	7.31E-2	5.42	–	5.74	0.94	0.12	0.086	10.3	–	0.40
EO7 <sub>IS</sub> <sup>200</sup>	P	0.1	12.09	10.07	0.32	0.29	0.00	1.71	–	2.37	0.72	0.00	0.003	10.3	–	0.51
FO4 <sub>IS</sub> <sup>200</sup>	P	0.1	12.14	3.75	3.64	2.88	8.98E-4	4.74	–	3.28	1.45	0.08	0.587	10.4	–	0.43
FO7 <sub>IS</sub> <sup>200</sup>	P	0.1	12.08	5.79	0.49	0.40	1.98E-2	5.81	–	3.43	1.69	0.04	0.010	10.3	–	0.50
EO4 <sub>IS</sub> <sup>200</sup>	A	0.01	12.07	9.40	0.27	0.17	9.92E-2	3.00	0.60	3.90	0.77	0.05	0.001	10.3	0.18	0.54
EO7 <sub>IS</sub> <sup>200</sup>	A	0.01	12.09	9.70	0.47	0.24	2.09E-1	2.40	0.48	4.39	0.55	0.06	0.002	10.32	0.24	0.49
FO4 <sub>IS</sub> <sup>200</sup>	A	0.01	12.14	6.77	0.98	0.72	1.86E-1	5.47	1.09	4.11	1.33	0.09	0.011	10.4	0.36	0.43
FO7 <sub>IS</sub> <sup>200</sup>	A	0.01	12.08	3.09	3.77	1.42	2.19	6.53	1.30	5.27	1.24	0.69	0.460	10.3	2.58	0.29
EO4 <sub>IS</sub> <sup>200</sup>	A	0.1	12.07	7.77	0.29	0.28	7.34E-4	5.02	1.00	3.72	1.35	0.00	0.002	10.3	0.01	0.53
EO7 <sub>IS</sub> <sup>200</sup>	A	0.1	12.09	8.27	0.80	0.62	5.66E-2	3.78	0.76	5.34	0.71	0.06	0.012	10.3	0.24	0.44
FO4 <sub>IS</sub> <sup>200</sup>	A	0.1	12.14	6.00	1.48	1.19	8.70E-4	5.81	1.16	3.10	1.88	0.01	0.041	10.4	0.05	0.44
FO7 <sub>IS</sub> <sup>200</sup>	A	0.1	12.08	3.82	3.82	3.28	2.55E-3	5.57	1.11	2.92	1.91	0.11	0.982	10.3	0.40	0.41

*Note.* All quantities as in Table 2.

**Table 4.** Relevant quantities at  $t = 13$  Gyr for the rotating models ( $k = 1$ ) with  $\sigma_{\text{es}} = 250 \text{ km s}^{-1}$  (aSF only).

Name	SF	$\eta_{\text{SF}}$	$M_{\text{inj}}$ ( $10^9 M_{\odot}$ )	$M_{\text{esc}}$ ( $10^9 M_{\odot}$ )	$M_{\text{gas}}$ ( $10^9 M_{\odot}$ )	$M_{\text{hot}}$ ( $10^9 M_{\odot}$ )	$M_{\text{c}}$ ( $10^9 M_{\odot}$ )	$M_{\text{new}}^*$ ( $10^9 M_{\odot}$ )	$M_{\text{inj}}^{\text{II}}$ ( $10^9 M_{\odot}$ )	$(t)_{*}^{\text{new}}$ (Gyr)	$M_{*}^{\text{new}} / (t)_{*}^{\text{new}}$ ( $M_{\odot} \text{ yr}^{-1}$ )	SFR ( $M_{\odot} \text{ yr}^{-1}$ )	$L_{\text{X}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$L_{\text{SNIIa}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$L_{\text{SNII}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$T_{\text{X}}$ (keV)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)
EO4 <sub>IS</sub> <sup>250</sup>	×	–	31.68	8.28	23.84	4.02	19.8	–	–	–	–	–	0.76	23.9	–	0.55
EO7 <sub>IS</sub> <sup>250</sup>	×	–	30.63	11.42	19.71	3.42	16.3	–	–	–	–	–	0.33	23.1	–	0.55
EO4 <sub>IS</sub> <sup>250</sup>	A	0.1	32.14	8.39	4.84	4.41	0.40	23.64	4.72	4.35	5.43	0.86	1.01	24.2	3.25	0.57
EO7 <sub>IS</sub> <sup>250</sup>	A	0.1	31.10	11.42	4.47	4.31	0.16	19.01	3.80	4.13	4.60	0.42	0.66	23.4	1.57	0.58
EO4 <sub>IS</sub> <sup>250</sup>	A	0.01	32.14	8.28	7.17	4.06	3.08	20.87	4.17	5.06	4.12	1.14	0.82	24.2	4.31	0.56
EO7 <sub>IS</sub> <sup>250</sup>	A	0.01	31.10	11.18	6.92	4.00	2.90	16.24	3.24	5.32	3.05	0.93	0.51	23.4	3.55	0.52


*Note.* All quantities are as in Table 2.

**Table 5.** Relevant quantities at  $t = 13$  Gyr for the mildly rotating models ( $k = 0.1$ ) with  $\sigma_{\text{es}} = 250 \text{ km s}^{-1}$  (aSF only).

Name	SF	$\eta_{\text{SF}}$	$M_{\text{inj}}$ ( $10^9 M_{\odot}$ )	$M_{\text{esc}}$ ( $10^9 M_{\odot}$ )	$M_{\text{gas}}$ ( $10^9 M_{\odot}$ )	$M_{\text{hot}}$ ( $10^9 M_{\odot}$ )	$M_{\text{c}}$ ( $10^9 M_{\odot}$ )	$M_{\text{new}}^*$ ( $10^9 M_{\odot}$ )	$M_{\text{inj}}^{\text{II}}$ ( $10^9 M_{\odot}$ )	$(t)_{*}^{\text{new}}$ (Gyr)	$M_{*}^{\text{new}} / (t)_{*}^{\text{new}}$ ( $M_{\odot} \text{ yr}^{-1}$ )	SFR ( $M_{\odot} \text{ yr}^{-1}$ )	$L_{\text{X}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$L_{\text{SNIIa}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$L_{\text{SNII}}$ ( $10^{40} \text{ erg s}^{-1}$ )	$T_{\text{X}}$ (keV)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)
EO4 <sub>k=0.1</sub> <sup>250</sup>	×	–	32.14	8.63	23.51	4.78	18.7	–	–	–	–	–	5.82	24.2	–	0.69
EO7 <sub>k=0.1</sub> <sup>250</sup>	×	–	31.10	12.94	18.17	3.44	14.7	–	–	–	–	–	2.20	23.4	–	0.69
EO4 <sub>k=0.1</sub> <sup>250</sup>	A	0.1	32.14	8.46	5.13	5.04	0.086	23.20	4.64	4.28	5.42	1.01	6.33	24.2	3.76	0.67
EO7 <sub>k=0.1</sub> <sup>250</sup>	A	0.1	31.10	13.06	3.63	3.56	0.065	18.01	3.60	4.03	4.47	0.45	2.78	23.4	1.76	0.68
EO4 <sub>k=0.1</sub> <sup>250</sup>	A	0.01	32.14	8.58	5.28	4.84	0.442	22.84	4.56	4.46	5.12	0.91	6.39	24.2	3.69	0.66
EO7 <sub>k=0.1</sub> <sup>250</sup>	A	0.01	31.10	12.96	4.14	3.48	0.656	17.50	3.50	4.22	4.14	0.63	2.51	23.4	2.32	0.68

*Note.* All quantities are as in Table 2.





**Figure 1.** Top panels: ISM X-ray luminosity  $L_X$  in the 0.3–8 keV band at 13 Gyr as a function of  $\sigma_{e8}$ , for the selection of 10 models from N14 re-simulated here (Table 1), plus two models with  $\sigma_{e8} = 250$  km s $^{-1}$  for the E0 counterpart, and  $k = 0.1$  (shown with thicker ellipses). The three panels (from left to right) refer to the same models without SF (from N14), with passive SF, and with active SF (with  $\eta_{SF} = 0.01$ ). Bottom panels: the same for the X-ray emission weighted temperature  $T_X$  in the 0.3–8 keV band at 13 Gyr. All  $L_X$  and  $T_X$  values are given in Tables 2–5. See Section 3.1 for more details.

with a lower limit for the ISM temperature of  $T = 10^4$  K. The total X-ray emission in the 0.3–8 keV band ( $L_X$ ), and the emission weighted temperature in the same band ( $T_X$ ), are calculated as volume integrals over the whole computational grid, using as weight the emissivity in the 0.3–8 keV band of a hot, collisionally ionized plasma (see N14 for more details).

### 3 RESULTS


Each simulation starts with the galaxy empty of gas, when the age of the original stellar population is 2 Gyr, and the ISM evolution is followed for the next 11 Gyr. Tables 2–5 list for all models the main quantities of interest at the end of the simulations; the tables also list the same quantities for the corresponding models without SF taken from N14 (hereafter ‘noSF’ models). First (Section 3.1), we present the main results focusing on the two sets of re-simulated models with largely differing values for the galaxy mass (i.e. the two sets with E0 counterpart of  $\sigma_{e8} = 200$  and 300 km s $^{-1}$ , the LM and HM models). In Section 3.2, we concentrate on the intermediate mass models (run only for the aSF), where this time  $k$  is equal to 1 or lower ( $k = 0.1$ ). In Sections 3.3 and 3.4, we discuss the consumption of cold gas mass and the formation of new stars. Finally, we investigate the relationship between the adopted recipe for SF (equation 1) and the Kennicutt–Schmidt relation (Section 3.5).

#### 3.1 Comparison between noSF and SF models

In a short summary, for noSF models, N14 found that rotation in LM galaxies favours the establishment of global winds,

with the consequent reduction of  $L_X$ ; in medium-to-high mass galaxies the conservation of angular momentum lowers the hot gas density in the central galactic region, leading again to a reduction of  $L_X$ , and also of  $T_X$  (because the external and colder regions weight more in the computation of  $T_X$ ). In LM galaxies, instead,  $T_X$  can become higher if rotation triggers a wind, due to the decrease of the ISM density, and the additional heating due to the high meridional velocities of the escaping material (see equation 7).


The first important result here is that when SF is added (both in the pSF and aSF modalities), the hot ISM evolution remains substantially similar to that found by N14 without SF, for models of the same mass. As a consequence of the insensitivity of the general behaviour to the addition of SF, the values of  $L_X$  and  $T_X$  for the models with SF, and their trends with the main galactic structural properties, are on average very similar to those found by N14. Therefore, the conclusions of N14 about the importance of shape and rotation in determining  $L_X$  and  $T_X$ , also as a function of galaxy mass, are confirmed. Fig. 1 demonstrates this result by plotting the final  $L_X$  and  $T_X$  values for the same rotating galaxy models evolved with noSF and with SF (with  $\eta_{SF} = 0.01$ ): the distribution of the points in the noSF and SF panels is very similar (and the analogous figure with the  $\eta_{SF} = 0.1$  models produces the same conclusions). In particular,  $L_X$  of HM models is only marginally sensitive to the presence of SF, while in LM models SF can introduce variations of  $L_X$ ; however, these keep within the (large) range of  $L_X$  values already found in N14 without SF. Thus, SF adds another cause of spread for  $L_X$ , at lower galaxy masses. Fig. 2 shows the resulting  $L_X$  versus  $T_X$  for the aSF models, compared with the same quantities derived recently from *Chandra* data for the hot gas of ETGs (Borson et al. 2011; Kim & Fabbiano 2015). From the latter two samples, we




**Figure 2.**  $L_X$  as a function of  $T_X$  for aSF models listed in Tables 1–5 (open ellipses, as in Fig. 1), compared with *Chandra* data for the hot gas of ETGs taken from Boroson, Kim & Fabbiano (2011) and Kim & Fabbiano (2015, grey squares with error bars). From the latter two samples, we selected only ETGs of morphological type later than E3, for a proper comparison with our models.

have selected only ETGs of morphological type later than E3, for a proper comparison with our models. Overall, the simulation results agree well with the observed X-ray properties of real ETGs (except perhaps for the most X-ray luminous models that may be slightly hotter than observed).

The reason for variations in  $L_X$  linked with SF in LM models is given by anticipations or delays in the most conspicuous features of their typical flow behaviour: N14 found (and the present simulations confirm) that the ISM in rotating models experiences periodic cooling episodes, where the gas injected by the stellar population accumulates until the radiative losses become catastrophic. In these episodes, the ISM quickly cools, emitting a large part of its internal energy as radiation; peaks in  $L_X$  and throats in  $T_X$  are then produced, as apparent in Fig. 5. The tenuous hot atmosphere left after a major cooling is then replenished by the new mass injection from the stellar population. In correspondence of the formation of these significant amounts of cold material (with short cooling times) SF is enhanced, and peaks in pace with  $L_X$ , as discussed in Section 3.4. On the contrary, very little variations of the flow due to SF are seen in HM galaxies. Figs 3–5 give particular examples of the general similarity in the hydrodynamical evolution of noSF and SF models, by plotting in parallel the ISM evolution for the noSF, pSF, and aSF cases; at the same time, the figures point out how this similarity is very close for HM models (see Fig. 4, for the HM model EO7<sub>IS</sub><sup>300</sup>), while some differences can be present in LM ones (see Fig. 3, for the LM model EO7<sub>IS</sub><sup>200</sup>). Notably, the hot ISM evolution in the HM models is practically identical with and without SF (see also Fig. 5): a massive cold disc forms early, and afterwards, even when it is mostly consumed by SF (see Section 3.3 below), around the disc the hot ISM evolution proceeds



**Figure 3.** Meridional sections of the heating over cooling time ratio for the LM EO7<sub>IS</sub><sup>200</sup> model, at a selection of representative times (indicated as galaxy ages in the top right of each panel). The heating time is  $t_{\text{heat}} = E/\dot{E}$  (where  $\dot{E}$  is the source term given in equation 7); the cooling time is defined in Section 2.2.1. From left to right, each column refers to the noSF, pSF, and aSF (with  $\eta_{\text{SF}} = 0.1$ ) models, respectively. Arrows describe the meridional velocity field; their length is proportional to the modulus of the gas velocity in the  $(z, R)$  plane, according to a scale shown in the bottom-left corner. See Section 3.1 for more details.



**Figure 4.** Meridional sections of the heating over cooling time ratio for the HM  $\text{EO7}_{\text{IS}}^{300}$  model. From top to bottom, each row refers to the noSF, pSF, and aSF (with  $\eta_{\text{SF}} = 0.1$ ) models, respectively. Arrows describe the velocity field, as in Fig. 3. Note the large cold disc, that does not completely disappear with SF. See Section 3.1 for more details.


almost unaltered by the energy and mass injection from SNII's. Less massive models, instead, are sensitive to the inclusion of SF, similarly to what found in previous studies where they showed to be very sensitive to any change in the galaxy properties (e.g. the mass distribution, the stellar population inputs, the stellar kinematics; Ciotti et al. 1991, N14). As shown by Figs 3 and 5, for the  $\text{EO7}_{\text{IS}}^{200}$  galaxy, at around 4 Gyr the noSF and the pSF models have already past their major cooling episode (see the peak in  $L_X$  in Fig. 5) followed by the onset of a wind, and the equatorial outflow becomes stronger and stronger thereafter (see how the purple regions become more and more extended in Fig. 3, starting from 4.6 Gyr, and the sharp and steady decline in  $L_X$  in Fig. 5). The aSF model, instead, has its major cooling catastrophe much later, as shown by the prominent dark green region in Fig. 3 at  $\simeq 8$  Gyr, which emits a conspicuous amount of radiation in the X-ray band, thus prompting the corresponding peak in  $L_X$  in Fig. 5. This delay in the major cooling episode, preceding the onset of a wind, is due to the newly formed stars that inject mass, and most importantly heat, in the ISM; thanks to this heat the galaxy can remain hot gas rich until  $\simeq 8$  Gyr, when finally the major cooling takes place. pSF, where mass is not injected instead, corresponds to an evolution very similar to that without SF.

A complementary illustration of the different behaviour between the HM and the LM galaxies is given by Fig. 5, that shows the  $L_X$  and  $T_X$  evolution for the same models of Figs 3 and 4, plus another set of three models representative of the LM class, the  $\text{FO7}_{\text{IS}}^{200}$  ones.

In LM models, a number of cooling episodes (peaks in  $L_X$ ) typically take place, possibly terminated by a major one that is followed by the onset of a wind and the clearing of the ISM from the galaxy (as in N14). The galaxy mass distribution in the  $\text{FO7}_{\text{IS}}^{200}$  models (middle panels of Fig. 5), is more concentrated than in the  $\text{EO7}_{\text{IS}}^{200}$  ones, thus the last major peak in  $L_X$  is delayed with respect to what happens for the  $\text{EO7}_{\text{IS}}^{200}$  models. Thus, in the  $\text{FO7}_{\text{IS}}^{200}$  galaxy, aSF produces a delay in the major cooling episode that is even longer than for the  $\text{EO7}_{\text{IS}}^{200}$  aSF model, to the point that it does not take place within the present epoch. pSF instead produces its anticipation with respect to what shown in the noSF case (Fig. 5), due to pSF subtracting gas and leaving a lower density region, radiating less and more easy to push out of the galaxy.

A final, general result is also that the evolution of a model with SF is more and more different from that of its corresponding noSF model, when increasing  $\eta_{\text{SF}}$ .

For illustration purposes, Fig. 6 shows a meridional section of the density distribution of the newly formed stars at the end of the simulations, for the models in Fig. 5. One can notice the more extended and massive disc in HM models, where it is also quite independent of the pSF or aSF scheme. On the contrary, in LM models, the disc is much less extended, and it can become larger (for the  $\text{EO7}_{\text{IS}}^{200}$  model) or smaller (for the  $\text{FO7}_{\text{IS}}^{300}$  model) when switching from the pSF to the aSF scheme; this is another evidence of the high sensitivity of the flow evolution to any change in the input parameters, at low galaxy masses.




**Figure 5.** Time evolution of the X-ray luminosity  $L_X$  and X-ray emission weighted temperature  $T_X$  for the  $EO7_{IS}^{200}$  (top panels),  $FO7_{IS}^{200}$  (middle panels),  $EO7_{IS}^{300}$  (bottom panels) models. The black, green, and red lines refer to the noSF model, and to the passive and active SF schemes, respectively, with  $\eta_{SF} = 0.1$ . The  $FO7_{IS}^{200}$  model has an evolution almost identical to that of  $EO7_{IS}^{300}$  shown here. See Section 3.1 for more details.

### 3.2 Intermediate mass ETGs

Here, we present the results for two representative models, of E4 and E7 shapes, with intermediate galaxy mass ( $\sigma_{e8} = 250 \text{ km s}^{-1}$  for the E0 counterpart, see Table 1), that have been run with the same kinematic configuration of the LM and HM models


presented previously ( $k = 1$ , isotropic rotator), and also with a lower rotation ( $k = 0.1$ ). For these, only aSF has been considered, with  $\eta_{SF} = 0.01$  and  $\eta_{SF} = 0.1$ . The final values for the most relevant quantities of these eight models are presented in Tables 4 and 5.



**Figure 6.** Meridional sections of the density distribution of the new stars formed up to 13 Gyr (in units of  $M_{\odot} \text{pc}^{-3}$ ), for the same models of Figs 5 and 8:  $\text{EO7}_{\text{IS}}^{200}$  (left-hand panels),  $\text{FO7}_{\text{IS}}^{200}$  (middle panels), and  $\text{EO7}_{\text{IS}}^{300}$  models (right-hand panels). Top and bottom panels refer, respectively, to pSF and aSF simulations, with  $\eta_{\text{SF}} = 0.1$ .

The E4 models with  $k = 1$  show the same smooth evolution of the HM models, i.e. no or very small oscillations in  $L_X$  and  $T_X$  (see the dashed lines in Fig. 7, similar to those in Fig. 5); in fact, the evolution of the hot gas is similarly not affected by SF. The E7 models with  $k = 1$ , instead, show oscillations in  $L_X$  and  $T_X$ , similar to those of the  $\text{EO7}_{\text{IS}}^{200}$  and  $\text{FO7}_{\text{IS}}^{200}$  models, but fewer and less pronounced (see the solid lines in Fig. 7, to be compared with those in Fig. 5). In the aSF model the final  $L_X$  is larger than for the noSF case, as found for the  $\text{FO7}_{\text{IS}}^{200}$  models (Section 3.1).

In models with rotational velocities reduced by a factor of 10 ( $k = 0.1$ ),  $L_X$  and  $T_X$  increase largely, as expected from the results



**Figure 7.** Time evolution of the X-ray luminosity  $L_X$  and X-ray emission weighted temperature  $T_X$  for the  $\text{EO4}_{\text{IS}}^{250}$  and  $\text{EO7}_{\text{IS}}^{250}$  models (top panels), and for the  $\text{EO4}_{k=0.1}^{250}$  and  $\text{EO7}_{k=0.1}^{250}$  models (bottom panels). For the aSF models,  $\eta_{\text{SF}} = 0.1$ . See Section 3.2 for more details.


of N14 showing a larger  $L_X$  and  $T_X$  in non-rotating galaxies ( $k = 0$ ). Again the E4 models show no oscillations in  $L_X$  and  $T_X$ , while the E7 models show numerous small oscillations in  $L_X$  and  $T_X$ , that increase in amplitude with time increasing (Fig. 7). The cold gas disc is much less extended: its radius is  $\lesssim 1$  kpc for  $k = 0.1$ , and  $\lesssim 10$  kpc for  $k = 1$  (slightly larger for flatter shape, and for lower  $\eta_{\text{SF}}$ ).

### 3.3 Mass exchange between the cold disc and the new stars

The second main result of the new simulations is that most of the cold gas disc is consumed by SF, both for pSF and aSF. This result, of general validity, is shown for the particular models of Fig. 5 in Fig. 8, where the cold gas mass  $M_c$  of the noSF models (black solid line) is at the end very close to that of the new stars  $M_*^{\text{new}}$  (dashed lines). For example, the  $\text{EO7}_{\text{IS}}^{200}$  model in the noSF case has a cold disc extending out to a 2.5 kpc radius, of  $M_c = 1.6 \times 10^9 M_{\odot}$ , while the corresponding pSF model at the end has no cold disc, and a mass in new stars of  $M_*^{\text{new}} \simeq 1.7 \times 10^9 M_{\odot}$  (a bit more gas has cooled in the pSF case than in the noSF case). Similarly, this same galaxy model, with aSF and  $\eta_{\text{SF}} = 0.1$ , at the end has a cold disc of just  $M_c \simeq 5.7 \times 10^7 M_{\odot}$ , and  $M_*^{\text{new}} \simeq 3.8 \times 10^9 M_{\odot}$ . This  $M_*^{\text{new}}$  is larger (approximately double) than for the pSF case, a fact that is partly accounted for by the injected mass from the new stars in the aSF model,<sup>2</sup> and mostly by the delay in the final cooling episode after which a galactic wind is established (Fig. 5); this delay keeps the galaxy gas-rich and star forming for a longer time (which is proved also by the lower escaped mass  $M_{\text{esc}}$  in the aSF than in the pSF case, see Table 3).

In general, the LM models with SF show a final  $M_c$  that can range from 0 to  $10^9 M_{\odot}$ , but is typically  $\lesssim 10^8 M_{\odot}$  (Fig. 9 and Table 3).


<sup>2</sup> We recall that, in the aSF scheme, the mass injected back into the ISM by the new stars can itself condense and produce new stars, that in turn contribute to the computation of  $M_*^{\text{new}}$ .



**Figure 8.** Left-hand panels: time evolution of the cold gas mass  $M_c$  (solid lines), and of the cumulative mass in newly formed stars  $M_*^{\text{new}}$  (dashed lines). Right-hand panels: time evolution of the SFR rates. The models are the same of Fig. 5. The black line refers to the noSF model, green, and red lines refer to pSF and aSF, respectively, with  $\eta_{\text{SF}} = 0.1$ . Note how  $M_c$  is much reduced in the SF models (green and red solid lines) with respect to the noSF models (black lines), and how at the same time  $M_*^{\text{new}}$  (dashed lines) reaches close to the noSF  $M_c$  values. See Section 3.3 for more details.

The final mass in the newly formed stellar disc is of the order of  $M_*^{\text{new}} \simeq \text{a few } \times 10^9 M_\odot$ ; thus, the  $M_*^{\text{new}}$  values reach almost those of the  $M_c$  of the corresponding noSF models (Table 3; Fig. 9). temperature below  $2 \times 10^4 \text{K}$ .

Also in the HM models, similarly to what found for the LM models,  $M_*^{\text{new}}$  at the end of the simulations is of the order of the  $M_c$  values of the noSF models. A larger massive disc remains in the HM models, though, even when including SF (e.g. Figs 8 and 9).



**Figure 9.** Top panels: final values of the cold gas mass, for the models in Tables 2–5 without SF (left-hand panel), and with aSF, for the two different  $\eta_{\text{SF}}$  values adopted here. Middle and bottom panels: final values of the stellar mass in newly formed stars  $M_*^{\text{new}}$ , of the SFR normalized to the original  $M_*$ , and of the mean formation time of the new stars  $(t)_*^{\text{new}}$ , calculated from the beginning of the simulation, for the same models with aSF in the top panels. All values are given in Tables 2–5. See Sections 3.3 and 3.4 for more details.

Their final  $M_c$  is  $\simeq 10^9 M_\odot$ , for  $\eta_{\text{SF}} = 0.1$  ( $\simeq 1/50$  of  $M_c$  without SF), and  $(8\text{--}12) \times 10^9 M_\odot$ , if  $\eta_{\text{SF}} = 0.01$  ( $M_c \simeq 1/5$  of  $M_c$  without SF), independently of pSF or aSF. As found for the LM galaxies, in the aSF models  $M_*^{\text{new}}$  is larger than in the pSF ones, and can be even larger than  $M_c$  of the noSF models (Figs 8 and 9). This time, though, the difference is lower, of  $\simeq 20$  per cent, and corresponds to secondary SF sustained by the material injected by the new stars. The final mass in the newly formed stellar disc is of the order of  $M_*^{\text{new}} \simeq \text{a few} \times 10^{10} M_\odot$  (Table 2). However, it must be noted that the structural properties of the HM models may not be very realistic: ETGs of this mass with the isotropic rotator kinematics are quite unlikely, since they are not observed to rotate this much (Emsellem et al. 2011). Real massive galaxies will have less rotation than adopted here with  $k = 1$ , and then they will have less cold gas and less new stars.

More realistic massive and rotating ETGs are those of Section 3.2, with aSF. For these, if  $k = 1$ , the final cold gas mass is again largely reduced when introducing SF, by a factor of  $\simeq 50$  (E4) and  $\simeq 100$  (E7), when  $\eta_{\text{SF}} = 0.1$ . At 13 Gyr,  $M_c$  is  $\text{few} \times 10^9 M_\odot$  for  $\eta_{\text{SF}} = 0.01$ , and  $\text{few} \times 10^8 M_\odot$  for  $\eta_{\text{SF}} = 0.1$ ;  $M_*^{\text{new}}$  is  $\simeq 2 \times 10^{10} M_\odot$ , for both  $\eta_{\text{SF}}$  values. If  $k = 0.1$ , the final  $M_c$  is reduced even more than for the  $k = 1$  models (Fig. 9), for both the E4 and E7 shapes:  $M_c$  is a  $\text{few} \times 10^8 M_\odot$  ( $\eta_{\text{SF}} = 0.01$ ), and  $\lesssim 10^8 M_\odot$  ( $\eta_{\text{SF}} = 0.1$ ). Note that the final cold mass  $M_c$ , without SF, is similarly large for the  $k = 1$

and 0.1 models (Tables 4 and 5; Fig. 9), but it is instead lower for the  $k = 0.1$  models than for the  $k = 1$  ones, when adding SF; one could conclude that in the  $k = 0.1$  models SF is more efficient, because it takes place in a cold disc which is smaller and denser, or that the evolution is different, if less gas overall cooled (the heating was more efficient). The second hypothesis is supported by the finding that, at the end of the simulations,  $M_*^{\text{new}}$  is similar for  $k = 1$  and 0.1 (e.g.  $2.4$  and  $2.3 \times 10^{10} M_\odot$  for the E4 models, respectively, with  $k = 1$  and 0.1, for  $\eta_{\text{SF}} = 0.1$ ), and similarly larger than  $M_c$  of the corresponding noSF models (e.g.  $2.0$  and  $1.9 \times 10^{10} M_\odot$ , for the E4 models, respectively, with  $k = 1$  and 0.1).

In conclusion, SF seems to be an important mechanism to solve the worrisome feature of massive cold gas discs in rotating ETGs without SF. In addition, we found the following trends, at the end of the simulations (see also the top two rows of panels of Fig. 9):  $M_c$  increases with  $M_*$ , and has a large spread for LM models, that increases for larger  $\eta_{\text{SF}}$ ;  $M_c/M_*$  is roughly constant with  $M_*$ , with a spread reaching down to values much lower than this constant for LM models; at any  $M_*$ ,  $M_c$  decreases when  $\eta_{\text{SF}}$  increases.  $M_*^{\text{new}}$  and  $M_*^{\text{new}}/M_*$  increase with  $M_*$ , with a spread for LM models, and are roughly independent of  $\eta_{\text{SF}}$ . This latter result means that more massive ETGs have been overall more efficient in forming stars from recycled material, considering their past  $\simeq 11$  Gyr.

### 3.4 The SF history

SF is not only responsible for the final mass budget of  $M_c$  and  $M_*^{\text{new}}$ , but also for the rate of SF (SFR), its evolution, and the (fiducial) age of  $M_*^{\text{new}}$ ; that is, for the SF history that we discuss here. SF shows a different evolution in LM and HM galaxies, in pace with their different ISM evolution (see Fig. 8). In LM galaxies, where major cooling episodes take place recursively, the SFR peaks suddenly in correspondence of these cooling events due to the increased density of cold gas, and declines right after, going eventually to practically zero either if the cold gas reservoir is completely consumed or the cooling episode is followed by a global wind (Section 3.1). Thus, in general LM galaxies have a larger and peaked SFR in their past (reaching  $1\text{--}2 M_\odot \text{ yr}^{-1}$ ), when the rate of stellar mass losses due to the original stellar population was much larger, and a low SFR ( $\lesssim 0.1 M_\odot \text{ yr}^{-1}$ ) at the present epoch. Typical values at 13 Gyr can be as low as zero, and at most as high as  $0.45 M_\odot \text{ yr}^{-1}$  (Table 3). These values compare well with the current rates estimated for the ATLAS<sup>3D</sup> sample, that range from  $\approx 0.01$  and  $3 M_\odot \text{ yr}^{-1}$ , with a median value of  $0.15 M_\odot \text{ yr}^{-1}$  for ETGs of  $\sigma_{e8}$  comparable to those of the LM models run here (Davis et al. 2014). HM galaxies, instead, show a more regular and steady production of cold gas, and so is their SFR (Fig. 8); at 13 Gyr, their SFR is  $2\text{--}3 M_\odot \text{ yr}^{-1}$  (Table 2; but we recall these models are not realistic). In the intermediate mass models (Section 3.2), the SFR is larger for the E4 models without oscillations in  $L_X$  than for the E7 ones. Overall, the SFR at 13 Gyr is similar for  $k = 1$  and 0.1: for  $k = 1$ , the SFR ranges from  $0.42$  to  $1.1 M_\odot \text{ yr}^{-1}$  (i.e. it is larger than for the LM models of same  $k = 1$ ); when  $k = 0.1$ , the SFR ranges from  $0.45$  to  $1.0 M_\odot \text{ yr}^{-1}$ . Fig. 9 summarizes the *present epoch* SFR behaviour, as a function of  $M_*$ , for the aSF models. To compare models of different mass, the plotted quantity is the SFR normalized to  $M_*$ . This quantity is  $\sim (1\text{--}4) \times 10^{-3} \text{ Gyr}^{-1}$ , quite independent of  $\eta_{\text{SF}}$ , for the intermediate mass and HM models; it can be largely varying for the LM models (from  $4 \times 10^{-4}$  to  $6 \times 10^{-3} \text{ Gyr}^{-1}$ , for  $\eta_{\text{SF}} = 0.1$ , and from  $8 \times 10^{-5}$  to  $\sim 10^{-3} \text{ Gyr}^{-1}$ , for  $\eta_{\text{SF}} = 0.01$ ). On average, the present epoch SFR/ $M_*$  increases with  $M_*$ , a result similar to that of the previous

Section 3.3, where we found that the integration over time of the SFR, that is  $M_*^{\text{new}}$ , and  $M_*^{\text{new}}/M_*$ , increase with  $M_*$ .

Tables 2 and 3 also list the mean formation time  $\langle t \rangle_*^{\text{new}}$  of the new stars since the beginning of the simulation ( $t_0 = 2$  Gyr), and the mean SFR (defined as  $M_*^{\text{new}}/\langle t \rangle_*^{\text{new}}$ ), both at 13 Gyr.  $\langle t \rangle_*^{\text{new}}$  as a function of time is defined as

$$\langle t \rangle_*^{\text{new}}(t) = \frac{1}{M_*^{\text{new}}(t)} \int_{t_0}^t (t' - t_0) \text{SFR}(t') dt', \quad (9)$$

where the SFR( $t$ ) is the volume integrated, instantaneous SFR. The values of  $\langle t \rangle_*^{\text{new}}$  at the *present epoch* are shown in Fig. 9 for the aSF cases. They range from 2.4 to 6.6 Gyr, for the LM models, with a tendency to be lower for the aSF than for the pSF, and for the larger  $\eta_{\text{SF}}$ . The HM and intermediate mass models have  $\langle t \rangle_*^{\text{new}}$  in a narrower range (from 4.2 to 5.5 Gyr), independent of the aSF or pSF, and are again lower for the larger  $\eta_{\text{SF}}$  (Fig. 9). Therefore, the new stars in LM models have a larger spread of formation times, and correspondingly a larger range of ages at the current epoch, than in larger mass models.

For the aSF, for  $\eta_{\text{SF}} = 0.01$ , the upper envelope of the more recent formation times (largest  $\langle t \rangle_*^{\text{new}}$  values) is  $\simeq 5.5$  Gyr, roughly independent of  $M_*$ , while the lower envelope (lowest  $\langle t \rangle_*^{\text{new}}$  values) increases with  $M_*$ . A similar trend is shown by the HM and intermediate mass models, for  $\eta_{\text{SF}} = 0.1$ , but shifted towards lower  $\langle t \rangle_*^{\text{new}}$  values ( $\lesssim 4.5$  Gyr); the spread of  $\langle t \rangle_*^{\text{new}}$  for LM models keeps instead large, extending to even lower  $\langle t \rangle_*^{\text{new}}$  values of  $\simeq 3$  Gyr. The decrease of  $\langle t \rangle_*^{\text{new}}$  for larger  $\eta_{\text{SF}}$ , i.e. for a more efficient SF, obtained here from hydrodynamical simulations provides support to the similar conclusion drawn in galaxy formation models focused on chemical evolution: these assume that the efficiency of SF is an increasing function of mass, which for them has – among others – the consequence that smaller galaxies continue to form stars for longer periods, while more massive galaxies host an older stellar population (e.g. Matteucci 1994).


It is interesting that  $\langle t \rangle_*^{\text{new}}$  decreases slightly, or remains similar, when switching from pSF to aSF; in fact, one could expect a more extended SF, due to the injection of material from the new stars, for the aSF. Evidently, the net results of the injection of both mass and energy is to produce the bulk of SF at roughly the same time, for the pSF and aSF schemes; only, the final mass in new stars is larger for aSF.

The mean age of the newly formed stars in Gyr is  $13 - t_0 - \langle t \rangle_*^{\text{new}}$ . This age is in any case larger than  $\sim 5.5$  Gyr, for the aSF.

### 3.5 The SF recipe and the Kennicutt–Schmidt relation

From equation (1) it follows that SF is favoured in regions of high density, and relatively short  $t_{\text{SF}}$ . These conditions are naturally present in low-temperature regions; accordingly, the simulations show that most of the SF takes place where the gas temperature is  $\lesssim 2 \times 10^4$  K, i.e. in the cold rotating disc.

Quite obviously, our results depend somehow on the adopted prescription for the SF, while a realistic, physically consistent, description of SF is not available. Yet, it is interesting to ask at what level our implementation of SF recovers some basic observational features of SF. More than half a century ago Schmidt (1959) conjectured that the rate of SF should vary as a power law of the gas density, and later Kennicutt (1998) suggested the parametrization in terms of mass surface densities  $\Sigma_{\text{SFR}} = A \Sigma_{\text{cold gas}}^N$ , valid for star-forming galaxies, with  $N \simeq 1.4$ – $1.5$  (Kennicutt & Evans 2012). We investigate here what scaling (if any) the adopted SF recipe of



**Figure 10.** Projected SFR versus mass surface density of the gas (at  $T < 2 \times 10^4$  K, for the blue symbols, and at  $T > 2 \times 10^4$  K, for the red ones), for two aSF models: EO7<sub>IS</sub><sup>200</sup> on the left, and EO7<sub>IS</sub><sup>250</sup> on the right. Full circles refer to  $\eta_{\text{SF}} = 0.01$  and crosses to  $\eta_{\text{SF}} = 0.1$ . The lower  $\eta_{\text{SF}} = 0.01$  coefficient for the SF recipe (equation 1) reproduces better the empirical Kennicutt–Schmidt scaling, that is shown by the solid line (with a slope of  $N = 1.4$ , and normalized as in Davis et al. (2014) for their ATLAS<sup>3D</sup> sample of molecular gas-rich ETGs). The blue dotted and dashed lines show the best fits for the cold gas only (coefficients in the frames in the lower right corners). See Section 3.5 for more details.

equation (1) produces in terms of  $\Sigma_{\text{SFR}}$  versus  $\Sigma_{\text{gas}}$ , when implemented in the simulations.

Fig. 10 shows for a few representative models at the end of the simulation the relationship between  $\Sigma_{\text{SFR}}$  and  $\Sigma_{\text{gas}}$ , calculated separately for the hotter ( $T > 2 \times 10^4$  K) and colder ( $T \leq 2 \times 10^4$  K) phases of the gas;  $\Sigma_{\text{SFR}}$  is the face-on projection of  $\dot{\rho}_{\text{SF}}$ . The figure shows for comparison a recent scaling for the Kennicutt–Schmidt relation, derived by Davis et al. (2014) for the ATLAS<sup>3D</sup> sample of molecular gas-rich ETGs (this has a slope of  $N = 1.4$ ). Fig. 10 tells that SF in the gas takes place basically in two regimes, and by far most of it is due to cold gas. The behaviour of the cold gas shows two remarkable features: the first is that a scaling similar to that of the Kennicutt–Schmidt relation is followed quite closely by the blue (cold gas) points; the second is that even the normalization is reasonably well reproduced by our scheme of ‘active SF’, when  $\eta_{\text{SF}} = 0.01$  (models with  $\eta_{\text{SF}} = 0.1$  correspond instead to a normalization that is too large). In particular, the cold gas in the  $\log \Sigma_{\text{SFR}} - \log \Sigma_{\text{gas}}$  plane follows a trend best fitted by a line of slope of  $\simeq 1.4$ – $1.5$ , a fact that can be explained as follows. SF in the cold gas is most likely regulated by the dynamical time-scale  $t_{\text{dyn}}$ , being  $t_{\text{cool}}$  very short at such temperatures and densities; thus, one roughly expects that  $\dot{\rho}_{\text{SF}} \propto \rho^{1.5}$ . A slope for the  $\Sigma_{\text{SFR}} - \Sigma_{\text{gas}}$  relation close to the empirical one then follows when neglecting the difference between volume and surface density, which is not unreasonable considering that the cold disc resides mainly in a thin region above and below the equatorial plane, extending vertically for just a couple of gridpoints ( $< 200$  pc; see also Kennicutt & Evans 2012). For the hotter gas, the red points in Fig. 10 are invariably fitted by a steeper line, with slope  $\simeq 2$ ; this is explained by a longer cooling time than the dynamical time, and then by  $\dot{\rho}_{\text{SF}} \propto \rho^2$ .

Overall, Fig. 10 proves that the recipe of equation (1) is very reasonable for SF, and, together with the chosen range of normalization for  $\eta_{\text{SF}}$ , it provides a scaling close to that observed for normal to starburst galaxies. These considerations add strength to the overall results obtained in this study.

## 4 CONCLUSIONS

In this work, we have explored the effects of SF on the evolution of rotating hot gas flows in ETGs; our previous (N14) high-resolution



2D hydrodynamical simulations of such flows, run for axisymmetric two-component models, were successful in producing the observed  $L_X$  and  $T_X$  of flat and rotating ETGs, but also revealed the formation of very massive cooled gas discs. To study the SF effects, we performed hydrodynamical simulations using the same numerical code of N14, where SF with a Salpeter IMF is inserted, following a simple recipe depending on the gas density, and the cooling and dynamical time-scales of the ISM (equation 1), with two possible values for the efficiency of SF ( $\eta_{\text{SF}} = 0.01$  and  $0.1$ ). We considered the new stellar generations to be passive (a pure sink of gas), or active (the new stars contribute mass and energy to the gas flow, via stellar winds and SNIIs). The new simulations have been run for a subsample of the N14 models, of high, intermediate and low galaxy masses, of E4 and E7 shapes, with the ordered stellar rotation described by the isotropic rotator, or with the rotational velocities reduced by a factor of 10.

It is found that subsequent generations of stars are formed from the cold gas that accumulates in the equatorial plane, and that most of the extended and massive cold disc found by N14 is consumed by this process. In particular, the main conclusions are summarized as follows.

(i) Remarkably, both for the passive and the active SF implementation, we confirm the results of the previous investigation without SF, concerning the trends of  $L_X$  and  $T_X$  of the hot ISM with galactic rotation. As found by N14,  $L_X$  is lowered by rotation: in low-mass ETGs this happens because rotation favours the establishment of global winds, and in medium-to-high mass ETGs because rotation lowers the hot gas density in the central galactic region. The average temperature  $T_X$  is also lower in intermediate/high mass ETGs, when rotation is important, due to the lower contribution of the central regions (usually hotter and denser in non-rotating systems), while it can be higher in low-mass ETGs, if rotation triggers a wind, due to the thermalization of meridional winds.

(ii) The robustness of the N14 results even after the addition of SF is explained differently for HM and LM galaxies. In HM galaxies, the evolution of the hot ISM is only marginally sensitive to the presence of SF, and  $L_X$  and  $T_X$  evolve smoothly as without SF: a massive cold disc forms early, it is mostly consumed by SF, and around it the hot ISM evolution proceeds almost unaltered by SF (even in presence of energy and mass injection by SNIIs). In less massive models, instead, the evolution of  $L_X$  is not smooth, but shows many peaks corresponding to major cooling episodes, as in N14. The time-occurrence of the peaks depends on SF, thus SF can induce variations of  $L_X$  at a certain chosen epoch; however, these keep within the already large range of  $L_X$  values typical of these masses, even without SF, being the gas flow very sensitive to many factors (as already found previously, e.g. Ciotti et al. 1991, N14).

(iii) For what concerns the global gas mass budget, at the end of the simulations the cold gas mass left in the equatorial disc  $M_c$  on average increases with  $M_*$ , with a larger spread for LM models; at fixed  $M_*$ ,  $M_c$  is lower for lower stellar ordered rotation, and for larger  $\eta_{\text{SF}}$ . In addition, the ratio  $M_c/M_*$  is roughly constant with  $M_*$ , with a spread reaching down to values much lower than this constant for LM models. In any case,  $M_c \lesssim 2 \times 10^9 M_\odot$  for galaxies with  $M_* \leq 4 \times 10^{11} M_\odot$ . Typical values for  $M_c$  are  $\lesssim$  a few  $\times 10^7 M_\odot$  for LM galaxies (with the maximum rotational level of  $k = 1$ ), with exceptions of  $M_c \simeq 10^9 M_\odot$ ; and a few  $\times 10^7 M_\odot$  (if  $k = 0.1$ ) to  $3 \times 10^9 M_\odot$  (if  $k = 1$ ) for ETGs of intermediate mass. These values compare well with those recently observed (Serra et al. 2012, 2014; Young et al. 2014). In particular, they can explain the observation

that massive, fast-rotating ETGs often have kinematically aligned gas, independent of environment (Davis et al. 2011, 2013).

(iv) The mass in newly formed stars  $M_*^{\text{new}}$ , and the ratio  $M_*^{\text{new}}/M_*$ , increase with  $M_*$ , again with a large spread for LM models, and are roughly independent of  $\eta_{\text{SF}}$ . This result means that more massive (rotating) ETGs have been overall more efficient in forming stars via recycling of their stellar mass losses, during the past  $\simeq 10$  Gyr. The mass in secondary generations of stars is  $(1 - 6) \times 10^9 M_\odot$  (if  $k = 1$ ) for LM models, and  $\simeq 2 \times 10^{10} M_\odot$  (for both  $k = 1$  and  $0.1$ ) for intermediate mass models. These should reside mostly in a disc, as most fast rotator ETGs possess, and be related to the birth of a younger, more metal-rich discy stellar component that is indeed observed (Krajnović et al. 2008; Cappellari et al. 2013). They should not be recognized as a ‘young’ stellar population in an absolute sense, though, since most of  $M_*^{\text{new}}$  formed a few Gyr ago, as can be evaluated from the evolution of their SFR.

(v) The time evolution of the SFR depends on the mass of the galaxy. LM galaxies ( $M_* = 1.25 \times 10^{11} M_\odot$ ) have a larger and peaked SFR in their far past (reaching  $1-2 M_\odot \text{ yr}^{-1}$ ), when the rate of stellar mass losses due to the original stellar population was much larger, and when major cooling episodes were frequent. In fact, the average age of the new stars of LM models ranges from 5 to 8 Gyr (for the aSF scheme). At the present epoch, their SFR is low ( $\lesssim 0.1 M_\odot \text{ yr}^{-1}$  typically), with the full range of values going from zero to  $\simeq 0.7 M_\odot \text{ yr}^{-1}$ . These results agree nicely with the low degree of SF and young stellar populations that is detected only in fast rotators, in the ATLAS<sup>3D</sup> sample (Kuntschner et al. 2010; Sarzi et al. 2013); also, they compare well with the current estimated rates, whose median value is  $\approx 0.15 M_\odot \text{ yr}^{-1}$  (Davis et al. 2014). More massive galaxies, instead, show a more regular and steady production of cold gas, and so is their SFR; the average age of their new stars ranges from 5.5 to 7.0 Gyr (in the aSF scheme). At 13 Gyr the SFR is  $\simeq (0.4-1) M_\odot \text{ yr}^{-1}$ , larger than for LM models, and somewhat larger than observed (e.g. McDermaid et al. 2015); this could be fixed by assuming a lower  $\eta_{\text{SF}}$ , or a lower ordered stellar rotation, at larger galaxy masses (Fig. 9). Thus, our models may still be rotating too much and producing too much cold gas, at galaxy masses  $> 2 \times 10^{11} M_\odot$ . The current SFR/ $M_*$ , instead, remains quite constant from LM to intermediate mass models, with a larger spread for LM models (as already found for other properties).

Finally, the SF recipe adopted for this work proved to be a reasonable one, given that it reproduces the slope of the Kennicutt–Schmidt relation, and even the normalization if  $\eta_{\text{SF}} \simeq 0.01$ .

Overall, we can suggest an origin (mostly) in the SF from cooling hot gas, for the presence of cold gas phases kinematically aligned with the stars, and for the low-level degree of SF, all features detected only in fast rotators in the ATLAS<sup>3D</sup> sample.

In the perspective of the galaxy evolution, we finally recall that the precise knowledge of the amount of gas flowing towards the galactic centre is of great importance for a proper study of feedback effects from the central black hole in rotating galaxies (e.g. Novak, Ostriker & Ciotti 2011; Gan et al. 2014). A major implication of this work is the fact that, if the gas produced by stellar evolution forms new stars, the amount of gas in principle available for fuelling the central supermassive black hole is much lowered. Of course, such fuelling would be impossible in presence of rotation and absence of viscosity. Disc instabilities, though, could break axial symmetry, and allow for the gas infall on to the black hole, possibly escaping SF. Phenomenologically, the effect of gravitational instabilities in the disc can be modelled as an effective gravitational viscosity (e.g. Bertin & Lodato 2001; Hopkins & Quataert 2011; Rafikov

2015), that favours accretion of cold gas towards the centre. Indeed, viscosity (that could be due for example to magnetorotational instability) is another ingredient not present in the current simulations, but that could affect the disc evolution. As the code is axisymmetric, we cannot follow the complex physics of (non-axisymmetric) disc instabilities that could be present. Although self-gravity of the gas is not considered in our simulations, yet for some models at a selection of times we computed the fiducial value of the radial profile of the stability parameter  $Q(R) = c_s k_R / \pi G \Sigma_c$ , where  $k_R$  is the local epicyclic frequency of the galaxy, and  $c_s$  and  $\Sigma_c$  are, respectively, the vertically mass-averaged values of the ISM sound velocity, and the surface density of the cold rotating gas. The  $Q$  values were computed for a layer  $\Delta z \simeq 200$  pc thick above and below the galaxy equatorial plane; these  $Q$  turned out to be quite independent of the adopted thickness value  $\Delta z$ , being dominated by the cold disc. These fiducial  $Q$  values maintained invariably larger than unity (indicating stability) when SF is allowed, while  $Q < 1$  on a central  $\simeq$ kpc-size region, in absence of SF. From this preliminary and qualitative analysis, it follows that SF and  $Q$ -instability are both present in the gaseous rotating discs. The relevant question is then if and what of the two processes is dominant in the disc. In a future work, we plan to extend further the present investigation by implementing in the code the cooperating effects of SF and mass discharge from the disc to the centre. In this new study, we will also consider the neglected late-time mass return from the newly born stellar population, that is the mass injected by stars of mass  $< 8 M_\odot$ ; this input could be modelled with the method described in Calura et al. (2014).

## ACKNOWLEDGEMENTS

We acknowledge Giuseppe Lodato and Jerry Ostriker for useful discussions, and Silvia Posacki for providing the galaxy models of N14. LC and SP were supported by the MIUR grant PRIN 2010-2011, project ‘The Chemical and Dynamical Evolution of the Milky Way and Local Group Galaxies’, prot. 2010LY5N2T.

## REFERENCES

Bertin G., Lodato G., 2001, *A&A*, 370, 342  
 Boroson B., Kim D.-W., Fabbiano G., 2011, *ApJ*, 729, 12  
 Calura F., Ciotti L., Nipoti C., 2014, *MNRAS*, 440, 3341  
 Cappellari M. et al., 2011, *MNRAS*, 413, 813  
 Cappellari M. et al., 2013, *MNRAS*, 432, 1862  
 Ciotti L., Ostriker J. P., 2007, *ApJ*, 665, 1038  
 Ciotti L., D’Ercole A., Pellegrini S., Renzini A., 1991, *ApJ*, 376, 380  
 Combes F., Young L. M., Bureau M., 2007, *MNRAS*, 377, 1795  
 David L. P., Forman W., Jones C., 1991, *ApJ*, 380, 39  
 Davis T. A. et al., 2011, *MNRAS*, 417, 882  
 Davis T. A. et al., 2013, *MNRAS*, 429, 534  
 Davis T. A. et al., 2014, *MNRAS*, 444, 3427  
 de Vaucouleurs G., 1948, *Ann. Astrophys.*, 11, 247  
 di Serego Alighieri S. et al., 2007, *A&A*, 474, 851  
 Dubois Y., Gavazzi R., Peirani S., Silk J., 2013, *MNRAS*, 433, 3297  
 Edge A. C. et al., 2010, *A&A*, 518, L47  
 Emsellem E. et al., 2011, *MNRAS*, 414, 888

Eskridge P. B., Fabbiano G., Kim D.-W., 1995, *ApJS*, 97, 141  
 Fabbiano G., 1989, *ARA&A*, 27, 87  
 Gan Z., Yuan F., Ostriker J. P., Ciotti L., Novak G. S., 2014, *ApJ*, 789, 150  
 Grossi M. et al., 2009, *A&A*, 498, 407  
 Hopkins P. F., Quataert E., 2011, *MNRAS*, 415, 1027  
 Kennicutt R. C., Evans N. J., 2012, *ARA&A*, 50, 531  
 Kennicutt R. C., Jr, 1998, *ApJ*, 498, 541  
 Khochfar S. et al., 2011, *MNRAS*, 417, 845  
 Kim D.-W., Fabbiano G., 2015, preprint (arXiv:1504.00899)  
 Ko J., Hwang H. S., Im M., Le Borgne D., Lee J. C., Elbaz D., 2014, *ApJ*, 791, 134  
 Krajnović D. et al., 2008, *MNRAS*, 390, 93  
 Kuntschner H. et al., 2010, *MNRAS*, 408, 97  
 Lagos C. d. P., Davis T. A., Lacey C. G., Zwaan M. A., Baugh C. M., Gonzalez-Perez V., Padilla N. D., 2014, *MNRAS*, 443, 1002  
 Li J.-T., Wang Q. D., Li Z., Chen Y., 2011, *ApJ*, 737, 41  
 McDermid R. M. et al., 2015, *MNRAS*, 448, 3484  
 McDonald M., Veilleux S., Rupke D. S. N., Mushotzky R., Reynolds C., 2011, *ApJ*, 734, 95  
 Martig M. et al., 2013, *MNRAS*, 432, 1914  
 Mathews W. G., 1989, *AJ*, 97, 42  
 Mathews W. G., Baker J. C., 1971, *ApJ*, 170, 241  
 Matteucci F., 1994, *A&A*, 288, 57  
 Mellier Y., Mathez G., 1987, *A&A*, 175, 1  
 Morganti R. et al., 2006, *MNRAS*, 371, 157  
 Naab T. et al., 2014, *MNRAS*, 444, 3357  
 Navarro J. F., Frenk C. S., White S. D. M., 1997, *ApJ*, 490, 493  
 Negri A., Ciotti L., Pellegrini S., 2014a, *MNRAS*, 439, 823  
 Negri A., Posacki S., Pellegrini S., Ciotti L., 2014b, *MNRAS*, 445, 1351 (N14)  
 Novak G. S., Ostriker J. P., Ciotti L., 2011, *ApJ*, 737, 26  
 O’Sullivan E., Forbes D. A., Ponman T. J., 2001, *MNRAS*, 324, 420  
 Oser L., Ostriker J. P., Naab T., Johansson P. H., Burkert A., 2010, *ApJ*, 725, 2312  
 Parriotti J. R., Bregman J. N., 2008, *ApJ*, 681, 1215  
 Pellegrini S., 2011, *ApJ*, 738, 57  
 Pellegrini S., 2012, in Kim D.-W., Pellegrini S., eds, *Astrophysics and Space Science Library*, Vol. 378, *Hot Interstellar Matter in Elliptical Galaxies*. Springer-Verlag, Berlin, p. 21  
 Pellegrini S., Held E. V., Ciotti L., 1997, *MNRAS*, 288, 1  
 Posacki S., Pellegrini S., Ciotti L., 2013, *MNRAS*, 433, 2259  
 Rafikov R. R., 2015, *ApJ*, 804, 62  
 Sarzi M. et al., 2006, *MNRAS*, 366, 1151  
 Sarzi M. et al., 2010, *MNRAS*, 402, 2187  
 Sarzi M. et al., 2013, *MNRAS*, 432, 1845  
 Sazonov S. Y., Ostriker J. P., Ciotti L., Sunyaev R. A., 2005, *MNRAS*, 358, 168  
 Schmidt M., 1959, *ApJ*, 129, 243  
 Serra P. et al., 2012, *MNRAS*, 422, 1835  
 Serra P. et al., 2014, *MNRAS*, 444, 3388  
 Suh H., Jeong H., Oh K., Yi S. K., Ferreras I., Schawinski K., 2010, *ApJS*, 187, 374  
 Tang S., Wang Q. D., 2005, *ApJ*, 628, 205  
 Thornton K., Gaudlitz M., Janka H.-T., Steinmetz M., 1998, *ApJ*, 500, 95  
 Yi S. K. et al., 2005, *ApJ*, 619, L111  
 Young L. M. et al., 2011, *MNRAS*, 414, 940  
 Young L. M. et al., 2014, *MNRAS*, 444, 3408

This paper has been typeset from a  $\text{\TeX}/\text{\LaTeX}$  file prepared by the author.