



HAL
open science

Venusian Thermosphere variability by IPSL Venus GCM

Antoine Martinez, Sebastien Lebonnois, Ehouarn Millour, Thomas Pierron,
Enora Moisan, Gabriella Gilli, Franck Lefèvre

► To cite this version:

Antoine Martinez, Sebastien Lebonnois, Ehouarn Millour, Thomas Pierron, Enora Moisan, et al..
Venusian Thermosphere variability by IPSL Venus GCM. EGU General Assembly 2022, May 2022,
Online, Unknown Region. 10.5194/egusphere-egu22-5271 . insu-03649793

HAL Id: insu-03649793

<https://insu.hal.science/insu-03649793v1>

Submitted on 23 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EGU22-5271, updated on 23 Apr 2022

<https://doi.org/10.5194/egusphere-egu22-5271>

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Venusian Thermosphere variability by IPSL Venus GCM

Antoine Martinez¹, Sebastien Lebonnois¹, Ehouarn Millour¹, Thomas Pierron¹, Enora Moisan¹, Gabriella Gilli^{2,4}, and Franck Lefevre³

¹Laboratoire de Météorologie Dynamique (LMD/IPSL), CNRS, Paris, France. (antoine.martinez@lmd.ipsl.fr)

²Instituto de Astrofísica de Andalucía (IAA-CSIC), Glorieta de la Astronomía s/n, Granada, España

³LATMOS, CNRS, Sorbonne Université, Université Versailles St Quentin, Paris, France

⁴Instituto de Astrofísica e Ciências do Espaço (IA), OAL, Tapada da Ajuda, PT1349-018 Lisboa, Portugal

For fifteen years, a Global Climate Model (GCM) has been developed for the Venus atmosphere at “Institut Pierre-Simon Laplace” (IPSL), in collaboration between LMD and LATMOS, from the surface up to 150 km altitude (Lebonnois et al., 2010; 2016). Recently, the vertical grid was extended from 10^{-5} Pa to 10^{-8} Pa (~180-250 km) and allows us to simulate the Venusian upper thermosphere. At the same time, improvements were made on the parameterization of non-LTE CO₂ near infrared heating rates, on the parameterization of non-orographic gravity waves and a tuning was performed on atomic oxygen production to improve the thermospheric densities and their effects (heating and cooling; Martinez et al., 2022; submitted).

This work focuses on validating the modeled thermospheric structure by comparison using data from the Pioneer Venus, Magellan and Venus Express missions which cover similar and complementary (equator and pole) regions at different periods of solar activity, typically above 130 km in altitude. In particular, we will discuss the importance of atomic oxygen in regulating the thermospheric temperature, the effect of the solar cycle on the upper thermosphere and the effect of non-orographic gravity waves on the diurnal temperature profile.

References:

Lebonnois, S., Hourdin, F., Eymet, V., Cresspin, A., Fournier, R., Forget, F., 2010. Superrotation of Venus’ atmosphere analyzed with a full general circulation model. *J. Geophys. Res. (Planets)* 115, 6006. <https://doi.org/10.1029/2009JE003458>.

Lebonnois, S., Sugimoto, N., Gilli, G., 2016. Wave analysis in the atmosphere of Venus below 100-km altitude, simulated by the LMD Venus GCM. *Icarus* 278, 38–51. <https://doi.org/10.1016/j.icarus.2016.06.004>.

Martinez et al. 2022, submitted to *Icarus*