



**HAL**  
open science

## Ultramafic Rock Carbonation: Constraints From Listvenite Core BT1B, Oman Drilling Project

A. Beinlich, O. Plümper, E. Boter, I. A. Müller, F. Kourim, M. Ziegler, Y.  
Harigane, R. Lafay, P. B. Kelemen

► **To cite this version:**

A. Beinlich, O. Plümper, E. Boter, I. A. Müller, F. Kourim, et al.. Ultramafic Rock Carbonation: Constraints From Listvenite Core BT1B, Oman Drilling Project. *Journal of Geophysical Research : Solid Earth*, 2020, 125, 10.1029/2019JB019060 . insu-03661300

**HAL Id: insu-03661300**

**<https://insu.hal.science/insu-03661300>**

Submitted on 6 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# JGR Solid Earth

## RESEARCH ARTICLE

10.1029/2019JB019060

### Special Section:

Ophiolites and Oceanic Lithosphere, with a focus on the Samail ophiolite in Oman

### Key Points:

- Large-scale carbonation resulted in pervasive replacement of serpentinite by the magnesite-quartz-fuchsite (listvenite) assemblage
- Clumped isotope thermometry indicates carbonation temperatures between 50 °C and 250 °C
- Carbonation textures indicate a transition from non-equilibrium spheroidal growth to near-equilibrium euhedral growth

### Supporting Information:

- Supporting Information S1
- Data Set S1
- Table S1
- Table S2
- Table S3
- Movie S1

### Correspondence to:

A. Beinlich,  
andreas.beinlich@uib.no

### Citation:

Beinlich, A., Plümper, O., Boter, E., Müller, I. A., Kourim, F., Ziegler, M., et al. (2020). Ultramafic rock carbonation: Constraints from listvenite core BT1B, Oman drilling project. *Journal of Geophysical Research: Solid Earth*, 125, e2019JB019060. <https://doi.org/10.1029/2019JB019060>

Received 14 NOV 2019

Accepted 10 APR 2020

Accepted article online 17 APR 2020

©2020. The Authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

## Ultramafic Rock Carbonation: Constraints From Listvenite Core BT1B, Oman Drilling Project

A. Beinlich<sup>1</sup> , O. Plümper<sup>2</sup> , E. Boter<sup>2</sup> , I. A. Müller<sup>3</sup> , F. Kourim<sup>3</sup> , M. Ziegler<sup>2</sup> , Y. Harigane<sup>4</sup> , R. Lafay<sup>5</sup> , P. B. Kelemen<sup>6</sup>, and the Oman Drilling Project Science Team

<sup>1</sup>Department of Earth Science, University of Bergen, Bergen, Norway, <sup>2</sup>Department of Earth Sciences, Utrecht University, Utrecht, The Netherlands, <sup>3</sup>Institute of Earth Sciences, Academia Sinica, Taipei, Taiwan, <sup>4</sup>Institute of Geology and Geoinformation, Geological Survey of Japan, The National Institute of Advanced Industrial Science and Technology, Ibaraki, Japan, <sup>5</sup>Géosciences Montpellier, Université Montpellier, Montpellier, France, <sup>6</sup>Lamont–Doherty Earth Observatory, Columbia University, Palisades, NY, USA

**Abstract** The occurrence of the quartz-carbonate alteration assemblage (listvenite) in ophiolites indicates that ultramafic rock represents an effective sink for dissolved CO<sub>2</sub>. However, the majority of earlier studies of ultramafic rock carbonation had to rely on the surface exposure of reaction textures and field relationships. Here we present the first observations on ultramafic rock alteration obtained from the 300 m deep BT1B drill hole, ICDP Oman Drilling Project, allowing for a continuous and high-resolution investigation. Hole BT1B recovered continuous drill core intersecting surface alluvium, 200 m of altered ultramafic rock comprising mainly listvenite and minor serpentinite bands at 90 and 180 m depth, and 100 m of the underlying metamorphic sole. Textural evidence suggests that the carbonation of fully serpentinized harzburgite commenced by non-equilibrium growth of spheroidal carbonate characterized by sectorial zoning resulting from radially oriented low-angle boundaries. In the serpentinite, carbonate spheroids are composed of alternating magnesite cores and dolomite rims, whereas texturally similar carbonate in the listvenite is composed of Fe-rich magnesite cores and Ca-Fe-rich magnesite rims. The distinct compositions and mineral inclusions indicate that the carbonation extent was controlled by fluid accessibility resulting in the simultaneous formation of limited carbonate in the serpentinite bands and complete carbonation in the listvenite parts of BT1B. The presence of euhedral magnesite overgrowing spheroidal carbonate in the listvenite suggests near-equilibrium conditions during the final stage of carbonation. The carbonate clumped isotope thermometry constrains carbonate crystallization between 50 °C and 250 °C, implying repeated infiltration of reactive fluids during ophiolite uplift and cooling.

## 1. Introduction

Alteration of the oceanic lithosphere by hydrothermal fluids is the primary driver of the hydrosphere-lithosphere volatile exchange and has significant consequences for rock physical properties and planetary-scale element fluxes. Furthermore, natural carbon uptake in thermodynamically stable carbonate in the oceanic lithosphere represents a means for permanently removing CO<sub>2</sub> from the atmosphere (Kelemen & Matter, 2008). The carbon uptake potential is particularly strong in the ultramafic part of the oceanic lithosphere that has high concentrations of divalent metal ions, required for carbonate mineral formation (Kelemen et al., 2011). Peridotite alteration by seawater typically results in the formation of Ca carbonate (calcite and aragonite) (e.g., Bonatti et al., 1980; Grozeva et al., 2017), while the alteration by metamorphic fluids frequently results in Mg-rich carbonate (dolomite and magnesite), accompanied by secondary silicate phases including talc, quartz, and fuchsite (Cr-muscovite) and additional Fe-oxide, -hydroxide, and/or sulfide phases. Altered ultramafic rocks consisting of the magnesite-quartz assemblage are termed listvenite and represent the end-product of CO<sub>2</sub>-fluid driven mafic, and ultramafic rock alteration as almost the entire inventory of Mg and Ca cations is incorporated into carbonate minerals (Halls & Zhao, 1995). Natural carbon uptake in ultramafic formations is significant as evidenced by abundantly exposed carbonate-bearing ophiolite complexes worldwide (Akbulut et al., 2006; Azer et al., 2019; Barnes & O'Neil, 1969; Barnes et al., 1978; Beinlich et al., 2010, 2012, 2018; Beinlich, John, et al., 2020; Belogub et al., 2017; Emam & Zoheir, 2013; Gahlan Hisham et al., 2018; Garcia del Real et al., 2016; Halls & Zhao, 1995; Hansen et al., 2005; Hinsken et al., 2017; Launay & Fontes, 1985; Menzel et al., 2018; Quesnel

et al., 2013; Tominaga et al., 2017; Tsikouras et al., 2006). One of the largest and most well-explored ophiolite complexes is the Samail ophiolite in the Sultanate of Oman and the United Arab Emirates (Chavagnac, Ceuleneer, et al., 2013; Chavagnac, Monnin, et al., 2013; Clark & Fontes, 1990; Falk et al., 2016; Falk & Kelemen, 2015; Kelemen & Matter, 2008; Lacinska & Styles, 2012; Lacinska et al., 2014; Mervine et al., 2014; Nasir et al., 2007; Neal & Stanger, 1983, 1985; Paukert et al., 2012; Stanger, 1985; Streit et al., 2012). In the Samail ophiolite, listvenite is present in the basal section above the underlying metamorphic sole. Earlier work has recognized the occurrence of calcite-, dolomite-, and quartz-dominated listvenite containing additional fuchsite  $\pm$  chlorite  $\pm$  spinel in the form of discontinuous planar lenses and fault-controlled vein systems (Nasir et al., 2007; Stanger, 1985; Wilde et al., 2002). More recently investigated listvenite from Wadi Mansah is characterized by magnesite as the dominant carbonate mineral consistent with peridotite alteration predominately driven by influx of CO<sub>2</sub> (Falk & Kelemen, 2015). In addition to the carbonated ultramafic basal section, the Samail ophiolite comprises a complete sequence of oceanic lithosphere from the upper mantle through to the continental crust all of which have recently been subject to comprehensive drilling and subsequent analysis of rock core, logging, hydrological, and microbial sampling in the course of the Oman Drilling Project (OmanDP, ICDP Exp. 5057) (Kelemen et al., 2020). To better constrain the conditions and reaction pathways driving complete peridotite carbonation at large scale, here we present the first results of microstructural investigation and carbonate clumped isotope thermometry on samples recovered from Hole BT1B of the OmanDP. BT1B recovered a 300-m-long continuous core across the basal thrust of the Samail ophiolite through 200 m of listvenite and intercalated serpentinite into an additional 100 m of the underlying metamorphic sole. Reaction textures are assessed across scales ranging from hand specimen to the nanoscale constraining critical reaction parameters and chemical equilibration during natural, large-scale ultramafic rock carbonation.

## 2. Geological Setting


The Samail Ophiolite, located in the Sultanate of Oman and the United Arab Emirates, is the most extensive and best-exposed cross-section through the oceanic lithosphere (Figure 1a). The ophiolite comprises an intact stratigraphy of oceanic crust composed of pelagic allochthonous sediments, overlying pillow basalts, and the sheeted dike complex through to gabbro and upper mantle peridotite (Coleman, 1981; Searle & Malpas, 1980). The basal thrust beneath the Samail nappe separates the ophiolite sequence from the underlying metamorphic sole consisting of metasedimentary phyllite, greenschist facies metabasalt, and minor amphibolite of the Haybi Complex (Searle & Malpas, 1980). The metamorphic sole is underlain by Permian to Late Cretaceous distal to deep-sea sediments, meta-volcanic rocks, and shelf-facies limestone of the Hawasina Group (Coleman, 1981; Nasir et al., 2007; Searle et al., 2015; Stanger, 1985). Syn- to post-emplacement tectonic tilting of the stratigraphy has been followed by unconformable deposition of Late Cretaceous to Early Miocene transgressive shallow marine sediments before Late Miocene to Quaternary uplift and erosion (Coleman, 1981; Searle et al., 2015; Searle & Malpas, 1980).

U-Pb dating of magmatic zircon and <sup>40</sup>Ar/<sup>39</sup>Ar ages of mica and hornblende from the metamorphic sole constrain the timing of thrusting of the Samail ophiolite onto the autochthonous Proterozoic to Mesozoic sedimentary Arabian continental margin between 95 and 80 Ma (Cowan et al., 2014; Gnos, 1998; Hacker, 1994; Hacker et al., 1996; Hacker & Gnos, 1997; Hacker & Mosenfelder, 1996; Lanphere, 1981; Rioux et al., 2012, 2013, 2016; Tilton et al., 1981; Warren et al., 2005). The recently obtained  $97 \pm 29$  Ma Rb-Sr isochron age of listvenite fuchsite, and the fact that listvenites in the Samail ophiolite are only found along the basal thrust and associated mélange zones, are consistent with syn-emplacement alteration of the ophiolite (Falk & Kelemen, 2015).

## 3. Materials and Methods

### 3.1. Macroscopic Sample Description and Context in Drill Core

Samples used for this study were collected from core BT1B in the course of shipboard core logging aboard D/V Chikyu in September 2017 (Kelemen et al., 2020). BT1B was drilled in March 2017 in Wadi Mansah (23°21.861' N, 58°20.149' E), has a total length of 300.1 m, and comprises upper alluvium (2%), serpentinite (8%), listvenite (56%; quartz + carbonate), and the underlying metamorphic sole (34%) (Figure 1b).



**Figure 1.** Geological context, down-hole stratigraphy, and representative core photographs of listvenite core BT1B. (a) Simplified geological map showing the location of hole BT1B in the context of the regional and local geology (after Nicolas & Boudier, 1995). (b) Downhole stratigraphy of core BT1B showing the relation between listvenite of distinct macroscopic appearance and the presence of relic serpentinite above the basal thrust separating the ophiolite from the underlying metamorphic sole. The blue dots indicate the location of half core photographs shown in Figure 1(c) (Kelemen et al., 2020). (c) Representative photographs showing the different macroscopic appearance of listvenite and serpentinite.

Serpentinite is present in two bands, one between 80 and 100 m depth (Upper Serpentinite) and another between 181 and 186 m (Lower Serpentinite) (Figure 1b). Serpentinite displays a dark green to black color in hand specimen, has a cryptocrystalline grain size, and is typically massive but intersected by abundant veins of light green serpentine and carbonate (Figure 1c). Carbonate-rich zones in the Upper Serpentinite occur between 80.54–80.73 m, 81.47–83.63 m, and 86.82–88.07 m and in gradational contacts between serpentinite and listvenite characterized by a distinct increase in carbonate vein abundance and color change to light green. The thickness of these zones ranges from 0.26 to 3.2 m (Kelemen et al., 2020).

Listvenite is the dominant rock type in the ultramafic part of BT1B and present as three massive bands between the alluvium and the Upper Serpentinite (6.02–80.82 m), between the Upper and Lower Serpentinite (100.2–181.3 m), and between the Lower Serpentinite and the basal thrust (185.5–196.6 m) (Figure 1b). Listvenite is fine-grained to cryptocrystalline and macroscopically heterogeneous with frequent color transitions on the centimeter to meter scale between dark and light red, orange, green, tan, and pale gray (Figure 1c). Red and orange listvenite is strongly brecciated in the drill core, and individual clasts show evidence for repeated brecciation and re-cementation such as truncation of clast-hosted veins at clast-matrix boundaries. Pale gray, tan, and green listvenite are characterized by the presence of cm-sized quartz-fuchsite intergrowths situated in a pale to dark gray matrix. Tan, pale gray, and green listvenite are mainly present between the Upper and Lower Serpentinite, are massive, and locally show foliation resembling ductile deformation.

The basal thrust (196.6–197.1 m) is developed as strongly foliated lower part of the lower listvenite unit and upper part of the underlying metamorphic sole separated by 0.42 m of brown to gray-green fault gouge. The metamorphic sole is composed of intercalated, foliated gray-green metabasalt and mafic metasediments. Hole BT1B did not reach the base of the metamorphic sole, but a recent review (Soret et al., 2017) reports that its thickness throughout the ophiolite “is commonly 10–100 m and does not exceed 500 m.” Falk and Kelemen (2015) report bulk compositions and Sr isotope ratios for phyllites and carbonates from the underlying Hawasina formation in outcrop within a few kilometers of Hole BT1B.

The study presented here is based on six serpentinite samples from the Upper Serpentinite and nine listvenite samples consisting of light red, dark red, and green listvenite from the Upper and Lower Listvenite (Figure 1).

### 3.2. Carbonate Microsampling for Stable Isotope Analyses

A subset of samples was microsampled for carbonate stable isotope analysis ( $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$ ) and clumped isotope thermometry. This sample subset includes matrix and vein magnesite, vein dolomite from the Upper Serpentinite (BT1B 43–02 and 44–03), and listvenite (Figure 1). The majority of listvenite carbonate samples are from the Lower Listvenite, comprising green, light red, and dark red listvenite, plus one additional light red listvenite from the Upper Listvenite. Listvenite matrix carbonate is mainly magnesite except for matrix dolomite from core sections BT1B 72–04 and BT1B 77–03. Listvenite vein dolomite was sampled from the Upper Listvenite (BT1B 32–02) and Lower Listvenite (BT1B 67–04) (see also supporting information Figure S1).

Carbonate was sampled by using a micro drill with 3.8 mm inner diameter and a handheld Dremel tool from thin-section billets at sites selected based on the petrography of the respective thin sections. The relatively large diameter of the drill compared to the typical grain size and vein diameter required sampling of homogeneous matrix areas and veins that were macroscopically devoid of crosscutting relationships. Carbonate samples were further crushed to a fine powder using an agate mortar and pestle prior to the analysis.

### 3.3. Analytical Methods

#### 3.3.1. Electron Microscopy

Electron microscopy was conducted on standard petrographic thin sections using an FEI Verios XHR FE-SEM at the Centre for Microscopy, Characterization and Analysis (CMCA), The University of Western Australia. A working distance of 4 mm and an acceleration voltage of 15 keV were used for BSE imaging. Additional TEM foils were prepared from spheroidal carbonate from serpentinite (BT1B 44–01) and listvenite (BT1B 60–03) by focused ion beam (FIB) milling at Utrecht University and CMCA using a FEI Helios Nanolab G3 CX DualBeam FIB-SEM at CMCA. Subsequent scanning transmission electron microscopy

(STEM) imaging and quantitative electron dispersive X-ray spectroscopy (EDS) element mapping were carried out using a FEI Titan G2 80–200 at CMCA and a FEI Talos F200X at Utrecht University. Quantitative mineral compositions and element distribution maps were acquired using the JEOL 8530F FE electron microprobe at CMCA using an acceleration voltage of 15 keV and a focused and 5  $\mu\text{m}$  defocused beam for silicate and carbonate minerals, respectively. The analytical procedure and application of reference materials follow the method described in Beinlich et al. (2018). Thin-section phase distribution mapping was conducted using a TESCAN Integrated Mineral Analyzer (TIMA) high-resolution Field Emission Scanning Electron Microscope (FESEM) in full liberation mode at the John De Laeter Centre, Curtin University. Operating conditions were 15 keV, 0.8 nA, and a working distance of 10 mm. Acquisition time per field was 227 seconds, and the map consists of 464 fields. Mineral abbreviations in the figures and tables follow the suggestions by Whitney and Evans (2010).

Crystallographic orientations of dolomite and magnesite in two samples BT1B 44–01 and BT1B 70–01 were measured by indexing Electron Backscatter Diffraction (EBSD) patterns acquired using a Hitachi SU3500 SEM equipped with an Oxford Instruments HKL EBSD Nordlys-Nano detector at the Geological Survey of Japan (GSJ). Before EBSD analysis, the thin sections were polished for several minutes with SYTON fluid using a Buehler VibroMet 2 to remove surface damage. Diffraction pattern acquisition was performed in low-vacuum mode using an accelerating voltage of 15 keV, a working distance of 18 mm, specimen tilt of 70°, and step size of 0.5  $\mu\text{m}$ . The orientation data were collected using the AZtec software and processed using the MTEX toolbox in Matlab™. All index data in this study represent points with a mean angular deviation (MAD) of <1°. Post-acquisition processing was done using a 10° segmentation angle for grain modeling, a cut-off grain size of 10 pixel, and the nearest-neighbor method to fill non-indexed areas.

### 3.3.2. X-Ray Microcomputed Tomography

Visualization of the carbonate spherule distribution in serpentinite sample BT1B 44–01 in three dimensions was done using a Zeiss Versa 520 X-ray microscope at CMCA. The X-ray source operated at 80 keV and 7 W and a Zeiss high-contrast, low-resolution 4X detector were used, achieving a spatial resolution of  $\sim 1.5 \mu\text{m}$ . The sample used was a hand-drilled 6 mm long and 3.8 mm diameter core from the thin-section billet used for petrography. Post-processing of the acquired 3D imaging data was done for a subvolume of the sample core using the Avizo software, version 9.4.

### 3.3.3. Stable Isotope Analysis

Stable isotope analyses were performed at the GeoLab, Utrecht University, The Netherlands. Before isotope analysis, the mineralogy of each sample powder was constrained by XRD (Table S1; Beinlich et al., 2020c). The duration of acid digestion was adjusted to either dolomite or magnesite according to the dominant carbonate species in the sample powder. The carbonate  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  values are reported relative to Vienna Pee Dee Belemnite (VPDB).

Dolomite samples were digested in 103% phosphoric acid at 70 °C for 20 minutes, and the released  $\text{CO}_2$  was continuously collected in a liquid nitrogen trap using a Kiel IV carbonate device, coupled to a 253 Plus isotope ratio mass spectrometer (both instruments from Thermo Scientific) and analyzed in Long-Integration Dual-Inlet mode (Müller, Fernandez, et al., 2017; with 600 seconds integration time per aliquot). The weight of individual aliquots of reference materials and unknown samples ranged between 75 and 95  $\mu\text{g}$ . The analyses were carried out in sequences consisting of 38 to 46 individual aliquots, of which 20 were carbonate reference materials. The reference materials used are calcite ETH-1 ( $\delta^{13}\text{C} = 2.20\text{‰}$ ,  $\delta^{18}\text{O} = 6.70\text{‰}$ ,  $\Delta_{47} = 0.258$ ), ETH-2 ( $\delta^{13}\text{C} = -10.0\text{‰}$ ,  $\delta^{18}\text{O} = -10.0\text{‰}$ ,  $\Delta_{47} = 0.256$ ) and ETH-3 ( $\delta^{13}\text{C} = 1.95\text{‰}$ ,  $\delta^{18}\text{O} = 7.18\text{‰}$ ,  $\Delta_{47} = 0.691$ ) (Bernasconi et al., 2018). Each sample and standard gas was measured against a  $\text{CO}_2$  reference gas ( $\delta^{13}\text{C} = -7.25\text{‰}$ ,  $\delta^{18}\text{O} = 1.65\text{‰}$ ), and each unknown sample was analyzed 10 to 14 times, spread over several measurement days.

Magnesite samples were digested offline, using 10–20 mg solid powder and 1–2 ml 103% phosphoric acid at 100 °C for 15–16 hours in individual, sealed vials using a custom-built vacuum line containing a cold trap with liquid nitrogen acetone slush ( $-96 \text{ °C}$ ) to remove  $\text{H}_2\text{O}$  trace quantities from the  $\text{CO}_2$  gas. Each measurement sequence consisted of six samples, of which three were identically treated carbonate reference materials (ETH-1, ETH-2, ETH-3). Each sample analysis, including reference materials, was replicated 4 to 10 times. In contrast to the dolomite analyses, the magnesite analyses were conducted using the Dual Inlet of

a Thermo Fisher Scientific MAT 253 in the traditional way by eight alternating reference gas-sample gas cycles (208 seconds sample gas integration time per measurement).

Clumped isotope analyses of dolomite are reported as  $\Delta_{47}$  in the absolute reference frame for 70 °C reaction temperature without projecting the values down to 25 °C acid digestion temperature due to the uncertainties in the dolomite specific temperature-dependent acid fractionation correction (see Müller, Violay, et al., 2017; Müller et al., 2019; Murray et al., 2016). The clumped isotope composition  $\Delta_{47}$  is the measured abundance of the  $^{13}\text{C}^{18}\text{O}^{16}\text{O}$  isotopologue ( $m/z$  47) in the carbonate acid digestion-released  $\text{CO}_2$  relative to an abundance that would be expected for a stochastic distribution of all isotopes. The deviation from the stochastic distribution is thermodynamically controlled (Eiler, 2007). The raw signal intensities were corrected for negative pressure baselines derived from the  $m/z$  44 beam in the mass spectrometer following the recipe described in Meckler et al. (2014) and Müller, Violay, et al. (2017). We then constructed from the measured raw  $\Delta_{47}$  values of the three ETH calcite standards and their accepted values an empirical transfer function to convert the raw  $\Delta_{47}$  values of the unknown samples into the absolute reference frame as described in Bernasconi et al. (2018). For the  $\Delta_{47}$  temperature estimates of dolomite, we used the mineral specific  $\Delta_{47}$ -temperature calibration of Müller et al. (2019) for 70 °C acid digestion temperature. For the dolomite  $\delta^{18}\text{O}$  values we applied the mineral-specific acid fractionation factor of 1.009926 for acid digestion at 70 °C (Rosenbaum & Sheppard, 1986).

In contrast to dolomite, magnesite samples were processed identically to the calcite samples since earlier magnesite clumped isotope studies did not observe any differences in acid fractionations of magnesite and calcite (García del Real et al., 2016) and more elaborate studies on the magnesite  $\Delta_{47}$  acid fractionation do not yet exist. The only published value for the magnesite  $\Delta_{47}$  acid fractionation (Müller, Violay, et al., 2017) may be erroneous due to incomplete isotopic re-equilibration during heating of the magnesite in this study. Thus, we use the calcite  $\Delta_{47}$ -temperature calibration of Kele et al. (2015) recalculated with the new IUPAC (International Union of Pure and Applied Chemistry) parameters in Bernasconi et al. (2018) to derive our estimates of the carbonate formation temperatures. To transfer the raw  $\Delta_{47}$  values of the magnesites into the absolute reference frame, we constructed an empirical transfer function from the measured raw  $\Delta_{47}$  values of all analyzed ETH-1,-2,-3 standards (runs DI-3 to DI-12; Beinlich et al., 2019) against their accepted values at 70 °C acid digestion temperature and applied the recalculated calcite  $\Delta_{47}$ -temperature calibration of Kele et al. (2015) for 70 °C acid digestion temperature (by subtracting an acid fractionation correction factor of 0.062‰ from the intercept of the calibration equation). For the magnesite  $\delta^{18}\text{O}$  values we applied the acid fractionation factor 1.009178 for acid digestion at 100 °C (Das Sharma et al., 2002).


## 4. Results

### 4.1. Microtexture and Phase Composition

#### 4.1.1. Serpentinite

Macroscopic description during core logging aboard D/V Chikyu distinguished “serpentinite” from “ophicarbonates” (Kelemen et al., 2020). However, subsequent microtextural analysis shows that also those samples that were macroscopically described as serpentinite are carbonate-bearing. Thus, samples previously classified as either serpentinite or ophicarbonates samples are grouped in this section. The increase in carbonate content, from the least-carbonated serpentinite to listvenite, is gradational. The distinction between carbonate-bearing serpentinite from listvenite is based on the absence of quartz in serpentinite and the lack of serpentine in listvenite.


The serpentinite with the smallest proportion of carbonate is dominated by lizardite and chrysotile (see Figures S2a, S2b, S3, S4) together with chromite, magnetite, bastite intergrowths after pyroxene, minor carbonate, and accessory  $\mu\text{m}$ -sized awaruite and is free of hematite and goethite. Primary olivine and pyroxene were not found in the investigated samples, and the serpentinite shows the typical mesh (hourglass) texture (Figures S2a and S2b). Carbonate occurs in veins and as spheroidal grains within the serpentine matrix. Crosscutting relationships indicate that carbonate vein formation was preceded by at least two generations of serpentine veining. Carbonate veins frequently follow or deflect into earlier serpentine veins and sometimes occur in subparallel sets with regular vein spacing of  $\sim 130\text{--}200\ \mu\text{m}$  (e.g., BT1B 44-03). The veins are composed of dolomite or magnesite or a combination of both. Magnesite veins show in most cases a



**Figure 2.** Representative back-scattered electron (BSE) images and quantitative element distribution maps of carbonate in serpentinite and listvenite. (a) Spheroidal magnesite–dolomite composite grains in the least carbonated serpentinite section 44–01. Quantitative element maps showing the chemical zonation of the carbonate. Ca-rich carbonate is identified as dolomite by Raman spectroscopy (not shown), whereas grain core regions are magnesite. FeO enrichment is confined to the magnesite. (b) Spheroidal carbonate in green listvenite core section 60–03. The carbonate growth morphology and chemical zonation is similar to that in serpentinite but Ca-rich carbonate is identified as Ca-magnesite that is also enriched in FeO. (c) Euhedral magnesite in green listvenite section 60–03. The core region of the euhedral magnesite grains resembles the morphology and composition of spheroidal magnesite shown in Figure 2b.

distinct chemical zonation comprising a Fe-rich central part and Fe-poor outer part at the contact with serpentine (Table S2; Beinlich et al., 2020b). The composition of serpentine after primary olivine is identical within the analytical uncertainty between samples BT1B 44–01 and BT1B 44–03. Serpentine formed after orthopyroxene (i.e., bastite) shows slightly lower MgO and higher FeO concentrations (Table S3; Beinlich et al., 2020a).


Abundant spheroidal carbonate is observed in serpentinite and ophicarbonate samples BT1B 44–01 and BT1B 43–02, respectively (Figure 2). These grains are characterized by their spherical shape and consistent grain diameters between 60 and 80  $\mu\text{m}$ . BSE imaging and element distribution maps show a distinct chemical zonation consisting of relatively large Mg-rich cores, surrounded by a zone of increased FeO and thin Ca-rich rims (Figure 2a). Raman spectroscopy (Figure S4) confirms that the Ca-rich zone of the spheroidal carbonate in serpentinite samples is dolomite. The chemical zonation is in some grains repetitive and defined by a second spherical dolomite zone (Figure 2a). The outer rim of the carbonate spheroids is in most cases composed of dolomite and less frequently by a thin rim of magnesite (Figure 3). Magnesite cores and dolomite rims are near their respective endmember compositions, while the outer magnesite core zones are characterized by FeO concentrations of up to 10 wt% (Figure 2a). Larger matrix carbonate clusters, composed of amalgamated carbonate spheroids, preserve the described chemical and mineralogical zonation (Figure 2a). Carbonate spheroids in the serpentinite frequently contain sub- $\mu\text{m}$  sized serpentine inclusions (Figure 3).



**Figure 3.** Details of the carbonate-serpentine interface in the least carbonated serpentinite core section 44–01. High angle annular dark-field image (HAADF) acquired in scanning transmission electron microscopy (STEM) mode showing a thin magnesite rim at the interface between the dolomite zone of a spheroidal carbonate and the surrounding serpentine matrix. The element distribution maps highlight the presence of abundant sub- $\mu\text{m}$  serpentine inclusions in carbonate.

#### 4.1.2. Listvenite

Listvenite samples are dominantly composed of carbonate, quartz, and relic chromite and generally lack precursor serpentine and primary magmatic silicate phases. Red listvenite contains additional fine-grained hematite and potentially goethite intergrown with carbonate and quartz. Matrix carbonate is dominated by magnesite, while carbonate veins are either magnesite or dolomite. Occasionally, the distribution of oxide strings in listvenite resembles that typically associated with a serpentine mesh (or hourglass) texture (Figures S2c and S2d). Based on their crosscutting relationships, thin, sometimes Fe-oxide-bearing magnesite veins (10–50  $\mu\text{m}$ ) represent in most cases the oldest vein generation. They are crosscut by thicker




**Figure 4.** Compositional details (STEM-HAADF) of the core region of a spheroidal magnesite grain in green listvenite section 60–03. The element distribution map indicates the presence of abundant, nm-sized SiO<sub>2</sub> inclusions.

magnesite veins (<1 mm), followed by quartz and/or quartz–carbonate veins (50–300 μm) and then by dolomite veins (<500 μm). The thicker magnesite veins occasionally contain euhedral magnesite showing an oscillatory variation in its Ca content. Additionally, coarse-grained magnesite veins frequently show a vein-internal chemical zonation, composed of near-endmember magnesite at the vein-matrix interface as well as high Fe and Ca concentrations in the vein center. Matrix quartz in the listvenite is in most cases fine-grained to cryptocrystalline and shows an interlocked texture. Less frequently, the quartz matrix appears to consist of clusters of sub-μm sized spheroidal quartz, creating an interstitial porosity (BSE images in Figures 2b and 2c).

Green listvenite samples (e.g., BT1B 52–01 and BT1B 60–03) contain additional fuchsite (Cr-muscovite, Table S3; Beinlich et al., 2020a) as intergrowths with quartz and associated with relic chromite. These samples also contain spheroidal carbonate (~80 μm grain diameter) that texturally resembles the matrix carbonate in serpentinite samples. Spheroidal carbonate in the listvenite occurs in the fine-grained quartz matrix as isolated grains and amalgamated clusters of individual spheroids. Listvenite carbonate spheroids also show a systematic compositional zonation with high-Mg concentrations in the core and elevated Ca and Fe concentrations in the rim (Figure 2b). However, the rims are composed of Ca- and Fe-rich magnesite instead of dolomite. Spheroidal magnesite is frequently overgrown by compositionally distinct euhedral magnesite (Figure 2c). Spheroidal carbonate in the listvenite contains abundant, fine-grained (≤100 nm) spherical silica inclusions, which may be quartz or amorphous silica (Figure 4). In addition, magnesite in the listvenite occasionally displays dendritic growth textures with initial Fe-richer branches and later Fe-poorer overgrowths (Figure S5).

#### 4.2. Carbonate Stable Isotope Composition and Clumped Isotope Thermometry

Dolomite and magnesite extracted from veins in serpentinite and listvenite and the listvenite matrix were analyzed for their δ<sup>13</sup>C (VPDB), δ<sup>18</sup>O (VPDB), and Δ<sub>47</sub> composition. Δ<sub>47</sub> values range between 0.275‰ and 0.548‰. The corresponding spread in formation temperatures for all samples is 45 ± 5 °C (1σ) to 247 ± 52 °C (Figure 5a, Table S1; Beinlich et al., 2020c). Listvenite matrix magnesite formation temperatures range between 52 ± 8 °C (BT1B 52–01) and 190 ± 31 °C (BT1B 68–02) and listvenite matrix dolomite



**Figure 5.** Stable O- and C-isotope systematics. (a) Down-hole plot of clumped isotope carbonate formation temperatures for listvenite and serpentinite samples. Colors and symbols refer to the legend shown in Figure 5c and differentiate between matrix and vein magnesite and dolomite. The downhole stratigraphy is identical to that shown in Figure 1. (b) Down-hole plot of magnesite and dolomite  $\delta^{18}\text{O}$  (VPDB). (c) Carbonate  $\delta^{18}\text{O}$  (VPDB) vs  $\delta^{13}\text{C}$  (VPDB) showing clustering at high oxygen and carbon isotope ratios, lower  $\delta^{18}\text{O}$  for some of the listvenite matrix and vein dolomite and lower  $\delta^{13}\text{C}$  for vein dolomite and magnesite from serpentinite sample BT1B 44–01.

temperatures between  $98 \pm 15$  °C (BT1B 72–04) and  $107 \pm 19$  °C (BT1B 77–03) (see also Figure S1). The dolomite and magnesite veins in the upper serpentinite yielded  $227 \pm 52$  °C (BT1B 43–02) and  $45 \pm 5$  °C (BT1B 44–03), respectively. In the listvenite, magnesite vein-matrix pairs indicate a higher temperature for the vein ( $T_{\text{vein}} - T_{\text{matrix}} = 11$  °C and 56 °C), whereas the matrix magnesite in BT1B 67–04 records a temperature 46 °C higher than the vein dolomite in the same sample.

Dolomite  $\delta^{13}\text{C}$  values range from  $-2.6\text{‰}$  to  $0.70\text{‰}$ , and magnesite  $\delta^{13}\text{C}$  values range from  $-3.8\text{‰}$  to  $0.60\text{‰}$ . Dolomite  $\delta^{18}\text{O}$  values range from  $-13.8\text{‰}$  to  $-6.2\text{‰}$ , and magnesite  $\delta^{18}\text{O}$  values range from  $-9.3\text{‰}$  to  $-6.9\text{‰}$  (Figures 5b and 5c, Table S1; Beinlich et al., 2020c). In the  $\delta^{13}\text{C}$  vs.  $\delta^{18}\text{O}$  space (Figure 5c), carbonate stable isotope compositions cluster in three groups characterized by equal  $\delta^{13}\text{C}$  but distinct  $\delta^{18}\text{O}$  and intermediate  $\delta^{18}\text{O}$  and low  $\delta^{13}\text{C}$ . Listvenite vein-matrix magnesite pairs (BT1B 60–03 and BT1B 68–02) show higher matrix  $\delta^{18}\text{O}$  of  $0.23\text{‰}$  and  $0.40\text{‰}$ , while  $\delta^{13}\text{C}$  values are  $0.26\text{‰}$  higher and  $-0.17\text{‰}$  lower in the matrix compared to the vein. Vein dolomite from BT1B 67–04 has a lower  $\delta^{18}\text{O}$  compared to the matrix magnesite ( $\Delta^{18}\text{O} = -5.9\text{‰}$ ) but slightly higher  $\delta^{13}\text{C}$  ( $\Delta^{13}\text{C} = 0.27\text{‰}$ ) (Figures 5b and 5c, Table S1; Beinlich et al., 2020c).


## 5. Discussion

### 5.1. Incipient Carbonation, Carbonate Nucleation, and Growth

Conceptually, the per-sample amount of carbonate can be related to reaction progress from incipiently altered serpentinite, containing only some carbonate, to fully carbonated listvenite. Along this conceptual reaction sequence, the carbonate growth texture evolves from isolated spheroids, through clusters of amalgamated carbonate spheroids, to spheroids overgrown by compositionally distinct and euhedral carbonate. The spheroidal carbonate texture in serpentinite (e.g., BT1B 44–03; Figure 2a) thus represents a snapshot of the incipient carbonation stage. The similar spheroid diameters imply nearly constant rates and durations of nucleation and growth. The presence of abundant carbonate veins, even in the serpentinite, suggests that the initial fluid infiltration was channelized and facilitated by tectonic deformation of the serpentinite host rock. This is further supported by 3-D textural analysis showing the existence of smaller veinlets connecting individual carbonate spheroids with larger veins (Figure 6; Movie S1). However, the 3-D textural analysis also shows that a relatively large number of carbonate spheroids occur isolated from carbonate veins. Consequently, further migration of reactive fluid must have occurred by diffusion through interconnected, fluid-filled pore space.

Analyses of TEM dark-field images suggest a porosity of  $2.68 \pm 1.02\%$  for the serpentinite. Most individual pore diameters are less than 100 nm (Figure 7). At this scale, nanofluidic transport phenomena can be significant and effectively govern metamorphic fluid-flow, mass transport, and hence carbon mineralization (Plümper, Botan, et al., 2017; Zheng et al., 2018). These transport phenomena are consistent with the relatively

even distribution of carbonate spheroids on the thin-section scale (Figures 8a and 8b), suggesting that the spatial distribution of carbonate nucleation sites and by implication also the pore fluid saturation state was homogeneous during nucleation. Indeed, the carbonate morphology is in agreement with a strong




**Figure 6.** 3-D reconstruction of X-ray tomography data acquired from the least carbonated serpentinite section 44–01. The BSE image shown on the left side shows an example of the reconstructed texture in two dimensions. The carbonate color code (blue/cyan) highlights connectivity between individual spheroids and the central vein. The pale green, translucent volume is filled by serpentine and omitted for clarity. The reconstructed volume consists of 83.0 vol.% serpentine, 5.4 vol.% isolated carbonate (blue), 8.2 vol.% vein carbonate and connected spheroids (cyan), and 3.4 vol.% oxide phases (magnetite and/or chromite). An animated version of the 3-D reconstruction can be found as a supplementary Movie S1.

driving force causing spherical growth instead of sub- to euhedral Mg-carbonate that is frequently observed in similar alteration systems elsewhere (Beinlich et al., 2012; Hansen et al., 2005; Sunagawa, 2005).

Based on the premise that spherulitic growth may result in morphologically similar carbonate and would indicate fast precipitation under far-from-equilibrium conditions (Gránásy et al., 2005; Shtukenberg et al., 2012), we further apply EBSD analysis to representative carbonate spheroids (Figure 9). This crystallographic examination indicates that spheroidal magnesite is composed of a single crystal and contains abundant, predominantly radial low-angle boundaries ( $2^{\circ}$ – $10^{\circ}$ ) that originate from small dolomite inclusions in the grain centers. High-angle boundaries are exclusively present in the dolomite (Figures 9a and 9b). Despite the resemblance with spherulitic carbonate, the fact that carbonate spheroids from BT1B are primarily single crystals is inconsistent with polycrystalline spherulitic growth. Instead, the shape and orientation of such magnesite grains are similar to sector zoning occasionally found in, for example, garnet and fluorite (e.g., Bosze & Rakovan, 2002; Stowell et al., 2011). Sectorial zoning is commonly related to compositionally distinct zones and ascribed to orientation-dependent differences in crystal growth rates. However, growth competition may also result in crystallographic misfit between adjacent sectors of identical composition, thus forming low-angle boundaries in chemically homogeneous single crystals that are unrelated to external deformation (see also Pearce et al., 2013).

## 5.2. Pervasive Carbonation and Listvenite Formation

Listvenite contains spheroidal and euhedral carbonate, both showing a systematic chemical zonation with Mg-rich and Ca-poor grain cores and Fe-Ca-rich rims. Spheroidal magnesite in the listvenite shows a gradual increase in MgO toward the core and a sharp transition to Ca- and Fe-rich magnesite in the rim. Theoretically, the textural similarity of carbonate in the listvenite and serpentinite may be interpreted as to indicate that spheroidal carbonate in the listvenite is inherited from the initial carbonation stage represented by spheroidal carbonate in the serpentinite. This scenario would involve dissolution of the dolomite rims and re-precipitation of Ca-bearing magnesite in a  $\text{CO}_2$ -rich reactive fluid. However, the core region of spheroidal carbonate in the listvenite has ~4–5 wt.% FeO, whereas the core region of spheroidal carbonate in the serpentinite has less than 2 wt.% FeO. Furthermore, listvenite carbonate spheroids contain abundant sub- $\mu\text{m}$  sized silica inclusions (Figure 4), whereas serpentinite magnesite contains serpentine inclusions (Figure 3). These characteristics suggest that the listvenite carbonate did not evolve from spheroidal carbonate in the serpentinite, but instead formed via direct replacement of serpentine, accompanied by the formation of quartz in the matrix. EBSD phase- and misorientation-maps show that spheroidal carbonates in the listvenite also consist of magnesite single crystals with radially orientated low-angle boundaries that originate from a central dolomite inclusion (Figures 9c and 9d). Thus, we infer that the listvenite carbonate spheroids also precipitated at relatively high supersaturation, resulting in growth competition between



**Figure 7.** Serpentinite porosity and pore diameter distribution. (a) STEM dark-field image of serpentine in sample BT1B 44-01 shows an example of sub- $\mu\text{m}$  sized porosity. The thresholded porosity of this image is 1.68%. (c) Pore diameter statistics and probability density (PD) based on a log-normal PD fit of the data shown in the histogram. Pore diameter data is based on the analysis of five binarized STEM dark-field images using the maximum entropy threshold in the Fiji distribution of ImageJ (Schindelin et al., 2012).

adjacent grain sectors and the spheroidal crystal habit as well as the less frequently observed dendritic texture of the magnesite (Figure S5) (Haque et al., 2016; Sunagawa, 2005).

Assuming that Ca is primarily sourced from the dissolution of clinopyroxene, the low CaO concentration in magnesite rims in the listvenite compared to the serpentinite is likely related to the larger total carbonate amount in the sample (dilution effect). High concentrations of FeO in rims of spheroidal magnesite and the euhedral magnesite overgrowths (Figures 2b and 2c) indicate increased availability of Fe during a relatively late stage of listvenite formation, that is, at the time when the carbonate rims and overgrowths formed relative to carbonate core zones. This availability of FeO is attributed to the breakdown of oxide phases (chromite, magnetite) in  $\text{CO}_2$ -rich alteration fluid during listvenite formation, as previously documented for sites in Canada and Norway by Hansen et al. (2005) and Tominaga et al. (2017). Perhaps, this occurred at the same time as crystallization of fuchsite, which formed from reactions involving spinel, silicates, and fluid. The transition from spheroidal to euhedral carbonate morphology reflects a driving force reduction during continued crystal growth (Sunagawa, 2005) and may be related to cooling and/or decrease in carbonate saturation.


### 5.3. Carbonation Temperature and Depth

The clumped isotope distribution in carbonate is thought to be independent of the isotopic composition of the fluid from which the carbonate precipitates (e.g., Eiler, 2007; Ghosh et al., 2006; Huntington et al., 2009), though this is a theoretical prediction that has not been rigorously tested over a broad range in fluid composition and temperature, particularly for dolomite and magnesite. Carbonate clumped isotope signatures can experience solid-state reordering during cooling, depending on the carbonate mineralogy and the cooling rate (Passey & Henkes, 2012). However, dolomite is relatively resistant to solid-state reordering at temperatures below 300  $^{\circ}\text{C}$ ; thus, the temperatures in our study likely reflect the dolomite formation temperatures (e.g., Bonifacie et al., 2017; Ryb et al., 2017). In the case of magnesite, solid-state reordering is less well studied. Nevertheless, its sluggish reaction kinetics with phosphoric acid and evidence for an equilibrium blocking temperature of approximately 490  $^{\circ}\text{C}$  (García del Real et al., 2016) suggest that also the magnesite clumped iso-

tope composition is resistant to isotopic reordering. Hence, we assume that the measured clumped isotope compositions of our carbonate samples reflect their formation temperatures.

The clumped isotope temperatures complement earlier thermometry work in the Samail ophiolite that described relatively low formation temperatures (clumped isotope temperatures between 23–43  $^{\circ}\text{C}$ ) for young carbonate veins in peridotite forming during present-day weathering (Streit et al., 2012), slightly higher temperatures for calcite veins in peridotite beneath a Cretaceous unconformity (25–60  $^{\circ}\text{C}$ , de Obeso & Kelemen, 2018), and yet higher temperatures for carbonate from outcrop samples of listvenite and the metamorphic sole within a few kilometers of Hole BT1B (listvenite 61–114  $^{\circ}\text{C}$ ,  $n = 15$ ; metamorphic sole 64–171  $^{\circ}\text{C}$ ,  $n = 4$ ) (Falk & Kelemen, 2015). Also, Falk and Kelemen (2015) documented the presence of intergrown quartz and antigorite, with and without talc and magnesite, in several samples from a gradational contact between listvenite and serpentinite about 2 km north of Hole BT1B. These mineral assemblages equilibrated with a relatively  $\text{H}_2\text{O}$ -rich fluid at temperatures of 80–120  $^{\circ}\text{C}$  (depending on the choice of thermodynamic data), broadly consistent with clumped isotope temperatures presented by Falk and Kelemen (2015).


For listvenite carbonate from BT1B analyzed in this study, the clumped isotope data yield a relatively wide range of formation temperatures, 52–247  $^{\circ}\text{C}$ . In detail, matrix magnesite from the Lower Listvenite (118.5 m



**Figure 8.** Phase distribution and carbonate nuclei density maps of serpentinite core section BT1B 44–01. (a) High-resolution phase distribution map on the thin-section scale showing the relation between spheroidal magnesite–dolomite composite grains, carbonate veins, and the serpentine matrix. The relative modal abundances of individual phases are 90.30 vol.% (serpentine), 5.90 vol.% (dolomite), 3.02 vol.% (magnesite), 0.42 vol.% (chromite), and 0.36 vol.% (magnetite). (b) Distribution density map of matrix carbonate evaluated per subsection of  $0.16 \text{ mm}^2$  by excluding vein carbonate (white overlay) and using the thresholding routines in the Fiji distribution of ImageJ (Schindelin et al., 2012). The color bar shows both carbonate area fraction and the corresponding number of nucleation sites assuming an average carbonate grain diameter of  $60 \mu\text{m}$ . The distribution of matrix carbonate grains shows clustering that is independent of the location of larger carbonate veins, indicating a relatively constant carbonate saturation state at the thin-section scale.

to 165 m depth) formed between  $52\text{--}190 \text{ }^\circ\text{C}$ , and listvenite matrix dolomite sampled near the basal thrust (178.7 m and 193.2 m depth) formed at temperatures of  $98\text{--}107 \text{ }^\circ\text{C}$ . Vein dolomite from the red Upper Listvenite formed at  $\sim 140 \text{ }^\circ\text{C}$  (67.4 m), whereas vein dolomite and magnesite from the relatively narrow Upper Serpentinite yielded one of the highest ( $227 \text{ }^\circ\text{C}$ ) and lowest ( $45 \text{ }^\circ\text{C}$ ) formation temperatures, respectively (Figure 5, Table S1; Beinlich et al., 2020d). Magnesite formation temperatures appear to increase with depth, whereas dolomite formation temperatures are relatively even within the uncertainty except for the high-temperature dolomite vein in the Upper Serpentinite ( $227 \text{ }^\circ\text{C}$ ; BT1B 43–02). Thus, our data show that the overall range in carbonate formation temperatures is wider than previously constrained by carbonate sampled in surface outcrops (Falk & Kelemen, 2015). Furthermore, the relatively erratic temperature distribution throughout core BT1B may reflect that the depth of carbonate formation varied nonsystematically with time, consistent with the repeated formation of deformation-induced high-permeability pathways allowing reactive fluid infiltration into different structural levels upon cooling of the system.

Assuming that cooling is related to ophiolite obduction, the obtained temperatures can be translated to carbonate formation depth ( $y$ ), using a simple equation for conductive heat transfer:




**Figure 9.** Phase and orientation maps using the Electron backscatter diffraction (EBSD) system of spheroidal carbonate from the least carbonated serpentinite (BT1B 44–01) and green listvenite (BT1B 70–01). Phase distribution (a) and misorientation to mean orientation (b) for a magnesite-dolomite composite grain in serpentinite. These data show that the sector zoning of the subgrain orientation may originate at dolomite inclusions near the grain center. Phase distribution (c) and misorientation to mean orientation (d) maps of spheroidal carbonate in listvenite sample 70–01 are similar to the observations for spheroidal carbonate in serpentinite. Black lines are magnesite-dolomite phase boundaries, and white lines indicate high angle dolomite–dolomite grain boundaries, and gray lines are low-angle ( $>2^\circ$ ) sub-grain boundaries in dolomite and magnesite.

$$y = \frac{q_0 - \sqrt{2T_0 H k \rho - 2H k \rho T + q_0^2}}{H \rho}, \quad (1)$$

where  $q_0$  is the surface heat flow ( $0.06 \pm 0.01 \text{ W}\cdot\text{m}^{-2}$  in forearc settings; Blackwell et al., 1990; Lewis et al., 1988),  $T_0$  is the temperature at the ocean floor (277 K),  $H$  is the current mean radioactive heat generation in the mantle ( $7.42 \times 10^{-12} \text{ W}\cdot\text{kg}^{-1}$ ) (Turcotte & Schubert, 2014),  $k$  is the thermal rock conductivity ( $2.9 \text{ W}\cdot\text{m}^{-1}\cdot\text{K}^{-1}$ ),  $\rho$  is the density of partially serpentinized peridotite ( $2900 \text{ kg}\cdot\text{m}^{-3}$ ), and  $T$  is the carbonate formation temperature (see also Plümper, King, et al., 2017).

The calculated depth is mainly dependent on chosen surface heat flow and thermal conductivity values. The  $2.9 \text{ W}\cdot\text{m}^{-1}\cdot\text{K}^{-1}$  thermal rock conductivity is based on measurements of serpentinized peridotite from the Mid-Atlantic Ridge (Kelemen et al., 2004), in agreement with the thermal properties of olivine and antigorite (Hofmeister, 1999; Osako et al., 2010). The paleo-surface heat flow is unknown but can be constrained based on pressure and temperature conditions of  $\sim 850^\circ\text{C}$  and 1 GPa recorded in the metamorphic sole underlying the ophiolite (Agard et al., 2016; Soret et al., 2017). The resulting surface heat flow ( $0.06 \pm 0.01 \text{ W}\cdot\text{m}^{-2}$ ) is consistent with these criteria and also with the inferred Permian age of the Samail mantle section and the  $\sim 92$  Ma mica Ar-Ar ages from the metamorphic sole (Hacker et al., 1996; Jaupart et al., 2015; van



**Figure 10.** Calculated equilibrium fluid  $\delta^{18}\text{O}$  (VSMOW) (Aharon, 1988; Horita, 2014) plotted against carbonate clumped isotope temperatures. The secondary abscissa shows depth inferred from the carbonate formation temperatures and simple heat conductivity calculation. The gray shaded area and transparent data points outline the range in fluid  $\delta^{18}\text{O}$  and temperature based on Falk and Kelemen (2015). Error bars reflect the propagated analytical uncertainty, standard deviation of the carbonate formation temperature, and the uncertainty of the mineral-fluid isotope fractionation calibration. Equilibrium fluid  $\delta^{18}\text{O}$  values calculated using various mineral-fluid oxygen isotope fractionation factors are shown in Figure S6.

Hinsbergen et al., 2019). The resulting geothermal gradient is  $25 \pm 0.6 \text{ }^\circ\text{C km}^{-1}$ . If our assumptions are correct, then the clumped isotope temperatures place the carbonation depth between  $\sim 10.1 \text{ km}$  at the highest temperature and  $1.7 \text{ km}$  at the lowest temperature, thus recording ophiolite uplift of  $\sim 8.4 \text{ km}$  (Figure 10; Figure S6).

#### 5.4. Implications for Reactive Fluid Composition

Ultramafic rock carbonation to listvenite is driven by the infiltration of  $\text{CO}_2$ -bearing reactive fluid into the ophiolite causing the dissolution of serpentine and precipitation of secondary Ca-Mg and Mg-carbonate in the serpentinite and additional quartz in the listvenite. Primary olivine was not found in core BT1B suggesting that serpentinization went to completion before  $\text{CO}_2$ -fluid infiltration, consistent with the mesh-textured distribution of oxides in listvenite (Figure S2). Hence, the carbonation reaction can be conceptualized, in Fe- and Ca-free space, as follows:



In the serpentinite, the reaction neither went to completion nor resulted in quartz/silica saturation of the alteration fluid, and  $\text{SiO}_{2,\text{aq}}$  was likely removed by the alteration fluid. In contrast, quartz/silica saturation was reached in the pervasively carbonated parts of BT1B that are now present as listvenite. The breakdown of relict clinopyroxene may have provided Ca for dolomite growth during limited carbonation of the serpentinite (e.g., in BT1B 44–01), whereas the significantly larger total amount of carbonate in the listvenite may have prevented dolomite precipitation instead of Ca-bearing magnesite. Additional replacement of primary

Cr-spinel results in the formation of fuchsite and implies the addition of potassium by the alteration fluid. Similarly altered ultramafic sequences have been described from several other ophiolite occurrences (e.g., Beinlich et al., 2012; Halls & Zhao, 1995; Hansen et al., 2005; Hinsken et al., 2017) and have been the subject of thermodynamic models linking mineral assemblages to fluid composition at a given alteration pressure and temperature (Beinlich, John, et al., 2020; Klein & Garrido, 2011; Klein & McCollom, 2013). However, the spheroidal carbonate texture in both the serpentinite and listvenite and the dendritic carbonate in the listvenite (Figures 2, 6, 9, and S5) are indicative of textural disequilibrium complicating constraints on the alteration fluid carbon concentration (i.e.,  $\text{XCO}_2$ ) using equilibrium thermodynamic modeling tools. Thus, we focus here on the implication of measured carbonate formation temperatures.

Assuming that the system had sufficient time to attain isotopic equilibrium between dissolved carbon species, the aqueous fluid, and the precipitating carbonate at the temperatures observed in this study, we can evaluate the fluid  $\delta^{18}\text{O}$  ( $\delta^{18}\text{O}_{\text{fluid}}$ ) based on the measured carbonate composition and available temperature-dependent fractionation factors (Aharon, 1988; Chacko & Deines, 2008; Horita, 2014; Matthews & Katz, 1977; Northrop & Clayton, 1966; Schauble et al., 2006; Schmidt et al., 2005; Zheng, 1999). The predicted oxygen isotope fractionation between magnesite and fluid ( $10^3 \ln \alpha^{\text{Mgs-fluid}}$ ) varies between 34‰ and 7.8‰ over the temperature range 45–247 °C. The strongest fractionation is predicted by Chacko and Deines (2008; magnesite- $\text{H}_2\text{O}$ ) at low temperature (34.9‰), which is distinctly stronger than the predicted fractionation by Zheng (1999) of  $\sim 27.3\%$ . The magnesite-fluid calibration by Aharon (1988; using magnesite 2) yields fractionation between 31‰ and 9.5‰ over the same temperature range. For dolomite, the predicted oxygen isotope fractionation varies between 22‰ and 7.2‰ for the temperature range 98–227 °C based on the most recent temperature relationship by Horita (2014). In the absence of experimental data on the temperature dependence of magnesite-fluid oxygen isotope fractionation, we here adopt the calibration by Aharon (1988) that is consistent with that by Zheng (1999) and applicable over the investigated temperature range, whereas the magnesite- $\text{H}_2\text{O}$  equation by Chacko and Deines (2008) is only applicable below 130 °C. Furthermore, for oxygen isotope fractionation between dolomite and fluid, we use the calibration by Horita (2014), calibrated over the temperature range 80–350 °C. By applying these

fractionation factors to the measured oxygen isotope composition of dolomite and magnesite samples that were used for clumped isotope thermometry defines an apparent trend from relatively elevated  $\delta^{18}\text{O}_{\text{fluid}}$  at a higher temperature (e.g., 12.2‰ at 247 °C) to relatively low  $\delta^{18}\text{O}_{\text{fluid}}$  at a lower temperature, for example, −9.9‰ at 45 °C. The different calibrations of the oxygen isotope fractionation factors result in a relative variation of  $\delta^{18}\text{O}_{\text{fluid}}$  of less than 3.7‰ at given temperature (Figure S6; see also García del Real et al., 2016).

As previously observed for the area around Hole BT1B (Falk & Kelemen, 2015; their Supplementary figure 8 and related text) and low-temperature carbonates in altered peridotite worldwide (García del Real et al., 2016), the resulting calculated  $\delta^{18}\text{O}_{\text{fluid}}$  displays a positive correlation with the corresponding carbonate formation temperature (Figure 10). This correlation may indicate carbonate precipitation from fluids that changed their oxygen isotope ratios and/or were derived from compositionally distinct sources. Low-temperature and low  $\delta^{18}\text{O}$  fluids may have been sourced from sediment compaction containing seawater-filled pore space during tectonic underthrusting (Falk & Kelemen, 2015; Kelemen et al., 2011), whereas high-temperature fluids with elevated  $\delta^{18}\text{O}$  may be released by subduction zone devolatilization reactions at the greenschist to amphibolite transition (Menzies et al., 2018). These different fluid sources can be reconciled with the measured carbonate compositions if the seawater  $\delta^{18}\text{O}$  remained fluid-buffered while trapped in the sedimentary pore space. Also, recent thermodynamic modeling has shown that magnesite-quartz assemblage requires fluid  $\text{CO}_2$  concentrations of ~2,500 ppm (at 100 °C and 5 kbar), that is, higher than that of low-pressure fluid saturated with calcite, suggesting that  $\text{CO}_2$  was primarily derived from subduction zone devolatilization (de Obeso et al., 2017; Kelemen et al., 2017).

Alternatively, the correlation between carbonate formation temperatures and calculated  $\delta^{18}\text{O}_{\text{fluid}}$  may be unrelated to the fluid composition and instead reflect a tendency toward departure from equilibrium of fluid-carbonate oxygen isotope exchange with decreasing temperature. Fluid-mineral oxygen isotope exchange may operate independently from the ordering of clumped isotopes in the carbonate mineral. Further experimental work on the relative rates of isotope exchange and ordering is required to fully investigate the apparent correlation of calculated  $\delta^{18}\text{O}_{\text{fluid}}$  and measured clumped isotope temperatures. However, independently determined fractionation factors are mutually consistent despite relative offsets (Figure S6), and a measurement artifact would be expected to cause a correlation between the measured carbonate clumped isotope composition and  $\delta^{18}\text{O}_{\text{carbonate}}$ .

## 6. Conclusions

Oman DP Hole BT1B intersects 200 m of listvenite containing two relic bands of the precursor serpentinite. The alteration to listvenite was pervasive, forming an assemblage consisting of Mg- and Ca-Mg-carbonate, quartz, minor fuchsite, and relic chromite. The presence of hematite and/or goethite in some parts of the core cannot be unambiguously linked to listvenite formation and may also result from a later overprint of already formed listvenite under oxidizing conditions. The least altered parts of BT1B are composed of serpentinite containing minor carbonate, while precursor olivine and pyroxene minerals could not be found during this study and earlier shipboard core logging. While we cannot rule out with certainty that carbonation was accompanied by serpentinization, the presence of oxides strings in carbonate resembling those typically found in serpentinized peridotite suggests that serpentinization went to completion prior to the carbonation.

The entire ultramafic part of the core is pervasively veined. Crosscutting relationships in the serpentinite indicate that carbonation was predated by at least two generations of serpentine veins. In the listvenite, crosscutting relationships indicate a general trend of early magnesite and minor dolomite veins being followed by the formation of a second set of broader magnesite veins that is subsequently crosscut by quartz and quartz-carbonate veins and then by dolomite veins. Whether the crosscutting relationships in the listvenite represent a network of fluid pathways during one alteration event or provide snapshots of consecutive fluid infiltration requires continued investigation beyond this contribution.

Carbonation of the serpentinite matrix resulted in the formation of abundant spheroidal magnesite-dolomite composite grains that show an even spatial and grain size distribution at the thin-section scale and appear to have nucleated on  $\mu\text{m}$ -sized, early dolomite. The textural analysis indicates that also listvenite formation commenced with the formation of spheroidal carbonate. However, the carbonate spheroids in the

#### Acknowledgments

The authors acknowledge the facilities and the scientific and technical assistance especially by Drs. M. Roberts, A. Suvorova, and M. Saunders of the Australian Microscopy & Microanalysis Research Facility at the Centre for Microscopy, Characterization & Analysis, The University of Western Australia, a facility funded by the University, State and Commonwealth Governments. This research used samples and/or data provided by the Oman Drilling Project. The Oman Drilling Project (OmanDP) has been possible through co-mingled funds from the International Continental Scientific Drilling Project (ICDP; Kelemen, Matter, Teagle Lead PIs), the Sloan Foundation-Deep Carbon Observatory (Grant 2014–3–01, Kelemen PI), the National Science Foundation (NSF-EAR-1516300, Kelemen lead PI), NASA-Astrobiology Institute (NNA15BB02A, Templeton PI), the German Research Foundation (DFG: KO 1723/21–1, Koepke PI), the Japanese Society for the Promotion of Science (JSPS no:16H06347, Michibayashi PI; and KAKENHI 16H02742, Takazawa PI), the European Research Council (ERC Adv: no.669972; Jamveit PI), the Swiss National Science Foundation (SNF:20FI21\_163073, Früh-Green PI), JAMSTEC, the TAMU-JR Science Operator, and contributions from the Sultanate of Oman Ministry of Regional Municipalities and Water Resources, the Oman Public Authority of Mining, Sultan Qaboos University, CRNS-Univ. Montpellier II, Columbia University of New York, and the University of Southampton. E.B. is grateful for receiving the Olaf Schuiling fond from Utrecht University, and A.B.'s participation in shipboard logging was supported by The Institute for Geoscience Research (TIGER). The clumped isotope analysis by I.A.M. and M.Z. was funded by the NWO VIDI grant WE.267002.1. O.P. has been supported by the ERC starting grant “nanoEARTH” (852069). TIMA phase mapping was undertaken using the Tescan Mira3 TIMA (ARC LE140100150) at the John de Laeter Centre, Curtin University. We thank M. Menzel and an anonymous reviewer for their insightful comments on the original manuscript.

listvenite are chemically and mineralogically distinct from those in the serpentinite. This suggests that the carbonate-bearing serpentinite and listvenite are unlikely to represent consecutive steps within the same reaction sequence but are the result of concomitant alteration, controlled by the accessibility of the alteration fluid, consistent with the preservation of serpentinite in BT1B.

The spheroidal and less commonly observed dendritic carbonate growth textures indicate fast precipitation from over-saturated fluid implying non-equilibrium conditions during serpentinite carbonation and the initial stage of listvenite formation. This may explain the pervasive nature of the alteration in those parts of BT1B that were accessible by the alteration fluid and also the lack of discernable talc-bearing transitional assemblages between the serpentinite and the listvenite that would be expected to form at chemical equilibrium in the temperature range defined by the clumped isotope thermometry. Spheroidal magnesite in the listvenite is frequently overgrown by euhedral magnesite, and both texturally distinct carbonate generations crystallized together with quartz. The formation of euhedral carbonate overgrowths in the listvenite reflects a driving force reduction and may be related to cooling and/or decrease in fluid saturation during the final stage of listvenite formation.

Clumped isotope thermometry of selected matrix and vein magnesite and dolomite indicates an overall carbonation temperature range of 45 to 247 °C. Carbonate formation temperatures show a nonsystematic distribution with sample depth in BT1B, indicating that reactive fluid could infiltrate the precursor ultramafic rock at different structural levels and times. Furthermore, the calculated oxygen isotope composition of the carbonate forming fluid varies between ~ -9.9‰ and ~12.1‰ and shows an apparent correlation with temperature. This correlation is consistent with previous observations in Oman (Falk & Kelemen, 2015) and suggests that carbonation took place as a multistage process driven by compositionally evolving reactive fluids that infiltrated the ophiolite during cooling.

#### Data Availability Statement

All data are available online. These include the processed carbonate stable isotope data and textural description of clumped isotope sampling spots (<https://doi.pangaea.de/10.1594/PANGAEA.914439>), the major element composition of carbonate phases (<https://doi.org/10.1594/PANGAEA.914455>), the major element composition of silicate phases (<https://doi.org/10.1594/PANGAEA.914440>), and the complete set of stable isotope data of all analyzed reference materials and unknown samples (<https://doi.pangaea.de/10.1594/PANGAEA.908649>).

#### References

- Agard, P., Yamato, P., Soret, M., Prigent, C., Guillot, S., Plunder, A., et al. (2016). Plate interface rheological switches during subduction infancy: Control on slab penetration and metamorphic sole formation. *Earth and Planetary Science Letters*, 451, 208–220. <https://doi.org/10.1016/j.epsl.2016.06.054>
- Aharon, P. (1988). A stable-isotope study of magnesites from the rum jungle uranium field, Australia: Implications for the origin of strata-bound massive magnesites. *Chemical Geology*, 69(1), 127–145. [https://doi.org/10.1016/0009-2541\(88\)90164-7](https://doi.org/10.1016/0009-2541(88)90164-7)
- Akbulut, M., & Pişkin, Ö., & Karayıit, A. İ (2006). The genesis of the carbonatized and silicified ultramafics known as listvenites: A case study from the Mihaliçcik region (Eskişehir), NW Turkey. *Geological Journal*, 41(5), 557–580. <https://doi.org/10.1002/gj.1058>
- Azer, M. K., Gahlan, H. A., Asimow, P. D., Mubarak, H. S., & Al-Kahtany, K. M. (2019). Multiple stages of carbonation and element redistribution during formation of ultramafic-hosted magnesite in Neoproterozoic ophiolites of the Arabian-Nubian shield, Egypt. *The Journal of Geology*, 127(1), 81–107. <https://doi.org/10.1086/700652>
- Barnes, I., & O'Neil, J. R. (1969). The relationship between fluids in some fresh alpine-type ultramafics and possible modern serpentinization, western United States. *GSA Bulletin*, 80(10), 1947–1960. [https://doi.org/10.1130/0016-7606\(1969\)80\[1947:TRBFIS\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1969)80[1947:TRBFIS]2.0.CO;2)
- Barnes, I., O'Neil, J. R., & Trescases, J. J. (1978). Present day serpentinization in New Caledonia, Oman and Yugoslavia. *Geochimica et Cosmochimica Acta*, 42(1), 144–145. [https://doi.org/10.1016/0016-7037\(78\)90225-9](https://doi.org/10.1016/0016-7037(78)90225-9)
- Beinlich, A., Austrheim, H., Glodny, J., Erambert, M., & Andersen, T. B. (2010). CO<sub>2</sub> sequestration and extreme mg depletion in serpentinized peridotite clasts from the Devonian Solund basin, SW-Norway. *Geochimica et Cosmochimica Acta*, 74(24), 6935–6964. <https://doi.org/10.1016/j.gca.2010.07.027>
- Beinlich, A., Austrheim, H., Mavromatis, V., Grguric, B., Putnis, C. V., & Putnis, A. (2018). Peridotite weathering is the missing ingredient of Earth's continental crust composition. *Nature Communications*, 9(1), 1–12. <https://doi.org/10.1038/s41467-018-03039-9>
- Beinlich, A., John, T., Vrijmoed, J. C., Tominaga, M., Magna, T., & Podladchikov, Y. Y. (2020). Instantaneous rock transformations in the deep crust driven by reactive fluid flow. *Nature Geoscience*, 13(4), 307–311. <https://doi.org/10.1038/s41561-020-0554-9>
- Beinlich, A., Plümper, O., Boter, E., Müller, I. A., Kourim, F., Ziegler, M., et al., & The Oman Drilling Project Phase 1 Science Party (2019). Clumped isotope data from hole BT1B, Oman Drilling Project. *PANGAEA*, n/a, n/a. <https://doi.org/10.1594/PANGAEA.908649>
- Beinlich, A., Plümper, O., Boter, E., Müller, I. A., Kourim, F., Ziegler, M., et al., & The Oman Drilling Project Phase 1 Science Party (2020a). Average compositions of serpentine and fuchsite from hole BT1B, Oman Drilling Project. *PANGAEA*. <https://doi.org/10.1594/PANGAEA.914440>

- Beinlich, A., Plümper, O., Boter, E., Müller, I. A., Kourim, F., Ziegler, M., et al., & The Oman Drilling Project Phase 1 Science Party (2020b). Average magnesite composition from selected serpentinite and listvenite samples from hole BT1B, Oman Drilling Project. *PANGAEA*. <https://doi.org/10.1594/PANGAEA.914455>
- Beinlich, A., Plümper, O., Boter, E., Müller, I. A., Kourim, F., Ziegler, M., et al., & The Oman Drilling Project Phase 1 Science Party (2020c). Carbonate stable and clumped isotope data, calculated carbonate formation temperatures and fluid equilibrium oxygen isotope composition from hole BT1B. *Oman Drilling Project. PANGAEA*. <https://doi.org/10.1594/PANGAEA.914439>
- Beinlich, A., Plümper, O., Hövelmann, J., Austrheim, H., & Jamtveit, B. (2012). Massive serpentinite carbonation at Linnajavri, N-Norway. *Terra Nova*, *24*(6), 446–455. <https://doi.org/10.1111/j.1365-3121.2012.01083.x>
- Belogub, E. V., Melekestseva, I. Y., Novoselov, K. A., Zabortina, M. V., Tret'yakov, G. A., Zaykov, V. V., & Yuminov, A. M. (2017). Listvenite-related gold deposits of the South Urals (Russia): A review. *Ore Geology Reviews*, *85*, 247–270. <https://doi.org/10.1016/j.oregeorev.2016.11.008>
- Bernasconi, S. M., Müller, I. A., Bergmann, K. D., Breitenbach, S. F. M., Fernandez, A., Hodell, D. A., et al. (2018). Reducing uncertainties in carbonate clumped isotope analysis through consistent carbonate-based standardization. *Geochemistry, Geophysics, Geosystems*, *28*95–2914. <https://doi.org/10.1029/2017gc007385>
- Blackwell, D. D., Steele, J. L., Frohme, M. K., Murphey, C. F., Priest, G. R., & Black, G. L. (1990). Heat flow in the Oregon Cascade Range and its correlation with regional gravity, curie point depths, and geology. *Journal of Geophysical Research*, *95*(B12), 19,475–19,493. <https://doi.org/10.1029/JB095iB12p19475>
- Bonatti, E., Lawrence, J. R., Hamlyn, P. R., & Breger, D. (1980). Aragonite from deep sea ultramafic rocks. *Geochimica et Cosmochimica Acta*, *44*(8), 1207–1214. [https://doi.org/10.1016/0016-7037\(80\)90074-5](https://doi.org/10.1016/0016-7037(80)90074-5)
- Bonifacie, M., Calmels, D., Eiler, J. M., Horita, J., Chaduteau, C., Vasconcelos, C., et al. (2017). Calibration of the dolomite clumped isotope thermometer from 25 to 350°C, and implications for a universal calibration for all (Ca, mg, Fe)CO<sub>3</sub> carbonates. *Geochimica et Cosmochimica Acta*, *200*, 255–279. <https://doi.org/10.1016/j.gca.2016.11.028>
- Bosze, S., & Rakovan, J. (2002). Surface-structure-controlled sectoral zoning of the rare earth elements in fluorite from long Lake, New York, and Bingham, New Mexico, USA. *Geochimica et Cosmochimica Acta*, *66*(6), 997–1009. [https://doi.org/10.1016/S0016-7037\(01\)00822-5](https://doi.org/10.1016/S0016-7037(01)00822-5)
- Chacko, T., & Deines, P. (2008). Theoretical calculation of oxygen isotope fractionation factors in carbonate systems. *Geochimica et Cosmochimica Acta*, *72*(15), 3642–3660. <https://doi.org/10.1016/j.gca.2008.06.001>
- Chavagnac, V., Ceuleneer, G., Monnin, C., Lansac, B., Hoareau, G., & Boulart, C. (2013). Mineralogical assemblages forming at hyperalkaline warm springs hosted on ultramafic rocks: A case study of Oman and Ligurian ophiolites. *Geochemistry, Geophysics, Geosystems*, *14*, 2474–2495. <https://doi.org/10.1002/ggge.20146>
- Chavagnac, V., Monnin, C., Ceuleneer, G., Boulart, C., & Hoareau, G. (2013). Characterization of hyperalkaline fluids produced by low-temperature serpentinization of mantle peridotites in the Oman and Ligurian ophiolites. *Geochemistry, Geophysics, Geosystems*, *14*, 2496–2522. <https://doi.org/10.1002/ggge.20147>
- Clark, I. D., & Fontes, J.-C. (1990). Paleoclimatic reconstruction in northern Oman based on carbonates from hyperalkaline groundwaters. *Quaternary Research*, *33*(3), 320–336. [https://doi.org/10.1016/0033-5894\(90\)90059-T](https://doi.org/10.1016/0033-5894(90)90059-T)
- Coleman, R. G. (1981). Tectonic setting for ophiolite obduction in Oman. *Journal of Geophysical Research*, *86*, 2497–2508. <https://doi.org/10.1029/JB086iB04p02497>
- Cowan, R. J., Searle, M. P., & Waters, D. J. (2014). Structure of the metamorphic sole to the Oman Ophiolite, Sumeini window and Wadi Tayyin: Implications for ophiolite obduction processes. *Geological Society, London, Special Publications*, *392*(1), 155–175. <https://doi.org/10.1144/sp392.8>
- Das Sharma, S., Patil, D. J., & Gopalan, K. (2002). Temperature dependence of oxygen isotope fractionation of CO<sub>2</sub> from magnesite–phosphoric acid reaction. *Geochimica et Cosmochimica Acta*, *66*(4), 589–593. [https://doi.org/10.1016/S0016-7037\(01\)00833-X](https://doi.org/10.1016/S0016-7037(01)00833-X)
- Eiler, J. M. (2007). “Clumped-isotope” geochemistry—The study of naturally-occurring, multiply-substituted isotopologues. *Earth and Planetary Science Letters*, *262*(3), 309–327. <https://doi.org/10.1016/j.epsl.2007.08.020>
- Emam, A., & Zoheir, B. (2013). Au and Cr mobilization through metasomatism: Microchemical evidence from ore-bearing listvenite, south Eastern Desert of Egypt. *Journal of Geochemical Exploration*, *125*, 34–45. <https://doi.org/10.1016/j.gexplo.2012.11.004>
- Falk, E. S., Guo, W., Paukert, A. N., Matter, J. M., Mervine, E. M., & Kelemen, P. B. (2016). Controls on the stable isotope compositions of travertine from hyperalkaline springs in Oman: Insights from clumped isotope measurements. *Geochimica et Cosmochimica Acta*, *192*, 1–28. <https://doi.org/10.1016/j.gca.2016.06.026>
- Falk, E. S., & Kelemen, P. B. (2015). Geochemistry and petrology of listvenite in the Samail ophiolite, Sultanate of Oman: Complete carbonation of peridotite during ophiolite emplacement. *Geochimica et Cosmochimica Acta*, *160*, 70–90. <https://doi.org/10.1016/j.gca.2015.03.014>
- Gahlan Hisham, A., Azer Mokhles, K., & Asimow Paul, D. (2018). On the relative timing of listwaenite formation and chromian spinel equilibration in serpentinites. *American Mineralogist*, *103*(7), 1087. <https://doi.org/10.2138/am-2018-6473>
- García del Real, P., Maher, K., Kluge, T., Bird, D. K., Brown, G. E., & John, C. M. (2016). Clumped-isotope thermometry of magnesium carbonates in ultramafic rocks. *Geochimica et Cosmochimica Acta*, *193*, 222–250. <https://doi.org/10.1016/j.gca.2016.08.003>
- Ghosh, P., Adkins, J., Affek, H., Balta, B., Guo, W., Schauble, E. A., et al. (2006). <sup>13</sup>C–<sup>18</sup>O bonds in carbonate minerals: A new kind of paleothermometer. *Geochimica et Cosmochimica Acta*, *70*(6), 1439–1456. <https://doi.org/10.1016/j.gca.2005.11.014>
- Gnos, E. (1998). Peak metamorphic conditions of garnet amphibolites beneath the Semail Ophiolite: Implications for an inverted pressure gradient. *International Geology Review*, *40*(4), 281–304. <https://doi.org/10.1080/00206819809465210>
- Gránásy, L., Pusztai, T., Tegze, G., Warren, J. A., & Douglas, J. F. (2005). Growth and form of spherulites. *Physical Review E*, *72*(1), 011605. <https://doi.org/10.1103/PhysRevE.72.011605>
- Grozeva, N. G., Klein, F., Seewald, J. S., & Sylva, S. P. (2017). Experimental study of carbonate formation in oceanic peridotite. *Geochimica et Cosmochimica Acta*, *199*, 264–286. <https://doi.org/10.1016/j.gca.2016.10.052>
- Hacker, B. R. (1994). Rapid emplacement of young oceanic lithosphere: Argon geochronology of the Oman Ophiolite. *Science*, *265*(5178), 1563–1565. <https://doi.org/10.1126/science.265.5178.1563>
- Hacker, B. R., & Gnos, E. (1997). The conundrum of Samail: Explaining the metamorphic history. *Tectonophysics*, *279*(1), 215–226. [https://doi.org/10.1016/S0040-1951\(97\)00114-5](https://doi.org/10.1016/S0040-1951(97)00114-5)
- Hacker, B. R., & Mosenfelder, J. L. (1996). Metamorphism and deformation along the emplacement thrust of the Samail ophiolite, Oman. *Earth and Planetary Science Letters*, *144*(3), 435–451. [https://doi.org/10.1016/S0012-821X\(96\)00186-0](https://doi.org/10.1016/S0012-821X(96)00186-0)
- Hacker, B. R., Mosenfelder, J. L., & Gnos, E. (1996). Rapid emplacement of the Oman ophiolite: Thermal and geochronologic constraints. *Tectonics*, *15*(6), 1230–1247. <https://doi.org/10.1029/96tc01973>

- Halls, C., & Zhao, R. (1995). Listvenite and related rocks - perspectives on terminology and mineralogy with reference to an occurrence at Cregganbaun, Co Mayo, republic-of-Ireland. *Mineralium Deposita*, 30(3–4), 303–313. <https://doi.org/10.1007/Bf00196366>
- Hansen, L. D., Dipple, G. M., Gordon, T. M., & Kellett, D. A. (2005). Carbonated serpentinite (listwanite) at Atlin, British Columbia: A geological analogue to carbon dioxide sequestration. *The Canadian Mineralogist*, 43, 225–239. <https://doi.org/10.2113/gscanmin.43.1.225>
- Haque, N., Cochrane, R. F., & Mullis, A. M. (2016). Rapid solidification morphologies in Ni<sub>3</sub>Ge: Spherulites, dendrites and dense-branched fractal structures. *Intermetallics*, 76, 70–77. <https://doi.org/10.1016/j.intermet.2016.06.012>
- van Hinsbergen, D. J. J., Maffione, M., Koornneef, L. M. T., & Guilmette, C. (2019). Kinematic and paleomagnetic restoration of the Semail ophiolite (Oman) reveals subduction initiation along an ancient Neotethyan fracture zone. *Earth and Planetary Science Letters*, 518, 183–196. <https://doi.org/10.1016/j.epsl.2019.04.038>
- Hinsken, T., Brocker, M., Strauss, H., & Bulle, F. (2017). Geochemical, isotopic and geochronological characterization of listvenite from the upper unit on Tinos, Cyclades, Greece. *Lithos*, 282, 281–297. <https://doi.org/10.1016/j.lithos.2017.02.019>
- Hofmeister, A. M. (1999). Mantle values of thermal conductivity and the Geotherm from phonon lifetimes. *Science*, 283(5408), 1699–1706. <https://doi.org/10.1126/science.283.5408.1699>
- Horita, J. (2014). Oxygen and carbon isotope fractionation in the system dolomite–water–CO<sub>2</sub> to elevated temperatures. *Geochimica et Cosmochimica Acta*, 129, 111–124. <https://doi.org/10.1016/j.gca.2013.12.027>
- Huntington, K. W., Eiler, J. M., Affek, H. P., Guo, W., Bonifacie, M., Yeung, L. Y., et al. (2009). Methods and limitations of ‘clumped’ CO<sub>2</sub> isotope (Δ<sub>47</sub>) analysis by gas-source isotope ratio mass spectrometry. *Journal of Mass Spectrometry*, 44(9), 1318–1329. <https://doi.org/10.1002/jms.1614>
- Jaupart, C., Labrosse, S., Lucazeau, F., & Mareschal, J. C. (2015). Temperatures, heat, and energy in the mantle of the earth. In G. Schubert (Ed.), *Treatise on Geophysics (Second Edition)*, (pp. 223–270). Oxford: Elsevier. <https://doi.org/10.1016/B978-0-444-53802-4.00126-3>
- Kele, S., Breitenbach, S. F. M., Capezzuoli, E., Meckler, A. N., Ziegler, M., Millan, I. M., et al. (2015). Temperature dependence of oxygen- and clumped isotope fractionation in carbonates: A study of travertines and tufas in the 6–95°C temperature range. *Geochimica et Cosmochimica Acta*, 168, 172–192. <https://doi.org/10.1016/j.gca.2015.06.032>
- de Obeso, J. C., & Kelemen, P. B. (2018). Fluid rock interactions on residual mantle peridotites overlain by shallow oceanic limestones: Insights from Wadi fins, Sultanate of Oman. *Chemical Geology*, 498, 139–149. <https://doi.org/10.1016/j.chemgeo.2018.09.022>
- de Obeso, J. C., Kelemen, P. B., Manning, C. E., Michibayashi, K., & Harris, M. (2017). *Listvenite formation from peridotite: Insights from Oman Drilling Project Hole BT1B and preliminary reaction path model approach*. American Geophysical Union, Fall Meeting 2017, Abstract #V43G-2941. New Orleans, Louisiana, USA.
- Kelemen, P. B., Kikawa, E., & Miller, J. (2004). Drilling mantle peridotite along the mid-Atlantic ridge from 14° to 16°N. *Proceedings of the Ocean Drilling Program, Initial Reports (Ocean Drilling Program, College Station, TX)*, 209, 1–139. <https://doi.org/10.2973/odp.proc.ir.209.2004>
- Kelemen, P. B., & Matter, J. (2008). In situ carbonation of peridotite for CO<sub>2</sub> storage. *Proceedings of the National Academy of Sciences*, 105(45), 17,295–17,300. <https://doi.org/10.1073/pnas.0805794105>
- Kelemen, P. B., Matter, J., Streit, E. E., Rudger, J. F., Curry, W. B., & Blusztajn, J. (2011). Rates and mechanisms of mineral carbonation in peridotite: Natural processes and recipes for enhanced, in situ CO<sub>2</sub> capture and storage. *Annual Review of Earth and Planetary Sciences*, 39, 545–576. <https://doi.org/10.1146/annurev-earth-092010-152509>
- Kelemen, P. B., Matter, J. M., Teagle, D. A. H., Coggon, J. A., & the Oman Drilling Science Team (2020). *Proceedings of the Oman Drilling Project*. College Station, TX: (International Ocean Discovery Program). <https://doi.org/10.14379/Oman.ph1-2.proc.2020>
- Kelemen, P. B., Godard, M., Johnson, K. T. M., Okazaki, K., Manning, C. E., Urai, J. L., et al., & The Oman Drilling Project Phase I Science Party (2017). *Peridotite carbonation at the leading edge of the mantle wedge: OmDP Site BT1*. American Geophysical Union, Fall Meeting 2017, Abstract #V24E-06. New Orleans, Louisiana, USA.
- Klein, F., & Garrido, C. J. (2011). Thermodynamic constraints on mineral carbonation of serpentinized peridotite. *Lithos*, 126(3), 147–160. <https://doi.org/10.1016/j.lithos.2011.07.020>
- Klein, F., & McCollom, T. M. (2013). From serpentinization to carbonation: New insights from a CO<sub>2</sub> injection experiment. *Earth and Planetary Science Letters*, 379, 137–145. <https://doi.org/10.1016/j.epsl.2013.08.017>
- Lacinska, A. M., & Styles, M. T. (2012). Silicified serpentinite—A residuum of a tertiary palaeo-weathering surface in the United Arab Emirates. *Geological Magazine*, 150(3), 385–395. <https://doi.org/10.1017/S0016756812000325>
- Lacinska, A. M., Styles, M. T., & Farrant, A. R. (2014). Near-surface diagenesis of ophiolite-derived conglomerates of the Barzaman formation, United Arab Emirates: A natural analogue for permanent CO<sub>2</sub> sequestration via mineral carbonation of ultramafic rocks. *Geological Society, London, Special Publications*, 392(1), 343–360. <https://doi.org/10.1144/sp392.18>
- Lanphere, M. A. (1981). K-Ar ages of metamorphic rocks at the base of the Samail Ophiolite, Oman. *Journal of Geophysical Research*, 86(B4), 2777–2782. <https://doi.org/10.1029/JB086iB04p02777>
- Launay, J., & Fontes, J.-F. (1985). Les sources thermales de Prony (Nouvelle-Calédonie) et leurs précipités chimiques. Exemple de formation de brucite primaire. *Géologie de la France*, 1, 83–100.
- Lewis, T. J., Bentkowski, W. H., Davis, E. E., Hyndman, R. D., Souther, J. G., & Wright, J. A. (1988). Subduction of the Juan de Fuca plate: Thermal consequences. *Journal of Geophysical Research*, 93(B12), 15,207–15,225. <https://doi.org/10.1029/JB093iB12p15207>
- Matthews, A., & Katz, A. (1977). Oxygen isotope fractionation during the dolomitization of calcium carbonate. *Geochimica et Cosmochimica Acta*, 41(10), 1431–1438. [https://doi.org/10.1016/0016-7037\(77\)90249-6](https://doi.org/10.1016/0016-7037(77)90249-6)
- Meckler, A. N., Ziegler, M., Millán, M. I., Breitenbach, S. F. M., & Bernasconi, S. M. (2014). Long-term performance of the Kiel carbonate device with a new correction scheme for clumped isotope measurements. *Rapid Communications in Mass Spectrometry*, 28(15), 1705–1715. <https://doi.org/10.1002/rcm.6949>
- Menzel, M. D., Garrido, C. J., López Sánchez-Vizcaíno, V., Marchesi, C., Hidas, K., Escayola, M. P., & Delgado Huertas, A. (2018). Carbonation of mantle peridotite by CO<sub>2</sub>-rich fluids: The formation of listvenites in the advocate ophiolite complex (Newfoundland, Canada). *Lithos*, 323, 238–261. <https://doi.org/10.1016/j.lithos.2018.06.001>
- Menzies, C. D., Wright, S. L., Craw, D., James, R. H., Alt, J. C., Cox, S. C., et al. (2018). Carbon dioxide generation and drawdown during active orogenesis of siliciclastic rocks in the southern Alps, New Zealand. *Earth and Planetary Science Letters*, 481, 305–315. <https://doi.org/10.1016/j.epsl.2017.10.010>
- Mervine, E. M., Humphris, S. E., Sims, K. W. W., Kelemen, P. B., & Jenkins, W. J. (2014). Carbonation rates of peridotite in the Samail Ophiolite, Sultanate of Oman, constrained through <sup>14</sup>C dating and stable isotopes. *Geochimica et Cosmochimica Acta*, 126, 371–397. <https://doi.org/10.1016/j.gca.2013.11.007>

- Müller, I. A., Fernandez, A., Radke, J., van Dijk, J., Bowen, D., Schweters, J., & Bernasconi, S. M. (2017). Carbonate clumped isotope analyses with the long-integration dual-inlet (LIDI) workflow: Scratching at the lower sample weight boundaries. *Rapid Communications in Mass Spectrometry*, *31*(12), 1057–1066. <https://doi.org/10.1002/rcm.7878>
- Müller, I. A., Rodríguez-Blanco, J. D., Storck, J.-C., do Nascimento, G. S., Bontognali, T. R. R., Vasconcelos, C., et al. (2019). Calibration of the oxygen and clumped isotope thermometers for (proto-)dolomite based on synthetic and natural carbonates. *Chemical Geology*, *525*, 1–17. <https://doi.org/10.1016/j.chemgeo.2019.07.014>
- Müller, I. A., Violay, M. E. S., Storck, J.-C., Fernandez, A., van Dijk, J., Madonna, C., & Bernasconi, S. M. (2017). Clumped isotope fractionation during phosphoric acid digestion of carbonates at 70°C. *Chemical Geology*, *449*, 1–14. <https://doi.org/10.1016/j.chemgeo.2016.11.030>
- Murray, S. T., Arienzo, M. M., & Swart, P. K. (2016). Determining the  $\Delta_{47}$  acid fractionation in dolomites. *Geochimica et Cosmochimica Acta*, *174*, 42–53. <https://doi.org/10.1016/j.gca.2015.10.029>
- Nasir, S., Al Sayigh, A. R., Al Harthy, A., Al-Khribash, S., Al-Jaaldi, O., Musllam, A., et al. (2007). Mineralogical and geochemical characterization of listwaenite from the Semail Ophiolite, Oman. *Chemie der Erde-Geochemistry*, *67*(3), 213–228. <https://doi.org/10.1016/j.chemer.2005.01.003>
- Neal, C., & Stanger, G. (1983). Hydrogen generation from mantle source rocks in Oman. *Earth and Planetary Science Letters*, *66*, 315–320. [https://doi.org/10.1016/0012-821X\(83\)90144-9](https://doi.org/10.1016/0012-821X(83)90144-9)
- Neal, C., & Stanger, G. (1985). Past and present serpentinisation of ultramafic rocks; an example from the Semail ophiolite nappe of northern Oman. In J. I. Drever (Ed.), *The Chemistry of Weathering, Nato ASI Series (C: Mathematical and Physical Sciences)*(Vol. 149). Dordrecht, The Netherlands: Springer.
- Nicolas, A., & Boudier, F. (1995). Mapping oceanic ridge segments in Oman ophiolite. *Journal of Geophysical Research*, *100*(B4), 6179–6197. <https://doi.org/10.1029/94jb01188>
- Northrop, D. A., & Clayton, R. N. (1966). Oxygen-isotope fractionations in systems containing dolomite. *The Journal of Geology*, *74*(2), 174–196. <https://doi.org/10.1086/627153>
- Osako, M., Yoneda, A., & Ito, E. (2010). Thermal diffusivity, thermal conductivity and heat capacity of serpentine (antigorite) under high pressure. *Physics of the Earth and Planetary Interiors*, *183*(1), 229–233. <https://doi.org/10.1016/j.pepi.2010.07.005>
- Passey, B. H., & Henkes, G. A. (2012). Carbonate clumped isotope bond reordering and geospeedometry. *Earth and Planetary Science Letters*, *351–352*, 223–236. <https://doi.org/10.1016/j.epsl.2012.07.021>
- Paukert, A. N., Matter, J. M., Kelemen, P. B., Shock, E. L., & Havig, J. R. (2012). Reaction path modeling of enhanced in situ CO<sub>2</sub> mineralization for carbon sequestration in the peridotite of the Samail Ophiolite, Sultanate of Oman. *Chemical Geology*, *330*, 86–100. <https://doi.org/10.1016/j.chemgeo.2012.08.013>
- Pearce, M. A., Timms, N. E., Hough, R. M., & Cleverley, J. S. (2013). Reaction mechanism for the replacement of calcite by dolomite and siderite: Implications for geochemistry, microstructure and porosity evolution during hydrothermal mineralisation. *Contributions to Mineralogy and Petrology*, *166*(4), 995–1009. <https://doi.org/10.1007/s00410-013-0905-2>
- Plümpner, O., Botan, A., Los, C., Liu, Y., Malthé-Sorensen, A., & Jamtveit, B. (2017). Fluid-driven metamorphism of the continental crust governed by nanoscale fluid flow. *Nature Geoscience*, *10*(9), 685–690. <https://doi.org/10.1038/ngeo3009>
- Plümpner, O., King, H. E., Geisler, T., Liu, Y., Pabst, S., Savov, I. P., et al. (2017). Subduction zone forearc serpentinites as incubators for deep microbial life. *Proceedings of the National Academy of Sciences*, *114*(17), 4324–4329. <https://doi.org/10.1073/pnas.1612147114>
- Quesnel, B., Gautier, P., Boulvais, P., Cathelineau, M., Maurizot, P., Cluzel, D., et al. (2013). Syn-tectonic, meteoric water-derived carbonation of the New Caledonia peridotite nappe. *Geology*, *41*(10), 1063–1066. <https://doi.org/10.1130/G34531.1>
- Rioux, M., Bowring, S., Kelemen, P., Gordon, S., Dudás, F., & Miller, R. (2012). Rapid crustal accretion and magma assimilation in the Oman-U.A.E. ophiolite: High precision U-Pb zircon geochronology of the gabbroic crust. *Journal of Geophysical Research*, *117*, B07201. <https://doi.org/10.1029/2012JB009273>
- Rioux, M., Bowring, S., Kelemen, P., Gordon, S., Miller, R., & Dudás, F. (2013). Tectonic development of the Samail ophiolite: High-precision U-Pb zircon geochronology and Sm-Nd isotopic constraints on crustal growth and emplacement. *Journal of Geophysical Research: Solid Earth*, *118*, 2085–2101. <https://doi.org/10.1002/jgrb.50139>
- Rioux, M., Garber, J., Bauer, A., Bowring, S., Searle, M., Kelemen, P., & Hacker, B. (2016). Synchronous formation of the metamorphic sole and igneous crust of the Semail ophiolite: New constraints on the tectonic evolution during ophiolite formation from high-precision U-Pb zircon geochronology. *Earth and Planetary Science Letters*, *451*, 185–195. <https://doi.org/10.1016/j.epsl.2016.06.051>
- Rosenbaum, J., & Sheppard, S. M. F. (1986). An isotopic study of siderites, dolomites and ankerites at high temperatures. *Geochimica et Cosmochimica Acta*, *50*(6), 1147–1150. [https://doi.org/10.1016/0016-7037\(86\)90396-0](https://doi.org/10.1016/0016-7037(86)90396-0)
- Ryb, U., Lloyd, M. K., Stolper, D. A., & Eiler, J. M. (2017). The clumped-isotope geochemistry of exhumed marbles from Naxos, Greece. *Earth and Planetary Science Letters*, *470*, 1–12. <https://doi.org/10.1016/j.epsl.2017.04.026>
- Schauble, E. A., Ghosh, P., & Eiler, J. M. (2006). Preferential formation of <sup>13</sup>C-<sup>18</sup>O bonds in carbonate minerals, estimated using first-principles lattice dynamics. *Geochimica et Cosmochimica Acta*, *70*(10), 2510–2529. <https://doi.org/10.1016/j.gca.2006.02.011>
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., et al. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. <https://doi.org/10.1038/nmeth.2019>
- Schmidt, M., Xeflide, S., Botz, R., & Mann, S. (2005). Oxygen isotope fractionation during synthesis of CaMg-carbonate and implications for sedimentary dolomite formation. *Geochimica et Cosmochimica Acta*, *69*(19), 4665–4674. <https://doi.org/10.1016/j.gca.2005.06.025>
- Searle, M. P., & Malpas, J. (1980). Structure and metamorphism of rocks beneath the Semail ophiolite of Oman and their significance in ophiolite obduction. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, *71*(4), 247–262. <https://doi.org/10.1017/S0263593300013614>
- Searle, M. P., Waters, D. J., Garber, J. M., Rioux, M., Cherry, A. G., & Ambrose, T. K. (2015). Structure and metamorphism beneath the obducting Oman ophiolite: Evidence from the Bani Hamid granulites, northern Oman mountains. *Geosphere*, *11*(6), 1812–1836. <https://doi.org/10.1130/GES01199.1>
- Shtukenberg, A. G., Punin, Y. O., Gunn, E., & Kahr, B. (2012). Spherulites. *Chemical Reviews*, *112*(3), 1805–1838. <https://doi.org/10.1021/cr200297f>
- Soret, M., Agard, P., Dubacq, B., Plunder, A., & Yamato, P. (2017). Petrological evidence for stepwise accretion of metamorphic soles during subduction infancy (Semail ophiolite, Oman and UAE). *Journal of Metamorphic Geology*, *35*(9), 1051–1080. <https://doi.org/10.1111/jmg.12267>
- Stanger, G. (1985). Silicified serpentinite in the Semail nappe of Oman. *Lithos*, *18*, 13–22. [https://doi.org/10.1016/0024-4937\(85\)90003-9](https://doi.org/10.1016/0024-4937(85)90003-9)
- Stowell, H., Zuluaga, C., Boyle, A., & Bulman, G. (2011). Garnet sector and oscillatory zoning linked with changes in crystal morphology during rapid growth, north cascades, Washington. *American Mineralogist*, *96*(8–9), 1354–1362. <https://doi.org/10.2138/am.2011.3759>

- Streit, E., Kelemen, P., & Eiler, J. (2012). Coexisting serpentine and quartz from carbonate-bearing serpentinized peridotite in the Samail Ophiolite, Oman. *Contributions to Mineralogy and Petrology*, *164*(5), 821–837. <https://doi.org/10.1007/s00410-012-0775-z>
- Sunagawa, I. (2005). *Crystals: Growth, Morphology and Perfection* (p. 295). New York: Cambridge University Press.
- Tilton, G. R., Hopson, C. A., & Wright, J. E. (1981). Uranium-lead isotopic ages of the Samail Ophiolite, Oman, with applications to Tethyan Ocean ridge tectonics. *Journal of Geophysical Research*, *86*(B4), 2763–2775. <https://doi.org/10.1029/JB086iB04p02763>
- Tominaga, M., Beinlich, A., Lima, E. A., Tivey, M. A., Hampton, B. A., Weiss, B., & Harigane, Y. (2017). Multi-scale magnetic mapping of serpentinite carbonation. *Nature Communications*, *8*(1), 1–10. <https://doi.org/10.1038/s41467-017-01610-4>
- Tsikouras, B., Karipi, S., Grammatikopoulos, T. A., & Hatzipanagiotou, K. (2006). Listwaenite evolution in the ophiolite melange of Iti Mountain (continental Central Greece). *European Journal of Mineralogy*, *18*(2), 243–255. <https://doi.org/10.1127/0935-1221/2006/0018-0243>
- Turcotte, D., & Schubert, G. (2014). *Geodynamics* (3rd ed.). Cambridge, U.K: Cambridge University Press.
- Warren, C. J., Parrish, R. R., Waters, D. J., & Searle, M. P. (2005). Dating the geologic history of Oman's Samail ophiolite: Insights from U-Pb geochronology. *Contributions to Mineralogy and Petrology*, *150*(4), 403–422. <https://doi.org/10.1007/s00410-005-0028-5>
- Whitney, D. L., & Evans, B. W. (2010). Abbreviations for names of rock-forming minerals. *American Mineralogist*, *95*(1), 185–187. <https://doi.org/10.2138/am.2010.3371>
- Wilde, A., Simpson, L., & Hanna, S. (2002). Preliminary study of tertiary hydrothermal alteration and platinum deposition in the Oman ophiolite. *Journal of the Virtual Explorer*, *6*, 7–13. <https://doi.org/10.3809/jvirtex.2002.00038>
- Zheng, X., Cordonnier, B., Zhu, W., Renard, F., & Jamtveit, B. (2018). Effects of confinement on reaction-induced fracturing during hydration of periclase. *Geochemistry, Geophysics, Geosystems*, *19*, 2661–2672. <https://doi.org/10.1029/2017gc007322>
- Zheng, Y.-F. (1999). Oxygen isotope fractionation in carbonate and sulfate minerals. *Geochemical Journal*, *33*(2), 109–126. <https://doi.org/10.2343/geochemj.33.109>