



HAL
open science

Pseudo-nitzschia sp. diversity and seasonality in the southern North Sea, domoic acid levels and associated phytoplankton communities

Alice Delegrange, A. Lefebvre, F. Gohin, Lucie Courcot, Dorothée Vincent

► To cite this version:

Alice Delegrange, A. Lefebvre, F. Gohin, Lucie Courcot, Dorothée Vincent. Pseudo-nitzschia sp. diversity and seasonality in the southern North Sea, domoic acid levels and associated phytoplankton communities. *Estuarine, Coastal and Shelf Science*, 2018, 214, pp.194-206. 10.1016/j.ecss.2018.09.030 . insu-03664848

HAL Id: insu-03664848

<https://insu.hal.science/insu-03664848>

Submitted on 3 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

***Pseudo-nitzschia* sp. diversity and seasonality in the southern North Sea, domoic acid levels and associated phytoplankton communities**

Delegrange A. ^{1,2,*}, Lefebvre Alain ³, Gohin Francis ⁴, Courcot L. ¹, Vincent D. ¹

¹ Univ. Littoral Côte d'Opale, Univ. Lille, CNRS, UMR 8187, LOG, Laboratoire d'Océanologie et de Géosciences, F-62930, Wimereux, France

² Ecole Supérieure du Professorat et de l'Éducation, Lille Nord de France, Communauté d'Universités et d'Établissements, F-59658, Villeneuve d'Ascq, France

³ IFREMER, LER/BL, 150 Quai Gambetta, F-62321, Boulogne-sur-Mer, France

⁴ IFREMER, DYNECO pelagos, Centre Bretagne, ZI de la Pointe du Diable, CS 10070, F-29280, Plouzané, France

* Corresponding author : A. Delegrange, email address : alice.delegrange@espe-lnf.fr

Abstract :

The diversity, toxicity and seasonality of *Pseudo-nitzschia* sp. were investigated from February to November 2012 in the southern Bight of the North Sea (SBNS) along the French coast. The identification of *Pseudo-nitzschia* species in this area was addressed for the first time in this study. Our results revealed a low species richness (3 distinct species) in association with moderate (102 pg mL⁻¹) to high (263 pg mL⁻¹) domoic acid (DA) levels in autumn and spring, respectively.

Pseudo-nitzschia succession corresponded to the dominance of *P. delicatissima* in April–May (86% of total diatoms) as a co-occurring species of the *Phaeocystis globosa* bloom. Following the *Phaeocystis* bloom (May–September), *P. pungens* dominated markedly over *P. fraudulenta* and *P. delicatissima* and was the only species present in autumn, although at low abundance (<1000 cell L⁻¹). The results of this study support the idea that *Pseudo-nitzschia* seasonality in the SBNS relies principally on temperature and nutrient availability (DIN and silicates), which, in turn, depend on locally fluctuating environmental conditions (rainfalls and winds). This study highlights the potential for the SBNS to be a potential risk area in regard to the possible impacts of DA on marine resources and the DA transfer through marine food webs. This is of particular concern since DA concentration in seawater was not systematically correlated to potentially toxic *Pseudo-nitzschia* abundance.

Highlights

► Low *Pseudo-nitzschia* sp. diversity (three species) in the southern North Sea. ► Domoic acid levels reached 263 pg mL⁻¹ in May 2012, 83% as dissolved DA. ► Strong species seasonality driven by temperature and nutrient availability (DIN and silicates). ► Autumnal DA concentrations could be attributed to *P. pungens*. ► DA concentration was not systematically correlated to *Pseudo-nitzschia* spp. abundance.

Keywords : *Pseudo-nitzschia*, Southern North Sea, Domoic acid, *Phaeocystis globosa*, Harmful algal blooms (HAB)

39 **1 Introduction**

40 The cosmopolitan diatom genus *Pseudo-nitzschia* sp. comprises more than 50 species
41 amongst which 24 are known to produce Domoic Acid (DA; Gai et al., 2018; Lundholm,
42 2018), a potent neurotoxin responsible for Amnesic Shellfish Poisoning (ASP). Since the first
43 reports of ASP in 1987 at Prince Edward Island, shellfish stocks have been systematically
44 monitored to prevent any outbreaks of ASP among humans. Since then, alert indicators such
45 as DA concentration and/or *Pseudo-nitzschia* sp. abundance monitoring have been shown to
46 be efficient tools, as no human poisoning has ever been reported (Trainer et al., 2012).
47 However, DA poisoning of marine mammals and seabirds can still occur when they feed on
48 DA-contaminated planktivorous prey (e.g. anchovies and sardines; Du et al., 2016; Gible et
49 al., 2018; Jensen et al., 2015; Lefebvre et al., 2002; Louw et al., 2018; Scholin et al., 2000;
50 Sierra-Beltran et al., 1997; Stauffer et al., 2012). The trend of increasing abundance of
51 *Pseudo-nitzschia* over the past decade has raised scientific awareness and concerns about
52 public health (e.g. Hernández-Fariñas et al., 2014; Lefebvre et al., 2014; Lundholm et al.,
53 2010; Parsons et al., 2002; Trainer et al., 2012) emphasizing the need to assess its dynamics,
54 and investigate potential DA production in exploited coastal areas. In the English Channel,
55 occurrence of DA in king scallop (*Pecten maximus*) has been reported several times from
56 south western regions such as the Bay of Seine and the Bay of Brest (Husson et al., 2016). In
57 the Eastern English Channel-North Sea (EEC-NS), reported ASP events were mainly
58 concentrated in the northern regions (Scotland, Norway, Denmark; Trainer et al., 2012)
59 although *Pseudo-nitzschia* sp. cells are present year-round in the southern regions (Bresnan et
60 al., 2015; Hernández-Fariñas et al., 2014; Schapira et al., 2008; Seuront et al., 2006). Thus,
61 reports of DA and shellfish closures were not recorded in the EEC-NS until 2014, when king
62 scallop stocks off the Bay of Somme (France) were contaminated with DA, which represented
63 the northernmost ASP alert of France (Lefebvre A., pers. com.). Existing studies and

64 monitoring surveys often limit the detection of *Pseudo-nitzschia* sp. to the genus level or to
65 morphological features measurable by optical microscopy, *i.e.* valve width, distinguishing
66 large species (width > 3 μm , *Pseudo-nitzschia* complex *seriata*), from narrow ones (width < 3
67 μm , *P. delicatissima* complex). This distinction by size categories has already made possible
68 seasonal studies within this genus (Bresnan et al., 2015; Brown and Bresnan, 2008; Fehling et
69 al., 2004; Thorel et al., 2017) as well as allowing for analysis of the short and long-term
70 changes in *Pseudo-nitzschia* with regard to environmental parameters (Díaz et al., 2014;
71 Husson et al., 2016; McKibben et al., 2015).

72 In the Eastern English Channel and Southern Bight of the North Sea, phytoplankton studies
73 have largely focused on spring bloom periods, revealing that diatoms and the
74 prymnesiophytes *Phaeocystis globosa* form the bulk of phytoplankton biomass (Bonato et al.,
75 2015; Schapira et al., 2008; Seuront et al., 2006). However, there is still a clear lack of
76 knowledge in this region regarding *Pseudo-nitzschia* species diversity and seasonality in
77 relation to environmental parameters and DA levels. Finally, studies defining *Pseudo-*
78 *nitzschia* species nutrient and physical requirements and biotic interactions at the regional
79 scale are scarce but crucial to predict proliferation and associated toxicity. Recently, Husson
80 et al. (2016) identified the environmental niche of *Pseudo-nitzschia* sp. in six distinct bays
81 from Brest to the Seine estuary. In this study, additional data focusing on *Pseudo-nitzschia*
82 seasonality and diversity in the EEC-NS ecosystem over an annual cycle is provided. The aim
83 is to define their importance in the seasonal phytoplankton succession and to estimate
84 associated DA concentrations with regard to environmental forcing and hydro-biological
85 conditions.

86 **2 Material and Methods**

87 **2.1 Sampling strategy**

88 The sampling station (~ 10 m depth) was located along the French coast of the southern North
89 Sea, in Dunkirk harbor (51°1'12'' N, 1°9'0''E, Figure 1). The inner part of this harbor forms
90 a small semi-enclosed embayment where semi-diurnal macrotidal tides prevail and enable
91 important water exchanges with coastal waters of the southern North Sea. Sampling was
92 conducted fortnightly from February 22 to November 14 2012 during flood (within 2 hours
93 before high tide) to consider only neritic plankton communities although benthic communities
94 could be re-suspended during high mixing periods.

95 Temperature (°C) and salinity were measured with an Aanderaa Instruments probe, turbidity
96 (Turb) with an Eutech instruments waterproof probe, pH with a HANNA pH probe, and
97 dissolved oxygen (dO₂, mg L⁻¹) with a Handy Polaris Oxyguard probe at each sampling date.
98 Seawater samples were collected at one meter depth using a Niskin bottle. For inorganic
99 nutrients, seawater samples (frozen at -20 °C until analysis) were analyzed by either
100 fluorimetry (NH₄⁺; Trilogy, Turner Designs; Holmes et al., 1999) or by the use of an
101 autoanalyzer (Alliance Integral Futura) for NO₂⁻, NO₃⁻, HPO₄²⁻, Si(OH)₄ following standard
102 protocols (Bendschneider and Robinson, 1952; Mullin and Riley, 1955; Murphy and Riley,
103 1962). The surface solar irradiance (ssi) was derived from the bi-directional reflectance
104 measured by MSG (METEOSAT Second Generation) and provided by the OSI SAF (Le
105 Borgne et al., 2006). Winds (hourly measurements) and rainfall (daily) data were obtained
106 from the Dunkirk Meteo France station (www.meteofrance.fr).

107 2.2 Phytoplankton standing stock and community composition

108 Phytoplankton biomass was assessed from chlorophyll *a* and pheopigment concentrations.
109 Seawater samples (250 to 500 mL) were filtered on glass fiber filters (Whatman GF/F) under
110 low vacuum and frozen at -20 °C until analysis (within 2 months after collection). Pigments
111 were extracted in 90 % acetone overnight at 4°C and chlorophyll *a*, and pheopigments
112 concentrations were estimated following Lorenzen (1966) using a pre-calibrated (chlorophyll

113 *a* from *Anacystis nidulans*, Sigma) fluorometer (Trilogy, Turner designs). For phytoplankton
114 community analyses, 250 mL samples were preserved in the field with lugol/glutaraldehyde
115 fixative (2 % final concentration, Verity et al., 2007) and stored at 4 °C in the dark until
116 analysis (within 3 months after collection). At the laboratory, 5 to 10 mL subsamples were
117 settled in Hydrobios counting chambers. Phytoplankton cells were identified and enumerated
118 using an inverted microscope (Nikon Eclipse TE2000-S, magnification ×200, ×400) under
119 phase contrast illumination. On average 942 ± 568 cells per sample were identified and
120 counted. For *Phaeocystis globosa*, total cells were enumerated without stage distinction (i.e.
121 isolated, colonial stages). Under optical microscopy, *Pseudo-nitzschia* cells were categorized
122 on the basis of cell width. More specifically, *Pseudo-nitzschia* cells with frustule width up to
123 $3\mu\text{m}$ were placed in the *P. delicatissima* complex while those with frustules above $3\mu\text{m}$ were
124 placed in the *P. seriata* complex as was done in several studies (Bresnan et al., 2015; Fehling
125 et al., 2004; Hasle and Syvertsen, 1997; Thorel et al., 2017).

126 When *Pseudo-nitzschia* sp. cell concentrations were higher than 10^3 cell L^{-1} , Scanning
127 Electron Microscopic (SEM) analyses were carried out for species identification, covering the
128 whole study period (April 4, May 2, May 16, June 1, September 5, October 3). In short, 10 to
129 100 mL of homogenized preserved samples were filtered on polycarbonate filters (24 mm
130 diam.; $0.8 \mu\text{m}$ porosity) and rinsed with Milli-Q water to eliminate salts. Filters containing
131 diatom frustules were acid cleaned (2 - 3 mL HCl 37%) in Teflon beakers overnight at room
132 temperature and rinsed in a Milli-Q water bath. Acid and water baths were filtered once again
133 using same filter to limit cell loss during the acidification and rinsing steps. Thereafter, filters
134 were placed on aluminium stubs with double sticky carbon tabs and air-dried during 12 h
135 under laminar flow. Lastly, dried filters were metallized under argon flow with Au/Pd during
136 90 s (Polaron SC 7620). Quantified measures based on literature review (e.g., valve length
137 and width, number of fibulae and of interstriae in $10 \mu\text{m}$) allowed for the discrimination of the

138 species within *Pseudo-nitzschia seriata* complex (width > 3 μm) and *P. delicatissima*
139 complex (width < 3 μm ; see Table 1), but species variety could not be inferred from our
140 study. At least 50 *Pseudo-nitzschia* sp. cells per sample were identified via SEM analyses.
141 When present in lower numbers, the entire filter was scanned for *Pseudo-nitzschia* frustules
142 and identification was carried out on available material. Dissolved and particulate domoic
143 acid (DA; pg mL^{-1}) concentrations were measured using ASP ELISA kits (Biosense, Bergen,
144 Norway) following the provided procedure. Total DA concentration (tDA *i.e.*, dissolved +
145 particulate DA) was measured from 5 mL of natural seawater samples stored at -20 $^{\circ}\text{C}$ until
146 analysis. For dissolved DA, collected seawater samples (5 mL) were filtered under low
147 vacuum pressure (< 50 mm Hg) on 0.2 μm cellulose acetate filters before being frozen (-20
148 $^{\circ}\text{C}$). On the day of the analysis, samples were thawed at 4 $^{\circ}\text{C}$ and dissolved DA samples were
149 directly processed. For total DA samples, sonication on ice was carried out in order to disrupt
150 cells, and samples were filtered on 0.2 μm cellulose acetate before the assay. Particulate DA
151 was obtained by subtracting dissolved DA from total DA. Given the limit of *Pseudo-nitzschia*
152 cell counts via microscopy, DA cell quota were computed only when *Pseudo-nitzschia* cell
153 abundance reached 10^3 cell L^{-1} (*i.e.* at least 5-10 *Pseudo-nitzschia* cells were counted).

154 2.3 Data analyses

155 In order to efficiently synthesize the environmental context of the survey and identify forcing
156 parameters, a Principal Component Analysis (PCA) was applied to the environmental data
157 matrix (11 parameters x 16 dates, Legendre and Legendre, 1998). Wind speed data from three
158 days before the sampling date were averaged. Rainfalls were summed over 3 days before the
159 sampling date in order to take into account watershed run-offs.

160 Nutrients concentrations of DIN (*i.e.* the sum of NH_4^+ , NO_2^- and NO_3^- concentrations), HPO_4^{2-}
161 and Si(OH)_4 concentrations were depicted in the PCA plots as N, P and Si, respectively.

162 Nutrient ratios (S/P, Si/N and Si/P) were considered as illustrative variables to highlight

163 potential nutrient limitations. Prior to the analysis, all data were tested for normality (Shapiro
164 test) and $\log(x+1)$ transformed.

165 *Pseudo-nitzschia* complex raw abundances were plotted on each PCA graph to give a picture
166 of species seasonality. Only variables significantly correlated to at least one principal
167 component (PC) of the PCA are presented and discussed (Spearman test, p-value <0.05).
168 Correlation coefficients will be referred to as r'_{PC1} , r'_{PC2} , r'_{PC3} for the first three principal
169 components of the PCA (Table 2). Additionally, correlations between *Pseudo-nitzschia*
170 complexes abundances and domoic acid concentrations have been assessed. Since data did not
171 fit the normality, correlations were tested using Spearman test (significance threshold was set
172 at p-value<0.05). Statistics were performed using R software packages (R Core Team, 2012)

173 **3 Results**

174 3.1 Environmental conditions

175 The present study began in the cold snap of February 2012 (temperature of minus 10°C) that
176 had not occurred in the area since 1991 (MeteoFrance data; regional bulletin,
177 www.meteofrance.fr). The following spring was cooler (-1.0 °C) and drier (half as much
178 rainfall) than average (Figure 2A, 2C). Stormy weather dominated the summer months (June-
179 July) and autumn. Autumn 2012 was characterized by strong southwestern and northeastern
180 winds (up to 100 km h⁻¹), excess precipitations (+30%) and a rapid drop in temperature.
181 Seawater surface temperature followed a typical seasonal evolution increasing from 4.9 °C in
182 February to 19.0 °C in August. By contrast, as measured by flood tide, salinity remained
183 stable with an average of 34.2 ± 0.3 (min: 33.4, max = 35.0; Figure 2A). The mean annual pH
184 value was 8.1 ± 0.3 and decreased continuously over the study period from February (8.6) to
185 November (7.6; Figure 2B). The mean annual dissolved oxygen concentration (dO₂) was $8.1 \pm$
186 1.3 mg L^{-1} with the highest values recorded in winter and late autumn, and the lowest (6.4 mg
187 L^{-1}) in late summer (Figure 2B). A slight increase in dO₂ concentrations coincided with the

188 spring bloom development in May. Turbidity averaged at 6.9 ± 3.2 NTU (Figure 2C). The
189 lowest turbidity values were measured in April and June (2.9 NTU) and the highest in May
190 and September (14.3 and 12.1 NTU, respectively).
191 With the exception of NH_4^+ , all concentrations of nutrients decreased rapidly to reach their
192 annual minima of 0.01, 0.35 and $0.30 \mu\text{mol L}^{-1}$ for $\text{NO}_2^- + \text{NO}_3^-$, HPO_4^{2-} and $\text{Si}(\text{OH})_4$,
193 respectively, in April (Figure 3). Thereafter, HPO_4^{2-} concentration varied between 0.5 and 1.0
194 $\mu\text{mol L}^{-1}$ and the maximal values followed storm events (July and November). $\text{NO}_2^- + \text{NO}_3^-$
195 concentrations remained stable below $5 \mu\text{mol L}^{-1}$ from April to October and increased to high
196 values at the end of the survey ($11.7 \mu\text{mol L}^{-1}$) although they did not reach the previous winter
197 maxima of $22.5 \mu\text{mol L}^{-1}$. Silicate concentrations decreased from 7.5 to 0.3 between February
198 and late March. A second minimal value ($0.2 \mu\text{mol L}^{-1}$) was recorded in June. The silicate
199 concentrations increased continuously during summer and reached $7.5 \mu\text{mol L}^{-1}$ at the end of
200 the survey. NH_4^+ concentrations were highly variable from February to June ranging between
201 1.0 and $3.8 \mu\text{mol L}^{-1}$. Thereafter, concentrations remained between 2.0 to $4.0 \mu\text{mol L}^{-1}$.

202 3.2 Phytoplankton standing stock, seasonality and domoic acid levels

203 The phytoplankton bloom, started at the end of February 2012 and lasted until early June. It
204 was characterized by three successive peaks in chlorophyll *a* concentrations (from 4.0 to 9.5
205 $\mu\text{g chl}a \text{ L}^{-1}$; Figure 4A), and phytoplankton abundance (from 8.2×10^5 to 3.6×10^6 cells L^{-1} ;
206 Figure 4B). From July to November, chlorophyll *a* concentrations ($2.3 \pm 0.7 \mu\text{g L}^{-1}$) along
207 with phytoplankton abundance ($1.9 \pm 0.7 \times 10^5$ cells L^{-1}) remained relatively stable and low.
208 Pheopigment concentrations followed the same pattern as chlorophyll *a* (Figure 5A) and
209 represented 22 to 49 % of total pigments. Four periods could be identified based on
210 phytoplankton community composition (Figure 4C): Period 1 (P1, February 20 – April 5) was
211 characterized by the dominance of small diatoms (5-20 μm in width/length; *Thalassiosira* sp.,
212 *Skeletonema costatum*, *Asterionellopsis glacialis* and *Chaetoceros* sp.), either as isolated cells

213 or in forming colonies, and phytoplankton abundance peaked at 8.2×10^5 cells L⁻¹. Period 2
214 (P2, April 5 to May 20) corresponded to the *P. globosa* bloom with abundances ranging from
215 1.8 to 3.6×10^6 cells L⁻¹ (Figure 4B). This bloom co-occurred with high abundances of
216 *Chaetoceros* sp. (8.7×10^4 cells L⁻¹, 64 % of the diatom community at the beginning of P2),
217 and *Pseudo-nitzschia* sp. (2.1×10^5 cells L⁻¹ equivalent to 25% and 86% of the phytoplankton
218 and the diatom communities, respectively; at the end of P2). Period 3 is the post-bloom period
219 (P3, from May 20 to September 8). Both fine walled summer diatoms such as *Guinardia* sp.
220 (*i.e.* *G. delicatula*, *G. flaccida*, and *G. striata*), and highly silicified spring diatoms such as
221 *Chaetoceros* sp. and *Thalassiosira* sp. composed the bulk of the diatom community in
222 association with small Cryptophytes. Period 4 (P4, from September 8 to November 16)
223 revealed the occurrence of intermediate sized diatoms (30 - 40 μ m in length/width; *Delphineis*
224 sp.) as well as diatoms in forming colonies (e.g., *Thalassiosira* sp.) which remained in low
225 number ($1.2 \pm 0.5 \times 10^5$ cells L⁻¹).

226 SEM analyses revealed a low diversity of *Pseudo-nitzschia* sp. in the area with three distinct
227 species namely *P. delicatissima* (belonging to the *P. delicatissima* complex), *P. fraudulenta*,
228 and *P. pungens* (both belonging to the *P. seriata* complex, see Table 1 and Figure 5). No
229 overlap between the two complexes was found in terms of species morphological features
230 since the width of *P. delicatissima* width was below 2 μ m, and the widths of *P. pungens* and
231 *P. seriata* were above 2.7 μ m (Table 1). *P. delicatissima* represented up to 99 % of SEM-
232 identified frustules during the *P. globosa* bloom period (P2, Mid-May) and outnumbered
233 other diatoms. By contrast, at the beginning of the post-bloom period (P3), the percentage of
234 *Pseudo-nitzschia* sp. to total diatoms decreased to 10 % of total diatoms. *P. pungens* markedly
235 dominated (54 % of the SEM-identified frustules) over *P. fraudulenta* and *P. delicatissima*
236 (33 % and 13 %, respectively). In autumn (P4), all the identified *Pseudo-nitzschia* sp.

237 frustules belonged to the *P. pungens* species ($< 10^3$ cells L⁻¹ representing $< 1\%$ of total
238 diatoms).
239 Domoic acid was mostly present as dissolved DA throughout the survey averaging $83 \pm 16 \%$
240 of the total DA (tDA). Particulate domoic acid (pDA) was detected in May (end of P2) and
241 was still measurable at the end of P4. pDA punctually represented 10 to 40 % of tDA (Figure
242 6). tDA concentration followed *Pseudo-nitzschia* sp. dynamics, particularly during the *P.*
243 *globosa* bloom and post-bloom periods (P2 and P3, Figure 7). tDA maxima (263 pg mL^{-1})
244 corresponded to the co-occurrence of *P. seriata* complex and *P. delicatissima* complex,
245 reaching 2.8×10^4 and 7.5×10^5 cells L⁻¹ in early May (P2), respectively. Thereafter, tDA
246 decreased to 118 and 110 pg mL^{-1} at the end of P2 and beginning of P3, which coincided with
247 the highest abundance of *P. delicatissima* complex and *P. seriata* complex, respectively. In
248 autumn, tDA concentrations remained high ($102 \pm 39 \text{ pg mL}^{-1}$) despite low abundances of *P.*
249 *seriata* complex (*i.e.* *P. pungens*), as suggested by the absence of correlation between tDA
250 and *Pseudo-nitzschia* abundances. When measurable, DA cell quota ranged between 0.016 pg
251 cell^{-1} in May and 7.9 pg cell^{-1} in September.

252 3.3 Abundance of *Pseudo-nitzschia* in relation to environmental factors

253 PCA confirmed the clear seasonal pattern of environmental variables and allowed to
254 synthesize their relation with the temporal variability of *Pseudo-nitzschia* (Figure 7). The first
255 three principal components (PC) explained 75.82 % of the total variance (PC1: 35.01 %, PC2:
256 28.77 %, PC3: 12.04 %). Considering coefficients in the linear combination of variables,
257 period P1 was mainly characterized by high DIN ($r'_{\text{PC1}} = 0.88$) and dO₂ ($r'_{\text{PC2}} = -0.75$)
258 concentrations, high N/P ratios ($r'_{\text{PC1}} = 0.66$; $r'_{\text{PC2}} = -0.50$) and low temperature and salinity
259 ($r'_{\text{PC2}} = 0.73$ and 0.64, respectively; Table 2). P2 (bloom period) was more related to high ssi
260 ($r'_{\text{PC1}} = -0.74$), pH ($r'_{\text{PC2}} = -0.64$), and increases in temperature and salinity ($r'_{\text{PC2}} = 0.73$ and
261 0.64), low nutrient stocks, particularly Si(OH)₄ and HPO₄²⁻ ($r'_{\text{PC1}} = 0.69$ and 0.87), and to a

262 lesser extent low Si/P ratio ($r'_{PC2}=0.44$, Table 2). Additionally, punctual storm events in May
263 characterized by high rainfalls ($r'_{PC3}=0.61$) and by strong winds ($r'_{PC3}=0.43$) led to a prompt
264 increase in turbidity (Figures 3 and 7B). High temperature, DIN limitations illustrated by low
265 N/P, and increasing Si/P ratio and Si(OH)_4 concentrations characterized periods P3 and P4
266 (Table 2).

267 The plotting of *Pseudo-nitzschia* abundance on the PCA graphs illustrated the importance of
268 high rainfall, increasing pH, ssi in the initiation of the *P. delicatissima* complex bloom
269 (Figures 7A and 7B). The seasonality of the *P. seriata* complex was essentially shown by
270 their high abundance in June (beginning of P3) and their disappearance in August and
271 September (end of P3; Figures 7C and 7D). The highest abundance of *P. seriata* complex was
272 observed at low nutrient levels (particularly DIN and Si(OH)_4), increasing surface solar
273 irradiance and high pH. From September to November 2012 (P4), rainfalls and wind speed
274 decreased, and *P. seriata* complex persisted at low abundance in increasing concentrations of
275 Si(OH)_4 , decreasing pH and concentrations of dO_2 . Finally, as illustrated by the distribution of
276 abundance values on the PCA plots, both complexes were present under Si(OH)_4 limitation
277 and the *P. seriata* complex seemed to tolerate higher variable environmental conditions (e.g.
278 nutrients) as compared to the *P. delicatissima* complex.

279 **4 Discussion**

280 4.1 Specificity of the 2012 spring bloom

281 In the southern North Sea - Eastern English Channel ecosystem, the phenology of
282 phytoplankton communities is generally characterized by a phytoplankton spring bloom
283 dominated by *Phaeocystis globosa* (Gentilhomme and Lizon, 1998; Lefebvre et al., 2011;
284 Schapira et al., 2008). In 2012, the spring bloom appeared to be of low magnitude and short
285 duration compared to other studies ($\sim 20 \mu\text{g L}^{-1}$ in Breton et al., 2000 and Kesaulya et al.,
286 2008 ; $\sim 50 \mu\text{g L}^{-1}$ in Lefebvre et al., 2011; Seuront et al., 2006), as exemplified by low

287 chlorophyll *a* values ($9.5 \mu\text{g L}^{-1}$, Figure 4A). This small bloom magnitude was confirmed by
288 the low *P. globosa* abundance (between 0.019 and $3.7 \times 10^6 \text{ cell L}^{-1}$ in this study vs 5.5 to
289 $100 \times 10^6 \text{ cell L}^{-1}$ in Gómez and Souissi, 2008; Seuront et al., 2006; Widdicombe et al., 2010)
290 and its low contribution to total phytoplankton abundance (50% vs $74\text{-}90\%$ between 1992 and
291 2011, Hernández-Fariñas et al., 2014). *P. globosa* dynamics can exhibit high interannual
292 variability (Lamy et al., 2006; Lefebvre et al., 2011) and the year 2012 appeared as a low
293 *Phaeocystis* year. The co-occurrence of the *P. delicatissima* complex and *P. globosa*
294 development (Figure 4; Period P2) is often described: *P. delicatissima* complex is a co-
295 occurring species of *P. globosa* bloom (Grattepanche et al., 2011; Hernández-Fariñas et al.,
296 2014; Monchy et al., 2012; Sazhin et al., 2007) and may serve as solid substrate during the
297 transitional phase from free-living cells to colonial stages (Peperzak and Gäbler-Schwarz,
298 2012; Rousseau et al., 2007; Verity et al., 2007).

299 4.2 Environmental factors drive *Pseudo-nitzschia* diversity and seasonality.

300 Three species of *Pseudo-nitzschia* were identified in 2012 in the Southern North Sea (*P.*
301 *delicatissima*, *P. pungens* and *P. fraudulenta*) by using SEM imagery and measurements.
302 These results are congruent with recent published data from adjacent areas (Eastern English
303 Channel - Thorel et al., 2017, Northern North Sea - Bresnan et al., 2015). Identification was
304 based on a literature review and best matches with species morphometric issues are specified
305 in Table 1. The identification of *Pseudo-nitzschia* species is a tricky issue since some species
306 can have very similar morphological characteristics (e.g. *P. delicatissima* with *P. dolorosa*, *P.*
307 *decipiens* or *P. fraudulenta* with *P. subfraudulenta*, Amato et al., 2007; Orsini et al., 2004;
308 Quijano-Scheggia et al., 2009). Further studies are therefore needed to confirm these
309 identifications using molecular tools (e.g. Pugliese et al. 2017).
310 The seasonality of *Pseudo-nitzschia* species within the Eastern English Channel – North Sea
311 (EEC-NS) ecosystem resulted in two distinct periods of abundance (spring and autumn)

312 during which a very small number of *Pseudo-nitzschia* species was observed. The three
313 identified species (*P. delicatissima*, *P. pungens*, and *P. fraudulenta*) are considered to be
314 cosmopolites (see Table 3; Hasle 2002 and Lelong et al. 2012 for reviews). They are also
315 known to have different but variable hydrological preferences depending on their ecosystem
316 inhabited, and genetic strains. Temperature and salinity values recorded in 2012 (4.9-19.0°C
317 and 33.6-35.0, respectively; Figure 2A) are consistent with the euryhaline features of *Pseudo-*
318 *nitzschia* spp. Most species are indeed able to grow at salinities comprised between 1 and >
319 30, occurring and growing more efficiently at high salinity (Cho et al., 2001; Husson et al.,
320 2016; Thessen et al., 2005). Prior studies have also demonstrated a wide temperature range for
321 *Pseudo-nitzschia* spp. (Dortch et al., 1997; Louw et al., 2016; Tas and Lundholm, 2016). In
322 particular, *P. delicatissima*, which was dominant in May (P2, 86% of the diatom community
323 and 99 % of identified frustules) seemed to better grow at temperatures ranging between 10-
324 12 °C (Table 3; Fehling et al. 2006). *P. pungens* and *P. fraudulenta* were commonly present
325 in summer (June) and autumn (October) when days were longer and temperatures higher
326 (Table 3). However, *P. pungens* seems to have a wider environmental tolerance than *P.*
327 *fraudulenta* (Klein et al., 2010) as it can peak at temperatures up to 21.0 °C (Terenko and
328 Terenko, 2012). The increasing relative contribution of *P. pungens* to *Pseudo-nitzschia*
329 species between June (54 %) and October (100 %) may be explained by two non-conflicting
330 hypotheses. Firstly, the *P. pungens* population may comprise different varieties exhibiting
331 different temperature preferences (e.g. Kim et al. 2015). Secondly, it can present varying
332 responses to temperature increases (Thessen et al., 2009).

333 *P. delicatissima* development began in February (P1) under high DIN concentrations, high
334 N/P ratios and low temperature (i.e. late winter conditions; Figure 7A, 7B). It is well known
335 that high abundances of *Pseudo-nitzschia* spp. are found in areas enriched with nutrients
336 (Parsons et al., 2002; Tas et al., 2016) which can originate from upwelling (Díaz et al., 2014),

337 mixing events and riverine inputs (Trainer et al., 2012). Paradoxically, the highest abundance
338 of *P. delicatissima* in May (P2) occurred after a decrease in DIN concentrations and rather
339 suggested nutrient limiting conditions. Pulses of DIN (Figure 3A) following rainfall events in
340 May (Figure 2C) could have favored the species rapid growth and bloom, a phenomenon
341 which has already been shown in estuarine systems (Tas et al., 2016; Tas and Lundholm,
342 2016). The results of this study also demonstrated that the maximum abundance of the *P.*
343 *delicatissima* complex occurred at high pH levels and dO₂ concentrations (P2, Figure 7).
344 Lundholm et al. (2004) suggested that high pH values (8.7 - 9.1) can inhibit the growth of
345 most *Pseudo-nitzschia* species. Such high values were never recorded during our study (7.6 to
346 8.6; Figure 3) and no significant relationships could be established between pH values and
347 *Pseudo-nitzschia* abundance. Rather than being associated solely with the *P. delicatissima*
348 complex, variability in dO₂ is likely associated with the photosynthetic activity of the whole
349 phytoplankton community including the *Phaeocystis* sp. bloom (P2).
350 High amount of particulate matter (i.e., high turbidity) linked to water mixing and river inputs
351 can limit light availability and consequently inhibit photosynthesis and growth. This light
352 attenuation can result in the limitation of both the spatial and temporal distribution of *Pseudo-*
353 *nitzschia* as demonstrated in estuarine and coastal systems (Dursun et al., 2016; Tas et al.,
354 2016). These studies are consistent with the *P. seriata* complex distribution pattern in the
355 Southern Bight of the North Sea (SBNS), the species being abundant when turbidity was low
356 (e.g. in June, Figure 2C, 7D). *P. delicatissima* could have benefited from the storm event
357 (high wind and rainfall) which occurred in May and particularly from the subsequent pulses of
358 DIN and Si(OH)₄ (Figures 3A and 7B).
359 The *P. seriata* species complex was the most abundant (beginning of P3) in low DIN and
360 SiOH₄ concentrations resulting in low Si/N ratios (figure 7C). These conditions may have
361 been caused by the lag effect of *P. delicatissima* and the associated *P. globosa* bloom having

362 already drawn down the nutrients. The *P. seriata* species complex was still persisting at low
363 abundance in the area from the end of the summer to the fall (end of P3 and P4). PCA plots
364 suggest that the growth of the *P. seriata* complex may benefit from nutrient pulses of silicate,
365 phosphate and DIN but that it tends to be limited at that time of the year by decreasing
366 temperature and irradiance (Figure 7C, 7D).

367 Silicate limitations were already proven to increase *Pseudo-nitzschia* abundance and toxicity
368 (Anderson et al., 2010; Thorel et al., 2017). For instance, *P. multiseriata* was able to
369 outcompete other phytoplankton species when silicate was low compared to nitrogen
370 (Sommer, 1994). Laboratory culture conditions conducive to DA production generally include
371 limitation by silicate or phosphate during the stationary phase (reviewed by Bates et al., 1989;
372 Trainer et al., 2012, 2008). This could explain this study findings of the high domoic acid
373 concentration in spring (P2, May, Figure 6) which coincided with low Si and DIN
374 concentrations along with low Si/P ratios (Figure 7A- C). However, no significant correlation
375 could be demonstrated between nutrients stocks and DA concentrations. Consequently, more
376 targeted studies (e.g., nutrient uptake experiments) are required in order to properly evaluate
377 the nitrogen nutritional status and nitrogen preference (i.e., $\text{NO}_2^- + \text{NO}_3^-$ vs NH_4^+ vs urea) for
378 *Pseudo-nitzschia* populations, and thus the potential influence of nitrogen on bloom dynamics
379 and toxicity in the EEC-NS.

380 4.3 Domoic acid concentrations and *Pseudo-nitzschia* species

381 The existence of toxigenic and non-toxigenic strains of the same species has been mentioned
382 recurrently for several *Pseudo-nitzschia* species (e.g. Sahraoui et al. 2011; Thessen et al.
383 2009; Villac et al. 1993) Sahraoui et al., 2011; Thessen et al., 2009; Villac et al., 1993). Total
384 DA concentrations measured during this study (max. 263 pg mL^{-1}) were higher than several
385 published values in the EEC-NS ecosystem (up to 7 pg mL^{-1} ; Downes-Tettmar et al. 2013;
386 Klein et al. 2010) but remained lower than recent measurements in the Bay of Seine (1.4×10^3

387 pg mL⁻¹, Thorel et al. 2017), and other ecosystems such as Monterey Bay (10×10³ pg mL⁻¹;
388 Walz et al. 1994), the Gulf of Mexico (3.3×10³ pg mL⁻¹, Liefer et al. 2013) or Luanda Bay,
389 Angola (1.4×10³ pg mL⁻¹, Blanco et al. 2010).

390 Although *Pseudo-nitzschia* abundance matched high DA concentration in spring, this
391 observation did not hold for the end of summer and autumn periods (Figure 6). Other species
392 (i.e. *Nitzschia navis-vargnica* and *N. bizertensis*) are known to produce DA in south-east Asia
393 and Tunisia, respectively (Bouchouicha-Smida et al., 2015; Kotaki et al., 2000; Lundholm and
394 Moestrup, 2000; Tan et al., 2016). Since neither species has been detected in North Atlantic
395 waters (including the English Channel and the North Sea) or been observed during this 2012
396 survey, it is likely that DA originate exclusively from *Pseudo-nitzschia* species.

397 Consequently, this result supports other studies where DA concentration in seawater
398 (particulate and dissolved) or in shellfish does not always depend on *Pseudo-nitzschia*
399 abundance (e.g. Bates et al. 1998; Bresnan et al. 2017; Klein et al. 2010; Lelong et al. 2012;
400 Walz et al. 1994). During P2 and P3, high tDA concentrations coincided with the co-
401 occurrence of two or three *Pseudo-nitzschia* species (i.e., *P. delicatissima* and *P. pungens*
402 during P2 and *P. delicatissima*, *P. pungens* and *P. fraudulenta* during P3). Since these three
403 species have been proven at least once to produce DA (Lelong et al., 2012), the formal
404 identification of the species responsible for DA concentration cannot be addressed with
405 certainty. Consistent with Downes-Tettmar et al. (2013), *P. pungens* was certainly toxic as it
406 was the only *Pseudo-nitzschia* sp. present in autumn when DA concentrations reached 101 pg
407 mL⁻¹. *P. delicatissima* and *P. fraudulenta* were demonstrated to be weakly toxigenic in
408 different ecosystems (Almandoz et al., 2017; Parsons et al., 2012; Rhodes et al., 1998).

409 Additionally, in the northern North Sea, *P. delicatissima* cells have not been observed to
410 produce DA (Scottish Waters, (Fehling et al., 2006, 2004). The DA cell quotas estimated for
411 the three samples in which particulate DA was associated with *Pseudo-nitzschia* abundance >

412 10^3 cell L^{-1} ranged between 0.016 and 7.9 pg. cell⁻¹. These DA cell quotas are comparable to
413 values obtained for batch cultures of *P. fraudulenta* (0.03 pg.cell⁻¹, Rhodes et al. 1998; 0.16
414 pg.cell⁻¹, Thessen et al. 2009) and *P. pungens* (0.2 to 10 pg. cell⁻¹; Lelong et al. 2012). They
415 are also congruent with DA cell quotas inferred from *in situ* surveys for *P. fraudulenta* (0.002
416 to 0.120 pg. cell⁻¹; Almandoz et al. 2017) and *P. delicatissima* (1.52 to 5.54 pg. cell⁻¹, Parsons
417 et al. 2012).

418 4.4 DA implications in the ecosystem and challenges for future surveys

419 In this study, the high proportion of dissolved domoic acid year round (~83%) may have two
420 main implications regarding phytoplankton communities succession and marine resources.

421 Dissolved domoic acid can indeed act as an effective Fe chelator, increasing its bioavailability
422 and providing a competitive advantage to *Pseudo-nitzschia* sp. over other phytoplankton
423 species for Fe assimilation (Prince et al., 2013; Rue and Bruland, 2001; Trick et al., 2010;
424 Zabaglo et al., 2016). Other studies suggest that DA synthesis requires available Fe (e.g.
425 Sobrinho et al. 2017). In both cases, accumulated iron in the polysaccharidic matrix of *P.*
426 *globosa* (Schoemann et al., 2005) could have benefitted *Pseudo-nitzschia* sp. growth and DA
427 production and could well explain its appearance every year, during and after the *P. globosa*
428 bloom.

429 Regarding marine resources, dissolved DA is reported to alter bivalve larvae growth (Liu et
430 al., 2007) and immunity (De Rijcke et al., 2015) and these effects could spread to population
431 recruitment and picking activities. Dissolved DA is also reported to inhibit herbivorous
432 feeding as shown for krill (Bargu et al., 2006) and copepod species (Tammilehto et al., 2015)
433 modifying matter and energy fluxes within the trophic web. These elements suggest the
434 potential for dissolved DA to impact marine resources throughout the year and to trigger long-
435 term effects regarding growth and recruitment.

436 Despite its absence in spring, particulate DA appeared at significant level (10-40%) over a
437 wide temporal window (May to December). During this period, *Pseudo-nitzschia* sp. could
438 provide a non-negligible food source from which particulate DA could be directly transferred
439 to higher trophic levels such as exploited filter feeders namely mussels or king scallops in the
440 EEC-NS (Ambiaud and Lefebvre, 2017). Phytoplankton consumers such as *Noctiluca*
441 *scintillans* and copepods could act as DA vectors/inducers (Escalera et al., 2007; Fock and
442 Greve, 2002; Leandro et al., 2010; Maneiro et al., 2005; Tammilehto et al., 2015) and were
443 shown to dominate the zooplankton community when particulate DA reached 40% of total
444 DA in the summer. Our data suggest that the SBNS could be an at-risk area regarding DA
445 contamination and/or exposure of marine resources to dissolved DA: the phytoplankton
446 abundance alert threshold was reached at least once for the *P. delicatissima* complex during
447 the 2012 survey (i.e. $>300\,000$ cells L^{-1} , (EC) No 853/2004 2004; (EC) No 854/2004 2004)
448 concomitantly with significant DA concentrations in seawater. However, high domoic acid
449 concentration was not always associated with high *Pseudo-nitzschia* abundance. This study at
450 the seasonal scale allowed the identification of risky periods regarding species diversity and
451 DA forms and concentrations in the SBNS ecosystem. Given the inter-annual variability of
452 *Pseudo-nitzschia* species occurrence (e.g. Thorel et al. 2017), long term monitoring is
453 fundamental to assess whether the species seasonality persist over years and whether it is
454 always associated to the same toxicogenic species and environmental parameters.

455 **Acknowledgments**

456 We acknowledge J. Mestre, B. Gomez and T. Raud for their technical support in the field and
457 at the laboratory. A PhD grant from the région Nord Pas-de-Calais and the Syndicat Mixte de
458 la Côte d'Opale (SMCO) attributed to A. Delegrange supported this work. This work was also
459 funded by a collaborative convention between former Aquanord S.A. and ULCO/LOG. We
460 thank Dr. H el ene H egaret and Dr. Philippe Soudant (IUEM - LEMAR) for relevant comments

461 and constructive suggestions on a former version of this manuscript. We also thank two
462 anonymous reviewers for their comments and suggestions improving our data presentation
463 and species description. Thanks are also due to Jennifer Morival
464 (jennifermorival.wixsite.com) for the complete English edition and correction of the
465 manuscript.

466 **References**

- 467 Almandoz, G.O., Fabro, E., Ferrario, M., Tillmann, U., Cembella, A., Krock, B., 2017. Species
468 occurrence of the potentially toxigenic diatom genus *Pseudo-nitzschia* and the associated
469 neurotoxin domoic acid in the Argentine Sea. *Harmful Algae* 63, 45–55.
470 <https://doi.org/10.1016/j.hal.2017.01.007>
- 471 Amato, A., Kooistra, W.H.C.F., Levialedi Ghiron, J.H., Mann, D.G., Pröschold, T., Montresor, M., 2007.
472 Reproductive Isolation among Sympatric Cryptic Species in Marine Diatoms. *Protist* 158, 193–
473 207. <https://doi.org/10.1016/j.protis.2006.10.001>
- 474 Anderson, C.R., Sapiano, M.R.P., Prasad, M.B.K., Long, W., Tango, P.J., Brown, C.W., Murtugudde, R.,
475 2010. Predicting potentially toxigenic *Pseudo-nitzschia* blooms in the Chesapeake Bay. *J. Mar.*
476 *Syst., GEOHAB Modeling* 83, 127–140. <https://doi.org/10.1016/j.jmarsys.2010.04.003>
- 477 Bargu, S., Lefebvre, K., Silver, M.W., 2006. Effect of dissolved domoic acid on the grazing rate of krill
478 *Euphausia pacifica*. *Mar. Ecol. Prog. Ser.* 312, 169–175. <https://doi.org/10.2307/24870068>
- 479 Bates, S.S., Bird, C.J., de Freitas, A.S.W., Foxall, R., Gilgan, M., Hanic, L.A., Johnson, G.R., McCulloch,
480 A.W., Odense, P., Pocklington, R., Quilliam, M.A., Sim, P.G., Smith, J.C., Subba Rao, D.V.,
481 Todd, E.C.D., Walter, J.A., Wright, J.L.C., 1989. Pennate diatom *Nitzschia pungens* as the
482 primary source of domoic acid, a toxin in shellfish from eastern Prince Edward Island,
483 Canada. *Can Fish Aquat Sci* 1203–1215.
- 484 Bates, S.S., Garrison, D.L., Horner, R.A., 1998. Bloom Dynamics and Physiology of Domoic-Acid-
485 Producing *Pseudo-nitzschia* Species, in: *Physiological Ecology of Harmful Algal Blooms*.
486 Anderson, D.M., Cembella A.D., and Hallegraeff G.M., Heidelberg, pp. 267–292.
- 487 Bendschneider, K., Robinson, R.J., 1952. A New Spectrophotometric Method for the Determination of
488 Nitrite in Sea Water.
- 489 Blanco, J., Livramento, F., Rangel, I.M., 2010. Amnesic shellfish poisoning (ASP) toxins in plankton and
490 molluscs from Luanda Bay, Angola. *Toxicon* 55, 541–546.
491 <https://doi.org/10.1016/j.toxicon.2009.10.008>
- 492 Bonato, S., Christaki, U., Lefebvre, A., Lizon, F., Thyssen, M., Artigas, L.F., 2015. High spatial variability
493 of phytoplankton assessed by flow cytometry, in a dynamic productive coastal area, in
494 spring: The eastern English Channel. *Estuar. Coast. Shelf Sci.* 154, 214–223.
495 <https://doi.org/10.1016/j.ecss.2014.12.037>
- 496 Bouchouicha-Smida, D., Bates, S.S., Lundholm, N., Lambert, C., Mabrouk, H.H., Sakka Hlaili, A., 2015.
497 Viability, growth and domoic acid toxicity of the diatom *Nitzschia bizertensis* following
498 filtration by the mussel *Mytilus* sp. *Mar. Biol.* 162, 0. <https://doi.org/10.1007/s00227-015-2758-x>
- 500 Bresnan, E., Fryer, R.J., Fraser, S., Smith, N., Stobo, L., Brown, N., Turrell, E., 2017. The relationship
501 between *Pseudo-nitzschia* (Peragallo) and domoic acid in Scottish shellfish. *Harmful Algae* 63,
502 193–202. <https://doi.org/10.1016/j.hal.2017.01.004>
- 503 Bresnan, E., Kraberg, A., Fraser, S., Brown, L., Hughes, S., Wiltshire, K.H., 2015. Diversity and
504 seasonality of *Pseudo-nitzschia* (Peragallo) at two North Sea time-series monitoring sites.
505 *Helgol. Mar. Res.* 1–12. <https://doi.org/10.1007/s10152-015-0428-5>

506 Breton, E., Brunet, C., Sautour, B., Brylinski, J.-M., 2000. Annual variations of phytoplankton biomass
507 in the Eastern English Channel: comparison by pigment signatures and microscopic counts. *J.*
508 *Plankton Res.* 22, 1423–1440. <https://doi.org/10.1093/plankt/22.8.1423>

509 Brown, L., Bresnan, E., 2008. Seasonal occurrence of *Pseudo-nitzschia* species in the west coast and
510 Shetland Isles, Scotland, in: Proceedings of the 12th International Conference on Harmful
511 Algae. Presented at the 12th International Conference on Harmful Algae, Moestrup, Ø. et al.,
512 Copenhagen, Denmark, pp. 165–167.

513 Cho, E.S., Park, J.G., Oh, B.C., Cho, Y.C., 2001. The application of species specific DNA-targeted probes
514 and fluorescently tagged lectin to differentiate several species of *Pseudo-nitzschia*
515 (*Bacillariophyceae*) in Chinhae Bay, Korea. *Sci. Mar.* 65, 207–214.

516 De Rijcke, M., Vandegheuchte, M.B., Vanden Bussche, J., Nevejan, N., Vanhaecke, L., De
517 Schamphelaere, K.A.C., Janssen, C.R., 2015. Common European harmful algal blooms affect
518 the viability and innate immune responses of *Mytilus edulis* larvae - ScienceDirect. *Fish*
519 *Shellfish Immunol.* 47, 175–181. <https://doi.org/10.1016/j.fsi.2015.09.003>

520 Díaz, P.A., Ruiz-Villarreal, M., Velo-Suárez, L., Ramilo, I., Gentien, P., Lunven, M., Fernand, L., Raine,
521 R., Reguera, B., 2014. Tidal and wind-event variability and the distribution of two groups of
522 *Pseudo-nitzschia* species in an upwelling-influenced Ría. *Deep Sea Res. Part II Top. Stud.*
523 *Oceanogr., Harmful Algal Blooms in Stratified Systems* 101, 163–179.
524 <https://doi.org/10.1016/j.dsr2.2013.09.043>

525 Dortch, Q., Robichaux, R., Pool, S., Milsted, D., Mire, G., Rabalais, N., Soniat, T., Fryxell, G., Turner, R.,
526 Parsons, M., 1997. Abundance and vertical flux of *Pseudo-nitzschia* in the northern Gulf of
527 Mexico. *Mar. Ecol. Prog. Ser.* 146, 249–264. <https://doi.org/10.3354/meps146249>

528 Downes-Tettmar, N., Rowland, S., Widdicombe, C., Woodward, M., Llewellyn, C., 2013. Seasonal
529 variation in *Pseudo-nitzschia* spp. and domoic acid in the Western English Channel. *Cont.*
530 *Shelf Res.* 53, 40–49. <https://doi.org/10.1016/j.csr.2012.10.011>

531 Du, X., Peterson, W., Fisher, J., Hunter, M., Peterson, J., 2016. Initiation and Development of a Toxic
532 and Persistent *Pseudo-nitzschia* Bloom off the Oregon Coast in Spring/Summer 2015. *PLOS*
533 *ONE* 11, e0163977. <https://doi.org/10.1371/journal.pone.0163977>

534 Dursun, F., Yurdun, T., Ünlü, S., 2016. The First Observation of Domoic Acid in Plankton Net Samples
535 from the Sea of Marmara, Turkey. *Bull. Environ. Contam. Toxicol.* 96, 70–75.
536 <https://doi.org/10.1007/s00128-015-1704-4>

537 (EC) No 853/2004, 2004. Regulation (EC) No 853/2004 of the European parliament and of the council
538 og 29 April 2004 laying down specific hygiene rules for on the hygiene of foodstuffs. *Off. J.*
539 *Eur. Union* 47.

540 (EC) No 854/2004, 2004. Regulation (EC) No 854/2004 of the European parliament and of the council
541 og 29 April 2004 laying down specific rules for the organization of official controls on
542 products of animal origin intended for human consumption. *Off. J. Eur. Union* 47.

543 Escalera, L., Pazos, Y., Moroño, Á., Reguera, B., 2007. *Noctiluca scintillans* may act as a vector of
544 toxigenic microalgae. *Harmful Algae* 6, 317–320. <https://doi.org/10.1016/j.hal.2006.04.006>

545 Fehling, J., Davidson, K., Bolch, C., Tett, P., 2006. Seasonality of *Pseudo-nitzschia*
546 spp. (*Bacillariophyceae*) in western Scottish waters. *Mar. Ecol. Prog. Ser.* 323, 91–105.

547 Fehling, J., Green, D.H., Davidson, K., Bolch, C.J., Bates, S.S., 2004. Domoic acid production by
548 *Pseudo-nitzschia seriata* (*Bacillariophyceae*) in Scottish waters. *J. Phycol.* 40, 622–630.
549 <https://doi.org/10.1111/j.1529-8817.2004.03200.x>

550 Fock, H., Greve, W., 2002. Analysis and interpretation of recurrent spatio-temporal patterns in
551 zooplankton dynamics: a case study on *Noctiluca scintillans* (*Dinophyceae*) in the German
552 Bight (North Sea). *Mar. Biol.* 140, 59–73. <https://doi.org/10.1007/s002270100685>

553 Gai, F.F., Hedemand, C.K., Louw, D.C., Grobler, K., Krock, B., Moestrup, Ø., Lundholm, N., 2018.
554 Morphological, molecular and toxigenic characteristics of Namibian *Pseudo-nitzschia* species
555 – including *Pseudo-nitzschia bucculenta* sp. nov. *Harmful Algae* 76, 80–95.
556 <https://doi.org/10.1016/j.hal.2018.05.003>

557 Gentilhomme, V., Lizon, F., 1998. Seasonal cycle of nitrogen and phytoplankton biomass in a well-
558 mixed coastal system (Eastern English Channel). *Hydrobiologia* 361, 191–199.
559 <https://doi.org/10.1023/A:1003134617808>

560 Gobble, C., Duerr, R., Bodenstein, B., Lindquist, K., Lindsey, J., Beck, J., Henkel, L., Roletto, J., Harvey,
561 J., Kudela, R., 2018. Investigation of a Largescale Common Murre (*Uria aalge*) Mortality Event
562 in California, USA, in 2015. *J. Wildl. Dis.* 54, 569–574. <https://doi.org/10.7589/2017-07-179>

563 Gómez, F., Souissi, S., 2008. The impact of the 2003 summer heat wave and the 2005 late cold wave
564 on the phytoplankton in the north-eastern English Channel. *C. R. Biol.* 331, 678–685.
565 <https://doi.org/10.1016/j.crv.2008.06.005>

566 Grattepanche, J.-D., Breton, E., Brylinski, J.-M., Lecuyer, E., Christaki, U., 2011. Succession of primary
567 producers and micrograzers in a coastal ecosystem dominated by *Phaeocystis globosa*
568 blooms. *J. Plankton Res.* 33, 37–50. <https://doi.org/10.1093/plankt/fbq097>

569 Hasle, G.R., 2002. Are most of the domoic acid-producing species of the diatom genus *Pseudo-*
570 *nitzschia* cosmopolites? *Harmful Algae* 1, 137–146. [https://doi.org/10.1016/S1568-](https://doi.org/10.1016/S1568-9883(02)00014-8)
571 [9883\(02\)00014-8](https://doi.org/10.1016/S1568-9883(02)00014-8)

572 Hasle, G.R., Syvertsen, E.E., 1997. Marine diatoms, in: *Identifying Marine Phytoplankton*. Tomas C.R.,
573 pp. 5–385.

574 Hernández-Fariñas, T., Soudant, D., Barillé, L., Belin, C., Lefebvre, A., Bacher, C., 2014. Temporal
575 changes in the phytoplankton community along the French coast of the eastern English
576 Channel and the southern Bight of the North Sea. *ICES J. Mar. Sci.* 71, 821–833.
577 <https://doi.org/10.1093/icesjms/fst192>

578 Holmes, Aminot, A., Kerouel, R., Hooker, B.A., Peterson, B.J., 1999. A simple and precise method for
579 measuring ammonium in marine and freshwater ecosystems. *Can. J. Fish. Aquat. Sci.* 56,
580 1801–1808.

581 Husson, B., Hernández-Fariñas, T., Le Gendre, R., Schapira, M., Chapelle, A., 2016. Two decades of
582 *Pseudo-nitzschia* spp. blooms and king scallop (*Pecten maximus*) contamination by domoic
583 acid along the French Atlantic and English Channel coasts: Seasonal dynamics, spatial
584 heterogeneity and interannual variability. *Harmful Algae* 51, 26–39.
585 <https://doi.org/10.1016/j.hal.2015.10.017>

586 Jensen, S.-K., Lacaze, J.-P., Hermann, G., Kershaw, J., Brownlow, A., Turner, A., Hall, A., 2015.
587 Detection and effects of harmful algal toxins in Scottish harbour seals and potential links to
588 population decline. *Toxicon* 97, 1–14. <https://doi.org/10.1016/j.toxicon.2015.02.002>

589 Kesaulya, I., Leterme, S.C., Mitchell, J.G., Seuront, L., 2008. The impact of turbulence and
590 phytoplankton dynamics on foam formation, seawater viscosity and chlorophyll
591 concentration in the eastern English Channel.

592 Kim, J.H., Park, B.S., Kim, J.-H., Wang, P., Han, M.-S., 2015. Intraspecific diversity and distribution of
593 the cosmopolitan species *Pseudo-nitzschia pungens* (Bacillariophyceae): morphology,
594 genetics, and ecophysiology of the three clades. *J. Phycol.* 51, 159–172.
595 <https://doi.org/10.1111/jpy.12263>

596 Klein, C., Claquin, P., Bouchart, V., Le Roy, B., Véron, B., 2010. Dynamics of *Pseudo-nitzschia* spp. and
597 domoic acid production in a macrotidal ecosystem of the Eastern English Channel
598 (Normandy, France). *Harmful Algae* 9, 218–226. <https://doi.org/10.1016/j.hal.2009.10.004>

599 Kotaki, Y., Koike, K., Yoshida, M., Thuoc, C.V., Huyen, N.T.M., Hoi, N.C., Fukuyo, Y., Kodama, M., 2000.
600 Domoic Acid Production in *Nitzschia* Sp. (bacillariophyceae) Isolated from a Shrimp-Culture
601 Pond in Do Son, Vietnam. *J. Phycol.* 36, 1057–1060. [https://doi.org/10.1046/j.1529-](https://doi.org/10.1046/j.1529-8817.2000.99209.x)
602 [8817.2000.99209.x](https://doi.org/10.1046/j.1529-8817.2000.99209.x)

603 Lamy, D., Artigas, L.F., Jauzein, C., Lizon, F., Cornille, V., 2006. Coastal bacterial viability and
604 production in the eastern English Channel: A case study during a *Phaeocystis globosa* bloom.
605 *J. Sea Res.* 56, 227–238. <https://doi.org/10.1016/j.seares.2006.04.003>

606 Le Borgne, P., Legendre, G., Marsouin, A., 2006. Validation of the OSI SAF radiative fluxes, in:
607 *Proceedings of the 2006 EUMETSAT Meteorological Satellite Conference, 12-16 June 2006.*
608 Presented at the EUMETSAT Meteorological Satellite Conference, Helsinki, Finland.

609 Leandro, L.F., Teegarden, G.J., Roth, P.B., Wang, Z., Doucette, G.J., 2010. The copepod *Calanus*
610 *finmarchicus*: A potential vector for trophic transfer of the marine algal biotoxin, domoic
611 acid. *J. Exp. Mar. Biol. Ecol.* 382, 88–95. <https://doi.org/10.1016/j.jembe.2009.11.002>
612 Lefebvre, A., Blondel, C., Cordier, R., Duquesne, V., Hébert, P., Vérin, F., 2014. Bulletin de surveillance
613 de la qualité du milieu marin littoral 2013. Départements du Nord, Pas-de-Calais et de la
614 Somme. Ifremer, Boulogne sur Mer, France.
615 Lefebvre, A., Guiselin, N., Barbet, F., Artigas, F.L., 2011. Long-term hydrological and phytoplankton
616 monitoring (1992–2007) of three potentially eutrophic systems in the eastern English
617 Channel and the Southern Bight of the North Sea. *ICES J. Mar. Sci. J. Cons.* 68, 2029–2043.
618 <https://doi.org/10.1093/icesjms/fsr149>
619 Lefebvre, K.A., Silver, M.W., Coale, S.L., Tjeerdema, R.S., 2002. Domoic acid in planktivorous fish in
620 relation to toxic *Pseudo-nitzschia* cell densities. *Mar. Biol.* 140, 625–631.
621 Legendre, P., Legendre, L.F.J., 1998. *Numerical Ecology*. Elsevier.
622 Lelong, A., Hégaret, H., Soudant, P., Bates, S.S., 2012. *Pseudo-nitzschia* (Bacillariophyceae) species,
623 domoic acid and amnesic shellfish poisoning: revisiting previous paradigms. *Phycologia* 51,
624 168–216. <https://doi.org/10.2216/11-37.1>
625 Liefer, J.D., Robertson, A., MacIntyre, H.L., Smith, W.L., Dorsey, C.P., 2013. Characterization of a toxic
626 *Pseudo-nitzschia* spp. bloom in the Northern Gulf of Mexico associated with domoic acid
627 accumulation in fish. *Harmful Algae* 26, 20–32. <https://doi.org/10.1016/j.hal.2013.03.002>
628 Liu, H., Kelly, M.S., Campbell, D.A., Dong, S.L., Zhu, J.X., Wang, S.F., 2007. Exposure to domoic acid
629 affects larval development of king scallop *Pecten maximus* (Linnaeus, 1758). *Aquat. Toxicol.*
630 81, 152–158. <https://doi.org/10.1016/j.aquatox.2006.11.012>
631 Lorenzen, C.J., 1966. A method for the continuous measurement of *in vivo* chlorophyll concentration.
632 *Deep Sea Res. Oceanogr. Abstr.* 13, 223–227. [https://doi.org/10.1016/0011-7471\(66\)91102-8](https://doi.org/10.1016/0011-7471(66)91102-8)
633 Louw, D.C., Doucette, G.J., Lundholm, N., 2018. Morphology and toxicity of *Pseudo-nitzschia* species
634 in the northern Benguela Upwelling System. *Harmful Algae* 75, 118–128.
635 <https://doi.org/10.1016/j.hal.2018.04.008>
636 Louw, D.C., Doucette, G.J., Voges, E., 2016. Annual patterns, distribution and long-term trends of
637 *Pseudo-nitzschia* species in the northern Benguela upwelling system. *J. Plankton Res.*
638 <https://doi.org/10.1093/plankt/fbw079>
639 Lundholm, N., 2018. Bacillariophyceae, in IOC-UNESCO Taxonomic Reference List of Harmful Micro
640 Algae [WWW Document]. URL <http://www.marinespecies.org/hab> (accessed 8.9.18).
641 Lundholm, N., Clarke, A., Ellegaard, M., 2010. A 100-year record of changing *Pseudo-nitzschia* species
642 in a sill-fjord in Denmark related to nitrogen loading and temperature. *Harmful Algae* 9, 449–
643 457. <https://doi.org/10.1016/j.hal.2010.03.001>
644 Lundholm, N., Hansen, P.J., Kotaki, Y., 2004. Effect of pH on growth and domoic acid production by
645 potentially toxic diatoms of the genera *Pseudo-nitzschia* and *Nitzschia*. *Mar. Ecol. Prog. Ser.*
646 273, 1–15. <https://doi.org/10.2307/24867469>
647 Lundholm, N., Moestrup, Ø.J., 2000. Morphology of the Marine Diatom *Nitzschia Navis-Varingica*, Sp.
648 Nov. (bacillariophyceae), Another Producer of the Neurotoxin Domoic Acid. *J. Phycol.* 36,
649 1162–1174. <https://doi.org/10.1046/j.1529-8817.2000.99210.x>
650 Maneiro, I., Iglesias, P., Guisande, C., Riveiro, I., Barreiro, A., Zervoudaki, S., Granéli, E., 2005. Fate of
651 domoic acid ingested by the copepod *Acartia clausi*. *Mar. Biol.* 148, 123–130.
652 <https://doi.org/10.1007/s00227-005-0054-x>
653 McKibben, S.M., Watkins-Brandt, K.S., Wood, A.M., Hunter, M., Forster, Z., Hopkins, A., Du, X.,
654 Eberhart, B.-T., Peterson, W.T., White, A.E., 2015. Monitoring Oregon Coastal Harmful Algae:
655 Observations and implications of a harmful algal bloom-monitoring project. *Harmful Algae*
656 50, 32–44. <https://doi.org/10.1016/j.hal.2015.10.004>
657 Monchy, S., Grattepanche, J.-D., Breton, E., Meloni, D., Sancier, G., Chabé, M., Delhaes, L., Viscogliosi,
658 E., Sime-Ngando, T., Christaki, U., 2012. Microplanktonic Community Structure in a Coastal
659 System Relative to a *Phaeocystis* Bloom Inferred from Morphological and Tag

660 Pyrosequencing Methods. PLoS ONE 7, e39924.
661 <https://doi.org/10.1371/journal.pone.0039924>
662 Mullin, J.B., Riley, J.P., 1955. The colorimetric determination of silicate with special reference to sea
663 and natural waters. *Anal. Chim. Acta* 12, 162–176. [https://doi.org/10.1016/S0003-](https://doi.org/10.1016/S0003-2670(00)87825-3)
664 [2670\(00\)87825-3](https://doi.org/10.1016/S0003-2670(00)87825-3)
665 Murphy, J., Riley, J.P., 1962. A modified single solution method for the determination of phosphate in
666 natural waters. *Anal. Chim. Acta* 27, 31–36. [https://doi.org/10.1016/S0003-2670\(00\)88444-5](https://doi.org/10.1016/S0003-2670(00)88444-5)
667 Orsini, L., Procaccini, G., Sarno, D., Montresor, M., 2004. Multiple rDNA ITS-types within the diatom
668 *Pseudo-nitzschia delicatissima* (Bacillariophyceae) and their relative abundances across a
669 spring bloom in the Gulf of Naples. *Mar. Ecol. Prog. Ser.* 271, 87–98.
670 <https://doi.org/10.3354/meps271087>
671 Parsons, M.L., Dortch, Q., Turner, R.E., 2002. Sedimentological evidence of an increase in *Pseudo-*
672 *nitzschia* (Bacillariophyceae) abundance in response to coastal eutrophication. *Limnol.*
673 *Oceanogr.* 47, 551–558.
674 Parsons, M.L., Okolodkov, Y.B., Aké-Castillo, J.A., 2012. Diversity and morphology of the species of
675 *Pseudonitzschia* (Bacillariophyta) of the National Park Sistema Arrecifal Veracruzano, SW Gulf
676 of Mexico. *Acta Bot. Mex.*
677 Peperzak, L., Gäbler-Schwarz, S., 2012. Current Knowledge of the Life Cycles of *Phaeocystis Globosa*
678 and *Phaeocystis Antarctica* (prymnesiophyceae)1. *J. Phycol.* 48, 514–517.
679 <https://doi.org/10.1111/j.1529-8817.2012.01136.x>
680 Prince, E.K., Imer, F., Pohnert, G., 2013. Domoic Acid Improves the Competitive Ability of *Pseudo-*
681 *nitzschia delicatissima* against the Diatom *Skeletonema marinoi*. *Mar. Drugs* 11, 2398–2412.
682 <https://doi.org/10.3390/md11072398>
683 Pugliese, L., Casabianca, S., Perini, F., Andreoni, F., Penna, A., 2017. A high resolution melting method
684 for the molecular identification of the potentially toxic diatom *Pseudo-nitzschia* spp. in the
685 Mediterranean Sea. *Sci. Rep.* 7, 4259. <https://doi.org/10.1038/s41598-017-04245-z>
686 Quijano-Scheggia, S.I., Garcés, E., Lundholm, N., Moestrup, Ø., Andree, K., Camp, J., 2009.
687 Morphology, physiology, molecular phylogeny and sexual compatibility of the cryptic
688 *Pseudo-nitzschia delicatissima* complex (Bacillariophyta), including the description of *P.*
689 *arenysensis* sp. nov. *Phycologia* 48, 492–509. <https://doi.org/10.2216/08-21.1>
690 R Core Team, 2012. R: A language and environmet for statistical computing. R Foundation for
691 Statistical Computing, Vienna, Austria.
692 REPHY, Laboratoire Environnement Ressources Boulogne sur Mer, 2017. Qualité du Milieu Marin
693 Littoral - Bulletin de la surveillance 2016. Ifremer.
694 Rhodes, L., Scholin, C., Garthwaite, I., 1998. *Pseudo-nitzschia* in New Zealand and the role of DNA
695 probes and immunoassays in refining marine biotoxin monitoring programmes. *Nat. Toxins* 6,
696 105–111. [https://doi.org/10.1002/\(SICI\)1522-7189\(199805/08\)6:3/4<105::AID-](https://doi.org/10.1002/(SICI)1522-7189(199805/08)6:3/4<105::AID-NT13>3.0.CO;2-9)
697 [NT13>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1522-7189(199805/08)6:3/4<105::AID-NT13>3.0.CO;2-9)
698 Rousseau, V., Chrétiennot-Dinet, M.-J., Jacobsen, A., Verity, P., Whipple, S., 2007. The life cycle of
699 *Phaeocystis*: state of knowledge and presumptive role in ecology. *Biogeochemistry* 83, 29–
700 47. <https://doi.org/10.1007/s10533-007-9085-3>
701 Rue, E., Bruland, K., 2001. Domoic acid binds iron and copper: a possible role for the toxin produced
702 by the marine diatom *Pseudo-nitzschia*. *Mar. Chem.* 76, 127–134.
703 [https://doi.org/10.1016/S0304-4203\(01\)00053-6](https://doi.org/10.1016/S0304-4203(01)00053-6)
704 Sahraoui, I., Bates, S.S., Bouchouicha, D., Mabrouk, H.H., Hlaili, A.S., 2011. Toxicity of *Pseudo-*
705 *nitzschia* populations from Bizerte Lagoon, Tunisia, southwest Mediterranean, and first
706 report of domoic acid production by *P. brasiliiana*. *Diatom Res.* 26, 293–303.
707 <https://doi.org/10.1080/0269249X.2011.597990>
708 Sazhin, A.F., Artigas, L.F., Nejtgaard, J.C., Frischer, M.E., 2007. The colonization of two *Phaeocystis*
709 species (Prymnesiophyceae) by pennate diatoms and other protists: a significant contribution
710 to colony biomass. *Biogeochemistry* 83, 137–145. [https://doi.org/10.1007/s10533-007-9086-](https://doi.org/10.1007/s10533-007-9086-2)
711 [2](https://doi.org/10.1007/s10533-007-9086-2)

- 712 Schapira, M., Vincent, D., Gentilhomme, V., Seuront, L., 2008. Temporal patterns of phytoplankton
713 assemblages, size spectra and diversity during the wane of a *Phaeocystis globosa* spring
714 bloom in hydrologically contrasted coastal waters. J. Mar. Biol. Assoc. U. K. 88, 649–662.
715 <https://doi.org/10.1017/S0025315408001306>
- 716 Schoemann, V., Becquevort, S., Stefels, J., Rousseau, V., Lancelot, C., 2005. Phaeocystis blooms in the
717 global ocean and their controlling mechanisms: a review. J. Sea Res., Iron Resources and
718 Oceanic Nutrients - Advancement of Global Environmental Simulations 53, 43–66.
719 <https://doi.org/10.1016/j.seares.2004.01.008>
- 720 Scholin, C.A., Gulland, F., Doucette, G.J., Benson, S., Busman, M., Chavez, F.P., Cordaro, J., DeLong, R.,
721 De Vogelaere, A., Harvey, J., Haulena, M., Lefebvre, K., Lipscomb, T., Loscutoff, S.,
722 Lowenstine, L.J., Marin Iii, R., Miller, P.E., McLellan, W.A., Moeller, P.D.R., Powell, C.L.,
723 Rowles, T., Silvagni, P., Silver, M., Spraker, T., Trainer, V., Van Dolah, F.M., 2000. Mortality of
724 sea lions along the central California coast linked to a toxic diatom bloom. Nature 403, 80–
725 84. <https://doi.org/10.1038/47481>
- 726 Seuront, L., Vincent, D., Mitchell, J.G., 2006. Biologically induced modification of seawater viscosity in
727 the Eastern English Channel during a *Phaeocystis globosa* spring bloom. J. Mar. Syst. 61, 118–
728 133. <https://doi.org/10.1016/j.jmarsys.2005.04.010>
- 729 Sierra-Beltran, A.P., Palafox-Urbe, M., Grajales-Montiel, J., Cruz-Villacorta, A., Ochoa, J.L., 1997. Sea
730 bird mortality at Cabo San Lucas, Mexico: evidence that toxic diatom blooms are spreading.
731 Toxicon 35, 447–453.
- 732 Sobrinho, B.F., de Camargo, L.M., Sandrini-Neto, L., Kleemann, C.R., Machado, E. da C., Mafra, L.L.,
733 2017. Growth, Toxin Production and Allelopathic Effects of Pseudo-nitzschia multiseriis
734 under Iron-Enriched Conditions. Mar. Drugs 15, 331. <https://doi.org/10.3390/md15100331>
- 735 Sommer, U., 1994. Are marine diatoms favoured by high Si:N ratios? Mar. Ecol. Prog. Ser. 115, 309–
736 315. <https://doi.org/10.2307/24849755>
- 737 Stauffer, B., Gellene, A., Schnetzer, A., Seubert, E., Oberg, C., Sukhatme, G., Caron, D., 2012. An
738 oceanographic, meteorological, and biological ‘perfect storm’ yields a massive fish kill. Mar.
739 Ecol. Prog. Ser. 468, 231–243. <https://doi.org/10.3354/meps09927>
- 740 Tammilehto, A., Nielsen, T.G., Krock, B., Møller, E.F., Lundholm, N., 2015. Induction of domoic acid
741 production in the toxic diatom Pseudo-nitzschia seriata by calanoid copepods. Aquat. Toxicol.
742 Amst. Neth. 159, 52–61. <https://doi.org/10.1016/j.aquatox.2014.11.026>
- 743 Tan, S.N., Teng, S.T., Lim, H.C., Kotaki, Y., Bates, S.S., Leaw, C.P., Lim, P.T., 2016. Diatom Nitzschia
744 navis-varingica (Bacillariophyceae) and its domoic acid production from the mangrove
745 environments of Malaysia. Harmful Algae 60, 139–149.
746 <https://doi.org/10.1016/j.hal.2016.11.003>
- 747 Tas, S., Dursun, F., Aksu, A., Balkis, N., 2016. Presence of the diatom genus Pseudo-nitzschia and
748 particulate domoic acid in the Golden Horn Estuary (Sea of Marmara, Turkey). Diatom Res.
749 31, 339–349. <https://doi.org/10.1080/0269249X.2016.1247020>
- 750 Tas, S., Lundholm, N., 2016. Temporal and spatial variability of the potentially toxic Pseudo-nitzschia
751 spp. in a eutrophic estuary (Sea of Marmara). J. Mar.
752 Biol. Assoc. U. K. 1–12. <https://doi.org/10.1017/S0025315416000837>
- 753 Terenko, L., Terenko, G., 2012. Dominant Pseudo-Nitzschia (Bacillariophyta) species in the Black Sea
754 (Ukraine). Bot. Lith. 18, 27–34. <https://doi.org/10.2478/v10279-012-0004-8>
- 755 Thessen, A.E., Bowers, H.A., Stoecker, D.K., 2009. Intra- and interspecies differences in growth and
756 toxicity of Pseudo-nitzschia while using different nitrogen sources. Harmful Algae, This issue
757 contains the special section on “Strains” 8, 792–810.
758 <https://doi.org/10.1016/j.hal.2009.01.003>
- 759 Thessen, A.E., Dortch, Q., Parsons, M.L., Morrison, W., 2005. Effect of Salinity on Pseudo-Nitzschia
760 Species (bacillariophyceae) Growth and Distribution1. J. Phycol. 41, 21–29.
761 <https://doi.org/10.1111/j.1529-8817.2005.04077.x>
- 762 Thorel, M., Claquin, P., Schapira, M., Le Gendre, R., Riou, P., Goux, D., Le Roy, B., Raimbault, V.,
763 Deton-Cabanillas, A.-F., Bazin, P., Kientz-Bouchart, V., Fauchot, J., 2017. Nutrient ratios

764 influence variability in *Pseudo-nitzschia* species diversity and particulate domoic acid
765 production in the Bay of Seine (France). *Harmful Algae* 68, 192–205.
766 <https://doi.org/10.1016/j.hal.2017.07.005>

767 Trainer, V.L., Bates, S.S., Lundholm, N., Thessen, A.E., Cochlan, W.P., Adams, N.G., Trick, C.G., 2012.
768 *Pseudo-nitzschia* physiological ecology, phylogeny, toxicity, monitoring and impacts on
769 ecosystem health. *Harmful Algae* 14, 271–300. <https://doi.org/10.1016/j.hal.2011.10.025>

770 Trainer, V.L., Hickey, B.M., Bates, S.S., 2008. Toxic diatoms. *Oceans Hum. Health Risks Remedies Sea*
771 219–237.

772 Trick, C.G., Bill, B.D., Cochlan, W.P., Wells, M.L., Trainer, V.L., Pickell, L.D., 2010. Iron enrichment
773 stimulates toxic diatom production in high-nitrate, low-chlorophyll areas. *Proc. Natl. Acad.*
774 *Sci.* 107, 5887–5892. <https://doi.org/10.1073/pnas.0910579107>

775 Verity, P.G., Whipple, S.J., Nejstgaard, J.C., Alderkamp, A.-C., 2007. Colony size, cell number, carbon
776 and nitrogen contents of *Phaeocystis pouchetii* from western Norway. *J. Plankton Res.* 29,
777 359–367. <https://doi.org/10.1093/plankt/fbm021>


778 Villac, M.C., Roelke, D.L., Villareal, T.A., Fryxell, G.A., 1993. Comparison of two domoic acid-producing
779 diatoms: a review, in: *Twelfth International Diatom Symposium, Developments in*
780 *Hydrobiology*. Springer, Dordrecht, pp. 213–224. [https://doi.org/10.1007/978-94-017-3622-](https://doi.org/10.1007/978-94-017-3622-0_23)
781 [0_23](https://doi.org/10.1007/978-94-017-3622-0_23)

782 Walz, P.M., Garrison, D.L., Graham, W.M., Cattey, M.A., Tjeerdema, R.S., Silver, M.W., 1994. Domoic
783 acid-producing diatom blooms in Monterey Bay, California: 1991-1993. *Nat. Toxins* 2, 271–
784 279. <https://doi.org/10.1002/nt.2620020505>

785 Widdicombe, C.E., Eloire, D., Harbour, D., Harris, R.P., Somerfield, P.J., 2010. Long-term
786 phytoplankton community dynamics in the Western English Channel. *J. Plankton Res.* 32,
787 643–655. <https://doi.org/10.1093/plankt/fbp127>

788 Zabaglo, K., Chrapusta, E., Bober, B., Kaminski, A., Adamski, M., Bialczyk, J., 2016. Environmental
789 roles and biological activity of domoic acid: A review. *Algal Res.* 13, 94–101.
790 <https://doi.org/10.1016/j.algal.2015.11.020>


791



792

793 Figure 1. A – Location of the study area. B – Location of the sampling station (black dot) in
 794 Dunkirk harbor (Southern Bight of the North Sea).


795



796

797 Figure 2. Time series of meteorological and hydrological parameters. A. Temperature (°C)
 798 and salinity, B. pH and dO₂ (mg L⁻¹), C. Turbidity (NTU) and cumulated rainfalls over three
 799 days (bars, mm) during the 2012 survey.


800



801

802 Figure 3. time series of nutrient concentrations ($\mu\text{mol L}^{-1}$). A. $\text{NO}_2 + \text{NO}_3$, NH_4^+ and total
 803 dissolved inorganic nitrogen ($\text{DIN} = \text{NO}_2 + \text{NO}_3 + \text{NH}_4^+$). B. HPO_4^{2-} . C. Si(OH)_4 , during the
 804 2012 survey.


805



806

807 Figure 4. Time series of A. Chlorophyll *a* (●) and pheopigment (■) concentrations ($\mu\text{g L}^{-1}$);
 808 B. Abundance (cell L^{-1}) of total phytoplankton (◆), diatoms (Δ), *Phaeocystis globosa* (□) and
 809 other phytoplankton groups (×); and C. relative abundance of diatom genus (%) during the
 810 2012 survey.


811



812

813


814 Figure 5. SEM micrographs of the three identified *Pseudo-nitzschia* species. A-B: *Pseudo-*
 815 *nitzschia delicatissima*; C-d: *Pseudo-nitzschia pungens*; E-F: *Pseudo-nitzschia fraudulenta*.
 816 Scale bars: A, C, E = 1 μ m, B, D, F = 10 μ m.



817

818 Figure 6: Time series of *Pseudo-nitzschia* complex *delicatissima* (◆) and *P. complex seriata*
 819 (\times) abundances (cell L⁻¹), and particulate and dissolved domoic acid (DA, pg mL⁻¹) during the
 820 2012 survey. Stars below the x axis indicate the dates of SEM analyses for *Pseudo-nitzschia*
 821 sp. diversity.

822



823

824 Figure 7: PCA plots of environmental variables. A and C. Plane defined by the two first axes
 825 of the PCA (PC1 and PC2) B and D. plane defined by the first (PC1) and third axis (PC3) of
 826 the PCA (B and D). Points correspond to sampling dates. The colors of the circles correspond
 827 to the periods (P1 to P4) determined from phytoplankton communities composition. Circles
 828 size is proportional to *Pseudo-nitzschia* abundance (cell L⁻¹) for *P. complex delicatissima* (A
 829 and B) and *P. complex seriata* (C and D). Black arrows represent variables significantly
 830 correlated to at least one represented PC.

Table 1. Morphological features of *Pseudo-nitzschia* species and comparison with literature values (min-max). * correspond to one frustule analysed and presenting outliers in poroids features (among >300 identified frustules)

Species	Width µm	Length µm	Central Interspace	Fibulae 10 µm ⁻¹	Interstriae 10 µm ⁻¹	Rows of poroids	Poroids 1 µm ⁻¹	Reference
<i>P. delicatissima</i>	1.3-2	N/A	+	22-29	33-43	2	8-11	Amato et al., 2005
	1.7-2.8	N/A	+	20-27	24-40	2	8-14	Fehling et al., 2006
	1.1-2	40-78	+	19-25	36-40	2	10-12	Skov et al., 1999
	1.0-1.5	42-66	+	20-23	env 40	2	10-12	Hasle et al., 1996
	1.7-2.3	17-48	+	18-36	37-44	2	7.5-13.4	Kaczmarek et al., 2008
	1.3-1.6	48-49	-	22-23	36-40	2	8-9	Klein et al., 2010
	1.5-2.0	19-76	N/A	19-26	35-40	2	12	Lundholm et al., 2006
	1.3-1.6	34-46	+	24-28	41-45	2	7-11	Moschandreou and Nikolaidis, 2010
	1.0-2.4	19-78	+	19-30	33-42	2	8-12.5	Quijano-Scheggia et al., 2010
	1.4-2.6	31-48	+	20-28	38-41	2	8-12	Fernandes et al 2014
	1.1-1.7	40.7-67.1	+	18-27	34-44	2(3*)	10-14(18*)	This study
<i>P. fraudulenta</i>	4.3-5.1	N/A	+	20-24	22-26	2-3	5-6	Fehling et al., 2006
	4.5-10	50-119	+	12-24	18-24	2-3	5-7	Skov et al., 1999
	5-6	73-117	+	19-23	19-23	2-3	4-5	Hasle et al., 1996
	3-4.5	63-99	+	15-17	14-15	2	N/A	Hernández-Becerril, 1998
	5.1-5.6	56.8-81	-	21-22	22-24	2-3	6-7	Klein et al., 2010
	4-7	71-101	N/A	18-30	19-28	2-3	6-7	Ljubešić et al., 2011
	4.4-5.8	48-80	+	22-24	21-24	2	5-6	Moschandreou and Nikolaidis, 2010
	2.9-7.1	38.9-131.5	+	12-24	18-24	2	4-7	Quijano-Scheggia et al., 2010
	5.2-6.2	64-84	+	20-23	22-23	2	5-6	Fernandes et al. 2014
	5.2-6.1	75.0-98.8	+	19-23	17-23	2-3	6-7	This study
<i>P. pungens</i>	3.1	N/A	-	12	12	2	3-4	Fehling et al., 2006
	2.4-5.3	74-174	-	9-16	9-16	1-2	3-4	Skov et al., 1999
	2.9-4.5	74-142	-	9-15	9-15	2	3-4	Hasle et al., 1996
	3-4	94-109	-	10-11	10-12	2	N/A	Hernández-Becerril, 1998
	2.9-3.3	61-106	-	11-13	11-14	2-3	2-4	Klein et al., 2010
	3.9-5.7	88-105	-	10-17	8-15	2	2-4	Ljubešić et al., 2011
	2.5-3.6	77-101	-	11-13	11-13	2	2-4	Moschandreou and Nikolaidis, 2010
	2-4.8	37-156	-	9-13	9-15	2	1-5	Quijano-Scheggia et al., 2010
	4-4.5	75-130	-	9-13	N/A	2	3-4	Stonik et al., 2001
	3.0-4.1	72-149	-	11-13	10-12	2	3-4	Fernandes et al 2014
		2.7-4.7	95.9-135.5	-	10-15	10-15	2-3	2-4

Table 2. Spearman correlation coefficient between principal components (PCx), environmental parameters, *Pseudo-nitzschia* complexes abundances and domoic acid concentration.

Parameters	PC1	PC2	PC3
<i>Main variables</i>			
Temperature (T)	-0.18	0.73**	-0.45 ⁺
Salinity (S)	-0.39	0.64**	0.13
pH	0.43 ⁺	-0.64**	0.29
dO ₂	0.22	-0.75***	0.26
Turbidity	0.498*	0.22	0.15
Rainfalls	-0.29	0.55*	0.61*
Windspeed	0.61*	0.51*	0.47
Surface Solar Irradiance (ssi)	-0.74**	-0.06	-0.15
DIN	0.88***	-0.29	-0.25
PO ₄ ²⁻	0.87***	0.28	-0.15
Si(OH) ₄	0.69***	0.37	-0.38
<i>Supplementary variables</i>			
Si/P	0.26	0.44	-0.38
Si/N	-0.23	0.69**	-0.19
N/P	0.66**	-0.50*	-0.20
<i>Pseudo-nitzschia and DA</i>			
<i>P. delicatissima</i> complex	-0.27	-0.33	0.47 ⁺
<i>P. seriata</i> complex	-0.31	-0.07	0.50*
tDA	-0.30	0.12	0.20
(* p-value < 0.05, ** p-value < 0.01, *** p-value < 0.001)			

Table 3. Environmental features of *Pseudo-nitzschia* spp. identified during the 2012 survey and literature values.

Species	Location	Temperature °C	Salinity	DIN µmol L ⁻¹	PO4 µmol L ⁻¹	Si µmol L ⁻¹	Abundance cells L ⁻¹	DA	Reference
<i>P. delicatissima</i>	Southern Adriatic Sea	8.79-29.91, R<0	R>0	no significant pattern			nd	nd	Caroppo et al., 2005
	Western Scotland	~8°C	nd	~3µM	~0.3µM	~2.5µM	1.6·10 ⁵	nd	Fehling et al., 2006
	Louisiana	nd	nd	no significant pattern			1·10 ⁶	5.5pg cell ⁻¹	Parsons et al., 2013
	NE Spanish Coast	R>0	36.8±1.1, R<0	R>0	nd	nd	17.6±5.2·10 ⁵	nd	Quijano-Scheggia et al., 2008
	Bay of Fundy	nd	nd	R>0	no significant correlation		nd	nd	Kaczmarek et al., 2007
	Gulf of maine	nd	nd	nd	nd	nd	nd	111-390 pg mL ⁻¹	Fernandes et al., 2014
	Black Sea	16-17.5	14.5-15.7	nd	nd	nd	1.5-3.1·10 ⁶	nd	
<i>P. delicatissima</i> complex	Western English Channel	déc-15	34.9±0.4, R>0	nd	R<0	nd	2,5·10 ⁵	0.2-0.4 pg mL ⁻¹	Downes-Tettmar et al., 2013
	Galicia – Spain	~ 13.1	~35.6	~2.1	~0.22	~3	6.6·10 ⁵	not detected	Díaz et al., 2014
	Southern Bight of the North Sea	9-12	34.2±0.2	3.5±2.5 decreasing	0.39±0.04	1.6±0.8	33·10⁵-7.5·10⁶	92-229pg mL⁻¹	This study
<i>P. pungens</i>	Louisiana	nd	nd	R<0	R>0	no corr	50000	nd	Parsons et al., 2013
	NE Spanish Coast	nd	nd	no pattern	nd	nd	nd	nd	Quijano-Scheggia et al., 2008
	Gulf of Maine	nd	nd	Nd	nd	nd	nd	40-1100 pg mL ⁻¹	Fernandes et al., 2014
	Bay of Fundy	nd	nd	R<0	R>0	nd	nd	nd	Kaczmarek et al., 2007
	Western English Channel	12-16	34.9±0.4	R<0	R<0	R<0	25000	0.2-0.4 pg mL ⁻¹	Downes-Tettmar et al., 2013
	Sea of Marmara, Turkey	11.6-22.8	21.6	nd	nd	1.8	4.8·10 ⁵	nd	Tas and Lundholm, 2016
	Sea of Marmara, Turkey	7.4-26.3	15.9-19.9	Nd	Nd	Nd	1.6-1.7·10 ⁴	1.0pg mL ⁻¹	Tas et al., 2016
<i>P. fraudulenta</i>	California	18.5-26.5	nd	0-3	nd	2.9-35.5	5-13 10 ⁵	detected	Gárate-Lizárraga et al., 2007
	North Western Mediterranean Sea	nd	R>0	R<0	nd	nd	nd	nd	Quijano-Scheggia et al., 2008
<i>P. seriata</i> complex including <i>P. pungens</i> , <i>P. fraudulenta</i>	Western Scotland	12-13	nd	~0.3	~0.25	~3	110000	nd	Fehling et al., 2006
	Southern Bight of the North Sea	15.4±0.3	34.50±0.03	2.4±1.8 decreasing	0.56±0.17	2.2±2.8 increasing	9.2·10⁵-2.8·10⁶	82-108pg mL⁻¹	This study

Min-max, mean ± standard deviation, R<0 and R>0: negative and positive correlation with diatom abundance, respectively., nd = no data

