



Major merging history in CANDELS. I. Evolution of the incidence of massive galaxy-galaxy pairs from $z = 3$ to $z \sim 0$

Tommy Wiklind, Mauro Stefanon, Paola Santini, Janine Pforr, Pablo G. Pérez-González, Hooshang Nayyeri, Bahram Mobasher, Yicheng Guo, Norman A. Grogin, Audrey Galametz, et al.

► To cite this version:

Tommy Wiklind, Mauro Stefanon, Paola Santini, Janine Pforr, Pablo G. Pérez-González, et al.. Major merging history in CANDELS. I. Evolution of the incidence of massive galaxy-galaxy pairs from $z = 3$ to $z \sim 0$. Monthly Notices of the Royal Astronomical Society, 2018, 475, pp.1549-1573. 10.1093/mnras/stx3260 . insu-03666274

HAL Id: insu-03666274

<https://insu.hal.science/insu-03666274>

Submitted on 12 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Major merging history in CANDELS. I. Evolution of the incidence of massive galaxy–galaxy pairs from $z = 3$ to $z \sim 0$

Kameswara Bharadwaj Mantha,^{1★} Daniel H. McIntosh,¹ Ryan Brennan,² Henry C. Ferguson,³ Dritan Kodra,⁴ Jeffrey A. Newman,⁴ Marc Rafelski,³ Rachel S. Somerville,² Christopher J. Conselice,⁵ Joshua S. Cook,¹ Nimish P. Hathi,³ David C. Koo,⁶ Jennifer M. Lotz,³ Brooke D. Simmons,⁷ Amber N. Straughn,⁸ Gregory F. Snyder,³ Stijn Wuyts,⁹ Eric F. Bell,¹⁰ Avishai Dekel,¹¹ Jeyhan Kartaltepe,¹² Dale D. Kocevski,¹³ Anton M. Koekemoer,³ Seong-Kook Lee,¹⁴ Ray A. Lucas,³ Camilla Pacifici,¹⁵ Michael A. Peth,¹⁶ Guillermo Barro,¹⁷ Tomas Dahlen,³ Steven L. Finkelstein,¹⁸ Adriano Fontana,¹⁹ Audrey Galametz,²⁰ Norman A. Grogin,³ Yicheng Guo,⁶ Bahram Mobasher,²¹ Hooshang Nayyeri,²¹ Pablo G. Pérez-González,²² Janine Pforr,^{23,24} Paola Santini,¹⁹ Mauro Stefanon²⁵ and Tommy Wiklund³

Affiliations are listed at the end of the paper

Accepted 2017 December 13. Received 2017 December 13; in original form 2017 July 29

ABSTRACT

The rate of major galaxy–galaxy merging is theoretically predicted to steadily increase with redshift during the peak epoch of massive galaxy development ($1 \leq z \leq 3$). We use close-pair statistics to objectively study the incidence of massive galaxies (stellar $M_1 > 2 \times 10^{10} M_\odot$) hosting major companions ($1 \leq M_1/M_2 \leq 4$; i.e. $<4:1$) at six epochs spanning $0 < z < 3$. We select companions from a nearly complete, mass-limited ($\geq 5 \times 10^9 M_\odot$) sample of 23 696 galaxies in the five Cosmic Assembly Near-Infrared Deep Extragalactic Legacy Survey fields and the Sloan Digital Sky Survey. Using 5–50 kpc projected separation and close redshift proximity criteria, we find that the major companion fraction $f_{\text{mc}}(z)$ based on stellar mass-ratio (MR) selection increases from 6 per cent ($z \sim 0$) to 16 per cent ($z \sim 0.8$), then turns over at $z \sim 1$ and decreases to 7 per cent ($z \sim 3$). Instead, if we use a major F160W flux-ratio (FR) selection, we find that $f_{\text{mc}}(z)$ increases steadily until $z = 3$ owing to increasing contamination from minor ($\text{MR} > 4:1$) companions at $z > 1$. We show that these evolutionary trends are statistically robust to changes in companion proximity. We find disagreements between published results are resolved when selection criteria are closely matched. If we compute merger rates using constant fraction-to-rate conversion factors ($C_{\text{merg,pair}} = 0.6$ and $T_{\text{obs,pair}} = 0.65$ Gyr), we find that MR rates disagree with theoretical predictions at $z > 1.5$. Instead, if we use an evolving $T_{\text{obs,pair}}(z) \propto (1+z)^{-2}$ from Snyder et al., our MR-based rates agree with theory at $0 < z < 3$. Our analysis underscores the need for detailed calibration of $C_{\text{merg,pair}}$ and $T_{\text{obs,pair}}$ as a function of redshift, mass, and companion selection criteria to better constrain the empirical major merger history.

Key words: galaxies: evolution – galaxies: high-redshift – galaxies: interactions – galaxies: statistics.

* E-mail: km4n6@mail.umkc.edu

1 INTRODUCTION

In a hierarchical universe, collisions between similar-mass galaxies (major mergers) are expected to occur, and many theoretical studies predict such merging plays an important role in the formation and evolution of massive galaxies. A key measurement for quantifying the role of major merging in galaxy development is the merger rate and its evolution during cosmic history. A host of studies have measured major merger rates at redshifts $z \leq 1.5$, primarily based either on close-pair statistics (e.g. Patton et al. 1997; Lin et al. 2004; Kartaltepe et al. 2007; Bundy et al. 2009), clustering statistics (e.g. Bell et al. 2006b; Robaina et al. 2010), and morphological disturbances and asymmetries (e.g. Lotz et al. 2008; Conselice, Yang & Bluck 2009). These studies have all found higher incidences of major merging at earlier look-back times and a strong to moderate decrease to the present epoch, in broad agreement with many theoretical predictions (e.g. Gottlöber, Klypin & Kravtsov 2001; Bower et al. 2006; Hopkins et al. 2010a). Despite these successes, large scatter (factor of 10) exists between even the most stringent individual constraints, owing to systematic uncertainties in different methodologies and merger time-scales. These issues are compounded for empirical estimates at the epoch of peak galaxy development ($z \sim 2-3$; ‘cosmic high noon’). Some early empirical estimates based on both methodologies found increasing major merger incidence at $z > 1.5$ (e.g. Bluck et al. 2009), but recent studies find a possible flattening or turnover in merger rates between $1 < z < 3$ (e.g. Ryan et al. 2008; Man, Zirm & Toft 2016; Mundy et al. 2017). These new empirical trends are in strong disagreement with recent theoretical models predicting that merger rates continue to rise from $z = 1$ to 3 and beyond (Hopkins et al. 2010a; Lotz et al. 2011; Rodriguez-Gomez et al. 2015). These discrepancies and the large variance between past measurements highlight the need for improved major merger constraints, especially during the critical high-noon epoch.

A host of selection-effect issues has plagued many previous attempts to constrain major merger statistics at high redshift, from low-number statistics and significant sample variance due to small-volume pencil-beam surveys, and rest-frame ultraviolet (UV) selections of both disturbed morphologies and close pairs. While the identification of close pairs is less prone to some systematics, the lack of statistically useful samples of spectroscopic redshifts or even moderately small-uncertainty photometric redshifts at $z > 1$ until very recently have limited the usefulness of this method. Moreover, the wildly varying close-companion selection criteria among previous studies is a plausible explanation for tensions between empirical merger rates and theoretical predictions (Lotz et al. 2011). In this study, we will address many of these shortcomings and systematically explore the impact of major close-companion selection criteria by analysing major companion fractions in a sample of 10 000 massive host galaxies (stellar mass $M_1 \geq 2 \times 10^{10} M_\odot$) from the *five Hubble Space Telescope (HST)* legacy fields in Cosmic Assembly Near-Infrared Deep Extragalactic Legacy Survey (CANDELS, Grogin et al. 2011; Koekemoer et al. 2011) and the Sloan Digital Sky Survey (SDSS). This comprehensive sample provides statistically useful major companion counts, down to a mass limit of $M_2 = 5 \times 10^9 M_\odot$, from rest-frame optical images over a large volume out to $z = 3$.

The hierarchical major merging of similar-mass haloes via gravitational accretion is the underlying physical driver of galaxy–galaxy major merging. Cosmological simulations predict that the major halo–halo merger rate rises steeply with redshift as $R \propto (1+z)^{2-3}$ (e.g. Fakhouri & Ma 2008; Genel et al. 2009; Fakhouri, Ma &

Boylan-Kolchin 2010), which is in agreement with simple analytical predictions based on Extended Press–Schechter theory of $R \propto (1+z)^{2.5}$ (Neistein & Dekel 2008; Dekel et al. 2013). Cosmologically motivated simulations of galaxy formation and evolution predict major galaxy–galaxy merger rates that follow $R \propto (1+z)^{1-2}$ over a wide redshift range (e.g. $z < 6$; Rodriguez-Gomez et al. 2015). While there is some debate on the increasing merger rate evolution among theoretical studies due to model dependencies (for review, see Hopkins et al. 2010a), some works claim flattening of merger rates with increasing redshift (e.g. Henriques et al. 2015), most agree with an increasing incidence (within a factor-of-two uncertainty). Not only are merger rates expected to be higher at early cosmic times, but major galaxy merging is predicted to play a crucial role in nearly all aspects of the formation and evolution of massive galaxies including build-up of spheroidal bulges and massive elliptical galaxies (Springel 2000; Khochfar & Burkert 2003, 2005; Naab, Khochfar & Burkert 2006; Cox et al. 2008), triggering and enhancement of star formation (SF) including nuclear starbursts (Sanders et al. 1988; Di Matteo et al. 2007, 2008; Martig & Bournaud 2008), and the fuelling of active galactic nuclei (AGNs, Hopkins et al. 2006; Younger et al. 2009; Narayanan et al. 2010; Hopkins et al. 2010a) and subsequent SF quenching (e.g. Di Matteo, Springel & Hernquist 2005; Hopkins et al. 2008).

Many empirical studies support the predictions that major merging may explain the documented build-up of massive and quenched (non-star-forming and red) galaxy number densities and their stellar content growth at $z < 1$ (e.g. Bell et al. 2006a; McIntosh et al. 2008; van der Wel et al. 2009), enhancement of SF activity (e.g. Jogee et al. 2009; Patton et al. 2011), and elevation of AGN activity (Treister et al. 2012; Weston et al. 2017; Hewlett et al. 2017). Despite this agreement, some studies find a weak major merging–SF connection and suggest mergers may not be the dominant contributor to *in situ* galactic SF (Robaina et al. 2009; Swinbank et al. 2010; Targett et al. 2011). Moreover, other studies find a lack of a merger–AGN connection (Grogin et al. 2005; Kocevski et al. 2012; Villforth et al. 2014, 2017). These conflicting observations lend support to theories that predict violent disc instabilities (VDI) due to the rapid hierarchical accretion of cold gas may be responsible for key processes like bulge formation and AGN triggering (Bournaud et al. 2011; Dekel & Burkert 2014). Indeed, a recent CANDELS study by Brennan et al. (2015) found the observed evolution of massive quenched spheroids at $z < 3$ is better matched to semi-analytic model (SAM) predictions that include both mergers and disc-instability prescriptions. Therefore, the role of major merging in galaxy evolution remains a critical open question. Hence, measuring the frequency and rate at which major mergers occur at different cosmic times using large, uniformly selected close-pair samples is a key step towards answering the role played by them in massive galaxy development.

Theoretical simulations predict that galaxies involved in major close pairs will interact gravitationally and coalesce over time into one larger galaxy, and thereby make them effective probes of ongoing or future merging. Many studies in the past have employed the close-pair method to estimate the frequency of major merging as a function of cosmic time. This typically involves searching for galaxies that host a nearby companion meeting a number of key criteria: (i) 2D projected distance, (ii) close redshift–space proximity, and satisfies a nearly-equal mass ratio (MR) M_1/M_2 between the host (1) and companion (2) galaxies. For each criterion, a wide range of choices is used in the literature. For projected separation R_{proj} , a search annulus is often employed with minimum and maximum radii. Common choices vary between $R_{\text{max}} \sim 30-140$ kpc (e.g.

Patton & Atfield 2008; de Ravel et al. 2009) and $R_{\min} \sim 0\text{--}14$ kpc (e.g. Bluck et al. 2009; Man et al. 2016). Depending on available redshift information, the choice of physical proximity criterion ranges from stringent spectroscopic velocity differences (commonly $\Delta v_{12} \leq 500 \text{ km s}^{-1}$; e.g. Lin et al. 2008) to a variety of photometric redshift z_{phot} error overlaps (e.g. Bundy et al. 2009; Man et al. 2012). To study similar-mass galaxy–galaxy mergers, previous studies have adopted stellar-MR selections ranging from $2 > M_1/M_2 > 1$ (or 2:1, e.g. De Propriis et al. 2007) to $5 > M_1/M_2 > 1$ (5:1, e.g. Lofthouse et al. 2017), with 4:1 being by far the most common MR criterion. In the absence of stellar-mass estimates, flux ratio (FR) F_1/F_2 is often used as a proxy for M_1/M_2 (e.g. Bridge et al. 2007). The wide range of adopted close-companion selection criteria lead to a large scatter in two decades of published pair-derived merger rates with redshift evolution spanning $R \propto (1+z)^{0.5-3}$ at $0 < z < 1.5$, and sometimes even indicating a flat or turnover in merger rates at $z > 1.5$ (Williams, Quadri & Franx 2011; Man et al. 2012; Man, Zirm & Toft 2016; Mundy et al. 2017). This large scatter in pair-derived merger rate constraints highlights the strong need for tighter constraints at cosmic high noon, and motivates a careful analysis of selection effects.

Numerical simulations of the gravitation interactions between merging galaxies can produce disturbed morphological features due to strong tidal forces (e.g. Barnes & Hernquist 1996; Bournaud & Duc 2006; Peirani et al. 2010). As such, morphological selections have also been used to empirically identify mergers. These selections are broadly divided into visual classifications (e.g. Darg et al. 2010; Kartaltepe et al. 2015), analysis of image-model residuals (e.g. McIntosh et al. 2008; Tal et al. 2009), and automated measures of quantitative morphology such as Gini-M20 (Lotz, Primack & Madau 2004) and Concentration-Asymmetry-Clumpiness metric system (Conselice 2003). Although morphology-based studies broadly find merger rates to be rising strongly with redshift as $(1+z)^{2-5}$ (e.g. López-Sanjuan et al. 2009; Wen & Zheng 2016), sometimes finding as high as 25 per cent–50 per cent of their sample as mergers (Conselice, Rajgor & Myers 2008b), there are significant study-to-study discrepancies where some studies find no merger rate evolution (e.g. Cassata et al. 2005; Lotz et al. 2008). Morphology-based selections depend on identifying relatively fainter disturbances than the galaxy, which makes this method prone to systematics. The cosmological surface brightness of galaxies falls off as $(1+z)^{-4}$, which can lead to a biased identification of faint merger-specific features as a function of redshift. In addition, most of these morphology-based merger rates are based on small-volume, pencil-beam surveys probing the rest-frame UV part of the spectrum, especially at $z > 1$. This can lead to overestimation of merger rates due to contamination from non-merging, high star-forming systems with significant substructure that can be confused as two merging galaxies. Recent theoretical developments suggest that VDI can also cause disturbances in the host galaxy morphology and mimic merger-like features (see Dekel, Sari & Ceverino 2009; Cacciato, Dekel & Genel 2012; Ceverino et al. 2015), which in principle may complicate the measurement of morphology-based merger rates. Thus, to robustly identify plausible merging systems out to high redshifts ($z \sim 3$) without having to rely on imaging-related systematics strongly, we resort to the close-pair method in this study. We acknowledge that the close-pair method has its limitations at high redshift where galaxies have large photometric redshift uncertainties, which may lead to incorrect merger statistics. In this study, we initially exclude the galaxies with unreliable redshifts from our analysis, but later add back a certain fraction of them by employing a statistical correction.

In this paper, we analyse galaxy–galaxy close pairs in a large sample of 5698 massive galaxies ($M_{\text{stellar}} \geq 2 \times 10^{10} M_{\odot}$) from the state-of-the-art CANDELS (Grogin et al. 2011; Koekemoer et al. 2011) of five highly studied extragalactic fields at six epochs spanning $z \sim 0.5\text{--}3.0$ (with a width $\Delta z = 0.5$). To simultaneously anchor our findings to $z = 0$, we take advantage of 4098 massive galaxies from the SDSS (York et al. 2000), Data Release 4 (DR4; Adelman-McCarthy et al. 2006) at $z \sim 0.03\text{--}0.05$, which is matched in resolution to CANDELS and probes an average of $\sim 1.3 \times 10^6 \text{ Mpc}^3$ per redshift bin. With the available data, we also perform rigorous analyses to understand the impact of different close-companion selection criteria on the derived results.

We structure this paper as follows: In Section 2, we provide a brief description of the CANDELS and SDSS data products (redshifts and stellar masses) and describe the selection of massive galaxies hosting major companions based on the stellar mass complete massive galaxy sample. In Section 3, we describe the calculation of major companion fraction and its redshift evolution including necessary statistical corrections. In Section 4, we discuss the impact of close-companion selection choices on the derived major companion fractions. In Section 5, we calculate the major merger rates based on the major companion fractions. We synthesize detailed comparisons of the companion fractions and merger rates to other empirical studies and theoretical model predictions, and also discuss plausible reasons and implications of disagreement between the observed and theoretical merger rates. We present our conclusions in Section 6. Throughout this paper, we adopt a cosmology of $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ($h = 0.7$), $\Omega_M = 0.3$, and $\Omega_{\Lambda} = 0.7$, and use the AB magnitude system (Oke & Gunn 1983).

2 DATA AND GALAXY SAMPLE

In this study, we analyse close galaxy–galaxy pairs selected from a large sample of massive galaxies from the CANDELS (Grogin et al. 2011; Koekemoer et al. 2011), spanning redshifts $0.5 \leq z \leq 3$, and subdivided into five epochs probing a volume of $\sim 1.3 \times 10^6 \text{ Mpc}^3$ each. We anchor our findings to $z \sim 0$ using a sample from the SDSS that is matched in volume and resolution to the CANDELS sample. To reliably track the major merging history since $z = 3$ using close-pair method, we start with a mass-limited sample of galaxies that will allow a complete selection of massive $> 2 \times 10^{10} M_{\odot}$ galaxies with major companions meeting our chosen stellar MR: $1 \leq M_1/M_2 \leq 4$ (1, 2 represent host and companion galaxies, respectively). In this section, we describe the relevant details of the data necessary to achieve this sample selection.

2.1 CANDELS : 5 legacy fields

2.1.1 Photometric source catalogues

The five CANDELS *HST* legacy fields – UDS, GOODS-S, GOODS-N, COSMOS, and EGS (for details, see Grogin et al. 2011) – have a wealth of multiwavelength data and cover a total area of $\sim 800 \text{ arcmin}^2$ ($\sim 0.22 \text{ deg}^2$). The CANDELS survey observations and image processing are described in Grogin et al. (2011) and Koekemoer et al. (2011), respectively. We use the photometric source catalogues from Galametz et al. (2013, UDS), Guo et al. (2013, GOODS-S), Barro et al., in preparation (GOODS-N), Nayyeri et al. (2017, COSMOS), and Stefanon et al. (2017, EGS). Each catalogue was generated with a consistent source detection algorithm using *SEXTRACTOR* applied to the F160W (*H* band) two-orbit depth CANDELS mosaic image produced for each field. These

Table 1. Galaxy sample information. Columns: (1) name of the field/survey; (2) the total number of sources in photometry catalogue before (after) applying good-source cuts described in Section 2.1.1; (3) the redshift range of interest in our study, used to select the mass-limited sample counts in (4) and (5), where the subsets with spectroscopic-redshift information are given in parenthesis.

Name (1)	Phot sources (2)	Redshift range (3)	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ (4)	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.3$ (5)
UDS	35 932 (33 998)	$0.5 \leq z \leq 3.0$	3019 (260)	1223 (141)
GOODS-S	34 930 (34 115)	$0.5 \leq z \leq 3.0$	2491 (892)	942 (403)
GOODS-N	35 445 (34 693)	$0.5 \leq z \leq 3.0$	2946 (494)	1133 (209)
COSMOS	38 671 (36 753)	$0.5 \leq z \leq 3.0$	3232 (11)	1307 (9)
EGS	41 457 (37 602)	$0.5 \leq z \leq 3.0$	2825 (199)	1093 (72)
CANDELS (Total)	186 435 (177 161)	$0.5 \leq z \leq 3.0$	14 513 (1856)	5698 (834)
SDSS-DR4 (1790 deg ²)	141 564	$0.03 \leq z \leq 0.05$	9183 (8524)	4098 (3859)

authors used profile template fitting (Laidler et al. 2007) to provide uniform photometry and spectral energy distributions (SEDs) for each galaxy at wavelengths spanning 0.4–1.6 μm , supplemented by ground-based data (for description, see Guo et al. 2013) and *Spitzer* photometry (3.6–8.0 μm) from the S-CANDELS survey (Ashby et al. 2015). Each photometric object was assigned a flag (Phot-Flag) to identify plausible issues using a robust automated routine described in Galametz et al. (2013). We use PhotFlag = 0 to remove objects with contaminated photometry due to nearby stars, image artefacts or proximity to the F160W coverage edges. This cut removes ~ 3 per cent–5 per cent of raw photometric sources depending on the field. We also use the stellarity index from *SEXTRACTOR* (Class_star ≥ 0.95) to eliminate bright star-like sources. We estimate that this additional cut removes active compact galaxies that makeup ~ 1.3 per cent of our total desired mass-limited sample. We note that including these galaxies has no significant impact on our conclusions. We tabulate the total raw and good photometric source counts for the five CANDELS fields in Table 1.

2.1.2 Redshifts and stellar masses

We use the CANDELS team photometric redshift and stellar mass catalogues available for each field. For the CANDELS UDS and GOODS-S fields, the redshifts are published in Dahlen et al. (2013), and the masses are found in Santini et al. (2015). For the remaining fields, we use the catalogues: GOODS-N (Barro et al., in preparation), COSMOS (Nayyeri et al. 2017), and EGS (Stefanon et al. 2017). As discussed extensively in Dahlen et al. (2013), photometric redshift probability distribution functions $P(z)$ were computed for each galaxy by fitting the SED data. This exercise was repeated by six participants (ID 4, 6, 9, 11, 12, and 13 in Dahlen et al. 2013) who performed SED fitting using different codes (EAZY, HYPERZ) and template sets (BC03, PEGASE, EAZY). Additional detailed discussion on individual code functionality and their respective fitting priors can be found in Dahlen et al. (2013). A team photometric redshift (z_{phot}) was computed for each source equal to the median of the six $P(z)$ peak redshifts. When compared to a known spectroscopic sample, these photometric redshifts have an outlier removed rms scatter $\sigma_z \sim 0.029$ (see Dahlen et al. 2013 for definition). Additionally, spectroscopic redshifts (z_{spec}) are also available for small subsets of galaxies in each field. The best available redshift z_{best} is catalogued as either the team z_{phot} or the good quality z_{spec} measurements when available, which are defined by the flag $q_{\text{zspec}} = 1$ (Dahlen et al. 2013). Note that the compilation of redshifts included in our analysis sample does not include grism redshifts. We limit our selection of massive galaxies to $0.5 \leq z_{\text{best}} \leq 3.0$, and we employ a redshift bin size $\Delta z = 0.5$ to probe evolution between 5 and 11 Gyr ago using

five roughly equal comoving volumes ranging between 7×10^5 and $1.3 \times 10^6 \text{ Mpc}^3$. We exclude redshifts $z_{\text{best}} < 0.5$ since this volume is ~ 10 times smaller ($\sim 1.3 \times 10^5 \text{ Mpc}^3$).

The stellar masses (M_{stellar}) were estimated for each source by fitting the multiband photometric data to SED templates with different stellar population model assumptions¹ fixed to the object’s z_{best} . The team stellar mass (see Santini et al. 2015; Mobasher et al. 2015) for each source is chosen as the median of the estimates based on the same assumptions of initial mass function (IMF, Chabrier 2003) and stellar population templates (Bruzual & Charlot 2003). Using the median mass estimate, we select a mass-limited ($M_{\text{stellar}} \geq 5 \times 10^9 M_{\odot}$) sample of 14 513 potential companion galaxies in a redshift range $0.5 \leq z_{\text{best}} \leq 3$ (for breakdown, see Table 1). As described in the next section, this provides a sample with high completeness.

2.1.3 $0.5 \leq z \leq 3.0$ sample completeness

We demonstrate the completeness of massive CANDELS galaxies with redshifts $0.5 \leq z \leq 3.0$ by adopting the method introduced in Pozzetti et al. (2010) (also see Nayyeri et al. 2017). Briefly, Pozzetti et al. computes a stellar-mass limit as a function of redshift, above which nearly all the galaxies are observable and complete. They do so by estimating the limiting stellar-mass ($M_{\text{stellar,lim}}$) distributions for the 20 per cent faintest sample population,² where $M_{\text{stellar,lim}}$ of a galaxy is the mass it would have if the apparent magnitude (H_{mag}) is equal to the limiting H -band magnitude (H_{lim}). We estimate the $M_{\text{stellar,lim}}$ by following Nayyeri et al. relation between the observed galaxy M_{stellar} and its $M_{\text{stellar,lim}}$ as $\log_{10}(M_{\text{stellar,lim}}) = \log_{10}(M_{\text{stellar}}) + 0.4(H_{\text{mag}} - H_{\text{lim}})$ (see Nayyeri et al. 2017) and use the published H -band 5σ limiting magnitudes (Grogin et al. 2011; Koekemoer et al. 2011; Galametz et al. 2013; Nayyeri et al. 2017; Stefanon et al. 2017).

In Fig. 1, we show the normalized cumulative distributions of $M_{\text{stellar,lim}}$ for the 20 per cent faintest CANDELS $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ galaxy samples³ in narrow ($\Delta z = 0.25$) redshift slices at

¹ Each model is defined by a set of stellar population templates, IMF, SFH, metallicity, and extinction law assumptions; see Mobasher et al. (2015).

² By considering the 20 per cent faintest galaxy sample of the apparent magnitude distribution at each redshift bin, only those galaxies with representative mass-to-light ratios close to the H_{lim} are used towards estimating the $M_{\text{stellar,lim}}$ (see Pozzetti et al. 2010, for additional details).

³ We compute the distributions independently for the five CANDELS fields and present the mean of them at each redshift slice. We find that the behaviour of individual field distributions is not significantly different from each other and with the mean distribution.




Figure 1. Stellar-mass completeness of the $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ CANDELS galaxy sample. We show the normalized cumulative distributions of the limiting stellar masses ($M_{\text{stellar,lim}}$) for the 20 per cent faintest galaxies of the desired mass-limited sample (dashed lines), colour-coded according to their respective redshift slices ($\Delta z = 0.25$) at $z \gtrsim 1$ (see Section 2.1.3 text for details). We show our desired mass limit in solid vertical black line and we mark the 90 per cent completeness in solid horizontal black line. At the highest redshift slice ($2.75 < z < 3$), we find that 90 per cent of the galaxies with $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ have their limiting stellar masses smaller than the desired major companion mass limit, implying that the CANDELS mass-limited sample of $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ galaxies is at least 90 per cent in the desired redshift range of this study $0.5 \leq z \leq 3.0$.

$z > 1$. At all redshift bins up to $z = 2.25$, we find that all the galaxies in our mass-limited sample have $M_{\text{stellar}} > M_{\text{stellar,lim}}$, which implies 100 per cent completeness. At redshifts $2.25 < z < 2.75$ and $2.75 < z < 3$, we find that the desired sample selection is > 95 per cent complete and 90 per cent complete, respectively. Additionally, we test the impact of surface brightness on the measured stellar-mass completeness by analysing the *effective* H -band surface brightness (SB_H) distributions of our desired mass-limited $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ galaxy sample at five redshift bins between $0.5 < z < 3$. We use an H -band surface brightness limit $SB_H = 26.45 \text{ mag arcsec}^{-2}$ based on the model–galaxy recovery simulations by Man et al. (2016) and find that 100 per cent and > 95 per cent of our desired galaxies have $SB_H < 26.45 \text{ mag arcsec}^{-2}$ at redshifts $0.5 < z < 2$ and $2 < z < 3$, respectively. This implies that even the population that constitutes *lowest* 10 per cent of the SB_H distribution (low surface brightness galaxies; hereby LSB galaxies) in our desired sample can be robustly detected up to $z = 3$. As the LSB galaxies only make up a small fraction (less than 10 per cent) of our desired mass-limited sample, we expect that a smaller completeness among these LSB galaxies will not have an significant impact on the close-pair statistics presented in this study. These tests permit us to robustly search for major companions associated with $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.3$ galaxies unaffected by significant incompleteness. We include the breakdown of $N_m = 5698$ massive galaxies per CANDELS field in Table 1.

2.2 SDSS

2.2.1 Redshifts and stellar masses

To anchor evolutionary trends to $z \sim 0$, we employ redshifts and stellar masses from Sample III of the SDSS Group Catalogue described in Yang et al. (2007). Briefly, this catalogue contains

$\sim 400\,000$ galaxies spanning a redshift range $0.01 < z < 0.2$ from the $\sim 4500 \text{ deg}^2$ sky coverage of the SDSS DR4 (Adelman-McCarthy et al. 2006). Yang et al. computed $(g - r)$ colour-based M_{stellar} estimates using the Bell et al. (2003) SED-fitting-based mass-to-light ratio calibrations and K -corrections from the New York University Value-Added Galaxy Catalog (Blanton et al. 2005). For consistency, these masses were corrected by -0.1 dex to convert from a ‘diet’ Salpeter IMF to a Chabrier (2003) IMF basis as in CANDELS. Besides the IMF, Bell et al. assumed similar exponentially declining star formation histories (SFHs) as the CANDELS team M_{stellar} participants, but used PEGASE stellar population models (Fioc & Rocca-Volmerange 1997) in contrast to Bruzual & Charlot (2003), respectively (for details, see Mobasher et al. 2015). However, de Jong & Bell (2007) explored the impact of these model assumptions and found that both PEGASE and Bruzual & Charlot (2003) yield similar results in terms of Bell et al. colour and mass-to-light ratio calibrations. In addition for a sample of galaxies with SDSS+GALEX photometry, Moustakas et al. (2013) found good agreement between SED fitting-derived stellar masses and independent SDSS photometry-based estimates. Hence, we conclude that the CANDELS and SDSS stellar mass estimates are not systematically different.

We select Sample III galaxies within a redshift range $0.03 \leq z \leq 0.05$ and sky area 1790 deg^2 ($RA = 100^\circ - 210^\circ$ and $Dec. = 17^\circ - 69^\circ$) to match the CANDELS sample in volume and resolution. Using these cuts, we find 9183 galaxies with $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$. We present the SDSS selection information in Table 1. We are aware of more recent data sets than the SDSS-DR4; e.g. the SDSS-DR7 (Abazajian et al. 2009) has an improvement in photometric calibration from 2 per cent (DR4) to 1 per cent (DR7). However, owing to the contribution from ~ 20 per cent random and ~ 25 per cent model dependent systematic uncertainties for Bell et al. M_{stellar} estimates, we argue that these small photometric improvements have no significant impact on our results. Hence, we use the SDSS-DR4 because it is readily available and it meets our volume and resolution requirements.

2.2.2 $z \sim 0$ sample completeness

The Yang et al. (2007) sample is magnitude-limited ($r < 17.77 \text{ mag}$) due to the SDSS spectroscopic target selection; this provides a ~ 90 per cent z_{spec} completeness (Strauss et al. 2002). Yang et al. included additional redshifts from supplementary surveys to improve the incompleteness due to spectroscopic fibre collisions (Blanton et al. 2003). As such, our $z \sim 0$ sample selection has 92.8 per cent z_{spec} completeness (Table 1). Nevertheless, several merger studies demonstrate that the SDSS spectroscopic incompleteness grows with decreasing galaxy–galaxy separation (McIntosh et al. 2008; Weston et al. 2017). We account for this issue and provide detailed corrections in Section 3.2. In addition, we demonstrate the stellar mass completeness of $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ galaxies by employing the method by Cebrián & Trujillo (2014). In Fig. 2, we show the stellar mass distributions for narrow redshift intervals ($\Delta z = 0.006$) at $z \leq 0.05$. We find that the $r < 17.77 \text{ mag}$ limit produces a turnover in counts at different masses as a function of redshift. At $z \leq 0.05$, the mass at which the distributions turn over (become incomplete) is well below our limit of $\log_{10}(M_{\text{stellar}}/M_{\odot}) = 9.7$. This indicates that our mass-limited sample is highly complete for selecting possible major companions in a complete sample of $N_m = 4098$ massive galaxies with $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.3$ and $0.03 \leq z \leq 0.05$ (see Table 1).




Figure 2. Stellar mass distributions for galaxies in the SDSS group catalogue (Yang et al. 2007) for four narrow redshift slices (see the legend). The stellar mass at which the mass distribution turns over owing to the $r < 17.77$ mag criteria for the SDSS spectroscopic target selection (Strauss et al. 2002) is given by the vertical dashed lines. The solid black line indicates our desired stellar mass limit of the companion galaxies ($M_{\text{stellar}} = 5 \times 10^9 M_{\odot}$). We find that companion galaxies are highly complete at $z \leq 0.05$.

2.3 Selection of massive galaxies hosting major companions

2.3.1 Projected separation

With our well-defined mass-limited samples for CANDELS and SDSS in hand, we start by identifying the massive ($M_{\text{stellar}} \geq 2 \times 10^{10} M_{\odot}$) galaxies hosting a major projected companion satisfying $1 \leq M_1/M_2 \leq 4$ and a projected physical separation of $5 \text{ kpc} \leq R_{\text{proj}} \leq 50 \text{ kpc}$. The choice of $R_{\text{proj}} \leq 50 \text{ kpc}$ is common in close-pair studies (Patton & Atfield 2008; Lotz et al. 2011; de Ravel et al. 2011) which is supported by the numerical simulation results showing that major bound companions with this separation will merge within $\lesssim 1 \text{ Gyr}$. Additionally, source blending from smaller separations ($\lesssim 1.2 \text{ kpc}$) can cause incompleteness at $z \gtrsim 0.04$ for SDSS and at $z \gtrsim 2.5$ for CANDELS. Thus, we adopt a lower limit of $R_{\text{proj}} = 5 \text{ kpc}$ ($\sim 4 \times$ the resolution), which also corresponds to the typical sizes of $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ galaxies at $2.0 \leq z \leq 2.5$. In summary, we find $N_{\text{proj}} = 318$ and $N_{\text{proj}} = 2451$ unique (i.e. duplicate resolved) massive galaxies hosting major projected companions in SDSS ($0.03 \leq z \leq 0.05$) and CANDELS (total of all five fields at $0.5 \leq z \leq 3.0$), respectively. We tabulate the breakdown of N_{proj} by redshift per each CANDELS field in Table 2.

2.3.2 Plausible physical proximity (SDSS)

We note that projected proximity does not guarantee true physical proximity as foreground and background galaxies can be projected interlopers. A common and effective method to define physical proximity is to isolate systems with a small velocity separation, which indicate that the host and companion galaxies are plausibly gravitationally bound. For the SDSS sample, we employ the common criteria $\Delta v_{12} = |v_1 - v_2| \leq 500 \text{ km s}^{-1}$ (e.g. Kartaltepe et al. 2007; Patton & Atfield 2008; Lin et al. 2008), where v_1 and v_2 are the velocities of the host and companion galaxies, respectively. Merger simulations find that systems that satisfy $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ typically merge within 0.5–1 Gyr (e.g. Conselice 2006). Other studies show that close-pair systems with $\Delta v_{12} > 500 \text{ km s}^{-1}$ are not likely

to be gravitationally bound (e.g. Patton et al. 2000; De Propriis et al. 2007). However, owing to spectroscopic-redshift incompleteness (see Section 2.2.2), we are only able to apply this velocity selection to a subset of galaxies from Section 2.3.1 which have spectroscopic redshifts. In doing so, we find $N_{\text{phys}} = 106$ massive galaxies hosting a major projected companion (in Section 2.3.1) meeting $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ criteria in the SDSS ($0.03 \leq z \leq 0.05$) sample. We describe the statistical correction for missing major companions due to spectroscopic incompleteness in Section 3.2.

2.3.3 Plausible physical proximity (CANDELS)

Most galaxies in the CANDELS catalogues do not have a spectroscopic redshift. Hence, we use a proximity method based on photometric redshifts and their uncertainties (σ_z) to select plausible, physically close companions (e.g. Bundy et al. 2009; Man, Zirm & Toft 2011; Man et al. 2016). As described in Section 2.1.2, each galaxy’s z_{phot} value is the median of the peak values (z_{peak}) of multiple photometric $P(z)$ distributions computed by the CANDELS team. However, the $P(z)$ data were not thoroughly analysed to derive z_{phot} errors for all of the CANDELS fields. Thus, we compute σ_z values from a single participant $P(z)$ data set that produces z_{peak} values that are consistent with the published team z_{best} values. This is necessary to achieve z_{phot} errors that are consistent with the z_{best} and stellar masses (calculated with z_{best}) that we use in this study. We find that the S. Wuyts⁴ photometric redshifts produced the best match to z_{best} (see Appendix) after testing all participant $P(z)$ data. The Wuyts $P(z)$ distributions for each CANDELS galaxy were computed using the photometric redshift code EAZY (Brammer, van Dokkum & Coppi 2008) and PEGASE (Fioc & Rocca-Volmerange 1997) stellar synthesis template models. We optimize⁵ the $P(z)$ for each galaxy and use this distribution to compute the uncertainty (σ_z) defined as the 68 per cent confidence interval of the photometric redshift z_{phot} (see Kodra et al., in preparation for details).

In Fig. 3, we show the photometric redshift uncertainties (σ_z) as a function of z_{best} for each galaxy in our sample ($M_{\text{stellar}} \geq 5 \times 10^9 M_{\odot}$). We find that the σ_z distributions in the CANDELS fields are qualitatively similar to each other. We find the σ_z distributions have small scatter up to $z \sim 1.5$ with their medians typically ranging between $0.02 \leq \sigma_{z,\text{med}} \leq 0.05$, and much larger scatter at $z \gtrsim 1.5$ with the medians ranging between $0.06 \leq \sigma_{z,\text{med}} \leq 0.08$. This large scatter is because the observed filters no longer span the 4000 Å break, which leads to larger uncertainties during template SED fitting (for additional details, see Kodra et al., in preparation). For each CANDELS field, we show the 80 per cent and 95 per cent outlier limits of the redshift normalized error [$\sigma_z/(1 + z_{\text{best}})$] distribution and present their values in Table 3. While the 95 per cent clipping limit rejects extreme outliers typically with $z_{\text{best}} > 1.5$, the 80 per cent limit does a reasonable job representing the upper envelope of the σ_z distribution at all redshifts. Therefore, to exclude galaxies with large z_{phot} errors, we elect to exclude those σ_z above the 80 per cent clipping limit. Hereafter, we define the large-error z_{phot} as *unreliable*.

⁴ Method 13 as specified in Dahlen et al. (2013).

⁵ We shift the $P(z)$ distributions and raise them to a power such that when compared to the test set of spectroscopic redshifts (z_{spec}), the 68 per cent confidence interval of the $P(z)$ should include z_{spec} 68 per cent of the time. A detailed description is given in Kodra et al. (in preparation).




Figure 3. Photometric redshift uncertainties (σ_z) as a function of z_{best} for galaxies with $M_{\text{stellar}} \geq 5 \times 10^9 M_{\odot}$ in each CANDELS field. The σ_z values are the 1σ photometric redshift errors from the optimized $P(z)$ distributions originally derived by S. Wuyts (see the text for details). In each panel, we show the 80 percent and 95 percent outlier clipped limits of the redshift normalized uncertainty $\sigma_z/(1 + z_{\text{best}})$ distribution in solid and dashed lines, respectively. We present these limits in Table 3.

For each galaxy in our CANDELS sample, we adopt the (Bundy et al. 2009, hereafter, B09) redshift proximity criteria given by:

$$\Delta z_{12}^2 \leq \sigma_{z,1}^2 + \sigma_{z,2}^2, \quad (1)$$

where $\Delta z_{12} = (z_{\text{best},1} - z_{\text{best},2})$ is the redshift difference of the host and companion galaxies, and $\sigma_{z,1}$ and $\sigma_{z,2}$ are their photometric redshift errors, respectively. It is important to note that projected pairs containing widely separated galaxies in redshift space that have large z_{phot} errors can satisfy equation (1). Hence, we apply the redshift proximity criteria only to those galaxies with reliable photometric redshifts. In summary, we select $N_{\text{phy}} = 504$ massive galaxies hosting major companions satisfying $5 \text{ kpc} \leq R_{\text{proj}} \leq 50 \text{ kpc}$, $1 \leq M_1/M_2 \leq 4$, and equation (1). We present the breakdown of N_{phy} in each redshift bin per CANDELS field in Table 2.

In Section 3.1, we describe a statistical correction to add back a subset of galaxies excluded because of an unreliable σ_z that could be statistically satisfying the redshift proximity criteria. Additionally, owing to the possibility that some companion galaxies may satisfy the close redshift proximity criterion by random chance, we discuss the statistical correction for random chance pairing in Section 3.3.3. We acknowledge the mismatch between the redshift proximity methods that we employ for the SDSS and CANDELS. In Section 4, we test the impact of this mismatch and find that it does not significantly impact our results and conclusions.

3 FREQUENCY OF MAJOR MERGING

To track the history of major merging, we start by analysing the fraction of massive ($M_{\text{stellar}} \geq 2 \times 10^{10} M_{\odot}$) galaxies hosting a

major companion selected in Section 2.3. The *major companion* fraction⁶ is

$$f_{\text{mc}}(z) = \frac{N_{\text{mc}}(z)}{N_{\text{m}}(z)}, \quad (2)$$

where at each redshift bin, N_{mc} is the number of massive galaxies hosting a major companion after statistically correcting the N_{phy} counts (see Sections 3.1 and 3.2), and N_{m} is the number of massive galaxies. The companion fraction f_{mc} is commonly used in the literature, and is the same as f_{merg} used by Man et al. (2016). We use the samples described in Section 2.3 to derive $N_{\text{mc}}(z)$ separately for the five redshift bins in CANDELS ($0.5 \leq z \leq 3.0$) and for the SDSS $z \sim 0$ anchor. We then discuss the application of a correction to account for galaxies satisfying the companion selection criteria by random chance. Finally, we characterize the companion fraction and use an analytical function to quantify the redshift evolution of f_{mc} during $0 < z < 3$.

3.1 Deriving $N_{\text{mc}}(z)$ for CANDELS

For redshifts $0.5 \leq z \leq 3$, we compute N_{mc} corrected for incompleteness owing to unreliable photometric redshifts as

$$N_{\text{mc}} = N_{\text{phy}} + C_1 N_{\text{proj,unreliable}}, \quad (3)$$

where N_{phy} is the number of massive galaxies with reliable z_{phot} values that host a major companion (Section 2.3.3), $N_{\text{proj,unreliable}}$ is the number of galaxies hosting major projected companions that are excluded because of unreliable z_{phot} values (Section 2.3.1), and C_1 is the correction factor used to statistically add back a subset of

⁶ While the companion fraction is related to the pair fraction, it is important to be clear that it is not the same (see Section 5.1).

Table 2. Detailed breakdown of variables involved in estimating the f_{mc} and $f_{\text{mc,c}}$ at five redshift bins between $0.5 \leq z \leq 3$. Columns: (1) name of the CANDELS field; (2) the CANDELS team z_{best} bin; (3) number count of massive ($\log M_{\text{stellar}}/M_{\odot} \geq 10.3$) galaxies; (4) number of massive galaxies hosting a major projected companion (Section 2.3.1), those of which that have unreliable photometric redshift values are shown in parenthesis; (5) number of massive galaxies with reliable z_{phot} that host a major projected companion satisfying redshift proximity [equation (8)] as described in Section 2.3.3; (6) the correction factor computed using equation (4); (7) the number of massive galaxies hosting a major companion after statistically correcting N_{phy} for incompleteness owing to unreliable z_{phot} values from equation (3); (8) correction factor to account for random chance pairing as described in Section 3.3.3; (9) the fraction of massive galaxies hosting a major companion (*major companion* fraction); and (10) random chance pairing corrected f_{mc} , as described in Section 3.3.3.

Name (1)	Redshift (2)	N_{m} (3)	N_{proj} ($N_{\text{proj, unreliable}}$) (4)	N_{phy} (5)	C_1 (6)	N_{mc} (7)	C_3 (8)	f_{mc} (per cent) (9)	$f_{\text{mc,c}}$ (per cent) (10)
UDS	$0.5 \leq z \leq 1$	256	132 (16)	44	0.38	50		20 ± 5	
	$1 \leq z \leq 1.5$	304	130 (23)	34	0.32	41		14 ± 4	
	$1.5 \leq z \leq 2$	290	130 (30)	25	0.25	33		11 ± 4	
	$2 \leq z \leq 2.5$	216	80 (31)	14	0.29	23		11 ± 4	
	$2.5 \leq z \leq 3$	157	82 (42)	3	0.07	6		4 ± 3	
GOODS-S	$0.5 \leq z \leq 1$	216	107 (21)	32	0.37	40		18 ± 5	
	$1 \leq z \leq 1.5$	252	116 (14)	34	0.33	39		15 ± 4	
	$1.5 \leq z \leq 2$	213	87 (22)	15	0.23	20		9 ± 4	
	$2 \leq z \leq 2.5$	138	56 (14)	3	0.07	4		3 ± 3	
	$2.5 \leq z \leq 3$	123	57 (15)	5	0.12	7		6 ± 4	
GOODS-N	$0.5 \leq z \leq 1$	333	195 (33)	56	0.35	67		20 ± 4	
	$1 \leq z \leq 1.5$	278	140 (27)	40	0.35	50		18 ± 4	
	$1.5 \leq z \leq 2$	209	99 (20)	15	0.19	19		9 ± 4	
	$2 \leq z \leq 2.5$	191	83 (17)	6	0.09	8		4 ± 3	
	$2.5 \leq z \leq 3$	122	61 (16)	5	0.11	7		6 ± 4	
COSMOS	$0.5 \leq z \leq 1$	448	244 (38)	40	0.19	47		11 ± 3	
	$1 \leq z \leq 1.5$	270	128 (37)	1	0.01	1		1 ± 1	
	$1.5 \leq z \leq 2$	350	157 (75)	15	0.18	29		8 ± 3	
	$2 \leq z \leq 2.5$	153	86 (54)	0	0.0	0		0 ± 0	
	$2.5 \leq z \leq 3$	86	36 (22)	6	0.43	15		18 ± 8	
EGS	$0.5 \leq z \leq 1$	224	120 (19)	21	0.21	25		11 ± 4	
	$1 \leq z \leq 1.5$	304	137 (19)	36	0.31	42		14 ± 4	
	$1.5 \leq z \leq 2$	331	171 (50)	39	0.32	55		17 ± 4	
	$2 \leq z \leq 2.5$	167	94 (18)	23	0.3	28		17 ± 6	
	$2.5 \leq z \leq 3$	67	35 (13)	2	0.09	3		5 ± 5	
All fields	$0.5 \leq z \leq 1$	1477	798 (127)	193	0.29	230	0.15	16 ± 2	13 ± 2
	$1 \leq z \leq 1.5$	1408	651 (120)	145	0.27	173	0.17	12 ± 2	10 ± 2
	$1.5 \leq z \leq 2$	1393	644 (197)	109	0.24	155	0.18	11 ± 2	9 ± 2
	$2 \leq z \leq 2.5$	865	399 (134)	46	0.17	63	0.25	7 ± 2	5 ± 2
	$2.5 \leq z \leq 3$	555	271 (108)	21	0.13	38	0.22	7 ± 2	5 ± 2

Table 3. Photometric redshift uncertainty outlier limits that are used to determine reliable z_{phot} values for each CANDELS field. Columns: (1) name of the CANDELS field; (2) and (3) the 80 per cent and 95 per cent outlier clipped limits of the redshift normalized uncertainty $\sigma_z/(1 + z_{\text{best}})$ distributions for galaxies in the mass-limited ($M_{\text{stellar}} \geq 5 \times 10^9 M_{\odot}$) sample for redshifts $0.5 \leq z \leq 3.0$ as shown in Fig. 3.

Name (1)	80 per cent limit (2)	95 per cent limit (3)
UDS	0.033	0.051
GOODS-S	0.038	0.058
GOODS-N	0.04	0.061
COSMOS	0.024	0.037
EGS	0.041	0.062

excluded galaxies that are expected to satisfy the redshift proximity criteria we employ. We estimate C_1 as

$$C_1 = \frac{N_{\text{phy}}}{N_{\text{proj}} - N_{\text{proj, unreliable}}} \quad (4)$$

To study $f_{\text{mc}}(z)$ from the overall sample and also its field-to-field variations, we calculate N_{mc} for five $\Delta z = 0.5$ bins separately for each CANDELS field and for the total sample using counts tabulated in Table 2. For example, in the $0.5 \leq z \leq 1$ bin, CANDELS contains 671 massive galaxies with reliable z_{phot} values hosting a major projected companion. This results in $C_1 = 193/671 = 0.29$ and a corrected count of $N_{\text{mc}} = 230$; i.e. we add back 29 per cent of the previously excluded galaxies at these redshifts. We tabulate C_1 and N_{mc} values in Table 2. We also note no significant difference in the computed N_{mc} values whether we use an 80 per cent or 95 per cent σ_z clipping limit (Section 2.3) to remove unreliable photometric redshifts.

3.2 Deriving $N_{\text{mc}}(z)$ for SDSS

To achieve an accurate low-redshift anchor for the fraction of massive galaxies hosting a major companion meeting our $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ velocity separation criterion, we calculate N_{mc}




Figure 4. The number of massive galaxies hosting a major companion in projected pairs as a function of projected separation from the SDSS: total (N'_{proj} ; bold line), the subset of galaxies with no spectroscopic-redshift information ($N'_{\text{proj,nospec}}$; dashed line), and the subset of galaxies satisfying $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ (N'_{phy} ; thin line). The *prime* signifies that the quantities are derived using a larger SDSS sample spanning $0.01 \leq z \leq 0.05$ and $\sim 4000 \text{ deg}^2$ (see Section 3.2) for nine R_{proj} bins between 5 and 50 kpc. The correction factor C_2 per R_{proj} bin (equation 6) is given by the red circles connected by a red solid line. The error bars represent 95 per cent binomial confidence of C_2 .

corrected for the SDSS spectroscopic incompleteness that varies with projected separation using $\Delta R_{\text{proj}} = 5 \text{ kpc}$ bins as follows:

$$N_{\text{mc}} = N_{\text{phy}} + \sum_{i=1}^9 (C_2 N_{\text{proj,nospec}})_i. \quad (5)$$

For each of nine bins between $R_{\text{proj}} = 5\text{--}50 \text{ kpc}$, we compute a correction factor $C_{2,i}$ necessary to add back a statistical subset of the $N_{\text{proj,nospec}}$ galaxies in the bin that lack spectroscopic redshifts but that we expect to satisfy the $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ criterion. Following the same logic as in equation (4), we estimate this correction at each ΔR_{proj} bin based on the counts of spectroscopic galaxies hosting a major projected companion and the fraction that satisfy $\Delta v_{12} \leq 500 \text{ km s}^{-1}$. Owing to our well-defined sample volume ($0.03 < z < 0.05$), the total sample of spectroscopic hosts with plausible physical companions (Section 2.3.2) is limited to $N_{\text{phy}} = 109$ over the nine separation bins. To reduce random errors from small number statistics, we use a larger redshift range ($0.01 \leq z \leq 0.05$) and SDSS footprint ($\sim 4000 \text{ deg}^2$), to calculate the correction factor at each ΔR_{proj} bin:

$$C_{2,i} = \left(\frac{N'_{\text{phy}}}{N'_{\text{proj}} - N'_{\text{proj,nospec}}} \right)_i. \quad (6)$$

In Fig. 4, we plot C_2 and the factors in equation (5) as a function of R_{proj} using this larger sample (emphasized with a *prime*). For example, in the $20 \leq R_{\text{proj}} \leq 25 \text{ kpc}$ separation bin, we find 119 massive galaxies hosting a major projected companion, of which 35 have small velocity separations and 44 lack spectroscopic redshifts. This results in a 47 per cent correction ($C_2 = 35/75 = 0.47$) at this separation. We find the probability for a small-separation pair ($R_{\text{proj}} = 5 \text{ kpc}$) to satisfy $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ (despite lacking the redshift information) is 85 per cent ($C_2 = 0.85$) and this rapidly decreases to ~ 30 per cent ($C_2 \sim 0.3$) at $R_{\text{proj}} = 30 \text{ kpc}$, and remains statistically constant between $R_{\text{proj}} = 30\text{--}50 \text{ kpc}$. This correction is important since the spectroscopic incompleteness $N'_{\text{proj,nospec}}/N'_{\text{proj}}$

ranges from >0.6 ($\sim 5\text{--}10 \text{ kpc}$) to 0.2 ($\sim 45\text{--}50 \text{ kpc}$) over the separations we probe, which is in agreement with trends published in fig. 2 from Weston et al. (2017).

3.3 Redshift evolution of major merging frequency

We use the corrected counts of massive galaxies hosting a major companion to compute the companion fraction (f_{mc} ; see equation 2) in the SDSS and CANDELS. We compare field-to-field variations of f_{mc} in CANDELS, quantify the redshift evolution of the f_{mc} from $z = 3$ to 0, and measure the impact of random chance pairing on the observed major companion fraction evolutionary trends.

3.3.1 Field-to-field variations

In Fig. 5(a), we plot $f_{\text{mc}}(z)$ for the combined CANDELS fields and compare this with the individual fractions from each field at each redshift; these are also tabulated in Table 2. Despite noticeable variations between the fractions derived from each CANDELS field owing to small-number statistics, we find fair agreement between multiple fields at each redshift. We note that the combined CANDELS sample and three individual fields (UDS, GOODS-S, and GOODS-N) show consistent trends with the highest merger fractions at $z \sim 1$, which then steadily decrease with increasing redshift. The EGS and COSMOS companion fractions exhibit different behaviour with redshift, the former peaks at $z \sim 2$ while the latter has no trend with redshift owing to a lack of galaxies hosting major companions in two different redshift bins. We compute the cosmic variance (σ_{CV}) on the combined CANDELS $f_{\text{mc}}(z)$ values using the prescription by Moster et al. (2011). For $\log_{10}(M_{\text{stellar}}/M_{\odot}) \gtrsim 10$ galaxies at $0.5 < z < 3.0$ populating the five CANDELS fields each with an area of 160 arcmin^2 such that their cumulative area matches that of the total CANDELS coverage ($5 \times 160 = 800 \text{ arcmin}^2$), we find that the number counts of galaxies hosting major companions have σ_{CV} ranging from 11 per cent ($z = 0.75$) to 18 per cent ($z = 2.75$).⁷ While most of the individual CANDELS-field fractions are consistent with the σ_{CV} within their large uncertainties (owing to small sample size), few f_{mc} values (e.g. at $z > 1.5$ for the COSMOS and EGS fields; see Fig. 5a) are significantly above the possible cosmic-variance limits.

3.3.2 Analytical fit to the major companion fraction evolution

To characterize the redshift evolution of the companion fraction during $0 < z < 3$, we anchor the combined CANDELS $f_{\text{mc}}(z)$ measurements to the SDSS-derived data point at $z \sim 0$. As shown in Fig. 5(a), the low-redshift fraction is $\sim 3 \times$ lower than the maximum $f_{\text{mc}} \sim 0.16$ value at $0.5 < z < 1$, which then decreases to $f_{\text{mc}} \sim 0.07$ at $2.5 < z < 3$. This suggests a turnover in the incidence of merging sometime around $z \sim 1$, in agreement with some previous studies (Conselice et al. 2003; Conselice, Rajgor & Myers 2008). Previous close-pair-based studies at $z \sim 0$ find fractions $f_{\text{mc}} \sim 2 \text{ per cent} \pm 0.5 \text{ per cent}$ (Patton et al. 2000; Patton & Atfield 2008; Domingue et al. 2009), but they used criteria that are different from our fiducial selection. Similarly, many empirical, close-pair-based studies in the literature broadly agree that f_{mc} rises at $0 < z < 1.5$ but with a range of evolutionary forms

⁷ We take into account that σ_{CV} is smaller for multiple, widely separated fields when compared to the σ_{CV} of a single contiguous field. For additional details, see Moster et al. (2011).




Figure 5. (a) The redshift evolution of the *major companion* fraction f_{mc} shown for the five CANDELS fields UDS (star), GOODS-S (left triangle), GOODS-N (right triangle), COSMOS (pentagon), and EGS (cross). The combined CANDELS fractions in five redshift bins (circles) and the SDSS low-redshift anchor (square) include 95 per cent binomial confidence limit error bars. To place our finding in the context of common, close-pair-based evolutionary trends found in the literature, we plot the shaded region (red) encompassing a common range of power-law slopes $f_{mc} = 0.06(1+z)^{1-2}$ at $0 < z < 1.5$. (b) The random chance corrected fractions ($f_{mc,c}$) for the five CANDELS Δz bins (open circles) are compared with the $f_{mc}(z)$ from (a). For $f_{mc,c}$, the binomial errors and scatter of C_3 (see Section 3.3.3) are added in quadrature. Best-fitting curves to the companion fraction (f_{mc}) evolution data [see equation (7) and Section 3.3.2 for details] are shown in solid (f_{mc}) and dashed ($f_{mc,c}$) lines, respectively. In the case of SDSS, since the correction $C_3 \sim 0.01$, we only plot f_{mc} for simplicity. From this figure, we conclude that the major companion fraction increases strongly from $z \sim 0$ to 1, and decreases steeply towards $z \sim 3$ (see the text for details).

$(1+z)^{1-2}$ owing to varying companion selection criteria (for discussion, see Lotz et al. 2011). After normalizing for these variations, we note that our SDSS-based f_{mc} is in good agreement with previous close-pair-based estimates, and our rising trend (see shaded region, Fig. 5a) between $0 < z < 1.5$ is in broad agreement with other empirical trends. In Section 5, we will present detailed comparisons to other studies by recomputing f_{mc} based on different companion selection choices that match closely with others.

All studies that measured redshift evolution of merger frequency at $0 < z < 1.5$, irrespective of the methodology, have used the power-law analytical form $f(z) \propto (1+z)^m$ to represent the best fit of $f(z)$. This functional form cannot be used to represent the observed rising and then decreasing $f_{mc}(z)$ for redshift ranges $0 < z < 3$. Therefore, following Conselice et al. (2008) and initially motivated by Carlberg (1990), we use a modified power-law exponential function given by

$$f_{mc}(z) = \alpha(1+z)^m \exp^{\beta(1+z)}. \quad (7)$$

As demonstrated in Fig. 5(a), this analytic function provides a good fit to the observed evolution. The best-fitting curve to the fractions derived from the SDSS and CANDELS measurements from our fiducial companion selection criteria has parameters $\alpha = 0.5 \pm 0.2$, $m = 4.5 \pm 0.8$, and $\beta = -2.2 \pm 0.4$. We note that we will apply this fitting function for different companion selection choices throughout our comparative analysis described in Sections 4 and 5.

3.3.3 Correction for random chance pairing

Finally, we note that a subset of massive galaxies hosting a major companion (N_{mc}) can satisfy the companion selection criteria by random chance. To account for this contamination, we apply a statistical correction and recompute the counts for the combined CANDELS sample per redshift bin as $N_{mc,c} = N_{mc}(1 - C_3)$ in each redshift bin. To compute C_3 , we generate 100 simulated Monte Carlo

(MC) randomized data sets.⁸ We define C_3 in each redshift bin as the ratio of massive galaxy number counts hosting major companions which satisfy our projected separation and photometric redshift proximity criteria in the MC data sets (i.e. by random chance) to the measured N_{mc} (Sections 3.1 and 3.2). For example, in redshift bin $1 < z < 1.5$, we find that 17 per cent of $N_{mc} = 173$ galaxies statistically satisfy the companion selection criteria by random chance. We tabulate C_3 values at each redshift for CANDELS in Table 2. We repeat this process for the SDSS ($0.03 < z < 0.05$) and find a very small correction of ~ 1 per cent ($C_3 \sim 0.01$). This demonstrates the very low probability for two SDSS galaxies to satisfy both the close projected separation and stringent spectroscopic-redshift proximity ($\Delta v_{12} \leq 500 \text{ km s}^{-1}$) criteria.

In Fig. 5(b), we compare the random chance corrected fractions $f_{mc,c} = N_{mc,c}/N_m$ at each redshift bin from CANDELS, to the uncorrected f_{mc} values copied from the left-hand panel. Owing to the insignificant 1 per cent correction at $z \sim 0$, we anchor both the corrected and uncorrected fits to the same SDSS data point. We find that $f_{mc,c}(z)$ follows the same evolutionary trend as $f_{mc}(z)$, in which the best-fitting curve rises to a maximum fraction at $z \sim 1$ and then steadily decreases to $z = 3$. At all redshifts, $f_{mc,c}$ is within the statistical errors of f_{mc} . The qualitatively similar redshift evolutionary trends of $f_{mc}(z)$ and $f_{mc,c}(z)$ is due to the nearly redshift independent amount of statistical correction for random pairing ($|df|/f_{mc} \sim 20$ per cent) at $1 < z < 3$. This is because of the nearly invariant angular scale in this redshift range, which results in similar random chance pairing probabilities. We note that some previous close-pair-based studies have applied this random chance

⁸ We generate these data sets by randomizing the positions of each galaxy in the $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ mass-limited sample and repeating the selection process in Section 2.3.

correction (Kartaltepe et al. 2007; Bundy et al. 2009), while others have not (e.g. Man et al. 2012).

4 IMPACT OF CLOSE-COMPANION SELECTION CRITERIA ON EMPIRICAL MAJOR COMPANION FRACTIONS

So far, we have discussed the derivation and redshift evolution of the major companion fraction f_{mc} based on our *fiducial* selection criteria described in Section 2.3. As illustrated in Table 5, previous studies have employed a variety of criteria to select companions. In this section, we study the impact of different companion selections on $f_{mc}(z)$ derived from our sample. We systematically vary each criterion (projected separation, redshift proximity, and stellar MR versus FR) individually, while holding the other criteria fixed to their fiducial values. Then, we compare each recomputed $f_{mc}(z)$ to the fiducial result from Fig. 5. In Section 3.3.3, we show that applying a statistical correction for random chance pairing will produce companion fractions that are ~ 20 per cent lower at each redshift interval between $0.5 \leq z \leq 3$. Therefore, we focus the following comparative analysis on uncorrected fractions and note that our qualitative conclusions are robust to whether or not we apply this correction.

4.1 Projected separation

To quantify the impact that changing *only* the criterion for companion projected separation will have on $f_{mc}(z)$, we compare fractions based on the fiducial $R_{proj} = 5\text{--}50$ kpc selection to those derived from two common criteria: $R_{proj} = 14\text{--}43$ kpc (e.g. Lin et al. 2008; Man et al. 2016) and $5\text{--}30$ kpc (e.g. Patton & Atfield 2008; Man et al. 2012). For each case, we hold all other companion selection criteria fixed such that we are strictly comparing fractions of major (stellar mass-selected) companions in close redshift proximity that are found within projected annular areas of $2/3$ and $1/3$ the fiducial selection window ($2475\pi \text{ kpc}^2$), respectively.

In Fig. 6, we plot the fiducial and non-fiducial $f_{mc}(z)$ data and their best-fitting curves. The key result from this figure is the observed redshift evolution of major companion fractions shown in Fig. 5 are qualitatively robust to changes in the projected separation criterion. For each case, we find the fractions increase from low redshift to $z \sim 1$, then turn over and decrease fairly steadily to $z = 3$. Quantitatively, the non-fiducial fractions are well fit by the same $f_{mc}(z)$ functional form as the fiducial results (see Table 4). Owing to the size of the best-fitting confidence intervals, the redshifts of the peak fractions are statistically equivalent. Larger samples of massive galaxy–galaxy pairs at redshifts $0.5 < z < 1.5$ are needed to place stringent constraints on the peak or turnover redshift.

Besides the overall f_{mc} trends with redshift, we find that both smaller projected separation criteria select smaller f_{mc} values than the fiducial selection does, as we expect. Despite being a factor of two smaller in projected area, the conservative $R_{proj} = 5\text{--}30$ kpc criterion coincidentally produces fractions that are statistically matched to those from the larger $R_{proj} = 14\text{--}43$ kpc selection criterion at most redshift intervals. Only the $0.5 < z < 1$ bin has unequal companion fractions between the two non-fiducial criteria. The coincidental finding of similar fractions using different projected separation criteria is consistent with an increased probability of physical companions at smaller projected separations (Bell et al. 2006b; Robaina et al. 2010).




Figure 6. Comparison of the redshift evolution of the major companion fractions based on three projected separation criteria: $R_{proj} = 5\text{--}50$ kpc (fiducial; black symbols, solid line), $R_{proj} = 14\text{--}43$ kpc (grey symbols, dashed line), and $R_{proj} = 5\text{--}25$ kpc (open symbols, dot-dashed line). Best-fitting curves for each $f_{mc}(z)$ spanning $0 < z < 3$ are a modified power-law exponential (equation 7; see Section 3.3.2 for details). The low-redshift fractions are from the SDSS (squares) and the $z > 0.5$ fractions are from CANDELS (circles). The error bar on each f_{mc} data point represents 95 per cent binomial confidence limit. The non-fiducial f_{mc} data points are offset by a small amount within each redshift bin for clarity.

4.2 Redshift proximity

Following a similar methodology as in Section 4.1, we quantify the impact that changing *only* the criterion for redshift proximity will have on $f_{mc}(z)$. As described in Section 2.3.2, we apply selections based on photometric redshifts and their uncertainties only to CANDELS data, and we anchor our evolutionary findings to stringent velocity-separation-based companion fractions at $z \sim 0$ from the SDSS. As such, we first compare CANDELS fractions ($0.5 < z < 3$) based on our fiducial B09 selection given in equation (1) to fractions based on two other related methods. Then, we demonstrate that the low-redshift fraction from the SDSS changes very little ($\Delta f / f_{fid} < 5$ per cent) if we select major companions using simulated photometric redshift errors that are similar in quality to the CANDELS σ_z values, rather than use our fiducial selection criterion ($\Delta v_{12} \leq 500 \text{ km s}^{-1}$).

In Fig. 7, we plot $f_{mc}(z)$ data and best-fitting curves for our fiducial selection and two additional methods. In each non-fiducial case, we hold the projected separation and MR selection criteria fixed to our fiducial choices. The first method is a simple overlapping of the host and companion galaxy 1σ photometric redshift errors

$$|\Delta z_{12}| \leq \sigma_{z,1} + \sigma_{z,2}. \quad (8)$$

This criterion is modified from Man et al. (2012) who used redshift overlap at the 3σ level. As with fiducial selection, we apply this criterion only to those galaxies with reliable photometric redshifts (see Section 2.3.2). Second, we incorporate available spectroscopic redshifts for the host and/or companion galaxies from CANDELS into a ‘hybrid’ of our fiducial B09 redshift proximity selection as follows:

- (i) if both galaxies have z_{spec} data, then the pair must satisfy $\Delta v_{12} \leq 500 \text{ km s}^{-1}$;

Table 4. Modified power-law exponential function parameter values for best-fitting models to major companion fraction evolution $f_{\text{mc}}(z)$ based on different selection criteria; our fiducial criteria are given in bold. Columns: (1) major companion definition – stellar MR $1 \leq M_1/M_2 \leq 4$, or H-band FR $1 \leq F_1/F_2 \leq 4$; (2) projected separation R_{proj} criterion; (3) redshift proximity method applied to CANDELS samples – equation (7) adopted from B09, equation (8) modified from Man et al. (2012), or ‘Hybrid’ version of B09 (see the text for details); and (4)–(6) best-fitting parameter values (α , m and β) and their 1σ confidence limits for the model function given in equation (7).

Major companion definition (1)	R_{proj} (2)	Redshift proximity (3)	α (4)	m (5)	β (6)
MR	5–50 kpc	B09	0.5 ± 0.2	4.6 ± 0.9	-2.3 ± 0.5
MR	14–43 kpc	B09	0.7 ± 0.5	6.2 ± 1.4	-3.1 ± 0.8
MR	5–30 kpc	B09	0.4 ± 0.1	5.4 ± 0.8	-2.6 ± 0.4
MR	5–50 kpc	modM12	0.5 ± 0.2	4.6 ± 0.9	-2.2 ± 0.5
MR	5–50 kpc	hybB09	0.4 ± 0.1	3.8 ± 0.7	-1.9 ± 0.4
FR	5–50 kpc	B09	0.1 ± 0.1	2.1 ± 1.2	-0.5 ± 0.6

Table 5. Compilation of companion selection criteria employed by previous empirical close-pair-based studies. Columns: (1) names of the studies – presented in the order of their mention in Fig. 10; (2) the choice of projected separation R_{proj} annulus; (3) the criterion to choose major companions, where $\mu_{12} = M_1/M_2$ is the stellar MR, FR is given by F_1/F_2 , and the luminosity ratio is given by L_1/L_2 ; (4) the choice of redshift proximity criterion to select companions in plausible close pairs, where Δz_{12} is the photometric redshift difference $z_1 - z_2$, Δv_{12} is the spectroscopic velocity difference, LOS is the statistical correction for line-of-sight projections, and $\text{CDF}(z_1, z_2)$ is the cumulative probability of the galaxies involved in a close pair; (5) the applied selection criterion to select the primary galaxy sample, which is either a stellar mass-limited or flux-limited selection; (6) the redshift range in which the study presented their findings. We divide the table into two parts, where studies based on MRs are above the solid-dashed line and vice versa. Newman et al. (2012) employed a redshift proximity where $\Delta z_{12}/1 + z_1 < 0.1$ for $z < 1$ and $\Delta z_{12}/1 + z_1 > 0.2$ when $z_1 > 1$. The dagger represents that the studies derived a ‘pair’ fraction rather than the companion fraction (f_{mc}), see Section 5.1 for details.

Study (1)	R_{proj} selection (2)	Major companion (3)	Redshift proximity (4)	Primary galaxy selection (5)	Redshift (6)
Ryan+08	5–30 kpc	$1 \leq \mu_{12} \leq 4$	2σ z_{phot} error overlap	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10$	$0.5 < z < 2$
Bundy+09†	5–30 kpc	$1 \leq \mu_{12} \leq 4$	$\Delta z_{12}^2 \leq \sigma_{z,1}^2 + \sigma_{z,2}^2$	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10$	$0.4 \leq z \leq 1.4$
Mundy+17†	5–30 kpc	$1 \leq \mu_{12} \leq 4$	$\text{CDF}(z_1, z_2)$	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10$	$0 < z < 2$
Williams+11	14–43 kpc	$1 \leq \mu_{12} \leq 4$	$ \Delta z_{12} /1 + z_1 < 0.2$	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.5$	$0.4 \leq z \leq 2$
Newman+12	14–43 kpc	$1 \leq \mu_{12} \leq 4$	See caption	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.7$	$0.4 \leq z \leq 2$
Man+16	14–43 kpc	$1 \leq \mu_{12} \leq 4$	Same as Newman+12	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.8$	$0 \leq z \leq 3$
Kartaltepe+07	5–30 kpc	$1 \leq L_1/L_2 \leq 4$	$\Delta v_{12} \leq 500 \text{ km s}^{-1}$	$M_V < -19.8 - 1.0z$	$0.1 \leq z \leq 1.2$
Bundy+09†	5–30 kpc	$1 \leq F_1/F_2 \leq 4$	LOS	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10$	$0.4 \leq z \leq 1.4$
Man+12	5–30 kpc	$1 \leq F_1/F_2 \leq 4$	3σ z_{phot} error overlap	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 11$	$0 \leq z \leq 3$
Bluck+09, +12	5–30 kpc	$1 \leq F_1/F_2 \leq 4$	Line-of-sight correction	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 11$	$1.7 \leq z \leq 3$
Lin+08	14–43 kpc	$1 \leq L_1/L_2 \leq 4$	$\Delta v_{12} \leq 500 \text{ km s}^{-1}$	$-21 < M_B + 1.3z < -19$	$z \leq 1.2$
López-Sanjuan+11†	14–43 kpc	$1 \leq L_1/L_2 \leq 4$	$\Delta v_{12} \leq 500 \text{ km s}^{-1}$	$M_V < -20 - 1.1z$	$z = 0.5, z = 0.8$
Man+16	14–43 kpc	$1 \leq F_1/F_2 \leq 4$	Same as Newman+12	$\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.8$	$0 \leq z \leq 3$

- (ii) or if only one galaxy has z_{spec} data, then z_{spec} must be within $z_{\text{phot}} \pm \sigma_z$ limit of the other galaxy;
- (iii) otherwise use B09 criterion.

As with changing the project separation for selecting close companions, the key takeaway from Fig. 7 is the redshift evolution trends for $f_{\text{mc}}(z)$ are robust to slight modifications in the redshift proximity criterion. Both non-fiducial companion fractions peak in the lowest-redshift CANDELS bin, the decline at $z > 1$ and their redshift evolution anchored to the SDSS $z \sim 0$ fraction are well fit by the same functional form with statistically equivalent best-fitting parameters (see Table 4). Besides the similar evolutionary trends, we notice that the modified Man et al. criterion results in f_{mc} values that are systematically ~ 11 per cent–24 per cent higher than the fiducial fractions between $0.5 < z < 3$. This is expected since simple photometric redshift error overlap (equation 1) is less stringent than a quadrature overlap; e.g. two galaxies with $\sigma_z = 0.06$ must have photometric redshifts within $\Delta z_{12} = 0.12$ compared to 0.085 (fiducial). On the other hand, the hybrid B09 method yields

9 per cent–17 per cent smaller major companion fractions than fiducial at $0.5 < z < 1.5$ owing to the availability of z_{spec} data for a fair fraction of CANDELS galaxies and the stringent Δv_{12} criterion for this subset, but nearly identical fractions at $z > 1.5$ for which the spectroscopic-redshift coverage becomes quite sparse.

We close our redshift proximity analysis by quantifying changes in the $f_{\text{mc}}(0)$ measurement from the SDSS data if we employ either the B09 or modified Man et al. photometric redshift error overlap criteria. We opt to generate simulated z_{phot} and σ_z for the SDSS sample such that these values mimic the quality of the CANDELS photometric data. First, we construct the combined $\sigma_z/(1 + z_{\text{phot}})$ distribution from all five CANDELS fields (see Fig. 3), apply a 3σ outlier clipping, and fit a normalized probability density function (PDF) to this distribution such that the area under the curve sums up to unity. Note that this PDF is the probability for each $\sigma_z/(1 + z_{\text{phot}})$ value and is different from $P(z)$ discussed in Section 2. For each galaxy, we set its SDSS redshift to z_{phot} and we assign it a σ_z value drawn randomly from the PDF distribution. Then, we repeat our close companion selections and find $f_{\text{mc}}(0) = 0.057$ (B09) and




Figure 7. Comparison of the redshift evolution of the major companion fractions $f_{mc}(z)$ based on three redshift proximity criteria applied to $z > 0.5$ galaxies from CANDELS: our fiducial (equation 1; black circles, solid line) adopted from B09; equation (8) (grey circles, dashed line) modified from Man et al. (2012); and a hybrid of the B09 criterion plus close velocity separation for the subset of CANDELS galaxies with spectroscopic redshifts (open circles, dot-dashed line). All three best-fitting curves (equation 7) to the $z > 0.5$ data are anchored to the same $f_{mc}(0)$ point (square symbol) based on SDSS companions with $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ velocity separation. The error bars are defined as in Fig. 6.

0.061 (modified Man et al.). The relative change in each fraction is roughly 3 percent compared to our fiducial selection of close velocity separation.

One can argue that actual z_{phot} data from the SDSS *ugriz* photometry will produce larger redshift errors and higher companion fractions. Yet, the motivation of this exercise is to quantify the difference between tight spectroscopic velocity separation versus photometric redshift proximity selection matched to the $z > 0.5$ data, not to make our $z \sim 0$ companion fraction more uncertain. We note that if we repeat this analysis by resampling each SDSS galaxy redshift from a PDF defined by its simulated σ_z , we find *smaller* major companion fractions since this effectively makes the redshift proximity worse (larger) and we demonstrated that companions have high probability of small velocity separations for the SDSS. Either way, all the SDSS f_{mc} anchors are significantly lower than the companion fractions at $z \sim 1$, which means the turnover trend in $f_{mc}(z)$ is a robust result.

4.3 Stellar mass versus flux ratios

Our fiducial choice for selecting major companions is the stellar MR criterion: $1 \leq M_1/M_2 \leq 4$. Some previous studies, in which galaxies lack stellar mass estimates, have used observed-band FRs of $1 \leq F_1/F_2 \leq 4$ as a proxy for selecting major companions (e.g. Bell et al. 2006b; Bluck et al. 2009). Bundy et al. (2004) and B09 speculated that FRs selection could preferentially lead to inflated companion counts especially at $z \gtrsim 2$ owing to increasing SF activity. Two recent studies found conflicting trends in $f_{mc}(z)$ estimates between $z \gtrsim 1$ and $z = 3$ (Man et al. 2012, 2016). Here, we quantify the impact that changing the major companion selection from 4:1 stellar MR to 4:1 FR has on companion fractions for massive galaxies in CANDELS and the SDSS while holding all other selection criteria fixed to fiducial values.




Figure 8. Comparison of the redshift evolution of major companion fractions $f_{mc}(z)$ based on our fiducial selection (stellar MR $1 \leq M_1/M_2 \leq 4$; black symbols, solid line) and based on *H*-band FRs ($1 \leq F_1/F_2 \leq 4$; open points; slightly offset in z -direction for clarity). Best-fitting curves to the FR fraction evolution are shown for modified power law (equation 7, dashed line) and a simple power law (dot-dashed line; $f_{mc}(z) = 0.07(1+z)^1$). The data symbols distinguishing CANDELS and SDSS fractions, and the error bars are the same as in Fig. 6. The FR selection results in larger fractions at $z > 1$ and an increasing power-law redshift dependence in contrast to the $f_{mc}(z)$ derived from stellar MRs.

In Fig. 8, we compare the fiducial MR-based $f_{mc}(z)$ to fractions based on *H* band (CANDELS) and *r* band (SDSS) 4:1 FRs. We find the FR-based major companion fractions follow starkly different evolutionary trends compared to MR-based fractions at $z \gtrsim 1$. At $0 < z < 1$, the FR and MR fractions both increase with redshift and agree within their 95 per cent confidence limits. However, FR produces increasingly larger fractions from $z \sim 1$ to $z = 3$ that diverge away from the fiducial $f_{mc}(z)$ trends and grow 1.5–4.5 times greater than MR-based fractions at these redshifts. We attempt to fit the FR-based $f_{mc}(z)$ with the same function form (equation 7) that we employ for MR fraction redshift evolution, but the best-fitting parameter values (see Table 4) for the steadily increasing fractions between $0 < z < 3$ are statistically consistent with a simple power law since the FR fractions do not peak nor turn over at $z > 1$. Therefore, we refit the FR fraction evolution with a power law and find $f_{mc}(z) = 0.07(1+z)^1$ for $0 < z < 3$.

To better understand the striking differences between the FR and MR $f_{mc}(z)$ trends, we analyse stellar MRs and FRs of CANDELS and SDSS close-pair systems that satisfy our fiducial 5–50 kpc separation and redshift proximity selections (see Section 2.3.2) in Fig. 9. At $z < 1$, we find good agreement between MR and FR values for a majority of major and *minor* ($M_1/M_2 > 4$) close-pair systems. The agreement suggests that *r* band (*H* band) FRs are a good approximation for stellar MRs at $z \sim 0$ ($0.5 < z < 1$). In detail, 80 per cent (72 per cent) of SDSS (CANDELS) pairs are considered major pairs according to *both* FR and MR criteria. We find 17 per cent (25 per cent) of FR-based major pairs have a $>4:1$ (minor) stellar MR, and only 3 per cent of MR-selected major pairs have minor FRs at $z < 1$.

In the bottom four panels of Fig. 9, we notice that the FR-based major companion selection suffers a steadily increasing contamination by minor companions according to CANDELS stellar MRs. This contamination rises from 40 per cent at $1 < z < 1.5$ to over




Figure 9. Stellar MR versus H -band FR for the close-pair systems satisfying our fiducial projected separation (5–50 kpc) and redshift proximity choices for SDSS ($\Delta v_{12} \leq 500 \text{ km s}^{-1}$; $z \sim 0$; top left) and CANDELS (B09; $0.5 \leq z \leq 3$; rest of the panels). We show the pairs that satisfy our fiducial MR criterion ($1 \leq M_1/M_2 \leq 4$) in large-black points, those having a $M_1/M_2 > 4$ but satisfy the FR criterion ($1 \leq |F_1/F_2| \leq 4$) in large-grey points, the remaining points in small-grey markers. To visualize the contamination from minor pairs ($M_1/M_2 > 4$), introduced by the FR criterion in each redshift panel, we show the percentage of pairs that satisfy either MR or FR selection that fall in each of the three quadrants. We notice that the contamination increases steadily from $z \sim 0$ to $z = 3$, causing inflated f_{mc} values observed in Fig. 8 (see Section 4.3 for discussion).

three quarters in the highest redshift interval. In contrast, our fiducial selection of major companions maintains a very small (~ 5 per cent) constant contamination of $F_1/F_2 > 4$ pairs between $1 < z < 3$. This analysis clearly demonstrates that FR selection results in inflated major companion counts at $z \gtrsim 1$, confirming the result that significantly different mass-to-light ratio properties of the companion galaxies at $z > 1$ may be the main cause of the contamination by minor companions (Man et al. 2016).

5 DISCUSSION

Our extensive analysis of the full CANDELS sample and a well-defined selection of SDSS galaxies in the preceding sections provides a new baseline for the evolving frequency of massive galaxies with close major companions (and of major mergers by extrapolation) spanning epochs from the start of cosmic high noon to the present day. In this section, we present a comprehensive comparison

of our measurements to those from previous studies by taking advantage of our analysis of varying the companion selection criteria from Section 4. We derive the empirical major merger rates based on a *constant* observability time-scale and find significant variation in the rates depending on our stellar MR or H -band FR choice. We discuss plausible reasons for these variations by recomputing merger rates based on theoretically motivated, redshift-dependent time-scale prescriptions. Finally, we describe the need for detailed calibrations of the complex and presumably redshift-dependent conversion factors required to translate from fractions to rates, which is key to improve major merger history constraints.

5.1 Comparison to previous empirical close-pair-based studies

We compare our CANDELS+SDSS major companion fractions from to the results from previous stellar mass and flux-selected, empirical close-pair-based studies tabulated in Table 5. We note




Figure 10. Comparison of major companion fractions from CANDELS+SDSS to those from previous studies that employed 4:1 stellar MR (a) and 4:1 FR (b) selections. In both panels, we outline the f_{mc} measurements data points as rectangles where their height represents the 95 percent binomial confidence limits per redshift bin (width of the rectangles). The data points of previous empirical studies are given in the panel keys (see Table 5). We compare fractions based on different projected separation criteria as follows: 5–30 kpc (solid-red line; open-red markers), and 14–43 kpc (blue-dashed line, filled-blue markers). We find our major companion fraction estimates are in good agreement with previous empirical constraints when the companion selection criteria are closely matched. We offset multiple fields from Mundy et al. (2017) by a small amount, and show the upper limits in filled markers with bold arrow for clarity.

that some studies quote fraction of pairs (f_{pair}), which is the fraction of close-pair systems among a desired massive galaxy sample of interest. This is different from the companion fraction f_{mc} . To correctly compare to previous studies which used a f_{pair} definition, we derive a simple conversion between the two definitions by separately computing the ratio of $f_{mc}/f_{pair} = 1.5$, which is constant for the CANDELS and SDSS samples at $0 < z < 1.5$. Overall, our $f_{mc}(z)$ results are in good agreement with previous studies when we properly convert the published pair fractions into companion fractions. We separate our comparisons into stellar MR- and FR-based studies in Fig. 10. For each set of comparisons, we further split them into matching 5–30 kpc and 14–43 kpc projected separation selections. We discuss each comparison with respect to any differences in the details of the redshift proximity selection.

5.1.1 Stellar mass-ratio-selected studies

We use the 5–30 and 14–43 kpc fractions from Section 4.1 (Fig. 6) and plot them in Fig. 10 alongside previous studies that employed a similar projected separation criterion and 4:1 stellar MR selection ($1 \leq M_1/M_2 \leq 4$). We start our discussion by comparing to the studies that employed a $R_{proj} = 5$ –30 kpc criterion. Ryan et al. (2008) studied major close pairs among galaxies more massive than $10^{10} M_{\odot}$ over the redshift range $0.5 < z < 2.5$, and found an unevolving fraction (~ 10 per cent). We agree with these estimates within their quoted uncertainties; however, we acknowledge that the consistency at $2 < z < 2.5$ is because of their large error bar (> 50 per cent). We also note that Ryan et al. used a $2\sigma_z$ overlap as their redshift proximity, which is less restrictive than our modified Man et al. (2012) redshift proximity criteria in equation (8). Based on our analysis in Section 4.2, this implies that the Ryan et al. fractions may be overestimated. Next, we compare to B09 who studied 5–30 kpc pairs among $\log_{10}(M_{stellar}/M_{\odot}) \geq 10$ galaxies between $0.4 < z < 1.4$ using a redshift proximity that we have adopted as our

fiducial choice. It is important to note that B09 elected a 4:1 K_s -band FR as their fiducial major companion selection criterion. However, in their analysis, they found that 80 per cent of their FR-selected companions also met $< 4:1$ stellar-MR major companion selection. To appropriately compare to B09 pair fractions, we include a 0.8 multiplicative factor to convert them to MR-based f_{mc} . We find that their estimates are on average smaller than our fractions, marginally agreeing at $0.7 < z < 1.4$, and disagreeing at $0.4 < z < 0.7$.

We now compare to the most recent study by Mundy et al. (2017), who analysed close pairs ($R_{proj} = 5$ –30 kpc) among a sample of $\log_{10}(M_{stellar}/M_{\odot}) \geq 10$ galaxies at $0 < z < 2$ from the UDS, VIDEO, GAMA, and COSMOS data sets (see Mundy et al. 2017, for details). In Fig. 10, we show all their pair fractions, except for an upper limit of 17.5 per cent from COSMOS at $1.5 < z < 2$ and convert them to f_{mc} . We find excellent agreement between GAMA-survey-based, spectroscopically derived fraction at $0 < z < 0.2$ and our SDSS-based fraction at $z \sim 0$. For the same GAMA sample, they also computed the fractions using photometric redshifts and found an average of ~ 5 per cent at $0 < z < 0.2$. We separately computed the SDSS f_{mc} using B09 redshift proximity and matching their $R_{proj} = 5$ –30 kpc selection, and find that our fraction $f_{mc}(0) \sim 2.8$ per cent is ~ 1.8 times smaller and marginally disagrees within their quoted uncertainties. At $0.2 < z < 1$, we find that Mundy et al. fractions are in good agreement with our $f_{mc}(z)$, except for their COSMOS-survey-based value at $0.2 < z < 0.5$, which is marginally smaller than our f_{mc} . At $1 < z < 1.5$, their fraction (~ 15 per cent) is in disagreement with our findings, while their upper limits are consistent with our results. Overall, we demonstrate good agreement with Mundy et al. (2017).

Now, we discuss our comparison to the studies that employed an $R_{proj} = 14$ –43 kpc separation selection. We start with Williams et al. (2011), who studied major ($< 4:1$ stellar-MR) close pairs among $\log_{10}(M_{stellar}/M_{\odot}) \gtrsim 10.5$ galaxies at $0.4 < z < 2$, and found a diminishing redshift evolution of the fractions. We find disagreement

with the Williams et al. fractions at $0.4 < z < 1.2$, where our f_{mc} are higher by 25 per cent, however, we show good agreement within their diminished fractions at $1.2 < z < 2$. We note that our sample is almost two times larger than their galaxy sample, and a likely reason for this discrepancy is that their uncertainties may have been underestimated. Next, we compare to Newman et al. (2012), who quantified major companion fraction among $\log_{10}(M_{\text{stellar}}/M_{\odot}) \gtrsim 10.7$ galaxies, and found that f_{mc} is flat (at ~ 10 per cent) at $0.4 < z < 2$. We find good agreement with their findings within the quoted uncertainties and in that redshift range.

Finally, we compare to a recent study by Man et al. (2016) who study a sample of $\log_{10}(M_{\text{stellar}}/M_{\odot}) \gtrsim 10.8$ galaxies to constrain major companion fractions at $0 < z < 3$ using two separate data sets, namely 3D-HST and UltraVISTA (see Man et al. 2016, for details). First, we find that our $f_{\text{mc}}(z)$ values are in excellent agreement with the Man et al. 3D-HST estimates over their redshift range, which is not surprising as 3D-HST covers 75 per cent of the CANDELS fields. We find that the Man et al. UltraVISTA fractions follow an increasing trend that is qualitatively similar to our $f_{\text{mc}}(z)$ at $0 < z < 1$, however, we notice that the former fractions are quantitatively smaller than the latter. At $z > 1$, we find that the UltraVISTA fractions are in good agreement with our results, except at $z \sim 2.25$, where our f_{mc} is two times smaller. Overall, we note that our companion fractions are in good agreement with Man et al. (2016) estimates. Moreover, Man et al. (2016) applied a parabolic fitting function to their UltraVISTA major companion fractions and report that their redshift evolution may be peaking around $z \sim 1-1.5$, in contrast to our fractions that peak around $0.5 < z < 1$ and diminish up to $z = 3$. This may be a consequence of more-massive galaxies experiencing growth earlier in cosmic time than the less-massive population, which is often referred to as the galaxy-downsizing phenomenon. In summary, we conclude that our stellar-MR-based major companion fractions $f_{\text{mc}}(z)$ are in good agreement with previous empirical close-pair-based estimates, once they are closely matched in selection criteria.

5.1.2 Flux-ratio-selected studies

We separately derive the major companion fractions for $R_{\text{proj}} = 5-30$ and $14-43$ kpc selections using H band (r band) FR selection for CANDLES (SDSS) and plot them in Fig. 10 alongside some published studies that used a similar projected separation and $<4:1$ flux (luminosity) ratio selection (see Table 5). First, we discuss the comparison to those studies that employ $R_{\text{proj}} = 5-30$ kpc. We start with Kartaltepe et al. (2007), who studied close pairs using a V -band luminosity-limited galaxy sample at $0.2 \leq z \leq 1.2$, and found that the companion fraction evolves as $f_{\text{mc}} \propto (1+z)^{3.1}$. We find that Kartaltepe et al. estimates are ~ 2 times smaller than our results at $0.5 < z < 1$, however, they start to converge at $z < 0.5$ and $z > 1$, and agree with our $f_{\text{mc}} \sim 10$ per cent at $z = 1.2$. We note that Kartaltepe et al. use a smaller separation annulus of $R_{\text{proj}} = 5-20$ kpc, and in addition they also apply a random chance pairing correction. Based on our analysis from Sections 3.3.3 and 4.1, we note Kartaltepe et al. fractions may be systematically smaller, and most plausibly are causing the observed discrepancy at $0.5 < z < 1.0$.

We now compare to B09, and remind our reader that they used a $4:1$ FR as their fiducial major companion criterion. We find good agreement with the Bundy et al. estimates at $0.7 < z < 1.4$, but find marginal disagreement at $0.4 < z < 0.7$. Next, we compare to Man et al. (2012), who analysed close pairs among $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 11$ galaxies using $4:1$ H -band FR criterion at $0 < z < 3$, and found fairly high fractions up to $z = 3$. We find good agreement with their

fractions over the full redshift range, however, we acknowledge that a proper comparison with our highest redshift bin results is not possible owing to their large uncertainty.

Finally, we compare to Bluck et al. (2009); Bluck et al. (2012), who analysed the incidence of close companions which satisfy a $4:1$ H -band FR selection among a GOODS NICMOS survey sample of $\log_{10}(M_{\text{stellar}}/M_{\odot}) > 11$ galaxies at $1.7 < z < 3$ (see Bluck et al. 2012, for details). Bluck et al. report that the fraction of galaxies hosting a nearby major companion evolves steeply with redshift as $f \propto (1+z)^3$ (dashed line in Fig. 10b), reaching up to $f = 0.19 \pm 0.07$ ($1.7 < z < 2.3$) and $f = 0.40 \pm 0.1$ ($2.3 < z < 3$), with a total fraction of $f = 0.29 \pm 0.06$. While we find consistent agreement at $1.7 < z < 2.3$, we note that their fractions are a factor of two higher at $2.5 < z < 3$. Similarly, we also find that their total fraction, which spans over $1.7 < z < 3$, is also larger by a factor of two when compared to our total f_{mc} at $2 < z < 3$. Although we qualitatively agree with the strongly increasing redshift evolution of Bluck et al., we note that our $f_{\text{mc}}(z)$ trend is shallower with a simple power-law slope of $m \sim 1$ (see Fig. 8). We argue that the steeply increasing fraction evolution by Bluck et al. may be a consequence of the increased contamination from minor companions by their FR selection (see Section 4.3). Also, significant field-to-field variance due to their small sample size and the use of a relatively less restrictive line-of-sight contamination correction may be contributing towards the observed (factor of two) disagreement to our fractions.

We now focus our comparisons to studies that employ an $R_{\text{proj}} = 14-43$ kpc selection (see Table 5). We start our comparison with Lin et al. (2008), who studied close pairs among luminosity-limited sample of galaxies at $0.12 < z < 1.1$. We find that our fractions are only marginally consistent with their results at $z < 0.5$ and disagree with them at $0.5 < z < 1$ as our fractions are ~ 2 times larger than their values. We note that Lin et al. used a stringent spectroscopic-redshift proximity criterion of $\Delta v_{12} \leq 500 \text{ km s}^{-1}$, which may be the reason for their smaller fractions and the observed discrepancy. Nevertheless, we notice that Lin et al. fractions start to converge with our estimates at $z > 1$, which may be due to the rising minor-companion contamination by FR selection leading to larger fractions. We now compare to López-Sanjuan et al. (2011), who study a spectroscopically confirmed sample of close pairs among B -band luminosity-limited galaxy sample to quantify pair fractions at two redshifts $z = 0.5$ and 0.8 . We find that our f_{mc} estimate at $z = 0.5$ is marginally higher than their fraction, but is in good agreement at $z = 0.8$. Also, we qualitatively agree with their increasing redshift evolutionary trend of the fractions at $0.5 < z < 0.8$.

Finally, we compare to Man et al. (2016), who also derived $4:1$ H -band FR-based companion fractions for $\log_{10}(M_{\text{stellar}}/M_{\odot}) \gtrsim 10.8$ galaxies spanning $0.1 < z < 3$ among the 3D-HST and UltraVISTA data sets. We find that our $f_{\text{mc}}(z)$ values trace very closely with the Man et al. 3D-HST estimates and are in excellent agreement within the uncertainties. We also find that Man et al. UltraVISTA estimate qualitatively agree with the increasing redshift evolution of our FR-selected $f_{\text{mc}}(z)$. However, we notice that their estimates at $z < 0.5$ are marginally smaller than our f_{mc} extrapolation between $z \sim 0$ and $0.5 < z < 1$. In summary, we conclude that our H -band FR-based $f_{\text{mc}}(z)$ are in good agreement with previous empirical studies once we match closely the choices of companion selection criteria.

5.2 Major merger rates

As outlined in the Introduction, empirical merger rates provide a fundamentally crucial measure of the importance of major merging

in the evolution of massive galaxies. Our new major companion fractions from CANDELS+SDSS allow us to place clearly defined new constraints on the evolution of major merger rates between $z = 3$ and 0. Here, we elect to focus on a straightforward comparison of merger rates derived from our major companion fractions with the largest systematic differences: stellar MR versus FR selection. Then we compare our merger rates to several recent empirical studies and cosmologically motivated predictions that employ similar companion selection criteria as we do.

The major merger rate based on close-companion (pair) statistics is defined as the number of major galaxy–galaxy merger events per unit time per selected galaxy (see Man et al. 2016):

$$R_{\text{merg,pair}} = \frac{C_{\text{merg,pair}} \times f_{\text{mc}}}{T_{\text{obs,pair}}} [\text{Gyr}^{-1}], \quad (9)$$

where $T_{\text{obs,pair}}$ is the average observability time-scale during which a galaxy–galaxy pair satisfies a given selection criterion, and the multiplicative factor $C_{\text{merg,pair}}$ attempts to account for the fact that not all such companions will merge within the $T_{\text{obs,pair}}$ interval (e.g. Lotz et al. 2011). It is clear that the $C_{\text{merg,pair}}$ and $T_{\text{obs,pair}}$ factors are *crucial* assumptions involved in converting observed companion fractions into merger rates. As such, these factors are a major source of systematic uncertainty in merger rate calculations. A few analyses of theoretical simulations have provided fairly broad guidelines for these key assumptions. Hopkins et al. (2010a) find that different merger time-scale assumptions can induce up to a factor-of-two uncertainty in the merger rates. Others have attempted to constrain the fraction-to-rate conversion factors and quote limits of $C_{\text{merg,pair}} = 0.4$ – 1 (Kitzbichler & White 2008; Jiang et al. 2014). Despite the need for improved constraints on these key merger rate factors, attempts to calibrate $C_{\text{merg,pair}}$ and $T_{\text{obs,pair}}$ in detail with regard to changing companion selection criteria and as a function of fundamental galaxy properties such as redshift and stellar mass, are lacking.

In practice, previous empirical studies have used merger rate factors that span a wide range. For example, some studies have adopted $C_{\text{merg,pair}} = 1$ for simplicity (e.g. Man et al. 2016), while others have employed an intermediate value of 0.6 (e.g. B09, Lotz et al. 2011) based on limits from simulations. Similarly, some empirical studies adopt their $T_{\text{obs,pair}}$ values based on analytical fitting function provided by Kitzbichler & White (2008), while others adopt values provided by Lotz et al. (2010) that range from 0.3 to 2 Gyr depending on the close-companion selection criteria. For the major merger rate calculations below, we make a simple assumption that these factors are constant over all redshifts and masses we probe. We adopt $C_{\text{merg,pair}} = 0.6$ and $T_{\text{obs,pair}} = 0.65$ Gyr, which is suitable for our fiducial $R_{\text{proj}} = 5$ – 50 kpc criterion (Lotz et al. 2010). Exploring the detailed impact of these assumptions is beyond the scope of this paper. However, in Section 5.3.2, we test two theoretically motivated redshift-dependent time-scale prescriptions to rederive the merger rates and discuss their implications.

In Fig. 11, we show the major merger rate evolution during $0 < z < 3$, which we derive from CANDELS+SDSS major companion fractions for stellar MR and FR selections with matched fiducial projected separation and B09 redshift proximity criteria. Holding aside the factor-of-two systematic uncertainty contribution from $T_{\text{obs,pair}}$ to merger rate calculations, we find both the MR- and FR-based merger rates evolutionary trends mimic their respective $f_{\text{mc}}(z)$ trends discussed in Sections 3.3 and 4.3, which is expected given our simplistic fraction-to-merger-rate conversion factors. We notice that both the MR and FR selections yield consistent major merger

rates at $z < 1$, where they increase strongly from ~ 0.05 ($z \sim 0$) to $\sim 0.15 \text{ Gyr}^{-1}$ ($0.5 < z < 1$). At $z > 1$, the different selections have divergent merger rates. The MR-based merger rates decline with redshift to $z = 3$, indicating a turnover at $z \sim 1$. In contrast, the inflated companion fractions owing to contamination by minor companions, the FR selection yields 1.5–4.5 times larger rates between $1 \lesssim z \lesssim 3$ with an increasing power-law trend $R_{\text{merg,pair}} \propto (1+z)^1$.

Next, we compare our empirical MR-based merger rates to previous empirical estimates (Fig. 11). For $z \lesssim 1.5$, we compare to Lotz et al. (2011) who compile close-pair fractions from several observational stellar-mass and luminosity-selected studies (magenta line) and found strongly increasing major merger rate evolution with redshift $R \propto (1+z)^{1.7-2.1}$, which qualitatively agrees with both our MR and FR rate evolutions over this redshift range. The Lotz et al. merger rate evolutions have systematically smaller normalization than ours because their compilation includes close-pair-based fractions that may be systematically smaller than our f_{mc} due to the use of different selection criteria. For example, Kartaltepe et al. (2007) and Lin et al. (2008) use a stringent proximity selections (see Section 5.1.2, and Table 5) and Patton & Atfield (2008) use a stringent major companion selection criterion.⁹ For merger rates at $0.5 < z < 3$, we compare our fiducial MR-based merger rate to a recent study of close pairs by Man et al. (2016). We find that our individual $R_{\text{merg,pair}}$ data points agree well with the Man et al. results (blue) within the measurement uncertainties at these redshifts. Yet, our standalone rising-diminishing merger rate evolution disagrees with the Man et al. flat trend. If we were to extend their rates to $z \sim 0$, we would disagree with them; however, we note that Man et al. do not attempt to anchor to a low redshift (for their 3D-HST sample), as such, claim a flat trend with redshift in agreement with Mundy et al. (2017). Recently, Mundy et al. (2017) in conjunction with Duncan et al. (in preparation) find that the Henriques et al. (2015) SAM mock light-cones accurately reproduce a flat pair-fraction trend that is consistent with the recent findings by Man et al. (2016), Snyder et al. (2017), and the results of this work. Employing an un-evolving close-pair observability time-scale assumption, Mundy et al. report a flat major merger rate evolution. Our fiducial MR-selected merger rates shown in Fig. 11 agree with the Mundy et al. result, which is not surprising owing to the assumption of constant observability time-scale.

Here, we discuss our comparison to theoretical major merger predictions. Since these are based on stellar MRs, we refrain from comparing them to our FR-based rates. Hopkins et al. (2010a) provide a comprehensive analysis of major galaxy–galaxy merging using Λ cold dark matter motivated simulation and SAM to derive major merger rate predictions, and quantified systematic error contribution from various theoretical model-dependent assumptions. We find that the Hopkins et al. major merger rate evolution of $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.3$ galaxies ($R = 0.04(1+z)^{1.35}$; green line in Fig. 11), within a factor-of-two uncertainty, agrees qualitatively and quantitatively with our MR-based rates up to $z < 1.5$. At $z > 1.5$, our fiducial MR-selected, diminishing merger-rate trend disagrees with the Hopkins et al. predictions. We also compare our merger rates to a recent theoretical prediction by Rodriguez-Gomez et al. (2015) from the Illustris numerical hydrodynamic simulation (Vogelsberger et al. 2014). We note that their $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 10.3$ galaxy merger rates

⁹ Patton & Atfield (2008) select major companions using a 2:1 luminosity ratio as opposed to our 4:1 criterion from a luminosity-limited galaxy sample at $z \sim 0$.




Figure 11. Comparison of CANDELS+SDSS galaxy–galaxy major merger rates $R_{\text{merg,pair}}(z)$ (number of mergers per galaxy per Gyr) for massive ($M_{\text{stellar}} \geq 2 \times 10^{10} M_{\odot}$) galaxies at $0 < z < 3$, to rates from previous empirical studies and theoretical model predictions. We show the $R_{\text{merg,pair}}(z)$ computed using major companion fractions (f_{mc}) based on fiducial projected separation (5–50 kpc) and redshift proximity (CANDELS: equation 1; SDSS: $\Delta v_{12} \leq 500 \text{ km s}^{-1}$) split into stellar MR (filled points; solid line) and FR (open points; dashed line) for CANDELS (circles) and SDSS (square). We employ simplistic assumptions for fraction-to-rate conversion factors $T_{\text{obs,pair}} = 0.65 \text{ Gyr}$ and $C_{\text{merg,pair}} = 0.6$, and show the variation of $R_{\text{merg,pair}}(z)$ for $C_{\text{merg,pair}} = 0.4\text{--}0.8$ in red shading. The error bars on the data points indicate their 95 per cent binomial confidence limits, solely based on the observed number counts. The solid (dashed) magenta line is the empirical merger rate evolution from Lotz et al. (2011) for stellar-mass (luminosity) limited selections, the empirical merger rate from Man et al. (2016) is shown in solid blue line along with its blue shaded uncertainty. We compare to theoretical galaxy–galaxy major merger rate predictions of $M_{\text{stellar}} \geq 2 \times 10^{10} M_{\odot}$ galaxies by Hopkins et al. (2010a, solid green lines), Rodríguez-Gomez et al. (2015, solid black), and to the analytical halo major merger rate prediction of $M_{\text{halo}} \sim 10^{12} M_{\odot}$ DM haloes by Neistein & Dekel (2008), Fakhouri et al. (2010), and Dekel et al. (2013) ($R \propto (1+z)^{2.5}$; dashed brown line). We strongly advise against trusting FR-based galaxy merger rates despite their agreement with the theoretical models, owing to the notable contamination from non-major mergers (see Section 4.3). Our fiducial MR-based empirical major merger rates show broad agreement up to $z \sim 1.5$, but demonstrate a strong tension at $z > 2$ when compared to previous empirical and theoretical constraints.

(black line in Fig. 11) follow an even stronger increasing redshift dependence ($R \propto (1+z)^{2.4\text{--}2.8}$) than the Hopkins et al. predictions, which also agrees qualitatively with our $f_{\text{mc}}(z)$ trend at $z < 1.5$. However, at $z < 1.5$, we note that the Rodríguez-Gomez et al. predictions are smaller than our empirical estimates. Similar to the conclusion from the Hopkins et al. comparison, our fiducial MR-based merger rates disagree with predictions from Rodríguez-Gomez et al. at $z > 1.5$. We present further discussion on the plausible reason for the observed discrepancy of flat versus rising merger rate evolution in Section 5.3.2.

Being the primary driver for galaxy–galaxy merging, halo–halo merger rates serve as a ceiling for both empirical measurements and theoretical galaxy merging predictions, and their qualitative trend is expected to mimic the galaxy merger rates closely. We find that our empirical MR-selected merger-rate trend broadly agrees with the analytical merger rates of $M_{\text{halo}} \sim 10^{12} M_{\odot}$ dark-matter (DM) haloes ($R \propto (1+z)^{2.5}$; brown line in Fig. 11) predicted by Neistein & Dekel (2008), Fakhouri et al. (2010), Dekel et al.

(2013) (also see Maulbetsch et al. 2007) at $z \lesssim 1$ but starts to deviate significantly at $z > 1$. Since the translation from a halo–halo merger rate to a galaxy–galaxy merger rate depends on the redshift-dependent mapping of galaxies on to their respective DM haloes (halo-occupation statistics), the galaxy merger rates are expected to follow a shallower redshift evolution than the ones set by the halo merger rates (for discussion, see Hopkins et al. 2010a). Although different assumptions of halo-occupation models may contribute up to a factor-of-two uncertainty in the theoretically predicted galaxy merger rates (e.g. Hopkins et al. 2010b), they all hint towards a rising incidence with increasing redshift. As such, a deviation of more than factor of five (e.g. see redshift bin $2 < z < 3$ in Fig. 11) between the constant time-scale-based fiducial MR-selected empirical rates (diminishing) and the analytical halo merger rate evolution (rising; $R \propto (1+z)^{2.5}$) is puzzling. Based on this analysis, we conclude that a straightforward MR-based estimation of major merger rates using a constant fraction-to-merger-rate observability time-scale ($T_{\text{obs,pair}} = 0.65 \text{ Gyr}$) are in disagreement with the theoretical




Figure 12. Comparing the redshift evolution of major merger rate of $M_{\text{stellar}} \geq 2 \times 10^{10} M_{\odot}$ galaxies based on our fiducial close-pair time-scale assumption $T_{\text{obs, pair}} = 0.65$ Gyr copied from Fig. 11 (solid red line, shading) to rates from different time-scale choices. We show the rates based on Jiang et al. (2014) scaling relation $T_{\text{obs, pair}} \propto H(z)^{-1/3}$ in red dotted line, and Snyder et al. (2017) relation $T_{\text{obs, pair}} \propto (1+z)^{-2}$ in red dot-dashed line (starting at $z_0 = 1$). We also plot the theoretical merger rate predictions shown in Fig. 11.

predictions at $z > 1.5$. We discuss the implications of these conclusions in Section 5.3.

5.3 Implications of the disagreement between simplistic empirical merger rates and model predictions

So far, we have demonstrated that major merger rates based on a rather straightforward MR-selection start to disagree with theoretical predictions of major galaxy–galaxy merging at $z \gtrsim 1.5$. This indicates there may be one or many redshift-dependent effects plaguing the key variables ($C_{\text{merg, pair}}$, f_{mc} , and $T_{\text{obs, pair}}$) in equation (9), causing the observed merger rates to disagree with simulations. Here, we first investigate for plausible systematic underestimation of f_{mc} values at $z \gtrsim 1.5$ as a reason for the observed discrepancy by testing if we are missing close companions by not choosing the full z_{phot} PDF when applying the B09 redshift proximity. Then, we discuss another possibility that the simplistic, constant value assumptions of $T_{\text{obs, pair}}$ may be incorrect, and test two theoretically motivated, redshift-dependent $T_{\text{obs, pair}}$ prescriptions to rederive merger rates (see Fig. 12). We also elaborate on the remnant formation times, which is a key difference between the merger rates measured in theoretical simulations and empirically calculated close-pair-based rates, and comment on how it may be affecting the data-theory comparison.

5.3.1 Are we missing companions by not accounting for the full z_{phot} PDFs?

One plausible reason for the $z > 2$ data-theory discrepancy may be that the adopted B09 redshift proximity criterion (equation 1) is missing plausible physical close-pair systems because we are using only the 1σ photometric uncertainties and not the full z_{phot} PDF. To test if this is the case we carry out a simple test, where we start by simplifying equation (1) as $\xi = \Delta z_{12} / \sqrt{\sigma_{z,1}^2 + \sigma_{z,2}^2}$; therefore, our fiducial criterion equates to $\xi \leq 1$. The distribution

of ξ values for physical close-pair systems is Gaussian, assuming σ_z values are Gaussian. Therefore, using $\xi \leq 1$ is equivalent to selecting 68 per cent of the full ξ distribution. Ideally, if one assumes the contribution from non-physical pairs (hereafter called as the background) per ξ bin is negligible, as much as 32 per cent of the close pairs (thereby companions) can be missed when using $\xi \leq 1$. For this exercise, we allow for a non-zero background to properly estimate the missing pairs at five redshift bins between $0.5 < z < 3.0$.

We begin with the cumulative distribution function (CDF) of ξ for all the close-pair systems selected using our fiducial $R_{\text{proj}} = 5$ –50 kpc projected separation criterion. To avoid errors from small number statistics, we do not limit close-pair systems with a stellar-MR selection. However, we require that both the host and companion galaxies to be brighter than $H = 25$ mag to exclude large ξ values from affecting the statistics. We now model CDF with a Gaussian integral function in conjunction with a constant background contribution, following the fitting formula: $N(\xi \leq c) = A / \sqrt{2\pi} \int_0^c \exp^{-x^2/2} dx + Bc$. Here, $N(\xi \leq c)$ is the number of close-pair systems that satisfy a cut-off significance (c) as $\xi \leq c$, and the variables A and B are the contributions from plausible physical close pairs and the background, respectively. For demonstrative purposes, we present the fits to the CDF and their best-fitting parameter values of A and B in Fig. 13 for two redshift bins between $1.5 < z < 3$. At each redshift bin, we define the ratio $(A/2)/N(\xi \leq 1)$ as the multiplicative correction factor to our fiducial f_{mc} that corrects for the missing close-pair systems due to Method I. When the background is negligible ($B \sim 0$), the correction factor is ~ 1.44 . However, if B is significant, then the correction factor may become less than unity. In Fig. 14, we plot the corrected f_{mc} alongside the fiducial fractions. We note that fractional change in the major companion fraction after applying the correction factor is less than 5 per cent at $0.5 < z < 2$. However, at $z > 2$, we find the relative contribution from the background starts to dominate the Gaussian contribution and therefore causes the fiducial f_{mc} to decrease by more than 10 per cent. Despite these changes, we note that the overall corrected $f_{\text{mc}}(z)$ values do not significantly deviate from the fiducial $f_{\text{mc}}(z)$ values. Therefore, we conclude that we are not missing close pairs (thereby companions), by using 68 per cent photometric-redshift errors for our fiducial redshift proximity criteria.

5.3.2 Evolving observability time-scale and remnant formation times

It is evident by now that the close-pair observability time-scale ($T_{\text{obs, pair}}$) is a key parameter that dictates the measurement of merger rates. Yet, a wide range of values can be found in the literature (see Section 5.2), often assumed to be constant over a wide range of redshifts. Therefore, with the growing evidence (Man et al. 2016; Mundy et al. 2017) for flat or possibly diminishing close-pair-based fractions at $1 < z < 3$, it is expected that a constant time-scale assumption-based merger rates to disagree with the continually rising merger-rate trends from the theoretical predictions. A possible implication of this disagreement is that the time-scale may be evolving with redshift, such that an ongoing merger can be observed as a close pair for a shorter period at earlier cosmic times. This may not be surprising as simulation-based merging-time-scale calibrations (e.g. Kitzbichler & White 2008; Jiang et al. 2014), motivated by dynamical friction time for a satellite DM halo to merge with the parent, are complex functions that depend on key merger properties such as the MR, galaxy mass, R_{proj} , and $H(z)^{-1/3}$ ($H(z) = H_0 E(z)$; H_0 is the present-day Hubble constant and $E(z)$ is the evolution of H_0 as a function of redshift). Moreover, the dynamical time-scale




Figure 13. The CDFs of $\xi = |\Delta z_{12}| / \sqrt{\sigma_{z,1}^2 + \sigma_{z,2}^2}$ (from equation 1) for projected pairs satisfying our fiducial $5 \text{ kpc} \leq R_{\text{proj}} \leq 50 \text{ kpc}$ selection (blue dashed, narrow line) as described in Section 5.3.1 for two redshift bins: $1.5 < z < 2$ (left) and $2.5 < z < 3$ (right). In each redshift-bin panel, following the equation $N(\xi \leq c) = A/\sqrt{2\pi} \int_0^c \exp^{-x^2/2} dx + Bc$, the best-fitting curve to the CDF is shown in narrow solid-red line. The breakdown of the best-fitting curve into Gaussian and background contributions are shown in bold solid-black and bold-dashed lines, respectively, and we print the best-fitting values (A and B) in each panel.




Figure 14. Comparing the redshift evolution of the fiducial major companion fraction $f_{\text{mc}}(z)$ (filled circles; copied from Fig. 5) to the fractions after correcting for full $\xi = |\Delta z_{12}| / \sqrt{\sigma_{z,1}^2 + \sigma_{z,2}^2}$ distribution usage (open circles; see Section 5.3.1 for details). For this exercise, we anchor both CANDELS $f_{\text{mc}}(z)$ to the same fiducial selection-based SDSS data point at $z \sim 0$. The error bars correspond to 95 per cent binomial confidence limits on the f_{mc} .

based on simple cosmological arguments is also redshift dependent, and can be approximated as $T \propto (1+z)^{-1.5}$ (see Snyder et al. 2017). However, as per the discussion in Snyder et al. (2017), any processes that can act faster than the dynamical time-scale and may impact the variables of close companion selection criteria, which in turn can change the $T_{\text{obs,pair}}$. In this context, the galaxies are undergoing a rapid transformation concerning their stellar-mass assembly, owing to rapid ongoing SF (Madau & Dickinson 2014) at $z > 1$. Snyder et al. found that this rapid mass assembly can shorten the close-pair

observability time-scale towards earlier redshifts, which evolves as $T \propto (1+z)^{-2}$ at $z > 1$. Moreover, they measured the close-pair fractions within the Illustris simulation (Vogelsberger et al. 2014) and inferred that the time-scale should evolve as $T \propto (1+z)^{-2}$ to match the intrinsic merger rates trend from Rodriguez-Gomez et al. (2015).

In Fig. 12, we recompute the merger rates for the fiducial MR-based companion fractions by adopting the two evolving time-scale assumptions $T_{\text{obs,pair}} \propto H(z)^{-1/3}$ and $T_{\text{obs,pair}} \propto (1+z)^{-2}$ from Jiang et al. (2014) and Snyder et al. (2017), respectively. Since Snyder et al. assert that the close-pair observability is mainly dynamical time dominated at $z < 1$, which results in an unevolving time-scale value, we choose to evolve the $T_{\text{obs,pair}}$ only at $z > 1$ when re-computing the merger rates. We find that the major merger rates based on Jiang et al. evolving time-scale still does not sufficiently agree with the theoretical predictions. On the other hand, when we use simulation-tuned Snyder et al. prescription of $T_{\text{obs,pair}} \propto (1+z)^{-2}$, we find that the merger rates agree closely to model predictions of Hopkins et al. (2010a), Rodriguez-Gomez et al. (2015), and closely mimic the Neistein & Dekel (2008) and Dekel et al. (2013) halo-halo merger rates. The fact that our evolving time-scale-based empirical merger rates agree with the theoretical predictions from Rodriguez-Gomez et al. imply a very good agreement between our $f_{\text{mc}}(z)$ and Snyder et al. fractions. Additionally, we argue that the disagreement between Henriques et al. (2015) and Mundy et al. (2017) flat merger rate evolution and the Hopkins et al. (2010a) and Rodriguez-Gomez et al. (2015) theoretical predictions (see Section 5.2) may be due to the unevolving close-pair observability time-scale assumption. These results imply that an evolving time-scale assumption may be more appropriate when measuring merger rates at $z > 1$, and it may be necessary to explain the discrepancy between a constant time-scale-based rates and theoretical model predictions.

Additionally, we also stress the difference in the measured quantities between the simulations and empirical close-pair-based studies. In theoretical simulations, a merger is counted towards the intrinsic

merger rate when the merging process is concluded, i.e. when the remnant of the progenitor galaxies is formed, and therefore it is an instantaneous measure. On the other hand, the close-pair method probes future merging systems, within some time frame after observing them. Snyder et al. found that the remnant formation time also evolves as $(1+z)^{-2}$, which is defined as the time prior to the formation of the remnant, when the merger is observed as a close pair. We follow section 4.1 in Snyder et al. and compute instantaneous remnant formation rates at $z > 1$, which is conceptually similar to the instantaneous merger rates quoted by the simulations. We find that remnant formation rate follows an increasing redshift trend, which is in qualitative agreement with finding by Snyder et al. (2017). We acknowledge that a comprehensive analysis of the empirical and theoretical measurement methods is necessary to make quantitative data-theory comparisons confidently. However, it is beyond the scope of the current study to fully address them. We reserve further discussions on this analysis to an accompanying paper (Paper 2 in the series; Mantha et al., in preparation).

5.4 Future work

So far, we have discussed plausible evolutionary observability time-scale prescriptions that may be necessary when deriving merger rates at $z > 1$. While this may be true, $C_{\text{merg,pair}}$ is another key parameter in equation (9) which may also depend on the companion selection criteria. In fact, recent studies (e.g. Lin et al. 2010; de Ravel et al. 2011) find that $C_{\text{merg,pair}}$ varies as a function of R_{proj} , redshift, and local overdensity (often represented by δ). Although, a redshift-dependent time-scale prescription sufficiently brings empirical and theoretical merger rates to agreement, it is possible that $T_{\text{obs,pair}}$ may be a simultaneous function of several key close-pair variables. As such, these conversion factors may be better represented as $C'_{\text{merg,pair}} = C_{\text{merg,pair}}(z, R_{\text{proj}}, \delta)$, $T'_{\text{obs,pair}} = T_{\text{obs,pair}}(z, R_{\text{proj}}, M_1/M_2, M_{\text{stellar}}, \delta)$, where $C'_{\text{merg,pair}}$ and $T'_{\text{obs,pair}}$ are the evolving prescriptions of C_{pair} and $T_{\text{obs,pair}}$, respectively. To fully quantify the interplay among the companion selection criteria and the fraction-to-merger-rate conversions, it is important to derive detailed calibrations for $C'_{\text{merg,pair}}$ and $T'_{\text{obs,pair}}$ using large-scale cosmological simulations. We will discuss this in a follow-up Paper 2 (Mantha et al. in preparation).

Furthermore, it has been speculated that stellar MR may be a poor representer of ongoing ‘significant’ mergers, owing to the dominant cold-gas contribution to the total baryonic mass (gas+stellar mass) of the galaxy at $z \sim 2-4$ (Lotz et al. 2011; Man et al. 2016). If this is true, a galaxy merger with $M_1/M_2 > 4$ may in fact be significant if the total baryonic (stellar+cold gas) MR of the galaxies is considered (i.e. $M_{1,\text{bar}}/M_{2,\text{bar}} \leq 4$). This may cause for many of the mergers that can be significant contributors to several aspects of galaxy evolution to be missed because of the selection bias. Empirical confirmation of this speculated selection bias deserves a dedicated analysis of its own, and we will discuss this in an accompanying paper (3 of this series).

SAMs are showing a great promise as the testing grounds for many galaxy evolution-related questions (e.g. Brennan et al. 2015; Pandya et al. 2016). Three studies (Somerville et al. 2008; Lu et al. 2014; Croton et al. 2016, Somerville et al., in preparation) have independently developed SAMs based on Bolshoi–Plank N -body simulation (Klypin et al. 2016) and produced mock data sets that mimic observational data in the five CANDELS fields. Additionally, multiple realizations of each CANDELS mock field are also made available. We will take advantage of the intrinsic merger history information and apply close-pair analysis on these mock data

sets to derive the necessary calibrations. With the help of multiple realizations per field, we will be able to quantify the impact of sample variance towards close-pair statistics and merger rates. Also, by introducing realistic, CANDELS-like random and systematic errors on to simulated z_{phot} and M_{stellar} quantities, we will carry out a comprehensive investigation of their impact on the measured merger rate evolution. These calibrations and analyses will help fill the gaps in our understanding of the merger rate measurements and will provide most up-to-date major merger rate constraints with future state-of-the-art telescopes.

6 CONCLUSIONS

In this paper, we analyse a large sample of nearly 9800 massive ($M_{\text{stellar}} \geq 2 \times 10^{10} M_{\odot}$) galaxies spanning $0 < z < 3$ using the five *HST* CANDELS fields (totaling $\sim 0.22 \text{ deg}^2$) and volume-matched region of the SDSS surveys to quantify the redshift evolution of major companion fraction $f_{\text{mc}}(z)$. We adopt a fiducial selection criteria of projected separation ($5 \leq R_{\text{proj}} \leq 50 \text{ kpc}$), redshift proximity $\Delta z_{12} \leq \sqrt{\sigma_{z,1}^2 + \sigma_{z,2}^2}$ (CANDELS) and $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ (SDSS), and stellar MR major companion selection criterion $1 \leq M_1/M_2 \leq 4$ (MR).

Our key result is the MR-based major companion fraction increases from 6 per cent ($z \sim 0$) to ~ 16 per cent ($0.5 < z < 1$) and decreases to ~ 7 per cent ($2.5 < z < 3$), indicating a turnover at $z \sim 1$ with power-law exponential (equation 7) best-fitting values $\alpha = 0.5 \pm 0.2$, $m = 4.6 \pm 0.9$, and $\beta = -2.3 \pm 0.5$. We perform comprehensive tests demonstrating that these evolutionary trends are robust to changes in companion selection criteria except for H -band FR selection, where FR-based $f_{\text{mc}}(z)$ agree with MR-based fractions up to $z \sim 1$, but disagree at $z > 1$ as they increase steadily with redshift as $f_{\text{mc}}(z) \propto (1+z)^1$ up to $z = 3$, confirming previous speculations (Bundy et al. 2004; Bundy et al. 2009). This disagreement is due to increasing contamination of FR selection by minor companions (MR > 4) from 40 per cent at $1 < z < 1.5$ to over three quarters at $2.5 < z < 3$, confirming the result by Man et al. (2016) that significantly different mass-to-light ratio properties of the companion galaxies at $z > 1$ may be the main cause for the contamination by minor companions.

We compute major merger rates for MR- and FR-based fractions by using a constant fraction-to-merger-rate conversion $C_{\text{merg,pair}} = 0.6$ and a non-evolving close-pair observability time-scale $T_{\text{obs,pair}} = 0.65 \text{ Gyr}$ (Lotz et al. 2010). We find that the MR-based rates follow an increasing trend with redshift up to $z \sim 1.5$, after which they decline up to $z \sim 3$. On the other hand, the FR-based rates follow a rising trend with redshift over our full redshift range $0 < z < 3$. We also recompute the MR-based rates using two additional theoretically motivated, redshift-dependent time-scales: $T_{\text{obs,pair}} \propto H(z)^{-1/3}$ (Jiang et al. 2014) and $T_{\text{obs,pair}} \propto (1+z)^{-2}$ (Snyder et al. 2017).

If recent cosmologically motivated merger simulations are representative of nature, our results indicate the strong need for improved understanding of how MR and FR estimates trace the galaxy mass (gas+stars) and halo MRs of merging systems as a function of redshift. This analysis underscores the strong need for detailed calibrations of these complex, presumably evolving prescriptions of $C_{\text{merg,pair}}$ and $T_{\text{obs,pair}}$ to constrain the empirical major merger rates confidently. In addition, $C_{\text{merg,pair}}$ and $T_{\text{obs,pair}}$ may be evolving not only as a function of redshift, but also stellar-mass, companion selection criteria ($R_{\text{proj}}, M_1/M_2$), and environment (local overdensity),

which we plan to pursue through a comprehensive analysis of close-pair statistics using mock data sets from the SAMs in Paper 2.

Finally, we recompute $f_{\text{mc}}(z)$ for different companion selections and we find good agreement with previous empirical estimates when we match their companion selection criteria. This confirms the report by Lotz et al. (2011) that part of the reason for the observed study-to-study differences is due to close-pair selection mismatch. Additionally, our comprehensive analysis of the impact of different selection criteria and statistical corrections on $f_{\text{mc}}(z)$ measurements yield the following results:

(i) The redshift evolution of major companion fractions is qualitatively robust to changes in projected separation criterion, with smaller R_{proj} resulting in lower fractions as expected. However, the 5–30 kpc selection (1/2 the area of 5–50 kpc selection) yields similar results to 14–43 kpc criterion owing to the increased probability of physical companions at smaller projected separations.

(ii) Moreover, the redshift evolution of major companion fractions is qualitatively robust to changes in redshift proximity criterion. While the modified Man et al. (Section 4.2) proximity-based fractions are systematically ~ 11 per cent–24 per cent higher than the fiducial values, the hybrid B09-based proximity (Section 4.2) yields 9 per cent–17 per cent smaller major companion fractions at $0.5 < z < 1.5$ and nearly identical values at $z > 1.5$ when compared to fiducial B09 (equation 1) based fractions, owing to the sparsity of spectroscopic-redshift coverage in CANDELS at these redshifts.

(iii) We recompute the SDSS $f_{\text{mc}}(0)$ evolutionary anchor using the modified Man et al. and B09 redshift proximity criteria applied to simulated z_{phot} errors similar to those for $z > 0.5$ CANDELS galaxies. We find the recomputed $z \sim 0$ companion fractions are statistically equivalent ($\Delta f / f_{\text{fid}} \sim 3$ per cent) to the fiducial $f_{\text{mc}}(0)$ values, which demonstrates a high probability of small velocity separations ($\Delta v_{12} \leq 500 \text{ km s}^{-1}$) for companions in 5–50 kpc projected pairs in the SDSS.

(iv) When major companion fractions are corrected for random chance pairing, we find negligible corrections (1 per cent) for SDSS owing to our stringent $\Delta v_{12} \leq 500 \text{ km s}^{-1}$ proximity selection, and approximately equal corrections of ~ 20 per cent at $0.5 < z < 3$ for CANDELS. Importantly, our key results regarding the evolutionary trends of major companion fractions are qualitatively the same with or without this statistical correction.

(v) We also demonstrate that the B09 redshift proximity method does not exclude close-pair systems owing to our use of a simple 68 per cent confidence limit to represent the full z_{phot} PDF.

(vi) Finally, we demonstrate that MR-selected merger rates using unevolving prescriptions of $C_{\text{merg,pair}}$ and $T_{\text{obs,pair}}$ agree with the theoretical predictions at $z < 1.5$, but strongly disagree at $z \gtrsim 1.5$. If we recompute our MR-based rates using an evolving, dynamical-time motivate time-scale by Jiang et al., we find the merger rates are still lower than and disagree with the theoretical merger rate predictions at $z > 1.5$. However, if we use a redshift-dependent time-scale $T_{\text{obs,pair}}(z) \propto (1+z)^{-2}$ motivated by Snyder et al. (2017), our MR-based rates agree with the theory at redshifts $0 < z < 3$.

ACKNOWLEDGEMENTS

This work is dedicated in memory of K. N. V. S. Kishore Babu, a dear friend who inspired KBM to pursue Astrophysics. We are grateful to Peter Behroozi, Mark Brodwin, Philip Hopkins, Kartheik Iyer, Allison Kirkpatrick, Viraj Pandya, Vicente Rodriguez-Gomez, Brett Salmon, and Raymond Simons for helpful comments and discussions during different manifestations of this work. Special thanks

to Daniel Shanabarger for carefully reading the manuscript, thanks to Gillen Brown, Cody Ciaschi, Bandon Decker, Benjamin Floyd, Ripon Saha, and Madalyn Weston for their valuable inputs during the University of Missouri Kansas City (UMKC) Galaxy Evolution Group discussions. DHM and KBM acknowledge support from the Missouri Consortium of National Aeronautics and Space Administration (NASA)’s National Space Grant College and Fellowship Program, and funding from the University of Missouri Research Board. RSS acknowledges support from the Downsborough family, and from the Simons Foundation. SL acknowledges support from the National Research Foundation of Korea grant, no. 2017R1A3A3001362, funded by the Korea government. This work is based on observations taken by the CANDELS Multi-Cycle Treasury Program with the NASA/ESA *HST*, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for Program number HST-GO-12060 was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555. This publication makes use of the SDSS. Funding for the creation and distribution of the SDSS Archive has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the US Department of Energy, the NASA, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. This publication also made use of NASA’s Astrophysics Data System Bibliographic Services, TOPCAT (Tools for OPERations on Catalogues And Tables, Taylor 2005), the core PYTHON package for the astronomy community (Astropy 1.2.1; Astropy Collaboration et al. 2013).

REFERENCES

- Abazajian K. N. et al., 2009, *ApJS*, 182, 543
- Adelman-McCarthy J. K. et al., 2006, *ApJS*, 162, 38
- Ashby M. L. N. et al., 2015, *ApJS*, 218, 33
- Astropy Collaboration et al., 2013, *A&A*, 558, A33
- Barnes J. E., Hernquist L., 1996, *ApJ*, 471, 115
- Bell E. F., McIntosh D. H., Katz N., Weinberg M. D., 2003, *ApJS*, 149, 289
- Bell E. F. et al., 2006a, *ApJ*, 640, 241
- Bell E. F., Phleps S., Somerville R. S., Wolf C., Borch A., Meisenheimer K., 2006b, *ApJ*, 652, 270
- Blanton M. R., Lin H., Lupton R. H., Maley F. M., Young N., Zehavi I., Loveday J., 2003, *AJ*, 125, 2276
- Blanton M. R. et al., 2005, *AJ*, 129, 2562
- Bluck A. F. L., Conselice C. J., Bouwens R. J., Daddi E., Dickinson M., Papovich C., Yan H., 2009, *MNRAS*, 394, L51
- Bluck A. F. L., Conselice C. J., Buitrago F., Grützbauch R., Hoyos C., Mortlock A., Bauer A. E., 2012, *ApJ*, 747, 34
- Bournaud F., Duc P.-A., 2006, *A&A*, 456, 481
- Bournaud F., Dekel A., Teyssier R., Cacciato M., Daddi E., Juneau S., Shankar F., 2011, *ApJ*, 741, L33
- Bower R. G., Benson A. J., Malbon R., Helly J. C., Frenk C. S., Baugh C. M., Cole S., Lacey C. G., 2006, *MNRAS*, 370, 645
- Brammer G. B., van Dokkum P. G., Coppi P., 2008, *ApJ*, 686, 1503
- Brennan R. et al., 2015, *MNRAS*, 451, 2933
- Bridge C. R. et al., 2007, *ApJ*, 659, 931
- Bruzual G., Charlot S., 2003, *MNRAS*, 344, 1000
- Bundy K., Fukugita M., Ellis R. S., Kodama T., Conselice C. J., 2004, *ApJ*, 601, L123
- Bundy K., Fukugita M., Ellis R. S., Targett T. A., Belli S., Kodama T., 2009, *ApJ*, 697, 1369
- Cacciato M., Dekel A., Genel S., 2012, *MNRAS*, 421, 818
- Carlberg R. G., 1990, *ApJ*, 359, L1

- Cassata P. et al., 2005, *MNRAS*, 357, 903
- Cebrián M., Trujillo I., 2014, *MNRAS*, 444, 682
- Ceverino D., Dekel A., Tweed D., Primack J., 2015, *MNRAS*, 447, 3291
- Chabrier G., 2003, *PASP*, 115, 763
- Conselice C. J., 2003, *ApJS*, 147, 1
- Conselice C. J., 2006, *ApJ*, 638, 686
- Conselice C. J., Bershadsky M. A., Dickinson M., Papovich C., 2003, *AJ*, 126, 1183
- Conselice C. J., Rajgor S., Myers R., 2008, *MNRAS*, 386, 909
- Conselice C. J., Yang C., Bluck A. F. L., 2009, *MNRAS*, 394, 1956
- Cox T. J., Jonsson P., Somerville R. S., Primack J. R., Dekel A., 2008, *MNRAS*, 384, 386
- Croton D. J. et al., 2016, *ApJS*, 222, 22
- Dahlen T. et al., 2013, *ApJ*, 775, 93
- Darg D. W. et al., 2010, *MNRAS*, 401, 1043
- de Jong R. S., Bell E. F., 2007, *Island Universes, Astrophysics and Space Science Proceedings*, Vol. 3. Springer, p. 107
- De Propriis R., Conselice C. J., Driver S. P., Liske J., Patton D., Graham A., Allen P., 2007, *ApJ*, 666, 212
- de Ravel L. et al., 2009, *A&A*, 498, 379
- de Ravel L. et al., 2011, preprint ([arXiv:1104.5470](https://arxiv.org/abs/1104.5470))
- Dekel A., Burkert A., 2014, *MNRAS*, 438, 1870
- Dekel A., Sari R., Ceverino D., 2009, *ApJ*, 703, 785
- Dekel A., Zolotov A., Tweed D., Cacciato M., Ceverino D., Primack J. R., 2013, *MNRAS*, 435, 999
- Di Matteo T., Springel V., Hernquist L., 2005, *Nature*, 433, 604
- Di Matteo P., Combes F., Melchior A.-L., Semelin B., 2007, *A&A*, 468, 61
- Di Matteo P., Bournaud F., Martig M., Combes F., Melchior A.-L., Semelin B., 2008, *A&A*, 492, 31
- Domingue D. L., Xu C. K., Jarrett T. H., Cheng Y., 2009, *ApJ*, 695, 1559
- Fakhouri O., Ma C.-P., 2008, *MNRAS*, 386, 577
- Fakhouri O., Ma C.-P., Boylan-Kolchin M., 2010, *MNRAS*, 406, 2267
- Fioc M., Rocca-Volmerange B., 1997, *A&A*, 326, 950
- Galamez A. et al., 2013, *ApJS*, 206, 10
- Genel S., Genzel R., Bouché N., Naab T., Sternberg A., 2009, *ApJ*, 701, 2002
- Gottlöber S., Klypin A., Kravtsov A. V., 2001, *ApJ*, 546, 223
- Grogin N. A. et al., 2005, *ApJ*, 627, L97
- Grogin N. A. et al., 2011, *ApJS*, 197, 35
- Guo Y. et al., 2013, *ApJS*, 207, 24
- Henriques B. M. B., White S. D. M., Thomas P. A., Angulo R., Guo Q., Lemson G., Springel V., Overzier R., 2015, *MNRAS*, 451, 2663
- Hewlett T., Villforth C., Wild V., Mendez-Abreu J., Pawlik M., Rowlands K., 2017, *MNRAS*, 470, 755
- Hopkins P. F., Somerville R. S., Hernquist L., Cox T. J., Robertson B., Li Y., 2006, *ApJ*, 652, 864
- Hopkins P. F., Cox T. J., Kereš D., Hernquist L., 2008, *ApJS*, 175, 390
- Hopkins P. F. et al., 2010a, *ApJ*, 715, 202
- Hopkins P. F. et al., 2010b, *ApJ*, 724, 915
- Jiang C. Y., Jing Y. P., Han J., 2014, *ApJ*, 790, 7
- Jogee S. et al., 2009, *ApJ*, 697, 1971
- Kartaltepe J. S. et al., 2007, *ApJS*, 172, 320
- Kartaltepe J. S. et al., 2015, *ApJS*, 221, 11
- Khochfar S., Burkert A., 2003, *ApJ*, 597, L117
- Khochfar S., Burkert A., 2005, *MNRAS*, 359, 1379
- Kitzbichler M. G., White S. D. M., 2008, *MNRAS*, 391, 1489
- Klypin A., Yepes G., Gottlöber S., Prada F., Heß S., 2016, *MNRAS*, 457, 4340
- Kocevski D. D. et al., 2012, *ApJ*, 744, 148
- Koekemoer A. M. et al., 2011, *ApJS*, 197, 36
- Laidler V. G. et al., 2007, *PASP*, 119, 1325
- Lin L. et al., 2004, *ApJ*, 617, L9
- Lin L. et al., 2008, *ApJ*, 681, 232
- Lin L. et al., 2010, *ApJ*, 718, 1158
- Lofthouse E. K., Kaviraj S., Conselice C. J., Mortlock A., Hartley W., 2017, *MNRAS*, 465, 2895
- López-Sanjuan C., Balcells M., Pérez-González P. G., Barro G., García-Dabó C. E., Gallego J., Zamorano J., 2009, *A&A*, 501, 505
- López-Sanjuan C. et al., 2011, *A&A*, 530, A20
- Lotz J. M., Primack J., Madau P., 2004, *AJ*, 128, 163
- Lotz J. M. et al., 2008, *ApJ*, 672, 177
- Lotz J. M., Jonsson P., Cox T. J., Primack J. R., 2010, *MNRAS*, 404, 575
- Lotz J. M., Jonsson P., Cox T. J., Croton D., Primack J. R., Somerville R. S., Stewart K., 2011, *ApJ*, 742, 103
- Lu Y. et al., 2014, *ApJ*, 795, 123
- Madau P., Dickinson M., 2014, *ARA&A*, 52, 415
- Man A. W. S., Zirm A., Toft S., 2011, preprint ([arXiv:1112.3764](https://arxiv.org/abs/1112.3764))
- Man A. W. S., Toft S., Zirm A. W., Wuyts S., van der Wel A., 2012, *ApJ*, 744, 85
- Man A. W. S., Zirm A. W., Toft S., 2016, *ApJ*, 830, 89
- Martig M., Bournaud F., 2008, *MNRAS*, 385, L38
- Maulbetsch C., Avila-Reese V., Colín P., Gottlöber S., Khalatyan A., Steinmetz M., 2007, *ApJ*, 654, 53
- McIntosh D. H., Guo Y., Hertzberg J., Katz N., Mo H. J., van den Bosch F. C., Yang X., 2008, *MNRAS*, 388, 1537
- Mobasher B. et al., 2015, *ApJ*, 808, 101
- Moster B. P., Somerville R. S., Newman J. A., Rix H.-W., 2011, *ApJ*, 731, 113
- Moustakas J. et al., 2013, *ApJ*, 767, 50
- Mundy C. J., Conselice C. J., Duncan K. J., Almaini O., Häußler B., Hartley W. G., 2017, *MNRAS*, 470, 3507
- Naab T., Khochfar S., Burkert A., 2006, *ApJ*, 636, L81
- Narayanan D. et al., 2010, *MNRAS*, 407, 1701
- Nayyeri H. et al., 2017, *ApJS*, 228, 7
- Neistein E., Dekel A., 2008, *MNRAS*, 388, 1792
- Newman A. B., Ellis R. S., Bundy K., Treu T., 2012, *ApJ*, 746, 162
- Oke J. B., Gunn J. E., 1983, *ApJ*, 266, 713
- Pandya V. et al., 2016, *MNRAS*, 472, 2054
- Patton D. R., Atfield J. E., 2008, *ApJ*, 685, 235
- Patton D. R., Pritchett C. J., Yee H. K. C., Ellingson E., Carlberg R. G., 1997, *ApJ*, 475, 29
- Patton D. R., Carlberg R. G., Marzke R. O., Pritchett C. J., da Costa L. N., Pellegrini P. S., 2000, *ApJ*, 536, 153
- Patton D. R., Ellison S. L., Simard L., McConnachie A. W., Mendel J. T., 2011, *MNRAS*, 412, 591
- Peirani S., Crockett R. M., Geen S., Khochfar S., Kaviraj S., Silk J., 2010, *MNRAS*, 405, 2327
- Pozzetti L. et al., 2010, *A&A*, 523, A13
- Robaina A. R. et al., 2009, *ApJ*, 704, 324
- Robaina A. R., Bell E. F., van der Wel A., Somerville R. S., Skelton R. E., McIntosh D. H., Meisenheimer K., Wolf C., 2010, *ApJ*, 719, 844
- Rodríguez-Gómez V. et al., 2015, *MNRAS*, 449, 49
- Ryan R. E., Jr, Cohen S. H., Windhorst R. A., Silk J., 2008, *ApJ*, 678, 751
- Sanders D. B., Soifer B. T., Elias J. H., Madore B. F., Matthews K., Neugebauer G., Scoville N. Z., 1988, *ApJ*, 325, 74
- Santini P. et al., 2015, *ApJ*, 801, 97
- Snyder G. F., Lotz J. M., Rodríguez-Gómez V., Guimarães R. d. S., Torrey P., Hernquist L., 2017, *MNRAS*, 468, 207
- Somerville R. S., Hopkins P. F., Cox T. J., Robertson B. E., Hernquist L., 2008, *MNRAS*, 391, 481
- Springel V., 2000, *MNRAS*, 312, 859
- Stefanon M. et al., 2017, *ApJS*, 229, 32
- Strauss M. A. et al., 2002, *AJ*, 124, 1810
- Swinbank A. M. et al., 2010, *MNRAS*, 405, 234
- Tal T., van Dokkum P. G., Nelan J., Bezanson R., 2009, *AJ*, 138, 1417
- Targett T. A., Dunlop J. S., McLure R. J., Best P. N., Cirasuolo M., Almaini O., 2011, *MNRAS*, 412, 295
- Taylor M. B., 2005, in Shopbell P., Britton M., Ebert R., eds, *ASP Conf. Ser. Vol. 347, Astronomical Data Analysis Software and Systems XIV*. Astron. Soc. Pac., San Francisco, p. 29
- Treister E., Schawinski K., Urry C. M., Simmons B. D., 2012, *ApJ*, 758, L39
- van der Wel A., Rix H.-W., Holden B. P., Bell E. F., Robaina A. R., 2009, *ApJ*, 706, L120
- Villforth C. et al., 2014, *MNRAS*, 439, 3342
- Villforth C. et al., 2017, *MNRAS*, 466, 812
- Vogelsberger M. et al., 2014, *MNRAS*, 444, 1518

- Wen Z. Z., Zheng X. Z., 2016, *ApJ*, 832, 90
 Weston M. E., McIntosh D. H., Brodwin M., Mann J., Cooper A., McConnell A., Nielsen J. L., 2017, *MNRAS*, 464, 3882
 Williams R. J., Quadri R. F., Franx M., 2011, *ApJ*, 738, L25
 Yang X., Mo H. J., van den Bosch F. C., Pasquali A., Li C., Barden M., 2007, *ApJ*, 671, 153
 York D. G. et al., 2000, *AJ*, 120, 1579
 Younger J. D., Hayward C. C., Narayanan D., Cox T. J., Hernquist L., Jonsson P., 2009, *MNRAS*, 396, L66

We find the redshifts based on S. Wuyts $P(z)$ s agree best with the CANDELS team z_{phot} and available z_{spec} values, where the median (of the CANDELS five fields) outlier fraction¹⁰ of 1.5 per cent and 3 per cent, respectively. This agreement is key to our analysis, and thus we choose the S. Wuyts σ_z values centred on their redshifts to estimate the σ_z for the CANDELS team z_{phot} values. For simplicity, in Fig. A1, we only show the S. Wuyts redshift estimates versus team z_{phot} .

APPENDIX: CHECKING CONSISTENCY OF CANDELS TEAM REDSHIFTS TO SINGLE PARTICIPANT ESTIMATES

In Section 2.3.2, we have discussed adopting the σ_z values based on single participant $P(z)$ as the photometric-redshift errors of z_{phot} to be able to apply the redshift proximity criterion successfully. To choose an appropriate participant for this analysis, we test the consistency of redshift estimates from six participants to the CANDELS team z_{phot} values for the sample of $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ galaxies.

¹⁰ It is defined as the fraction of $\log_{10}(M_{\text{stellar}}/M_{\odot}) \geq 9.7$ galaxies that are outliers. We consider a galaxy to be an outlier if $(z_{\text{participant}} - z_{\text{phot}})/(1 + z_{\text{phot}}) > 0.1$.




Figure A1. Comparison between the single participant (S. Wuyts) $P(z)$ -based redshifts (z_{wuyts}) to the CANDELS team z_{phot} values for $\log_{10} M_{\text{stellar}}/M_{\odot} \geq 9.7$ galaxies (black squares) and those with spectroscopic redshifts (red squares) for five CANDELS fields. In each panel, the top portion of the figure compares both redshift estimates with one-to-one correspondence (red line), and the bottom portion visualizes the redshift normalized scatter defined as $\zeta = (z_{\text{wuyts}} - z_{\text{phot}})/(1 + z_{\text{phot}})$ centred on zero. We find that the S. Wuyts redshift estimates best match to the CANDELS team values with a median outlier fraction of 1.5 per cent and 3 per cent for z_{phot} and z_{spec} samples, respectively (see the text in Appendix for details).

- ¹Department of Physics and Astronomy, University of Missouri-Kansas City, Kansas City, MO 64110, USA
- ²Department of Physics and Astronomy, Rutgers, The State University of New Jersey, NJ 08854-8019, USA
- ³Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA
- ⁴Department of Physics and Astronomy and PITT PACC, University of Pittsburgh, Pittsburgh, PA 15260, USA
- ⁵School of Physics and Astronomy, University of Nottingham, Nottingham, NG7 2RD, UK
- ⁶University of California, 1156 High Street, Santa Cruz, CA 95064, USA
- ⁷Oxford Astrophysics, Denys Wilkinson Building, Oxford OX1 3RH, UK
- ⁸Astrophysics Science Division, Goddard Space Flight Center, MD 20771, USA
- ⁹Department of Physics, University of Bath, Claverton Down, Bath BA2 7AY, UK
- ¹⁰The University of Michigan, 300E West Hall, Ann Arbor, MI 48109-1107, USA
- ¹¹Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel
- ¹²School of Physics and Astronomy, Rochester Institute of Technology, Rochester, NY 14623, USA
- ¹³Department of Physics and Astronomy, Colby College, Waterville, ME 04961, USA
- ¹⁴Center for the Exploration of the Origin of the Universe, Seoul National University, Seoul, Korea

- ¹⁵Goddard Space Flight Center, Code 665, Greenbelt, MD 20771, USA
- ¹⁶Department of Physics and Astronomy, The Johns Hopkins University, Baltimore, MD 21218, USA
- ¹⁷University of California at Berkeley, 501 Campbell Hall #3411, Berkeley, CA 94720-3411, USA
- ¹⁸Department of Astronomy, The University of Texas at Austin, Austin, TX 78712, USA
- ¹⁹INAF-Osservatorio Astronomico di Roma Via Frascati 33, F-00078, Monte Porzio Catone (RM), Roma, Italy
- ²⁰Max Planck Institute für Extraterrestrische Physik, D-85741 Garching bei München, Germany
- ²¹Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA
- ²²Departamento de Astrofísica, Facultad de CC. Físicas, Universidad Complutense de Madrid, Ciudad Universitaria, E-28040 Madrid, Spain
- ²³Aix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, F-13388, Marseille, France
- ²⁴Scientific Support Office, Directorate of Science and Robotic Exploration, European Space Research and Technology Centre (ESA/ESTEC), Keplerlaan 1, NL-2201 AZ Noordwijk, The Netherlands
- ²⁵Leiden Observatory, Leiden University, Niels Bohrweg 2, NL-2333 CA Leiden, Netherlands

This paper has been typeset from a \LaTeX file prepared by the author.