



**HAL**  
open science

## Assessing orbital vs. volcanic control on carbon cycle during the Early Cretaceous

Mathieu Martinez, Beatriz Aguirre-Urreta, Marina Lescano, Guillaume Dera, Julieta Omarini, Maisa Tunik, Tomas Frederichs, Heiko Pälike, Luis O'Dogherty, Roque Aguado, et al.

► **To cite this version:**

Mathieu Martinez, Beatriz Aguirre-Urreta, Marina Lescano, Guillaume Dera, Julieta Omarini, et al.. Assessing orbital vs. volcanic control on carbon cycle during the Early Cretaceous. vEGU21, 2021, à renseigner, Unknown Region. 10.5194/egusphere-egu21-11871 . insu-03669248

**HAL Id: insu-03669248**

**<https://insu.hal.science/insu-03669248>**

Submitted on 31 May 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EGU21-11871, updated on 11 May 2021  
<https://doi.org/10.5194/egusphere-egu21-11871>  
EGU General Assembly 2021  
© Author(s) 2021. This work is distributed under  
the Creative Commons Attribution 4.0 License.



## Assessing orbital vs. volcanic control on carbon cycle during the Early Cretaceous

**Mathieu Martinez**<sup>1</sup>, Beatriz Aguirre-Urreta<sup>2</sup>, Marina Lescano<sup>2</sup>, Guillaume Dera<sup>3</sup>, Julieta Omarini<sup>4,5</sup>, Maisa Tunik<sup>4</sup>, Tomas Frederichs<sup>6</sup>, Heiko Pälike<sup>6</sup>, Luis O'Dogherty<sup>7</sup>, Roque Aguado<sup>8</sup>, Miguel Company<sup>9</sup>, and Jose Sandoval<sup>9</sup>

<sup>1</sup>Univ. Rennes, Géosciences Rennes, Vandoeuvre-les-Nancy, France (mathieu.martinez@univ-rennes1.fr)

<sup>2</sup>Instituto de Estudios Andinos Don Pablo Groeber, CONICET & Universidad de Buenos Aires, Ciudad Universitaria, pabellón 2, 1428 Buenos Aires, Argentina

<sup>3</sup>Géosciences Environnement Toulouse (GET), Université Paul Sabatier Toulouse 3, CNRS UMR 5563, IRD, Toulouse, France

<sup>4</sup>Instituto de Investigación en Paleobiología y Geología, CONICET & Universidad de Río Negro, Sede Alto Valle, 8332 General Roca, Río Negro, Argentina

<sup>5</sup>IIPG, CONICET. Av. J. A. Roca 1242, General Roca, Río Negro, Argentina

<sup>6</sup>MARUM: Center for marine Environmental Sciences, Leobener Strasse 8, Universität Bremen, 28359 Bremen, Germany

<sup>7</sup>Departamento Ciencias de la Tierra, Universidad de Cádiz, CASEM, 11510 Puerto Real, Spain

<sup>8</sup>Departamento de Geología y CEACTEMA, Universidad de Jaén, Campus Científico-Tecnológico de Linares, 23700 Linares, Spain

<sup>9</sup>Departamento de Estratigrafía y Paleontología, Universidad de Granada, 18002 Granada, Spain

The interval from the Valanginian to the Barremian stages (137–121 Ma; Early Cretaceous) is punctuated by several episodes of environmental changes, accompanied by shifts in weathering intensity on the continents and changes in the Tethyan neritic carbonate production. We synthesize here the astrochronology of two recent studies performed in the Neuquén basin, Vocontian Basin and Subbetic Domain (Aguirre-Urreta et al., 2019; Martinez et al., 2020), anchored to CA-ID-TIMS U-Pb ages, which conclusions have been included in the Geologic Time Scale 2020 (Gale et al, in press). We applied this time scale to a compilation of carbon-isotope ratio from belemnites and proxies of detrital supply in the Tethyan area (Vocontian Basin and Subbetic Domain). From this compilation, we show that the episodes of environmental changes are paced by a 2.4-Myr cycle and, with a lower amplitude, a 1.2-Myr cycle. In addition, the new time scale shows the synchronicity between the Weissert Event and the Parana-Etendeka Large Igneous Province. In the series of carbon-isotope ratios measured on belemnite rostra, the amplitude of the 2.4-Myr cycle is twice higher during the Valanginian than in the Late Barremian and three times higher than in the Hauterivian and Early Barremian, suggesting that the activity of the Parana-Etendeka Large Igneous Province amplified the initial orbital forcing to trigger the environmental changes observed during the Mid-Valanginian.

Reference:

Aguirre-Urreta, B., Martinez, M., Schmitz, M., Lescano, M., Omarini, J., Tunik, M., Kuhnert, H.,

Concheyro, A., Rawson, P.F., Ramos, V.A., Reboulet, S., Noclin, N., Frederichs, T., Nickl, A.-L., Pälke, H., 2019. Interhemispheric radio-astrochronological calibration of the time scales from the Andean and the Tethyan areas in the Valanginian–Hauterivian (Early Cretaceous). *Gondwana Research* 70, 104-132. <https://doi.org/10.1016/j.gr.2019.01.006>.

Gale, A.S., Mutterlose, J., Batenburg, S., in press. Chapter 27: The Cretaceous Period, in: Gradstein, F.M., Ogg, J.G., Schmitz, M.D., Ogg, G.M. (Eds.) *Geologic Time Scale 2020*. Elsevier BV, Amsterdam, The Netherlands, pp. 1023–1086.

Martinez, M., Aguado, R., Company, M., Sandoval, J., O'Dogherty, L., 2020. Integrated astrochronology of the Barremian Stage (Early Cretaceous) and its biostratigraphic subdivisions. *Global and Planetary Change* 195, 103368. <https://doi.org/10.1016/j.gloplacha.2020.103368>.