

Breakdown curves of $\text{CH}_2^{(+)}$, $\text{CH}_3^{(+)}$, and $\text{CH}_4^{(+)}$ molecules

I. Construction and application to electron collisions and UV photo-dissociation (*Corrigendum*)

T. IdBarkach¹, M. Chabot¹, K. Béroff², S. Della Negra¹, J. Lesrel¹, F. Geslin¹, A. Le Padellec³,
T. Mahajan², and S. Díaz-Tendero⁴

¹ Institut de Physique Nucléaire d'Orsay (IPNO), CNRS, Univ. Paris Sud, Université Paris-Saclay, 91406 Orsay, France
e-mail: chabot@ipno.in2p3.fr

² Institut des Sciences Moléculaires d'Orsay (ISMO), CNRS, Univ. Paris Sud, Université Paris-Saclay, 91405 Orsay, France

³ Université de Toulouse, UPS-OMP, IRAP, 31028 Toulouse cedex 4, France

⁴ Departamento de Química, Módulo 13, Condensed Matter Physic Center (IFIMAC), Institute for advanced Research in Chemical Sciences (IAdChem), Universidad Autónoma de Madrid, 28049 Madrid, Spain

A&A, 628, A75 (2019), <https://doi.org/10.1051/0004-6361/201935760>

Key words. astrochemistry – ISM: molecules – molecular processes – errata, addenda

A typographical error was made in Eq. (1) of the paper, which expresses the relation between the kinetic energy and the radial distribution for the detector, based on the isotropic hypothesis of the collision. The formula should be written as:

$$f(E, r) = \begin{cases} \frac{r}{\sqrt{1-(r/r_{90}(E))^2}} & \text{if } r < r_{90}(E) \\ 0 & \text{else} \end{cases} \quad (1)$$

We thank Prof. Xavier Urbain for notifying us of the error.