



HAL
open science

Detection of extreme events from IASI observations

Adrien Vuvan, Anne Boynard, Pascal Prunet, Dominique Jolivet, Olivier Lezeaux, Patrice Henry, Claude Camy-Peyret, Pierre-François Coheur, Lieven Clarisse, Cathy Clerbaux

► **To cite this version:**

Adrien Vuvan, Anne Boynard, Pascal Prunet, Dominique Jolivet, Olivier Lezeaux, et al.. Detection of extreme events from IASI observations. 5th IASI International Conference, Dec 2021, Evian, France. insu-03676741

HAL Id: insu-03676741

<https://insu.hal.science/insu-03676741v1>

Submitted on 24 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Introduction

- The 3 IASI instruments on-board the Metop satellites have been sounding the atmospheric composition since 2006 allowing the monitoring of atmospheric chemistry (Clerbaux, et al 2009) and the detection of exceptional events such as fires (Coheur et al., 2009; R'Honi et al., 2013), volcanic eruptions (Clarisse et al., 2008) or pollution events (Boynard et al, 2014).
- The early detection of extreme events is key to take appropriate decisions regarding protection of inhabitants and the environment.
- With IASI providing global observations twice a day in near real time, a new way for the systematic and continuous detection of exceptional atmospheric events to support operational decisions is possible.

Objectives

- Using and improving a method for the detection and characterization of extreme events, which relies on the principal component analysis method (Atkinson, N. C., 2010, 2011; Chefdeville S., 2010).
- Creating a record of extreme events (volcanic eruption and fires) based on IASI data.
- Characterizing and classifying fire, volcanic eruption and pollution events.

Method

The Principal Component method

- The reference database is generated using nearly 120 000 spectra randomly selected in 2013 over the entire globe.
- From the reference database, eigenvectors are created to statistically depict the atmospheric variability during a full year, around different conditions of acquisition over the entire globe.
- By projecting raw apodized spectra (y) on the eigenvectors, reconstructed spectra \tilde{y} are obtained, which are supposed to contain only information on the standard atmospheric variability, except for anomalous events.
- The residual normalized by the IASI noise (N) is calculated as follows: $r = N^{-1}(y - \tilde{y})$.

| Spectral bands (cm ⁻¹) | Peaks (Spectroscopic Database) | Molecule ID |
|------------------------------------|----------------------------------------------------------------------------------------|-------------------------------|
| 667.250 - 667.750 | 667.5 cm ⁻¹ (Q-branch CO ₂) | CO ₂ |
| 711.500 - 713.500 | 713 cm ⁻¹ (Q-branch of HCN v ₂) | HCN |
| 729.250 - 730.000 | 729.25 cm ⁻¹ (C ₂ H ₂ Q-branch v ₂) | C ₂ H ₂ |
| 763.000 - 763.750 | 763 cm ⁻¹ (v ₂ HNO ₂) | HNO ₂ |
| 821.750 - 822.250 | 822 cm ⁻¹ (C ₂ H ₂ Q-branch v ₂) | C ₂ H ₂ |
| 853.500 - 854.250 | 854 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 867.750 - 868.250 | 868 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 878.500 - 880.000 | 879 cm ⁻¹ (v ₂ HNO ₂) | HNO ₂ |
| 887.250 - 888.250 | 888 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 891.750 - 892.250 | 892 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 895.500 - 896.750 | 896 cm ⁻¹ (v ₂ HNO ₂) | HNO ₂ |
| 908.000 - 909.000 | 908.3 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 931.750 - 933.750 | 930 cm ⁻¹ (transition NH ₃ Q-branch) | NH ₃ |
| 949.000 - 950.500 | 949 cm ⁻¹ (v ₂ band vibration liaison CH ₄) | C ₂ H ₆ |
| 966.000 - 968.000 | 967 cm ⁻¹ (transition NH ₃ Q-branch) | NH ₃ |
| 991.750 - 993.500 | 992.8 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 1007.750 - 1008.250 | 1008 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 1034.000 - 1034.750 | 1034 cm ⁻¹ (CH ₃ OH Q-branch) | CH ₃ OH |
| 1046.250 - 1047.250 | 1047 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 1065.750 - 1066.250 | 1066 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 1075.750 - 1076.250 | 1076 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 1084.500 - 1085.750 | 1085 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 1103.000 - 1104.250 | 1104 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 1104.500 - 1105.750 | 1105 cm ⁻¹ (v ₂ band Q-branch) | HCOOH |
| 1180.500 - 1184.750 | 1184 cm ⁻¹ (v ₂ band Q-branch) | CH ₃ COOH |
| 1121.500 - 1122.750 | 1122 cm ⁻¹ (NH ₃ Q-branch) | NH ₃ |
| 1325.750 - 1326.250 | 1326 cm ⁻¹ (v ₂ HNO ₂) | HNO ₂ |
| 1344.500 - 1346.500 | 1345 cm ⁻¹ (SO ₂ v ₂ band) | SO ₂ |
| 1370.500 - 1372.000 | 1371 cm ⁻¹ (estimate absorption in the SO ₂ v ₂ band) | SO ₂ |
| 1375.750 - 1377.000 | 1376 cm ⁻¹ (absorption in the SO ₂ v ₂ band) | SO ₂ |
| 1710.750 - 1711.500 | 1711 cm ⁻¹ (Q branch HNO ₂) | HNO ₂ |
| 1776.750 - 1777.250 | 1777 cm ⁻¹ (HCOOH v ₂ band) | HCOOH |
| 2032.500 - 2087.000 | 2050.2 - 2069.65 cm ⁻¹ (v ₂ OCS branch) | OCS |
| 2111.000 - 2112.250 | 2111.50 cm ⁻¹ (P-branch CO) | CO |
| 2123.000 - 2124.250 | 2123.75 cm ⁻¹ (P-branch CO) | CO |
| 2130.000 - 2132.250 | 2131.75 cm ⁻¹ (P-branch CO) | CO |
| 2157.750 - 2158.725 | 2158.00 cm ⁻¹ (R-branch CO) | CO |
| 2164.750 - 2166.000 | 2165.75 cm ⁻¹ (R-branch CO) | CO |
| Not defined | Unknown | UNKNOWN |

Table.1 : Definition of molecule indicators using known absorption peaks for the characterisation of PCA detections.

The MIN_MAX_RESIDUALS method

- For a given granule, the extreme values of the residuals are saved for each IASI channel for the entire spectral domain (645cm⁻¹ to 2760cm⁻¹) → we obtain an array called « MIN_MAX_RESIDUAL » (Fig.1).
- By analyzing the signal in each IASI channel of the MIN_MAX_RESIDUALS, we compare the strongest peaks to pre-defined micro-windows where specific molecular signatures are expected (Table.1).
- The method is applied on a granule basis to allow the detection in real time.




Fig.1 : Example of strongest features observed in Australian fires residuals during the day (2020/01/01) from the MIN_MAX_RESIDUALS.

Results: analysis of the year 2020

Volcanic eruption events




Fig.3 : Spatial distribution of PCA detection pixels for indicators SO₂ (left) and for indicators HNO₃ (right) for the year 2020.




Fig.4 : Spatial distribution of IASI L2 SO₂ total column for the year 2020.




Fig.6 : Maximum absolute intensity of PCA signal for the 2020 year with referenced eruptions in black.

References

- Atkinson N. C. et al.: Potential for the use of reconstructed IASI radiances in the detection of atmospheric trace gases, <https://doi.org/10.5194/amt-3-991-2010>, 2010.
- Boynard A. et al.: First simultaneous space measurements of atmospheric pollutants in the boundary layer from IASI: a case study in the North China, <https://doi.org/10.1002/2013GL058333>, 2014.
- Chefdeville S.: Analyse de trois années d'« outliers » dans les mesures de l'instrument IASI : détection et étude d'événements extrêmes, rapport de stage Master 1, 2010.
- Clarisse L. et al.: Tracking and quantifying volcanic SO₂ with IASI, the September 2007 eruption at Jebel at Tair, <https://doi.org/10.5194/acp-8-7723-2008>, 2008.
- Clerbaux C. et al.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, Atmos, <https://doi.org/10.5194/acp-9-6041-2009>, 2009.
- Coheur P.-F. et al.: IASI measurements of reactive trace species in biomass burning plumes, <https://doi.org/10.5194/acp-9-5655-2009>, 2009.
- R'Honi Y. et al.: Exceptional emissions of NH₃ and HCOOH in the 2010 Russian wildfires, <https://doi.org/10.5194/acp-13-4171-2013>, 2013.
- Rothman L.S. et al.: The HITRAN2012 molecular spectroscopic database, <https://doi.org/10.1016/j.jqsrt.2013.07.002>, 2012.

Fire events




Fig.7 : Spatial distribution of PCA detection pixels for indicators CO, NH₃, HCOOH, HCN, C₂H₂ and C₂H₄ for the year 2020.

- The PCA method is able to detect volcanic eruptions with the indicators for HNO₃ and SO₂ (Fig. 3).
- The PCA method detect more volcanic eruptions than the L2 data (Fig. 4).
- An excellent agreement is found between SO₂ PCA and L2 data. This allow potentially a fast quantification of this molecule (Fig. 5).
- Record of volcanic eruption events for 2020 with PCA (Fig. 6).

- Indicators for CO, NH₃, HCOOH, HCN, C₂H₂ and C₂H₄ are used for the detection of fires and show different molecular species emitted by fires in Australia, Siberia, Brazil, Africa and California (Fig.7).

- The method allows to detect extreme events, especially in case of the presence of large concentrations.
- Multiple indicators can detect same events, allowing a characterization and a classification of an extreme event.

Future work

- Extending the record of volcanic eruption events based on IASI L1 data for the entire IASI period.
- Validating the method for fire events (comparison with Level 2 data).
- Characterizing and classifying the events.

Acknowledgements

IASI is a joint mission of EUMETSAT and the Centre National d'Etudes Spatiales (CNES, France). This study was supported by a CIFRE grant. The AERIS data infrastructure provides access to the Level 2 IASI data.