

The Pristine survey - XVIII. C-19: tidal debris of a dark matter-dominated globular cluster?

Raphaël Errani, Julio F. Navarro, Rodrigo Ibata, Nicolas Martin, Zhen Yuan, David S. Aguado, Piercarlo Bonifacio, Elisabetta Caffau, Jonay I. González Hernández, Khyati Malhan, et al.

▶ To cite this version:

Raphaël Errani, Julio F. Navarro, Rodrigo Ibata, Nicolas Martin, Zhen Yuan, et al.. The Pristine survey - XVIII. C-19: tidal debris of a dark matter-dominated globular cluster?. Monthly Notices of the Royal Astronomical Society, 2022, 514, pp.3532-3540. 10.1093/mnras/stac1516. insu-03707832

HAL Id: insu-03707832 https://insu.hal.science/insu-03707832

Submitted on 11 Apr 2023 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Monthly Notices of the ROYAL ASTRONOMICAL SOCIETY

MNRAS **514**, 3532–3540 (2022) Advance Access publication 2022 June 2

The *Pristine* survey – XVIII. C-19: tidal debris of a dark matter-dominated globular cluster?

Raphaël Errani,^{1*} Julio F. Navarro,² Rodrigo Ibata,¹ Nicolas Martin[®],¹ Zhen Yuan[®],¹ David S. Aguado,^{3,4} Piercarlo Bonifacio[®],⁵ Elisabetta Caffau,⁵ Jonay I. González Hernández,^{6,7} Khyati Malhan,⁸ Rubén Sánchez-Janssen,⁹ Federico Sestito[®],² Else Starkenburg,¹⁰ Guillaume F. Thomas^{6,7} and Kim A. Venn[®]²

¹Université de Strasbourg, CNRS, Observatoire astronomique de Strasbourg, UMR 7550, F-67000 Strasbourg, France

¹⁰Kapteyn Astronomical Institute, University of Groningen, Landleven 12, NL-9747 AD Groningen, the Netherlands

Accepted 2022 May 27. Received 2022 May 20; in original form 2022 March 3

ABSTRACT

The recently discovered C-19 stellar stream is a collection of kinematically associated metal-poor stars in the halo of the Milky Way lacking an obvious progenitor. The stream spans across an arc of ~15° in the sky, and orbit-fitting suggests an apocentric distance of ~20 kpc and a pericentre of ~10 kpc. The narrow metallicity dispersion of stars with available spectra, together with light element abundance variations, suggests a globular cluster (GC) origin. The observed metallicity ([Fe/H] \approx -3.4), however, is much lower than that of any known GC. In addition, the width and velocity dispersion of the stream are similar to those expected from disrupting dwarf galaxies, and substantially larger than the tidal debris of GCs able to disrupt on C-19's orbit. We propose here an unconventional model where the C-19 progenitor is a dark matter-dominated stellar system with GC-like abundance patterns. We use *N*-body simulations to show that the tidal disruption of a ~100 pc King-model stellar component embedded in a ~20 km s⁻¹ cuspy cold dark matter halo yields debris consistent with C-19's observed width and velocity dispersion. The stellar component of the progenitor is fully disrupted, and is spread over two distinct streams, one corresponding to C-19 and another possibly hiding behind the Galactic plane. If such companion stream were found, it would suggest that dark matter-dominated dwarfs may also develop GC-like enrichment patterns, a finding that would inform our theoretical understanding of the formation of multiple populations in GCs and dwarf galaxies alike.

Key words: globular clusters: general – Galaxy: kinematics and dynamics – galaxies: dwarf – galaxies: evolution – dark matter.

1 INTRODUCTION

The number of known stellar streams in the Milky Way (MW) has increased dramatically over the last few decades thanks to discoveries made by the Sloan Digital Sky Survey (e.g. Odenkirchen et al. 2001; Belokurov et al. 2006; Koposov et al. 2012), the Dark Energy Survey (e.g. Shipp et al. 2018), and the *Gaia* satellite (e.g. Malhan, Ibata & Martin 2018; Ibata, Malhan & Martin 2019; Ibata et al. 2021), among others.

Stellar streams form through tidal stripping of globular clusters (GCs) or dwarf galaxies (DGs) as they orbit in the gravitational potential of the MW. Since GCs are in general denser and physically smaller than dwarfs, they produce narrow, kinematically cold stellar

streams (e.g. the Pal-5 stream; Odenkirchen et al. 2001). Dwarfs, on the other hand, are expected to produce thicker and kinematically hotter streams (e.g. the massive Sagittarius stream; Ibata et al. 2001; Newberg et al. 2002). Stream kinematic properties are particularly useful when inferring the progenitor properties of 'orphan' streams, where no bound remnant of the original system can be identified.

Kinematic signatures, however, are inconclusive on their own. One reason is the recent discovery of 'ultra-faint dwarfs' (UFDs, usually defined as DGs with $L < 10^5 L_{\odot}$), whose structural properties (size, velocity dispersion, and luminosity) overlap those of some GCs (Simon 2019). Indeed, it often isn't clear whether a system is a GC or a UFD unless a fairly precise estimate of its dynamical mass can be made: objects dominated by dark matter are considered UFDs, whereas self-gravitating ones are regarded as GCs.

This litmus test, however, cannot be applied to 'orphan streams' as they are not self-gravitating systems, so other diagnostics need to

²Department of Physics and Astronomy, University of Victoria, Victoria, BC V8P 5C2, Canada

³Dipartimento di Fisica e Astrofisica, Univerisità degli Studi di Firenze, via G. Sansone 1, I-50019 Sesto Fiorentino, Italy

⁴INAF/Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125 Firenze, Italy

⁵GEPI, Observatoire de Paris, Université PSL, CNRS, 5 Place Jules Janssen, F-92195 Meudon, France

⁶Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

⁷Dpto. Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain

⁸ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

⁹UK Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

^{*} E-mail: errani@unistra.fr

be considered. A popular one makes use of the ubiquitous presence of 'multiple populations' (MPs) in massive, old GCs. These MPs show a narrow dispersion in metallicity (as measured by [Fe/H]), but wide variations in the light-element abundances of individual stars (Gratton, Sneden & Carretta 2004; Gratton et al. 2019). These variations are also often correlated (or anti-correlated) between elements such as C, N, O, Na, Mg, and Al, signalling a complex nucleosynthetic process that operated during GC formation and whose origin has not yet been fully elucidated (Bastian & Lardo 2018).

UFDs, on the other hand, appear to have formed their stars differently, with available data indicating a fairly broad dispersion in [Fe/H] (σ _[Fe/H]) (Leaman 2012; Willman & Strader 2012), and no GC-like correlations between light-element abundances (Tolstoy, Hill & Tosi 2009; Simon 2019). It is therefore common practice to associate narrow σ _[Fe/H] and wide variations in light-element abundances with GCs, although even here important exceptions apply: For example, the well-studied ω Cen GC has a fairly broad σ _[Fe/H] (Norris & Da Costa 1995), and younger, more metal-rich GCs appear to lack MPs (Milone et al. 2020), and some UFDs, like Grus-1,¹ have unresolved dispersion in the metallicity (and velocity) of the stars measured so far (Walker et al. 2016; Ji et al. 2019).

Stream kinematic properties are also inconclusive as progenitor diagnostics because thin, cold streams originating from GCs may be thickened and 'heated' if formed in Galactic subhaloes prior to accretion into the MW (Carlberg 2018, 2020; Malhan, Valluri & Freese 2021). Such heating may also occur because of the gravitational effect of dark matter substructures in the MW halo (Ibata et al. 2002; Johnston, Spergel & Haydn 2002; Peñarrubia 2019), or because of the dispersal of stream stars in axisymmetric (Erkal, Sanders & Belokurov 2016) as well as non-axisymmetric or timeevolving potentials (Price-Whelan et al. 2016).

The recent detection of C-19, a stream of metal-poor stars recently discovered in the MW halo by combining the power of *Gaia* proper motions with the photometric metallicities of the Pristine survey, has renewed interest in this topic (Martin et al. 2022). Interestingly, C-19 has kinematic properties that suggests a DG origin [i.e. a width and velocity dispersion of \approx 180 pc (for a heliocentric distance of 20 kpc) and \approx 6 km s⁻¹, respectively; see Yuan et al. 2022], but the element abundances of the few stars measured so far indicate a narrow spread in metallicity ($\sigma_{\text{[Fe/H]}} < 0.18$; Martin et al. 2022) and light-element abundance variations akin to those in GCs. At the same time, at [Fe/H] ≈ -3.4 , C-19 is more metal-poor than any other DG (Chiti et al. 2021) or known GC (Beasley et al. 2019), so elucidating the properties of its progenitor is particularly intriguing.

We explore here an unconventional model, where C-19's progenitor is envisioned as a dark matter-dominated DG with GC-like enrichment patterns. As we show below, this model circumvents a major difficulty afflicting GC progenitor models: no known GC can disrupt fully on C-19's orbit and leave a stream as thick and as kinematically hot as observed. Either the stream has been heated after formation by some of the external effects mentioned earlier, or C-19's progenitor was unlike any GC or dwarf system that has survived to the present.

This contribution is meant to explore the consequences of assuming that C-19's progenitor was a dark matter-dominated dwarf, and to identify features of this scenario that may falsify it. Our intention is not to deny the possibility that the C-19 stream may have indeed originated from a conventional GC (with an unconventional dynamical heating history), nor to survey the full range of possible origins for C-19, but rather to explore the possibility that the most metal-poor stellar systems to form in the early Universe might have differed systematically from the examples (GCs or UFDs) that have survived to the present day. Should C-19's progenitor be actually a dark matter-dominated low-surface-brightness UFD, as proposed here, this would be extremely useful for theoretical models of the origins of MPs in GCs, many of which assume that the high density environment of present-day GCs is critical for the existence of MPs.

The paper is structured as follows. We introduce the numerical modelling of the stream in Section 2, and discuss our main results in Section 3. In particular, we explore the evolution of systems modelled after three well-known GCs (NGC 2419, Pal-5, and Pal-14) placed on C-19's current orbit, and discuss whether it is possible to match the properties of C-19 without leaving behind at the same time a luminous bound remnant (Section 3.1). We then explore the stripping of a cold dark matter-dominated DG model in Section 3.2. We end with a brief summary of our main conclusions in Section 4.

2 C-19 MODELLING

2.1 Observational summary

The extreme metallicity of C-19 stars {[Fe/H] = -3.38 ± 0.06 (stat.) ± 0.20 (syst.); Martin et al. 2022} is well below the metallicity of the second most metal-poor stream discovered so far, the Phoenix stream ([Fe/H] ≈ -2.7 ; Wan et al. 2020), and well below the most metal-poor DG, Tucana 2 ([Fe/H] ≈ -2.8 ; Chiti et al. 2021). It is also more metal-poor than all known GCs, implying that its progenitor was a unique system by any account.

As mentioned above, its width ² and velocity dispersion³ (177 \pm 25 pc and 6.2^{+2.0}_{-1.4} km s⁻¹, respectively; see Yuan et al. 2022) suggest a DG origin rather than a GC one. We show this in Fig. 1, where we compare the width and velocity dispersion of C-19 (red symbols with errorbars) with a compilation⁴ of half-light radii, $R_{\rm h}$, and line-of-sight velocity dispersions, $\sigma_{\rm los}$, of GCs (filled grey triangles) and DGs (filled blue circles) with $L < 10^7 L_{\odot}$. The same figure also includes data for other streams in the halo of the MW (blue crosses for DG streams, and grey crosses for GC streams). The GC stream

⁴Data for DGs is as compiled in McConnachie (2012) (version 2021 January, width updated data for Antlia 2, Crater 2 (Ji et al. 2021), Tucana (Taibi et al. 2020), Tucana 2 (Chiti et al. 2021), And 19 (Collins et al. 2020), and And 21 (Collins et al. 2021)), and for GCs as in Harris (1996) (version 2010 December, and updated half-light radii and velocity dispersions for Pal-5 from Kuzma et al. 2015; Gieles et al. 2021; for NGC 2419 from Baumgardt et al. 2009, and for Pal-14 from Hilker 2006; Jordi et al. 2009). Width and velocity dispersions for stellar streams are from the S⁵ survey (Li et al. 2021b, table 1), with stream width defined as one standard deviation of a Gaussian profile. In addition, data is shown for the Sagittarius stream [width and velocity dispersion from Ibata et al. (2020) and Gibbons, Belokurov & Evans (2017), respectively], the Palomar-5 stream (Bonaca et al. 2020; Kuzma et al. 2015), the GD-1 stream (Koposov, Rix & Hogg 2010; Gialluca, Naidu & Bonaca 2021), and the C-19 stream [both width and dispersion from Yuan et al. (2022), for a heliocentric distance of 20 kpc].

¹Ji et al. (2019) identify Grus-1 as a UFD based on its member stars being deficient in neutron-capture elements (indicative for DGs, see e.g. Frebel & Norris 2015), while Walker et al. (2016) refer to Grus-1 more agnostically as an 'ultrafaint stellar system'.

²The width of the stream is defined as one standard deviation in right ascension α at fixed declination δ for a heliocentric distance of 20 kpc, multiplied by $\cos(\delta)$.

³The transverse velocity dispersion is estimated as the dispersion in line-ofsight velocity with respect to the fitted orbit at fixed declination δ .



Figure 1. Half-light radii $R_{\rm h}$ and line-of-sight velocity dispersions $\sigma_{\rm los}$ of DGs (blue, filled circles) and GCs (grey, filled triangles) with luminosities $L < 10^7 L_{\odot}$, compared against the characteristic width and velocity dispersion (see footnotes 2, 3) of the C-19 stream (red errorbars). Indicative values of width and velocity dispersion are also shown for other stellar streams (crosses, in blue for streams with DG progenitors, and in grey for those with GC progenitors). N-body models explored in this work are marked in red. References for the data shown are given in footnote 4. A line of constant crossing time $T_{\rm cross} \sim 2\pi R_{\rm h}/\sigma_{\rm los}$ that equals the circular orbital time at pericentre $T_{\text{peri}} \approx 2\pi r_{\text{peri}}/(240 \,\text{km}\,\text{s}^{-1})$ is shown in grey. Systems that fall to the right of this line are likely to experience significant tidal stripping on the C-19 orbit. The two dashed diagonal lines correspond to self-gravitating Plummer models (equation 3) with fixed total masses of $M_{\star} = 6 \times 10^3 \,\mathrm{M_{\odot}}$ (the minimum stellar mass of C-19, see the text), and of $10^7 \, M_{\odot}$. The characteristic width and velocity dispersion of the C-19 stream place it closer to the regime of DGs than to the regime of GCs.



Figure 2. Metallicity [Fe/H] and luminosity *L* of the DGs (blue, filled circles) and GCs (grey, filled triangles) shown in Fig. 1. The C-19 progenitor (shown in red, with errorbars in [Fe/H] corresponding to ± 0.2 dex systemic uncertainty) appears unique in its properties when compared against known DGs and GCs.

closest in width and velocity dispersion to C-19 is the 'Atlas and Aliqa Uma' stream, which is likely perturbed from interaction with the Sagittarius dwarf (Li et al. 2021a).

Fig. 2, on the other hand, compares the metallicity of C-19 and its inferred luminosity $\sim 3 \times 10^3 L_{\odot}$ (more appropriately, its lower limit, as derived by Martin et al. 2022) with DGs and GCs. Clearly, regardless of its luminosity, the progenitor of C-19 has no known parallel in our Local Group.

Aside from its width and velocity dispersion, another C-19 property useful for comparison with our models is its angular extent. This is estimated to be $\sim 15^{\circ}$ in the sky, with only one star candidate beyond that region (Yuan et al. 2022), and no identifiable bound remnant. We shall take this to imply that the detected stream contains the bulk of the stars in the disrupted progenitor (assumed fully disrupted). Since streams grow longer and sparser as they orbit the Galactic halo, this, in turn, suggests that the stream is relatively 'young', in the sense that its disruption has happened relatively recently (i.e. in the most recent past few orbital periods).

Because the stream's pericentric distance of $r_{\rm peri} \sim 10 \,\rm kpc}$ and apocentric distance $r_{\rm apo} \sim 20 \,\rm kpc}$ indicate an orbital period of just $T_{\rm orb} \sim 300 \,\rm Myr$, this implies that either (i) the progenitor has been recently accreted into the MW halo, or that (ii) the stellar component of C-19 was deeply embedded in a more massive (dark) halo that has only recently been sufficiently stripped to affect its stars. Since (i) is disfavoured by the relatively short orbital time and small apocentric distance, we take this observation to support, at least in principle, the premise that C-19's progenitor could have been more massive and extended in the past.

2.2 Numerical methods

We summarize in this section the *N*-body models and numerical tools used in this work.

2.2.1 Host galaxy model

We model the MW using a static, axisymmetric, analytical potential with a circular velocity of $V_c(R_{\odot}) = 240 \text{ km s}^{-1}$ at the solar circle, $R_{\odot} = 8.3 \text{ kpc}$. We use the parametrization of Errani & Peñarrubia (2020), which in turn is derived from the McMillan (2011) model.

In this parametrization, the dark matter halo is modelled using a spherical NFW (Navarro–Frenk–White) profile with virial⁵ mass $M_{200} = 1.15 \times 10^{12} \,\mathrm{M_{\odot}}$, virial radius $r_{200} = 192 \,\mathrm{kpc}$, and concentration parameter c = 9.5. The bulge is modelled using a Hernquist (1990) sphere with total mass $M_{\rm b} = 2.1 \times 10^{10} \,\mathrm{M_{\odot}}$ and scale length $a_{\rm b} = 1.3 \,\mathrm{kpc}$. Thick and thin discs are axisymmetric Miyamoto & Nagai (1975) potentials with masses $M_{\rm thick} = 2.0 \times 10^{10} \,\mathrm{M_{\odot}}$ and $M_{\rm thin} = 5.9 \times 10^{10} \,\mathrm{M_{\odot}}$, and horizontal and vertical scale lengths $a_{\rm thick} = 4.4 \,\mathrm{kpc}$, $a_{\rm thin} = 3.9 \,\mathrm{kpc}$ and $b_{\rm thick} = 0.92 \,\mathrm{kpc}$, $b_{\rm thin} = 0.31 \,\mathrm{kpc}$, respectively.

2.2.2 Orbit

We use the potential described above to fit an orbit that matches approximately the sky location $\{\alpha, \delta\}$, proper motions $\{\mu_{\alpha}*, \mu_{\delta}\}$, and radial velocities v_{los} (if available) of C-19 member candidates identified by Yuan et al. (2022). Our orbit fitting assumes a solar motion relative to the local standard of rest as in Schönrich, Binney & Dehnen (2010).

The fitted orbit has a pericentre of $r_{\text{peri}} = 11 \text{ kpc}$, an apocentre of $r_{\text{apo}} = 23 \text{ kpc}$, and an orbital period of $T_{\text{orb}} = 0.34 \text{ Gyr}$. The orbit is shown in Fig. 3 in heliocentric Galactic coordinates (l, b). Fig. 4 shows the orbit in Galactocentric Cartesian coordinates, with the Sun at $(x_{\odot}, y_{\odot}, z_{\odot}) = (-R_{\odot}, 0, 0)$.

⁵Virial quantities are computed within spheres of mean density equal to 200 times the critical density for closure, $\rho_{\rm crit} = 3H_0^2/8\pi G$, and are indicated with a '200' subscript.



Figure 3. Orbit fitted to the C-19 members in heliocentric galactic coordinates *l*, *b*, integrated backward (black line) and forward (grey line) in time from the apocentre closest to the C-19 stream. Apo- and pericentres are marked along the orbit. Stream members with radial velocity measurements are shown in red; those where only proper motion measurements are available are shown in blue. The blue shaded area corresponds to the segment with right ascension $345^{\circ} < \alpha < 360^{\circ}$ and declination $-15^{\circ} < \delta < 40^{\circ}$ shown in Figs 7 and 9. The current orbital pole ($l \approx -140^{\circ}$, $b \approx -6^{\circ}$) is marked by a cross.



Figure 4. Orbit fitted to C-19 member (black solid for integration backward in time, grey for forward integration), like Fig. 3, but in galactocentric Cartesian coordinates. Apo- and pericentres are marked along the orbit. The Sun is located at $(x_{\odot}, y_{\odot}, z_{\odot}) = (-R_{\odot}, 0, 0)$. The bottom right-hand panel shows a zoom-in on the main stream segment, with C-19 member velocities shown as vectors. For those member stars without radial velocity measurement (shown in blue), the radial velocity of the orbit is used in addition to the measured proper motions to compute the velocity vector shown.

The bottom right-hand panel of Fig. 4 shows a zoom-in on the main stream segment, with velocities shown as vectors. For member stars without measured radial velocities (shown in blue), the radial velocity of the orbit at their location is used to obtain the velocity vectors in Cartesian coordinates.

2.3 Globular cluster models

We model GCs as self-gravitating Plummer spheres with isotropic velocity dispersion,

$$\rho_{\rm P}(r) = \frac{3M_{\star}}{4\pi a^3} \left[1 + (r/a)^2 \right]^{-5/2},\tag{1}$$

Table 1. Projected half-light radii R_h , line-of-sight velocity dispersions σ_{los} , and total masses M_{\star} of the GC models used in this work. Masses are estimated using equation (3).

Model	Equivalent GC	$R_{\rm h}~({\rm pc})$	$\sigma_{\rm los}~({\rm kms^{-1}})$	$M_{\star}~({ m M}_{\odot})$	
GC-1	NGC 2419	19.0 ^a	4.14 ± 0.48^{a}	5.1×10^5	
GC-2	Pal-5	18.7^{b}	1.2 ± 0.3^{c}	4.2×10^{4}	
GC-3	Pal-14	27.8^{d}	0.38 ± 0.12^e	6.3×10^3	

^aBaumgardt et al. (2009).

^bGieles et al. (2021).

^cKuzma et al. (2015).

^dHilker (2006).

^eJordi et al. (2009).

with total mass M_{\star} and scale radius *a*. For spherical Plummer models, the projected half-light radius coincides with scale radius, $R_{\rm h} = a$.

The line-of-sight velocity dispersion profile equals (see e.g. Dejonghe 1987, equation 43)

$$\sigma_{\rm los}^2(R) = \frac{GM_{\star}}{a} \frac{3\pi}{64} \left[1 + (R/a)^2 \right]^{-1/2} , \qquad (2)$$

which implies that the total mass, M_{\star} , may be estimated from the projected half-light radius, $R_{\rm h}$, and the central line-of-sight velocity dispersion, $\sigma_{\rm los}(0)$,

$$M_{\star} \approx 6.8 R_{\rm h} \sigma_{\rm los}^2(0)/G \,. \tag{3}$$

The importance of Galactic tides on the evolution of a cluster may be gauged by comparing its inner crossing time, $T_{\rm cross} \sim 2\pi R_{\rm h}/\sigma_{\rm los}(0)$, with the circular orbital time at pericentre, $T_{\rm peri} \approx 2\pi r_{\rm peri}/(240 \,{\rm km \, s^{-1}})$.

The latter is shown in Fig. 1 as a solid grey line. Systems on or to the right of this line are expected to be severely affected by tides on such orbit; systems well to the left should be able to survive relatively unscathed for many orbital periods.

As may be seen from Fig. 1, most GCs are too dense to disrupt on an orbit like that of C-19, with $r_{\text{peri}} = 11$ kpc; only Pal-14, the least dense known MW GC, is likely to disrupt fully on such orbit. With this caveat, we explore three different self-gravitating GC models, with parameters chosen as *GC-1*, *GC-2*, and *GC-3* in Fig. 1. These models span a fairly wide range of velocity dispersions and radii, and hence also of stellar mass, as listed in Table 1. Model *GC-1* is a fairly dense and massive ($M_{\star} \approx 5.1 \times 10^5 \text{ M}_{\odot}$) GC, with parameters resembling those of NGC 2419. Model *GC-3*, on the other hand, is a fluffy, low mass ($M_{\star} \approx 6 \times 10^3 \text{ M}_{\odot}$) cluster resembling Pal-14. Model *GC-2* has intermediate properties, resembling those of Pal-5.

The *N*-body realizations have $N = 10^6$ particles each and are generated using the code of Errani & Peñarrubia (2020), available online.⁶

2.4 Dark matter-dominated dwarf galaxy-like model

Our DG-like model assumes that the stellar component of C-19's progenitor may be modelled as spherical King-like models (King 1962) embedded in a Navarro–Frenk–White (hereafter, NFW; Navarro, Frenk & White 1996, 1997) halo. Stars are assumed to be subdominant gravitationally, and can therefore be modelled as massless tracers of the NFW halo, as detailed below.

The NFW dark matter subhalo model is given by

$$\rho_{\rm NFW} = \rho_{\rm s} \left(r/r_{\rm s} \right)^{-1} \left(1 + r/r_{\rm s} \right)^{-2},\tag{4}$$

⁶https://github.com/rerrani/nbopy

where ρ_0 and r_s denote the scale density and radius, respectively. The corresponding circular velocity curve $V_c = \sqrt{GM(< r)/r}$ reaches its maximum of $V_{\rm mx} \approx 1.65 r_{\rm s} \sqrt{G\rho_{\rm s}}$ at a radius $r_{\rm mx} \approx 2.16 r_{\rm s}$. The *N*-body realization consists of 10⁷ particles. The NFW density profile is exponentially tapered beyond 10 $r_{\rm s}$ to prevent the divergence of the total mass.

We choose the virial mass of the subhalo to approximately correspond to the minimum halo mass able to form stars imposed by the hydrogen cooling limit (HCL) at an early redshift, chosen here to be z = 5 (Benítez-Llambay et al. 2019). The motivation for this choice is mainly one of simplicity: Given C-19's extreme metallicity, its progenitor is likely one of the oldest systems to form in the Universe, and therefore if dark matter dominated, it should populate some of the lowest mass systems where efficient energy dissipation of the collapse energy may occur. More specifically, we chose a maximum circular velocity of $V_{\rm mx} = 20 \,\rm km \, s^{-1}$, and a radius of maximum circular velocity $r_{\rm mx} = 1.7 \,\rm kpc$ in order to match the z = 5 mean mass–concentration relation (Ludlow et al. 2016). The mass enclosed within $r_{\rm mx}$ equals $M_{\rm mx} \approx 1.6 \times 10^8 \,\rm M_{\odot}$ and the virial mass of the subhalo is $M_{200} \approx 3 \times 10^8 \,\rm M_{\odot}$.

Stars are embedded in this subhalo as massless tracers, and can therefore be modelled by attaching to each dark matter particle a 'stellar probability' that depends on their initial binding energy. We express the binding energy in dimensionless units,

$$\mathcal{E} = 1 - E/\Phi_0 \,, \tag{5}$$

where $\Phi_0 \equiv \Phi(r=0)$ corresponds to the potential minimum of the dark matter halo. In these units, the most-bound state is $\mathcal{E} = 0$, and $\mathcal{E} = 1$ corresponds to the boundary between bound and unbound. For NFW profiles, the potential minimum is directly related to the maximum circular velocity, $\Phi_{0,\text{NFW}} = -4\pi G \rho_s r_s^2 \approx -4.64 V_{\text{mx}}^2$ (for details, see e.g. Errani et al. 2021). For the tapered NFW models used in this work, we have $\Phi_0 \approx -4.32 V_{\text{mx}}^2$.

We select the stellar energies so that the stars form a deeply embedded cored profile within the NFW halo dark matter cusp, with a (2D) half-light radius of $R_h \approx 85$ pc, and a central line-ofsight velocity dispersion of $\sigma_{los} \approx 8.1$ km s⁻¹. We accomplish this by simply assigning a uniform 'stellar probability' to all dark matter particles between a specified minimum, \mathcal{E}_{min} , and maximum, \mathcal{E}_{max} , initial binding energy (see top panel of Fig. 5). Probabilities are set to zero for dark matter particles outside that range.

This choice results in a stellar profile well described by a sharply truncated King (1962) surface brightness profile with concentration parameter $c_{\rm K} = 1$ and King tidal radius $R_{\rm t,K} = 210 \,\rm pc$ (see Table 2 for a summary of the model parameters).

Fig. 5 shows the resulting stellar density and velocity dispersion profiles (shown in red) compared with the dark matter (in blue). Note that because the stars are assumed to be subdominant, the stellar mass of the galaxy can be rescaled arbitrarily. Our DG-like model for the C-19 progenitor assumes a total stellar luminosity consistent with the minimum inferred from the minimum integrated luminosity of C-19, i.e. $3 \times 10^3 L_{\odot}$ (Martin et al. 2022). Using a mass-to-light ratio for old stellar populations of ~2 (Maraston 2005; Woo, Courteau & Dekel 2008), this corresponds to a stellar mass of ~6 $\times 10^3 M_{\odot}$.

2.5 N-body code

We use the particle-mesh code SUPERBOX (Fellhauer et al. 2000) to evolve the *N*-body particles of the GC and DG models in the specified MW potential. The code uses two comoving grids, centred on the *N*-body model, with 128³ grid cells each. The sizes of the comoving grids are chosen relative to the size of the *N*-body model, resulting for



Figure 5. Initial conditions of the DG-like model. Dark matter properties are shown in blue, those of stars in red. The total stellar luminosity is scaled to $3 \times 10^3 L_{\odot}$. The top panel shows the initial energy distributions of dark matter and stars, as well as the energy distribution of those particles which remain bound after 0.8, 1.8, and 2.7 Gyr, corresponding to the snapshots shown in Fig. 8. The initial (2D) surface brightness profile is shown in the bottom left, together with the projected initial dark matter density (arbitrary normalization). The stellar profile is cored, with a projected half-light radius of $R_h \approx 85$ pc. A King (1962) surface brightness profile with concentration parameter $c_{\rm K} = 1$ and King core radius $R_{\rm K} = 210$ pc is shown, for reference, as a black dotted curve. The stellar line-of-sight velocity dispersion profile is shown in the bottom right, with a central line-of-sight velocity of $\sigma_{\rm los} \approx$ $8.1 \,{\rm km \, s^{-1}}$.

Table 2. Properties of the dark matter subhalo model used in this work. The maximum circular velocity V_{mx} is chosen to approximately correspond to the minimum halo mass imposed by the HCL (Benítez-Llambay et al. 2019). The radius of maximum circular velocity r_{mx} matches the z = 5 mean mass–concentration relation (Ludlow et al. 2016). Half-light radius and line-of-sight velocity dispersion correspond to the massless stellar tracer with energies $0.08 \le \mathcal{E}_{\star}0.12$ as shown in Fig. 5.

Model	r _{mx} (kpc)	$V_{\rm mx}~({\rm kms^{-1}})$	$M_{ m mx}~({ m M}_{\odot})$	$R_{\rm h}~({\rm kpc})$	$\sigma_{\rm los}~({\rm kms^{-1}})$
DG	1.7	20.0	1.6×10^8	0.085	8.1

Plummer (NFW) models in cell spacings of $\Delta x \approx 10 a/128$ ($\Delta x \approx 10 r_{\rm mx}/128$) for the medium resolution grid and $\Delta x \approx a/128$ ($\Delta x \approx r_{\rm mx}/128$) for the high resolution grid.

The equations of motions are solved using a leap-frog-scheme with a time-step $\Delta t = T_a/200$ for Plummer models, where T_a denotes the period of a circular orbit at radius *a*, and with a time-step $\Delta t = T_{mx}/400$ for NFW models, where T_{mx} denotes the period of a circular orbit at r_{mx} . Using these choices, a circular orbit with a radius equal to the grid resolution $r = \Delta x$ is resolved, in the case of Plummer models, with ≈ 120 time-steps, and in the case of NFW models, with ≈ 16 steps.



Figure 6. NGC 2419-like model *GC-1*, evolved for 10 Gyr, in heliocentric Galactic coordinates. The position of the cluster remnant is indicated by an arrow. Shown also are MW GCs (Baumgardt et al. 2019) with apo- and pericentric distances roughly compatible with C-19. No GC lies on the C-19 orbit (black curve), and all GCs are considerably more metal-rich than C-19 (see colour coding).

3 RESULTS

We begin by analysing the streams produced by the C-19 GC progenitors described in Section 2.3, before considering dark matter-dominated progenitors in the following subsection.

3.1 Self-gravitating GC model progenitors

3.1.1 NGC 2419-like progenitor (GC-1)

The *GC-1* model represents a GC with structural properties similar to that of NGC 2419. It has an initial velocity dispersion (~4 km s⁻¹) slightly below that measured for the C-19 stream (~6 km s⁻¹), but it is much more massive ($M_{\star} \approx 5 \times 10^5 \text{ M}_{\odot}$) than the lower limit on the stellar mass inferred for C-19 ($M_{\star} \gtrsim 6 \times 10^3 \text{ M}_{\odot}$). More importantly, NGC 2419 has a half-light radius of only ~19 pc, implying a relatively high density compared with the inner mean density of the host Galaxy model at its pericentric radius (i.e. it lies well to the left of the grey line in Fig. 1).

Because of this, *GC-1* is barely affected by Galactic tides, losing only 15 per cent of its stars over 10 Gyr of evolution. This model would predict a massive bound remnant on the same orbit as C-19, at odds with observations, where no known cluster has been found to share C-19's orbit. We show this in Fig. 6, which shows stars in grey from the GC-1 progenitor in heliocentric Galactic coordinates l, b. The location of the bound luminous remnant (with stellar mass $\sim 4 \times 10^5 \,\mathrm{M_{\odot}}$) is indicated by an arrow.

The figure also shows the positions of all known MW GCs with apo- and pericentric distances in a fairly wide range around the corresponding C-19 values (i.e. $10 \text{ kpc} < r_{apo} < 30 \text{ kpc}$ and pericentric distances $5 \text{ kpc} < r_{peri} < 20 \text{ kpc}$, taken from Baumgardt et al. 2019). No GC shown in Fig. 6 lies closer than 4° from C-19's current orbital path. In addition, as mentioned earlier, the metallicity of all known GCs is far in excess of that measure for C-19 members, making them incompatible with being potential progenitors to the C-19 stream. We conclude that GC-1 is not viable as a possible progenitor of the C-19 stream.



Figure 7. GC model *GC-3* with a Pal-14-like progenitor on the C-19 orbit, evolved for 10 Gyr. The top panel compares the line-of-sight velocities of the *N*-body model as a function of declination δ against the observational data (red points with errorbars). A histogram, summing over all stars with declination $15^{\circ} < \delta < 40^{\circ}$, is shown on the right, comparing the C-19 data (red) against the *N*-body model (grey). The bottom panel compares right ascension α as a function of declination δ (red and blue points correspond to member stars with and without line-of-sight velocity measurements, respectively). The model completely disrupts, and after 5 Gyr roughly matches the C-19 stream in length and width. However, the velocity dispersion of the simulated stream is much lower than that of the C-19 member stars.

3.1.2 Pal 5-like progenitor (GC-2)

The *GC*-2 model has properties similar to those of Pal-5, a relatively fluffy MW GC, substantially less dense and less massive than NGC 2419. As may be seen from Fig. 1, this cluster model still has a characteristic crossing time $T_{\rm cross} \sim 2\pi R_{\rm h}/\sigma_{\rm los}$ shorter than the circular orbital time $T_{\rm peri} \approx 2\pi r_{\rm peri}/(240 \,{\rm km \, s^{-1}})$ at pericentre, and, as a result, we would not expect *GC*-2 to be fully disrupted by tides.

Indeed, our results show that *GC-2* loses ~65 per cent of its initial mass after completing ~30 orbits in about 10 Gyr. This implies that *GC-2* would predict that a bound remnant with total luminosity $\sim 10^4 L_{\odot}$ should remain visible at present, also at odds with observations. We hence conclude that, just like for GC-1, the GC-2 model is not a viable progenitor of C-19.

3.1.3 Pal 14-like progenitor (GC-3)

Finally, we consider a much lower density GC model (*GC-3*), inspired by Pal-14, the least dense GC known in the Galaxy. When placed on C-19's orbit, this system is tidally destroyed after about a dozen of orbits. In order to match C-19's observed length and width, we are forced to study the remnant after just ~5 Gyr of evolution. The comparison with C-19, shown in Fig. 7, shows that *GC-3* is able to approximately match the width (and length) of the observed stream without leaving behind a bound luminous remnant. However, the velocity dispersion of this stream (~1.4 km s⁻¹) is much lower than that observed for C-19 (~6 km s⁻¹), even if analysed just after disruption. Since streams generally lengthen and become kinematically colder as they evolve (see e.g. Helmi & White 1999), it is unlikely that this model can reproduce C-19 even if analysed at a later time. While 'fanning' on polar orbits in axisymmetric potentials



Figure 8. Distribution of dark matter (grey) and stars (colour) for the DG-like model at different times after accretion, in galactocentric Cartesian coordinates. At T = 0.8 Gyr (left-hand panel), some dark matter has been stripped, but the stellar component is largely unaffected by tides. At T = 1.8 Gyr (second panel from left), a stellar stream has formed, but a bound stellar cluster remains centred on the remnant dark matter subhalo. At T = 2.6 Gyr (third panel from left), the stellar cluster has been completely dispersed, and no bound stars remain within the dark matter subhalo. This scenario is a possible model for the formation of the C-19 stream, with the leading stream corresponding to C-19. Observed stream members are shown using red and blue points (with distances adjusted), in good agreement with the leading stream of the disrupted DG-like model. Note that while no bound stellar remnant remains, the dark matter subhalo does not fully disrupt. The right-hand panel (T = 2.7 Gyr) shows a similar situation, but here, the trailing tidal stream matches the C-19 data. The centre of mass of the bound dark matter remnant is shown using an arrow in all panels. A video of this simulation is available on the journal website.

may increase the width of streams over time (see e.g. Erkal et al. 2016), even after 10 Gyr of evolution, the velocity dispersion of the *GC-3* stream remains below that of C-19. Earlier accretion times are in conflict with the observed length of the C-19 stream: Evolving the *GC-3* model for \sim 10 Gyr results in a stream more than twice as long as C-19. We conclude that this model does not provide a good approximation to C-19's data either.

3.2 Dark matter-dominated dwarf galaxy-like model

The failure of the GC progenitors studied above to reproduce C-19 is not entirely unexpected, and may be traced to the assumption that these systems were initially self-gravitating. Indeed, GC progenitors with initial stellar mass similar to that inferred for C-19 must lie close to the grey dashed $6 \times 10^3 M_{\odot}$ line in Fig. 1. To disrupt fully on C-19's orbit, GC progenitors must also lie to the right of the solid grey line, indicating the circular orbit time at a pericentre of 11 kpc.

These restrictions put potential C-19 progenitors quite far from C-19's locus in the half-light radius versus velocity dispersion plane. Indeed, tentatively⁷ applying equation (3) to the measured width and dispersion of the C-19 stream suggests a dynamical mass of $\sim 10^7 \, M_{\odot}$ (see dashed grey line in Fig. 1).

This mass corresponds roughly to that enclosed within $\sim 200 \text{ pc}$ of a halo at the HCL boundary (see Section 2.4), a fact that motivates the choice of stellar component parameters described in Section 2.4, which are shown by the red symbol labelled '*DG*' in Fig. 1. Dark matter-dominated models have an advantage over self-gravitating GC models, for they allow us to construct systems where the stellar component may fully disrupt, leaving behind no self-bound stellar remnant, even if a small dark matter clump may remain. We explore these issues in more detail below.

As an illustration, we choose here to simulate the tidal evolution of the dwarf-like model for roughly eight pericentric passages, or \sim 3 Gyr. Tides strip mass gradually and 'outside-in' in terms of initial binding energy (Errani et al. 2021). This is shown in the top panel of Fig. 5, where thin dashed lines outline the sharp energy truncation experienced by the halo at various stages of evolution (i.e. after 0.8, 1.8, and 2.7 Gyr, respectively). Note that after 2.7 Gyr of evolution, there is no bound stellar remnant left, although a dense clump of dark matter (consisting of the innermost initial dark matter cusp) remains bound (see also Errani & Navarro 2021).

Fig. 8 shows projections of the dwarf-like system at those stages of the evolution. The dark matter surface density ρ_{2D} is shown in grey, and stellar surface brightness Σ in colour. At t = 0.8 Gyr (or after two pericentric passages, left-hand panel), some dark matter has been stripped, but the stellar component is still largely unaffected by tides. At t = 1.8 Gyr (second panel from left), a stellar stream has formed, and a bound stellar remnant remains centred on the dark matter subhalo (marked by the arrows in each panel). At t = 2.6 Gyr and t = 2.7 Gyr (third panel from left, and right-hand panel), the stellar component has been completely dispersed, and no bound stellar progenitor is detectable, consistent with observations.

This scenario leads to the formation of two fairly symmetric streams, each of which, at t = 2.6 Gyr and t = 2.7 Gyr, respectively, matches fairly well the width and velocity dispersion of the C-19 stream. This is illustrated in Fig. 9 for the trailing stream at t = 2.7 Gyr; the top panel compares line-of-sight velocities of the simulation model against C-19 member stars, while the bottom panel shows the positions in the sky. Adjacent histograms sum over the main stream segment, defined as all stars with $15^{\circ} \le \delta \le 40^{\circ}$.

What about the other stream? In the case where C-19 is identified with the trailing stream (i.e. t = 2.7 Gyr in Fig. 8), this would predict the presence of a second, nearly symmetric stream segment that is currently not observed. However, if C-19 were identified with the leading stream (i.e. at t = 2.6 Gyr), then the bulk of the trailing stream would be at very low Galactic latitudes, and could have easily escaped detection. At least from the kinematic perspective, this scenario provides an attractive explanation for the unexpected width and velocity dispersion of C-19.

The times quoted here are for illustrative purpose only: Matching models with earlier (or later) accretion time may be constructed through small modifications to the initial stellar binding energies. As the angular separation between leading and trailing stream increases

⁷Equation (3) was derived for self-gravitating Plummer spheres. Here, we apply it tentatively to the width and velocity dispersion of the C-19 stream, assuming that these values are a rough proxy for size and velocity dispersion of the C-19 progenitor at the time when the stream formed.



Figure 9. Comparison of the DG-like model against velocities (top panel) and positions (bottom panel) of C-19 member stars (as in Fig. 7). The width and dispersion of the *N*-body model match that of the C-19 stream members. The arrow shows the position of the centre of mass of the bound dark matter remnant.

over time, the discovery of a second companion stream to C-19, sharing its orbital pole⁸ ($l \approx -140^\circ$, $b \approx -6^\circ$, see Fig. 3), might allow to constrain the time of accretion of the C-19 progenitor on to the MW. If the C-19 progenitor is still in the process of disrupting, leading and trailing streams may appear as a single, connected structure.

4 SUMMARY AND CONCLUSIONS

The C-19 stellar stream is the most metal-poor stream known to date, with a mean metallicity of [Fe/H] ≈ -3.4 . As only a very small spread in metallicity ($\sigma_{\text{[Fe/H]}} \leq 0.18$) is observed, it is tempting to associate C-19 with a tidally disrupted GC. However, the unusual width and velocity dispersion of the stream suggest a possible DG origin for the stream. We have explored here scenarios where the progenitor of C-19 is either a self-gravitating GC or a dark matter dominated stellar system similar in structure to dwarf spheroidal galaxies.

Our main conclusions may be summarized as follows:

(i) The C-19 stellar stream has a width and a velocity dispersion exceeding those of known GC streams, and those structurally place C-19 closer to the regime of DGs than that of GC streams.

(ii) Self-gravitating GC progenitors similar in structure to NGC 2419 and Pal-5 are too dense to be fully disrupted on C-19's orbit over a period of 10 Gyr. These may be ruled out by the lack of a bound luminous remnant associated with C-19.

(iii) A self-gravitating progenitor similar to the Pal-14 GC matches the integrated luminosity of C-19 and disrupts fully, but yields a stream too kinematically cold to match C-19's observed velocity dispersion.

(iv) A dark matter-dominated DG-like progenitor, where stars follow a King-like profile embedded in a low-mass cuspy cold dark

⁸While the pole of the C-19 orbit precesses over time, the angular separation between the orbital poles of leading and trailing streams grows at a much lower pace.

matter model near the HCL, is able to reproduce the width and velocity dispersion of C-19, leaving behind no bound stellar remnant.

(v) The DG progenitor model predicts the existence of a second tidal stream, similar to C-19 but at Galactic latitudes likely coinciding with the Galactic plane, hindering its discovery. The discovery of such a stream, symmetric in structure and luminosity to C-19, would strongly favour the dwarf-like progenitor model.

Although the dynamical evidence seems to favour a DG-like progenitor for C-19, it is difficult to completely rule out GClike progenitors. Indeed, different dynamical processes have been discussed in the literature that could increase the width and velocity dispersion of GC streams beyond the values expected from stripping in smooth tidal fields such as the one considered here. These include the gravitational effect of (dark) matter clumps in the MW halo (Ibata et al. 2002; Johnston et al. 2002; Peñarrubia 2019), the dispersal of stream stars in non-axisymmetric or time-evolving potentials (Price-Whelan et al. 2016), tidal evolution of the GC within a subhalo prior to accretion on to the MW (Carlberg 2018, 2020; Malhan et al. 2021). as well as collisional processes within the GC that might facilitate tidal disruption (Gieles et al. 2021). What observations could help us distinguish a GC stream that has been thickened and 'heated' from the DG-like progenitor we explored here? A non-exhaustive list of diagnostics include the following:

(i) Signs of collisional processes in the progenitor, like masssegregation of C-19 member stars along the orbit, might give dynamical hints on the progenitor dynamics and allow to distinguish a (collisional) GC from a (collision-less) dwarf-like system.

(ii) Our DG-progenitor model predicts two rather symmetric tidal streams. The detection of a second stream segment, with a luminosity and structure similar to that of the current C-19 segment, would be an indicator in favour of the dwarf-like model, in contrast to the more irregular streams of pre-accretion stripped GCs (see e.g. Malhan et al. 2021).

(iii) A better understanding of the origin of light-element abundance patterns in GCs and in low-metallicity UFDs, together with improved observational constraints on a larger number of C-19 stars, might allow to use them as discriminating diagnostics of C-19's progenitor. Is it actually possible for DGs to develop GC-like light-element enrichment patterns while keeping a narrow spread in metallicity?

If the hypothesis of a dark matter-dominated progenitor holds, then C-19 is allowing a glimpse at galaxy evolution at the faintest end of the galaxy luminosity function, and would suggest that the earliest, most metal-poor stellar systems formed may differ substantially from the more metal-rich globulars and dwarfs that have survived to the present day.

ACKNOWLEDGEMENTS

RE, RI, and ZY acknowledge funding from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (grant agreement no. 834148). DA acknowledges support from the ERC Starting Grant H2020/808240. JIGH acknowledges financial support from the Spanish Ministry of Science and Innovation (MICINN) project PID2020-117493GB-I00. ES acknowledges funding through VIDI grant 'Pushing Galactic Archaeology to its limits' (with project number VI.Vidi.193.093), which is funded by the Dutch Research Council (NWO). KAV is grateful for funding through the Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grants and CRE- ATE programs. ZY acknowledges support from the French National Research Agency (ANR) funded project 'Pristine' (ANR-18-CE31-0017).

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Bastian N., Lardo C., 2018, ARA&A, 56, 83
- Baumgardt H., Côté P., Hilker M., Rejkuba M., Mieske S., Djorgovski S. G., Stetson P., 2009, MNRAS, 396, 2051
- Baumgardt H., Hilker M., Sollima A., Bellini A., 2019, MNRAS, 482, 5138
- Beasley M. A., Leaman R., Gallart C., Larsen S. S., Battaglia G., Monelli M., Pedreros M. H., 2019, MNRAS, 487, 1986
- Belokurov V. et al., 2006, ApJ, 642, L137
- Benítez-Llambay A., Frenk C. S., Ludlow A. D., Navarro J. F., 2019, MNRAS, 488, 2387
- Bonaca A. et al., 2020, ApJ, 889, 70
- Carlberg R. G., 2018, ApJ, 861, 69
- Carlberg R. G., 2020, ApJ, 889, 107
- Chiti A. et al., 2021, Nat. Astron., 5, 392
- Collins M. L. M., Tollerud E. J., Rich R. M., Ibata R. A., Martin N. F., Chapman S. C., Gilbert K. M., Preston J., 2020, MNRAS, 491, 3496
- Collins M. L. M. et al., 2021, MNRAS, 505, 5686
- Dejonghe H., 1987, MNRAS, 224, 13
- Erkal D., Sanders J. L., Belokurov V., 2016, MNRAS, 461, 1590
- Errani R., Navarro J. F., 2021, MNRAS, 505, 18
- Errani R., Peñarrubia J., 2020, MNRAS, 491, 4591
- Errani R., Navarro J. F., Ibata R., Peñarrubia J., 2021, MNRAS, 511, 6001
- Fellhauer M., Kroupa P., Baumgardt H., Bien R., Boily C. M., Spurzem R., Wassmer N., 2000, New Astron., 5, 305
- Frebel A., Norris J. E., 2015, ARA&A, 53, 631
- Gialluca M. T., Naidu R. P., Bonaca A., 2021, ApJ, 911, L32
- Gibbons S. L. J., Belokurov V., Evans N. W., 2017, MNRAS, 464, 794
- Gieles M., Erkal D., Antonini F., Balbinot E., Peñarrubia J., 2021, Nat. Astron., 5, 957
- Gratton R., Sneden C., Carretta E., 2004, ARA&A, 42, 385
- Gratton R., Bragaglia A., Carretta E., D'Orazi V., Lucatello S., Sollima A., 2019, A&AR, 27, 8
- Harris W. E., 1996, AJ, 112, 1487
- Helmi A., White S. D. M., 1999, MNRAS, 307, 495
- Hernquist L., 1990, ApJ, 356, 359
- Hilker M., 2006, A&A, 448, 171
- Ibata R., Irwin M., Lewis G. F., Stolte A., 2001, ApJ, 547, L133
- Ibata R., Bellazzini M., Thomas G., Malhan K., Martin N., Famaey B., Siebert A., 2020, ApJ, 891, L19
- Ibata R. et al., 2021, ApJ, 914, 123
- Ibata R. A., Lewis G. F., Irwin M. J., Quinn T., 2002, MNRAS, 332, 915
- Ibata R. A., Malhan K., Martin N. F., 2019, ApJ, 872, 152
- Ji A. P., Simon J. D., Frebel A., Venn K. A., Hansen T. T., 2019, ApJ, 870, 83

- Ji A. P. et al., 2021, ApJ, 921, 20
- Johnston K. V., Spergel D. N., Haydn C., 2002, ApJ, 570, 656
- Jordi K. et al., 2009, AJ, 137, 4586
- King I., 1962, AJ, 67, 471
- Koposov S. E., Rix H.-W., Hogg D. W., 2010, ApJ, 712, 260
- Koposov S. E. et al., 2012, ApJ, 750, 80
- Kuzma P. B., Da Costa G. S., Keller S. C., Maunder E., 2015, MNRAS, 446, 3297
- Leaman R., 2012, AJ, 144, 183
- Li T. S. et al., 2021a, ApJ, 911, 149
- Li T. S. et al., 2021b, ApJ, 928, 18
- Ludlow A. D., Bose S., Angulo R. E., Wang L., Hellwing W. A., Navarro J. F., Cole S., Frenk C. S., 2016, MNRAS, 460, 1214
- McConnachie A. W., 2012, AJ, 144, 4
- McMillan P. J., 2011, MNRAS, 414, 2446 Malhan K., Ibata R. A., Martin N. F., 2018, MNRAS, 481, 3442
- Malhan K., Valluri M., Freese K., 2021, MNRAS, 501, 179
- Maraston C., 2005, MNRAS, 362, 799
- Martin N. F. et al., 2022, Nature, 601, 45
- Milone A. P. et al., 2020, MNRAS, 491, 515
- Miyamoto M., Nagai R., 1975, PASJ, 27, 533
- Navarro J. F., Frenk C. S., White S. D. M., 1996, ApJ, 462, 563
- Navarro J. F., Frenk C. S., White S. D. M., 1997, ApJ, 490, 493
- Newberg H. J. et al., 2002, ApJ, 569, 245
- Norris J. E., Da Costa G. S., 1995, ApJ, 447, 680
- Odenkirchen M. et al., 2001, ApJ, 548, L165
- Peñarrubia J., 2019, MNRAS, 484, 5409
- Price-Whelan A. M., Johnston K. V., Valluri M., Pearson S., Küpper A. H. W., Hogg D. W., 2016, MNRAS, 455, 1079
- Schönrich R., Binney J., Dehnen W., 2010, MNRAS, 403, 1829
- Shipp N. et al., 2018, ApJ, 862, 114
- Simon J. D., 2019, ARA&A, 57, 375
- Taibi S., Battaglia G., Rejkuba M., Leaman R., Kacharov N., Iorio G., Jablonka P., Zoccali M., 2020, A&A, 635, A152
- Tolstoy E., Hill V., Tosi M., 2009, ARA&A, 47, 371
- Walker M. G. et al., 2016, ApJ, 819, 53
- Wan Z. et al., 2020, Nature, 583, 768
- Willman B., Strader J., 2012, AJ, 144, 76
- Woo J., Courteau S., Dekel A., 2008, MNRAS, 390, 1453
- Yuan Z. et al., 2022, MNRAS, 514, 1664

SUPPORTING INFORMATION

Supplementary data are available at MNRAS online.

C19_model.mp4

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a TEX/LATEX file prepared by the author.