



Enhanced Carbonate Counter Pump and upwelling strengths in the Indian sector of the Southern Ocean during MIS 11

Margaux Brandon, Stéphanie Duchamp-Alphonse, Elisabeth Michel, Amaelle Landais, Gulay Isguder, Patricia Richard, Nicolas Pige, Franck Bassinot, Samuel L. Jaccard, Annachiara Bartolini

► To cite this version:

Margaux Brandon, Stéphanie Duchamp-Alphonse, Elisabeth Michel, Amaelle Landais, Gulay Isguder, et al.. Enhanced Carbonate Counter Pump and upwelling strengths in the Indian sector of the Southern Ocean during MIS 11. *Quaternary Science Reviews*, 2022, 287, pp.107556. 10.1016/j.quascirev.2022.107556 . insu-03710162

HAL Id: insu-03710162

<https://insu.hal.science/insu-03710162>

Submitted on 20 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Enhanced Carbonate Counter Pump and upwelling strengths in the Indian sector of the Southern Ocean during MIS 11

Brandon Margaux ^{1,2,*}, Duchamp-Alphonse Stéphanie ¹, Michel Elisabeth ², Landais Amaëlle ², Isguder Gulay ², Richard Patricia ², Pige Nicolas ³, Bassinot Franck ², Jaccard Samuel L. ⁴, Bartolini Annachiara ⁵

¹ Université Paris-Saclay, CNRS, GEOPS, 91405, Orsay, France

² Université Paris-Saclay, CNRS, CEA, UVSQ, Laboratoire des Sciences du climat et de l'environnement, 91191, Gif-sur-Yvette, France

³ Univ Lyon, UCBL, ENSL, UJM, CNRS, LGL-TPE, F-69622, Villeurbanne, France

⁴ Institute of Earth Sciences, University of Lausanne, 1015, Lausanne, Switzerland

⁵ Muséum National d'Histoire Naturelle, Département Origines & Évolution, CR2P MNHN, CNRS, Sorbonne Université, 8 rue Buffon CP38, 75005, Paris, France

* Corresponding author : Margaux Brandon, email address : margaux.brandon@locean.ipsl.fr

Abstract :

While numerous studies have highlighted the central role of Southern Ocean (SO) dynamics in modulating rapid increases in atmospheric CO₂ concentrations during deglaciations, fewer studies have yet focused on the impact of the Biological Carbon Pump - and more specifically the Carbonate Counter Pump (CCP) - in contributing to increase the CO₂ concentration in oceanic surface waters and thus, in the atmosphere. Here, we present micropaleontological (coccolith, planktonic foraminifera) and geochemical (CaCO₃, CaXRF, $\delta^{13}\text{CN}$, pachyderma) constraints from sediment core MD04-2718 retrieved in the Polar Front Zone of the Indian Ocean covering the time interval spanning Marine Isotope Stage (MIS) 12 to MIS 10 (440,000–360,000 years). We compare our results with published records from the SO to reconstruct past changes in CCP and upwelling dynamics and understand their leverage on the ocean-atmosphere partitioning of CO₂. We demonstrate that the sharp increase in atmospheric pCO₂ during Termination V was likely associated with enhanced deep-water ventilation in the SO, that promoted the release of previously sequestered CO₂ to the ocean surface as the westerly wind belt and the frontal system migrated southwards. Enhanced CCP is observed later, during MIS 11, and is likely the consequence of higher sea surface temperature and higher nutrient availability due to the reinvigoration of SO upwelling leading to increased coccolith (and to a lesser degree, planktonic foraminifera) production and export. The low eccentricity signal recorded during MIS 11 might have additionally strengthened the CCP, exerting a specific control on Gephyrocapsa morphotypes. In addition to the strong global biological productivity and higher carbon storage on land, these synergistic mechanisms may have permitted to shape the distinctive 30 ka-long pCO₂ plateau characteristic of MIS 11.

Highlights

► Multiproxy approach to reconstruct Carbonate Counter Pump strength and upwelling dynamics. ► Increase in $p\text{CO}_2$ during Termination V coincides with enhanced upwelling. ► Strong Carbonate Counter Pump during MIS 11 in the Indian Southern Ocean. ► Enhanced upwelling and Carbonate Counter Pump helped shape the $p\text{CO}_2$ plateau.

Keywords : Southern Ocean, Marine Isotope Stage 11, Coccolith, Planktonic foraminifera, Geochemistry, Carbonate Counter Pump, Upwelling, Atmospheric CO_2 concentration

1. Introduction

The Earth's climate system of the past 800 000 years (800 ka) has been characterised by glacial-interglacial cycles that are tightly coupled to pCO₂ changes (Lüthi et al., 2008; Bereiter et al., 2015). Causes underlying these changes, especially in conjunction with deglaciations (glacial Terminations) that registered pCO₂ rises of about 50-100 ppm, are still under debate. It is recognised that complex, synergistic interactions involving changes in sea-ice cover, surface water stratification and marine productivity in the SO together with the dynamics of the westerly winds may have played a major role in modulating the efficiency of the physical and biological carbon pumps (François et al., 1997; Stephens and Keeling, 2000; Sigman and Boyle, 2000; Hodell et al., 2003; Toggweiler et al., 2006; Jaccard et al., 2013; Ferrari et al., 2014; Abelmann et al., 2015; Galbraith & Jaccard, 2015; Menviel et al., 2018; Gottschalk et al., 2019; Stein et al., 2020; Sigman et al., 2021). The reinvigoration of SO upwelling has been held responsible for a substantial part of the abrupt pCO₂ rise observed during Terminations (Hodell et al., 2003; Anderson et al., 2009; Skinner et al., 2010; Siani et al., 2013, Jaccard et al., 2016; Rae et al., 2018; Gottschalk et al., 2020; Sigman et al., 2021) while an increase in the efficiency of the Biological Carbon Pump (BCP) might have accounted for up to 40-50 ppmv CO₂ drawdown during glacial maxima (Hain et al., 2010; Martinez-Garcia et al., 2009; Jaccard et al., 2013). However, due to a lack of high-resolution primary productivity reconstructions and, more generally, to the scarcity of multiproxy approaches (Duncan et al., 2016; Duchamp-Alphonse et al., 2018), it remains difficult to apprehend the spatio-temporal patterns underpinning changes in the BCP and compare them to the SO upwelling dynamics to understand their impact on pCO₂. Furthermore, most of the studies aiming to reconstruct changes in the strength and efficiency of the BCP focus on the export of phytoplankton-derived organic carbon to the deep ocean (i.e., the Soft Tissue Pump (STP)) that directly lowers atmospheric CO₂ (Martin et al., 1990; Martinez-Garcia et al., 2009; Martinez-Garcia et al., 2014; Thöle et al., 2019). They typically overlook the export of phytoplankton- and zooplankton-derived calcium carbonate (CaCO₃), which raises surface water CO₂

(i.e., the CCP) (Baumann et al., 2004; Barker et al., 2006; Broecker and Clark, 2009; Salter et al., 2014; Rembauville et al., 2016), and contributes to the CaCO_3 /organic carbon ratio in the biogenic particle flux to the sea-floor (namely the rain ratio), especially in the SO (Duchamp-Alphonse et al., 2018; Krumhardt et al., 2020). This omission may relate to the fact that calcareous phytoplankton is in part represented by coccoliths, which detailed analysis requires sophisticated tools such as SYRACO (SYstème de Reconnaissance Automatique des Coccolithes), a software using automated optical microscope that allows the robust determination of coccolith mass and thus, coccolithophore calcification rates, based on the optical properties (birefringence) of calcite under cross-polarized light (Beaufort et al., 2005; 2014). It is also probably linked to the lack of well-preserved carbonate fraction in sediment cores of the SO, making the reconstruction of biogenic carbonate export in the past challenging (Barker et al., 2006; Jaccard et al., 2013; Thöle et al., 2019).

Amongst the interglacials of the past 800 ka, MIS 11 is the longest (~30 ka) as well as one of the warmest (Tzedakis et al., 2022 and references therein). Following the cold glacial MIS 12, MIS 11 begins at the end of Termination V (TV), the first Termination recording high amplitude increases in both pCO_2 (~ 90 ppmv) (Bereiter et al., 2015; Nehrbass-Ahles et al., 2020) and Antarctic air temperatures (> 12°C) (Jouzel et al., 2007). MIS 11 occurs during an eccentricity minimum (Laskar et al., 2004) and coincides with strong global biosphere productivity (Brandon et al., 2020), a major phase in coral reef expansion (Husson et al., 2018), and a climax in coccolithophore (*Gephyrocapsa* sp.) production (Barker et al., 2006; Beaufort et al., 2022). Recent studies document the increase in coccolith abundances over this time interval in the Pacific and Atlantic sectors of the SO (Flores et al., 2003; Saavedra-Pellitero et al., 2017a; 2017b). However, no studies exist in the Indian sector of the SO yet, and none focus on the CCP and its impact on atmospheric pCO_2 based on combined coccolith and planktonic foraminifera carbonate export data.

Here, we provide coccolith and planktonic foraminifera abundance and mass from core MD04-2718 retrieved in the Indian sector of the SO. Using these data, we demonstrate the biogenic origin of the sedimentary CaCO_3 at this site and calibrate the carbonate signal to reconstruct high resolution changes in CCP strength across the time interval covering MIS 12 – MIS 10 (440-360 ka), including TV and MIS 11. Combined with reconstructed summer SST and $\delta^{13}\text{C}_{\text{N. pachyderma}}$ data obtained from the same core, as well as previously published micropaleontological and geochemical data from the Pacific and Atlantic sectors of the SO as well as geochemical data from EPICA Dome C (EDC), our study further highlights the synergistic contribution of changing global productivity and terrestrial carbon storage, SO ventilation and CCP strength on atmospheric CO_2 during this key period of the past 800 ka.

2. Regional oceanic setting

The SO is delimited by the Antarctic continent to the South and by the Subtropical front (STF) to the North (Fig. 1a). The eastward-flowing Antarctic Circumpolar Current (ACC) connects the Atlantic, Pacific and Indian sectors of the SO. This current, driven by Coriolis and the Southern Hemisphere westerlies, is characterised by the largest mass transport of all ocean currents, averaging 150 Sv ($10^6 \text{m}^3 \text{s}^{-1}$) in the Indian Ocean (Park et al., 2009). Its Northern and Southern Boundaries (NB and SB) are often delimited around 40°S and 60°S respectively, but their exact location depends on bottom topography (Sokolov and Rintoul, 2009, Park et al., 2019) and varies latitudinally (Freeman et al., 2016). Three hydrological fronts generally characterised by stronger flows, are defined in the ACC: the Subantarctic Front (SAF), the Polar Front (PF) and the Southern Antarctic Circumpolar Front (SACCF) (Fig. 1). They help defining the Subantarctic Zone (SAZ) between the STF and the SAF, the Polar Front Zone (PFZ) between the SAF and the PF, and the Antarctic Zone (AAZ) delimited by the PF to the North, and the SB to the South (Orsi et al., 1995, Park et al., 2019). Within the Indian sector of the SO, the Kerguelen Plateau that extends from ~ 46°S to 63°S and between ~ 62°E and 85°E, is the second largest submarine plateau in the world. Due to its great meridional extent and topography, it constitutes a major bathymetric constrain to the ACC flow and thus exerts a significant impact on both ACC transport and front patterns (Park et al., 2009; Sokolov and Rintoul, 2009). In this context, the PF is located alongside the southern coast of the Kerguelen archipelago (Park et al., 2014). The circulation in the Southern Ocean is also regulated by the upwelling of CO_2 - and nutrient-rich Circumpolar Deep Waters (CDW) that resurface at the Antarctic divergence in the AAZ (Fig. 1c). These waters represent the source for the Subantarctic Mode Water (SAMW) and the Antarctic Intermediate Water (AAIW) that flow northward. Today, the change in the strength of upwelling affects the partitioning of CO_2 between the ocean subsurface and the atmosphere on decadal timescales (Lovenduski et al., 2008; Landschützer et al., 2015; Gruber et al., 2019).

In the SO, primary productivity is primarily steered by bottom-up control involving both physical and biogeochemical processes, which are spatially and temporally heterogeneous. Today, the presence/absence of sea ice together with the progressive temperature decrease polewards subdivide the SO into distinct provinces characterised by specific functional assemblages (Boyd, 2002; Constable et al., 2014). In ecosystems that remain ice-free throughout the year, i.e., within the Permanently Open Ocean Zone (POOZ), primary productivity is commonly co-limited by light and micronutrient availability (Boyd et al., 2002; Moore et al., 2013). While the residual surface ocean macronutrient concentrations in the SO are among the highest globally (Pollard et al., 2006) largely due to the large-scale upwelling of CDW at the Antarctic divergence, primary productivity is relatively low, owing to the scarcity of bioavailable iron (Fe) imposes on phytoplankton growth (Martin et al.,

1990; Pollard et al., 2009). In all cases, phytoplankton that flourishes south of the PF is dominated by diatoms, with blooms generally occurring during austral summer (Rigual-Hernandez et al., 2015). Phytoplankton that thrives north of the PF is more heterogeneous and represented by nano- and picoplankton of different origins. One of the most striking patterns associated with phytoplankton in this area, is the significant coccolithophore blooms that occur during spring and summer between ~40 and ~60°S, and that forms the so-called Great Calcite Belt that enriches surface waters in CaCO₃ (Balch et al., 2011; 2016; Rembauville et al., 2016). Within the Indian sector of the SO, near the Kerguelen plateau, this Great Calcite Belt is observed in the PFZ (Smith et al., 2017) and the northern part of the AAZ (between Kerguelen and Heard and McDonald Islands) and may contribute ~ 60 to 85 % of the CCP (Blain et al., 2007; Rembauville et al., 2016; Smith et al., 2017; Rigual-Hernandez et al., 2020). Relatively high primary productivity may also be observed locally, near the PF and the Crozet and Kerguelen plateaus, where oceanic current and/or bottom topography enriches surface waters with iron (Blain et al., 2007; Sokolov and Rintoul, 2007; Pollard et al., 2009; Quéguiner et al., 2013; Tagliabue et al., 2014).

3. Material and methods

3.1 Material core

The 50-meter-long marine sediment core MD04-2718 (48°53,31S; 65°57,42E, 1428 m water depth) was retrieved from the NW flank of the Kerguelen Plateau in 2004, during the MD136/VIGO expedition aboard the R.V. Marion Dufresne (Fig. 1, Michel and Waelbroeck, 2017) within the PFZ. Surface water properties are largely controlled by CO₂- and nutrient- rich waters upwelling in the Antarctic Zone, that are advected northward by the Ekman transport. Sediment recovered at this site is primarily siliciclastic with alternations of nannofossil (calcareous) and diatom (opal) oozes. MIS 11 is clearly identifiable and represented by calcareous-rich ooze, between 11.5 and 13.1 m core depth.




Figure 1 Core location and study area. **a, b** Location of the studied core (MD04-2718, red dot) as well as cores from previous studies (ODP sites 1088 and 1090 (Hodell et al., 2003) and PS75/059-2 (Saavedra-Pellitero et al., 2017a; 2017b) (black dots). Oceanic fronts are from Park et al. (2019). NB: Northern Boundary of the ACC, SAZ: Subantarctic Zone, SAF: Subantarctic Front, PFZ: Polar Front Zone, PF: Polar Front, AAZ: Antarctic Zone, SACCF: Southern Antarctic Circumpolar Front, SB: Southern Boundary of the ACC. **c** Salinity (psu) section at 66°E using WOA18 (Zweng et al., 2018) on ODV software (Schlitzer, 2018). SAMW: Subantarctic Mode Water, AAIW: Antarctic Intermediate Water, CDW: Circumpolar Deep Water, AABW: Antarctic Bottom Water.

3.2 Geochemical analyses

Planktonic foraminifera $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ compositions (expressed in ‰) were measured at the LSCE (Laboratoire des Sciences du Climat et de l'Environnement), based on 67 *Neoglobobulimina* *pachyderma* senestral (200-250 μm fraction, Supplementary Fig. 1) samples, subsampled every 3-4 cm between 11.03 and 13.3 m depth (except for 13.07, 13.10 and 13.13 m, which are barren of *N. pachyderma* s). For each sample, 30 specimens were hand-picked, cleaned using methanol,

ultrasonicated, weighted, and analysed using an IRMS GV ISOPRIME mass spectrometer coupled with an extraction line. The measurements are reported versus the Vienna Pee Dee Belemnite standard (VPDB) with respect to NBS-19 standard, with a mean external reproducibility (1σ) for carbonate standards of $\pm 0.04\text{ ‰}$ and 0.06 ‰ for $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, respectively. The reproducibility of isotopic measurements on *N. pachyderma* s. are 0.06 and 0.1 ‰ , for $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, respectively.

High-resolution, relative sedimentary calcium (Ca) concentrations were measured every 1 cm using an Aavatech profiling X-ray Fluorescence (XRF) core scanner at ETH Zurich, Switzerland with a 10s integration at 10 kV and 500 μA . In parallel, sedimentary CaCO_3 concentrations were determined on 26 samples at GEOPS Laboratory with a $\pm 2\%$ precision, using the vacuum-gasometric technique described in Duchamp-Alphonse et al., (2018) where the amount of CO_2 degassed during the reaction of $100\text{ mg} \pm 2\text{ mg}$ of dried sediments with 6N HCl is converted into CaCO_3 percentages. Comparison between Ca counts and discrete CaCO_3 contents show a highly significant statistical correlation ($r^2=0.97$), and Ca_{XRF} signal is therefore directly converted into CaCO_3 content (Jaccard et al., 2009; 2010; Duchamp-Alphonse et al., 2018; Supplementary Fig. 2).

3.3 Micropaleontological analyses

3.3.1 Coccoliths

Coccolith abundance and mass were determined at the GEOPS (GEOscience Paris Saclay) Laboratory. Smear slides were prepared for 68 samples, based on the protocol detailed in Duchamp-Alphonse et al., (2018), after Beaufort et al., (2014). For each sample, the abundance and mass of individual coccoliths (with an average of 6,172 coccoliths per sample (176 to 13,233)) were automatically determined with the help of the SYRACO software using a Leica DM6000B microscope. SYRACO detects and classifies most of the coccoliths (*Gephyrocapsa caribbeanica*, small *Gephyrocapsa*, *G. oceanica*, *G. muelleriae*, *Calcidiscus leptoporus* and *Coccolitus pelagicus*) present on the slides. Coccolith mass was also directly obtained using SYRACO based on a quasi-linear relationship between their brightness (birefringence in grey scale) and their thickness under cross-polarized light. Because this method applies to coccoliths thinner than $1.55\text{ }\mu\text{m}$, we interpret only thickness and mass data for *Gephyrocapsa* specimens, as *C. leptoporus* and *C. pelagicus* display third-order interferences. The presence of relatively large detrital particles may lead to an underestimation of small specimens including *Gephyrocapsa* sp. and an overestimation of larger ones including *C. pelagicus*. To correct for these potential biases, the abundance of each morphotype was verified manually on 18 samples covering the full spectrum of coccolith abundance observed from MIS 12 to

MIS 10. Visual counts were performed on the same view fields analysed by SYRACO and were compared to the results obtained by the software. *Gephyrocapsa* sp. abundances from SYRACO counts were calibrated using the correlation coefficients between the two datasets, while *C. pelagicus* abundances have been obtained visually for the entire sample set because of the confusion made by SYRACO between detrital particles and this species. *C. leptoporus* abundances did not differ significantly between SYRACO and visual counts were not corrected.

The number of coccoliths per gram of sediment (absolute coccolith abundance – ACA) was determined for each morphotype/species using the equation of Grelaud et al., (2009) as follows:

$$ACA = \frac{N_c * S_f}{N_o * S_o * W_s}$$

where N_c represents the total coccolith number per sample, S_f , the flat beaker area (3117 mm²), N_o , the number of view fields (165), S_o , the view field area (0.01 mm²) and W_s , the dry sediment weight.

The coccolith calcite mass per gram of sediment (per g. sed) was determined using the same equation, where N_c is replaced by M_c , that represents the coccolith calcite mass per view fields. *Gephyrocapsa* calcite mass (CM_{gephy}) was obtained based on the estimation of specimen abundance and mean mass as follows:

$$CM_{Gephy} = ACA * mean\ mass_{Gephy}$$

3.3.2 Planktonic foraminifera

The abundance and planktonic foraminifera species composition were determined at the LSCE with a binocular microscope, using the > 150 µm fraction of 57 samples ranging between 11.30 m and 13.3 m depth. For each sample, an average of 450 planktonic foraminifera were counted. The absolute planktonic foraminifera abundance (AFA) per gram of sediment was determined as follows:

$$AFA = \frac{N_f * 2^{split}}{W_{dry}}$$

where N_f represents the number of planktonic foraminifera counted per sample (> 300), split is the number of splits performed on the sediment before counting ($0 \leq split \leq 6$), and W_{dry} , the dry sediment sample weight.

The planktonic foraminifera calcite mass was determined at the LSCE using mean weights of the major planktonic foraminifera species from the >150 µm fraction. The foraminifera calcite mass is

thus underestimated as it does not consider juveniles planktonic foraminifera or smaller species such as *G. uvula*. Yet the weight variations of this fraction are considered representative of the weight variations of the total planktic foraminifera fraction. The calcite mass produced by planktonic foraminifera, $CaCO_{3plank.foram}$, is (in mg/g):

$$CaCO_{3plank.foram} = \frac{(\sum_i (\sum_j m_{ji} * size_{ji}) * X_i) * AFA}{10^7}$$

Where m_{ji} represents the mean weight in size range j of specie i (in μg), $size_{ji}$ the percentage of population in size range j for specie i , and X_i the percentage of the specie i in the sample. The error on the mass will be:

$$\frac{1}{10^7} * (\sum_i X_i * (\sum_j size_{ji} * err_{m_{ij}} + \sum_j err_{size_{ji}} * m_{ij}) + \sum_i err_{X_i} * (\sum_j size_{ji} * m_{ji}))$$

where err is the error on each of the variables.

The size distribution, considering eight intervals (150-200 μm , 200-250 μm , 250-315 μm , 315-355 μm , 355-400 μm , 400-450 μm , 450-500 μm and >500 μm), and the mean weight for each size has been determined at six depths for the eight species representing > 99 % of the assemblage. The mean weight has been determined from 15 weighings of 1 to 4 individuals (to reach weight between 10 to 40 μg), to decrease the weight error to 19 % $[(1/(2(n-1)))^{0.5}]$. The eight species include *Neogloboquadrina pachyderma* sinistral and dextral, *Globigerinita glutinata*, *Turborotalia quinqueloba*, *Globigerina bulloides*, *Globorotalia inflata*, *Globorotalia crassaformis* and *Globorotalia truncatulinoides*. We determine mean weights of : $3.55 \pm 0.32 \mu g$ for *N. pachyderma*, $9.42 \pm 2.15 \mu g$ for *G. bulloides*, 3.48 ± 0.42 for *G. glutinata*, $13.9 \pm 2.37 \mu g$ for *G. crassaformis* and $22.4 \pm 5.39 \mu g$ for *G. inflata*. We assumed a similar weight to *G. glutinata* for *T. quinqueloba*. For *G. truncatulinoides*, we assumed a mean weight similar to *G. crassaformis*. For *N. pachyderma*, 90 to 97 % of the population ranges in the minimum measured size range (150-200 μm), and between 71 to 100 % for *G. glutinata* and *T. quinqueloba*. For the other species the size range distribution varies with depth with the main size range varying from 150-200 μm to 250-315 μm for *G. bulloides*, *G. crassaformis* and *G. truncatulinoides* and from 150-200 μm to 315-355 μm for *G. inflata*. The error on the calcite mass of the planktonic foraminifera depends thus mainly on the error on the population fraction in the different size range for the larger species. The error on the calcite mass of the planktonic foraminifera also considers the variations in planktonic foraminifera weight and size with depth observed in the core, that can be linked with the changes in atmospheric CO_2 concentration (Barker and Elderfield, 2002; Moy et al., 2009; Hennehan et al., 2017). For the 57 samples, the calcite mass of the planktonic foraminifera and its error have been calculated from the counted abundance of the

different species and their mean size and mean weight calculated from the six depths. The error on this calcite mass varies from 26 to 57 % and is linearly linked with the abundance of *G. bulloides* ($r^2=0.91$). Although large, the error remains negligible compared to the observed variations that are almost a factor of a thousand.

Summer SST estimates were determined on the same 57 samples as those used for planktonic foraminifera assemblages based on two independent approaches: the Modern Analog Technique and the percentage of *N. pachyderma* s. The former one relies on the use of a database of modern foraminifera assemblage (Haddam et al., 2016) and the PaleoAnalog software (Therón et al., 2004) to find the closest modern analogues for the fossil assemblage (Prell, 1985; Haddam et al., 2016). The latter one is based on the linear relation between SST and the percentage of *N. pachyderma* s. in the total planktonic foraminifera assemblage following the equation of Govin et al. (2009):

$$T^{\circ}\text{C} (\% \text{ pachy s.}) = -0.087 (\% \text{ pachy s.}) + 11.339$$

No SST could be determined between 13.13 and 13.07 m depth as the number of planktonic foraminifera was insufficient (<300) for a robust temperature estimate. During glacial periods the number of foraminifera per gram of sediment decreases drastically and, in these three levels, the >150 μm fraction contains almost exclusively ice rafted detritus and radiolarians.

4. Results

4.1 Geochemical signals

Planktonic foraminifera $\delta^{18}\text{O}$ values range between 2.2 ‰ and 3.8 ‰ (Fig. 2a). After a rapid decrease from 3.7 to 2.2 ‰ between 13.00 and 12.33 m, values gradually increase to reach around 3.6 ‰ at 11.33 m. The $\delta^{13}\text{C}$ signal ranges from 0.75 ‰ to 1.67 ‰ (Fig. 2b). $\delta^{13}\text{C}$ values are characterised by a slightly increasing plateau between 12.80 and 12.23 m and two sharp decreases to the minimum value of 0.8 ‰ from 13.23 to 13.03 m and from 11.57 to 11.33 m.

Sedimentary CaCO_3 concentrations range between 5 % and 99 % (Fig. 2c). Values never exceed 20 % between 13.30 and 13.10 m and between 11.30 and 11.03 m, while they average 85 % between 12.67 and 11.90 m. The most striking trends are observed between 13.10 and 12.66 m and 12.03 and 11.41 m, where CaCO_3 concentrations undergo a drastic increase to maximum values of 99 % at 12.65 m and a two-fold decreasing trend to reach minimum values of 7 % at 11.38.




Figure 2 Bulk carbonate signals and stable isotope records of planktonic foraminifera *N. pachyderma s.* from sediment core MD04-2718. **a** $\delta^{18}\text{O}_{\text{N. pachyderma s}}$ (‰), **b** $\delta^{13}\text{C}_{\text{N. pachyderma s}}$ (‰), **c** CaCO_3 (%) derived from Ca_{XRF} signal, the envelope representing the error on the calculation of CaCO_3 .

4.2 Micropaleontological signals

Coccolith assemblages are predominantly composed of *Gephyrocapsa* sp. i.e., specimens from the Noelaerhabdaceae family whose proportions exceed 98.5 %, reaching maximum values (99.9 %) between 12.87 and 11.33 m depths (Supplementary Fig. 3). *C. pelagicus* and *C. leptoporus* are present as well, but in much lower proportions. They are mainly represented by the subspecies *C. pelagicus* ssp. *braarudii* (10-14 μm) and *C. leptoporus* (5-8 μm) and range between 0 and 0.9 %, and 0 and 0.6 % of the assemblages, respectively. The number of total coccoliths per gram of sediments

(per g. sed.) ranges from $0.017 \cdot 10^9$ (at 13.10 m) to $40 \cdot 10^9$ (at 12.10 m) and is mainly represented by the number of *Gephyrocapsa* sp. per g. sed. that fluctuates between $0.016 \cdot 10^9$ and $40 \cdot 10^9$ (Fig. 3b and 3c).




Figure 3 Coccolith and $\delta^{18}O_{N. pachys}$ signals between 1103 and 1330 cm of the sediment core MD04-2718. **a.** $\delta^{18}O_{N. pachys}$ (‰), abundances of : **b** coccoliths (number of coccoliths $\cdot 10^9$ per g sed.), **c** *Gephyrocapsa* (number $\cdot 10^9$ per g sed.), **d** *C. leptoporus* (number $\cdot 10^6$ per g sed.), **e** *C. pelagicus* (number $\cdot 10^6$ per g sed.), **f** *Gephyrocapsa* mean mass (pg). Envelopes in **b**, **c**, **d** and **f** show the error on SYRACO abundance for each specie and for *Gephyrocapsa* mean mass.

335 After an increasing trend from 12.90 to 12.67 m, values show a plateau around $22.5 \cdot 10^9$
 336 specimens per g. sed. between 12.67 and 11.80 m and decreases to reach values as low as $0.35 \cdot 10^9$
 337 specimens per g. sed. at 11.17 m. *C. leptoporus* abundance fluctuates between $0.06 \cdot 10^6$ and
 338 $21.6 \cdot 10^6$ specimens per g. sed. and its distribution generally follows that of *Gephyrocapsa* sp. (and
 339 thus of total coccoliths), with an increasing trend between 13.07 and 12.67 cm followed by a long-
 340 term decreasing trend up to 11.47 m, interrupted by a transient a peak at 12.10 m (Fig. 3d). *C.*
 341 *pelagicus* abundance varies between 0 and $1.6 \cdot 10^6$ coccoliths per g. sed (Fig. 3e). It is totally absent
 342 from the sediment between 12.73 and 11.54 m and is thus only present between 13.30 and 12.77 m
 343 and between 11.53 and 11.03 m, where it reaches values as high as 1.6 and $0.77 \cdot 10^6$ specimens per
 344 g. sed, respectively. *Gephyrocapsa* sp. mean mass varies between 1.03 pg and 2.24 pg (Fig. 3f). After
 345 reaching maximum values (from 2.07 to 2.24 pg) between 13.23 and 13.13 m, data are consistent
 346 with a long-term decrease to reach minimum values around 11.40 m depth.

347 The planktonic foraminifera assemblages are mainly composed of *N. pachyderma* sinistral and
 348 *G. bulloides* with proportions ranging between 0 % and 99.8 % and between 0 and 75.1 %,
 349 respectively. The other species, represented by *G. glutinata*, *G. inflata*, *G. crassaformis*, *N.*
 350 *pachyderma* dextral and *T. quinqueloba* never exceed 6.8 %. (Supplementary Fig. 4). The total
 351 number of planktonic foraminifera fluctuates between 0 and $18.5 \cdot 10^3$ specimens per g. sed (Fig. 4b).
 352 No specimen is found between 13.13 and 13.07 m, and the total abundance of planktonic
 353 foraminifera never exceeds $5.7 \cdot 10^3$ between 13.30 and 12.9 m and between 11.5 and 11.3 m.
 354 Between 12.9 and 11.5 depth the number of *N. pachyderma* s. varies between 0.3 and $12.4 \cdot 10^3$
 355 specimens per g. sed (Fig. 4c), and significantly influences the total number of planktonic
 356 foraminifera which generally shows the same trends. Its abundance shows an overall plateau around
 357 $3 \cdot 10^3$ specimens per g. sed, and 2 maxima ($12.4 \cdot 10^3$ and $10.9 \cdot 10^3$ specimens per g. sed) at 12.70
 358 and 11.73 m, respectively. The number of *G. bulloides* fluctuates between 2.10 and $4.7 \cdot 10^3$
 359 specimens per g. sed (Fig. 4d). After reaching maximum value at 12.70 m, it depicts a long-term
 360 decreasing trend to the minimum abundance of 2.1 specimens per g. sed. at 11.3 m. The number of
 361 *N. pachyderma* d. varies between 0.14 and 548.6 specimens per g. sed. and follows the general same
 362 trends as those of total planktonic foraminifera and *N. pachyderma* s. (Fig. 4e). *G. glutinata*
 363 abundance ranges between 0.2 and 551 specimens per g. sed (Fig. 4f). It depicts rather scattered
 364 values with two peaks (512 and 551 specimens per g. sed.) at 12.70 and 12.33 m. *G. crassaformis*
 365 abundance fluctuates between 0 and 259.5 specimens per g. sed, with values averaging 150
 366 specimens per g. sed between 12.70 and 12.13 m (Fig. 4g). *T. quinqueloba* abundance fluctuates
 367 between 0 and 219.4 specimens per g. sed, and no specific trend may be highlighted (Fig. 4h). Finally,

G. inflata abundance ranges between 0 and 106 specimens per g. sed (Fig. 4i). Similarly, to *G. crassaformis*, no or very few specimens are found between 12.23 and 11.60 m.

The reconstructed summer SSTs fluctuate between 1.7 and 9.9 °C (Fig. 4j). The maximum temperature is reached after a drastic warming amounting ~8 °C between 13.27 and 13.00 m. It is followed by temperatures fluctuating between 4.6 and 8.3°C from 12.93 to 12.70 m, and a plateau around 8°C between 12.67 and 12.27 m. Then, summer SSTs gradually decrease and reach down to 3 °C between 12.03 and 11.30 m.




Figure 4 Planktonic foraminifera and $\delta^{18}\text{O}_{N.pachys}$ signals between 1103 and 1330 cm of the sediment core MD04-2718. **a** Signal of $\delta^{18}\text{O}_{N.pachys}$ (‰), **b** Number of total planktonic foraminifera (per g sed.), Number (per g. of sed.) of: **c** *N. pachyderma s.*, **d** *G. bulloides*, **e** *N. pachyderma d.*, **f** *G. glutinata*, **g** *G. crassaformis*, **h** *T. quinqueloba*, **i** *G. inflata*, **j** Reconstructed summer SST (°C) using *N. pachyderma s.* percentages (blue curve) and based on Modern Analog Technique method (green curve).

5. Age model

The age model of core MD04-2718 is based on the planktonic foraminifera $\delta^{18}\text{O}$ record and sedimentary CaCO_3 signal obtained on the uppermost 30 m of the core (Supplementary Fig. 5). These records helped identifying glacial-interglacial cycles and determining that the uppermost 20 m of the

core covered the last 800 ka. It also allowed us to define the depth interval 13.00 to 11.00 m as representing the MIS 12 – MIS 10 time-interval and confirm our assumption that the nannofossil ooze between 11.5 and 13.1m was deposited during MIS 11 (see section 3.1). Numerous studies showed that the correlation between reconstructed SSTs in the Southern Ocean and air temperature reconstructed from Antarctic ice cores was helpful to better constrain sediment core age model (e.g., Pichon et al., 1992; Sowers et al. 1993; Waelbroeck et al., 1995; Thöle et al., 2019; Ai et al., 2020). Here, we used the record of δD of the EPICA Dome C ice core as a proxy for Antarctic surface temperature (Jouzel et al., 2007). As such, for the interval MIS 12 – 10, the age model of core MD04-2718 was refined based on the graphical alignment of our reconstructed summer SST with record of the δD record on the AICC2012 timescale (Bazin et al., 2013), assuming an in-phase relationship (Fig. 5a). Using Analyseries software (Paillard et al., 1996), we set 5 tie points which helped us identifying the 11.03- 13.3 m core section as representing the time interval between 438.4 and 360.2 ka i.e., between the end of MIS 12 and the beginning of MIS 10. Over this period, the sedimentation rate averages 3.1 cm.ka⁻¹ with lowest values (1 cm.ka⁻¹) being recorded during MIS 12, and highest ones (4.9 cm.ka⁻¹) being observed at the beginning of MIS 11 (Fig. 5b). Given these sedimentation rates, the temporal resolution of our record ranges from ~ 0.6 to 3.8 ka for micropaleontological and stable isotope data and of 0.2 to 2 ka for the CaCO₃ signal.




Figure 5 a Age model of MD04-2718 based on the correlation between MD04-2718 summer SST reconstructions and EDC δD signal on AICC2012 timescale. Blue curve is the reconstructed summer SST using *N. pachyderma* s. percentages. Green curve is the reconstructed summer SST based on Modern Analog Technique method. Black curve is the δD measured in EDC ice core (Jouzel et al., 2007) on the AICC2012 timescale (Bazin et al., 2013). Black dots are the tie points between SST and δD **b** Estimated sedimentation rate of MD04-2718 core. Black curve represents the depth versus the calibrated age (ka BP on AICC2012) and the grey shaded area represent the variation of sedimentation rate with time.

6. Discussion

6.1 CaCO_3 as a record of CCP strength

In pelagic environments of the SO, sedimentary carbonate concentration and/or coccolith abundances have been assumed to reflect biogenic carbonate export production - and by inference the strength of the CCP - based on the paradigm that in modern environments, and more particularly in the Great Calcite Belt, coccolithophores represent an important – if not dominant – part of exported mineralized organisms (Salter et al., 2014; Rembauville et al., 2016). However, the bulk sedimentary carbonate signal may include detrital or diagenetic phases and within the biogenic fraction, contributions of planktonic foraminifera and pteropods, the latter often considered negligible (Salter et al., 2014; Rembauville et al., 2016). Therefore, special attention must be devoted to assessing the significance of bulk sedimentary carbonate signals before making interpretations in terms of planktonic carbonate productivity and, by inference, of the CCP strength. It requires to carefully select the sedimentary cores that robustly reflect biogenic carbonate accumulation and to consider the potentially obfuscating impact of diagenesis.

Core MD07-2418 was retrieved at 1,428 m water depth i.e., well above the modern lysocline located at ~4,300 m (Kolla et al., 1976; Howard and Prell, 1994). The area only receives carbonate-free terrigenous material sourced from volcanic rocks of the Crozet and Kerguelen plateau (Nougier, 1970). Thus, the sedimented carbonate material directly relates to local biogenic carbonate production, for which the diagenetic overprint and potential detrital contribution are minimal. These assumptions are corroborated by Scanning Electron Microscopy (SEM) observations highlighting that the sedimentary carbonate fraction is predominantly constituted by very well-preserved foraminifera and coccoliths, sometimes represented as intact coccospheres (Fig. 6). In addition, we measured the relative proportion of fragmented planktonic foraminifera shells versus intact foraminifers, a well-characterised preservation index (Vazquez-Riveiros et al., 2010; Flores et al., 2003) (Supplementary Fig. 6). This foraminifera fragmentation index varies between 0 and 15.2 % with values higher than 3 % observed only between 426 and 417 ka. These results clearly indicate that planktonic foraminiferal preservation is generally good (Le and Shackleton, 1992).

Dissolution processes could result in differential sedimentary preservation in the coccolith assemblages as well, preferentially preserving resistant specimens. However, the assemblage is mainly constituted by well-preserved *Gephyrocapsa* sp., as seen on SEM observations. Furthermore, within this genus, small *Gephyrocapsa* sp. (<3 μm) follow the exact same trends as the larger ones,

which suggests that differential dissolution does not significantly affect coccolith assemblages in sediments from core MD04-2718 (Supplementary Fig. 6).




Figure 6 Visual estimation of coccolith preservation of sediment core MD04-2718 based on Tescan VEGA II SEM observation (Museum National d'Histoire Naturelle, MNHN, Paris) showing good preservation **a** at 389.1 ka and **b** at 405.8 ka.

At site MD04-2718, *Gephyrocapsa* calcite mass varies between 0.03 mg/g and 54.62 mg/g. The lowest values are recorded during MIS 12 (0.03 mg/g) and MIS 10 (0.43 mg/g) and the highest ones are documented during MIS 11. In detail, *Gephyrocapsa* calcite mass increases from 0.03 to 36.68 mg/g at the beginning of MIS 11, averages 30 mg/g during MIS 11, and decreases gradually from 382 ka to 372 ka to reach low values of 0.43 mg/g during glacial MIS 10. The *Gephyrocapsa* calcite mass record closely resembles the one of *Gephyrocapsa* sp. abundance, whereas no relationship may be observed with *Gephyrocapsa* mean mass (Fig. 7 and Supplementary Fig. 7). Moreover, while the variability in *Gephyrocapsa* sp. mean mass along the records are of a factor of two, the observed variations in abundance are more than a factor of two thousand. We therefore posit that changes in *Gephyrocapsa* calcite mass are mainly the consequence of changes in *Gephyrocapsa* abundance rather than in shell mean mass linked with shift in calcification or morphotypes.—The planktonic foraminifera (>150 µm) calcite mass varies between 0 and 93.9 mg/g. Briefly, it is characterised by null or low values during MIS 12 (< 7.2 mg/g) and MIS 10 (< 1.08 mg/g) and generally higher values during MIS 11 and the beginning of MIS 10 (> 20 mg/g). Maximum values are observed around 412.7 ka i.e., at the beginning of MIS 11. According to the significant linear correlation that exists between the *Gephyrocapsa* sp. + planktonic foraminifera calcite mass and sedimentary CaCO₃ content ($r^2 = 0.93$), it appears that both plankton groups constitute the major component of the biogenic carbonate signal. *Gephyrocapsa* sp. seems to have a relatively larger contribution on the sedimentary

CaCO₃ content as shown by its greater correlation to CaCO₃ ($r^2 = 0.89$) than planktonic foraminifera calcite mass itself ($r^2 = 0.83$) (Supplementary Fig. 2). Such results are consistent with the major control of coccoliths and more particularly morphotypes of the Noelaerhabdaceae family exert on PIC export today (Rembauville et al., 2016; Patil et al., 2020).

In the following, changes in the downcore CaCO₃ signal at site MD04-2718 will be interpreted in terms of changes in planktonic carbonate export, and by inference changes in CCP strength.




Figure 7 Origin of the carbonate fraction. **1** Correlation between CaCO₃ and Gephyrocapsa + planktonic foraminifera calcite mass, **2 a** planktonic foraminifera calcite mass (mg/g), **b** Gephyrocapsa mean mass (pg), **c** Gephyrocapsa calcite mass (mg/g), **d** CaCO₃ (%). Smoothed curves use a three-point average using IGOR software. The shaded areas represent the error on the curve. The dotted blue lines represent the beginning and end of MIS 11 from right to left respectively, and the dotted red line represent the middle of the rising slope of CO₂ from glacial to interglacial.

6.2 Increased CCP during MIS 11 as a response to the southward migration of westerly winds and ACC front positions

As represented by the variation in sedimentary CaCO₃ content, it appears that the CCP was strengthened during MIS 11 compared to MIS 12 and MIS 10 (Fig. 8). By analogy to the modern SO (Baumann et al., 1999; Zondervan et al., 2007) and the Pacific sector of the SAZ during MIS 12-MIS 10

(Schaefer et al., 2005; Ullermann, 2015; Saavedra-Pellitero et al., 2017a; 2017b), it is probable that such recorded pattern is related to substantial changes in SST and nutrient conditions. In addition to reconstructed SSTs, the abundances of *C. pelagicus* and *C. leptoporus* can be used to indirectly assess past changes in sea surface nutrient and temperature conditions. Indeed, while *C. pelagicus* seems to proliferate under colder conditions than *C. leptoporus* (Cobianchi et al., 2012), both these species appear to have affinities with relatively nutrient-rich waters (Andruleit, 1997; Balestra et al., 2004; Amore et al., 2012). This behaviour has been well documented on glacial/interglacial timescales in the SO (Flores et al., 1999; Findlay and Flores, 2000; Marino et al., 2009; Saavedra-Pellitero et al., 2017a; 2017b) to characterize frontal zone dynamics (Flores et al., 1999; Findlay and Giraudeau, 2000; Flores et al., 2003; Baumann et al., 2004; Malinverno et al., 2015; Marino et al., 2008; Saavedra-Pellitero et al., 2017a; 2017b). At core site MD04-2718, reconstructed SSTs range between ~2-4°C during glacial MIS 12 and MIS 10, and ~8°C during MIS 11 (Fig. 8). In parallel, relatively fertile conditions are inferred during TV and MIS 11 by the coccolith species *C. pelagicus* and *C. leptoporus* that peak between 426 and 415 ka and between 412 and 392 ka, respectively. The CCP was thus stronger during the relatively warm and nutrient-rich MIS 11 interglacial compared to the relatively cold and nutrient-depleted MIS 12 and MIS 10. The CCP trend shows however some temporal lag with the SST and nutrient records, highlighting other mechanisms in potentially regulating the variability of CCP strength. This is particularly obvious during TV (430-424 ka), when the CCP remains relatively weak despite drastic increases in SSTs and nutrient conditions. At that time, decreasing $\delta^{13}\text{C}_{\text{N.pachyderma}}$ values and increasing mean mass of *Gephyrocapsa* are recorded (Fig. 8). The rapid $\delta^{13}\text{C}$ decrease during the deglaciation is also documented in the South Pacific (Saavedra-Pellitero et al., 2017b) and South Atlantic (Hodell et al., 2003) (Fig. 8). This isotopic signal is consistent with the resumption of SO upwelling, which supplies abundant nutrients, previously sequestered in the ocean interior owing to greater water-column stratification, to the productive ocean surface (François et al., 1997; Spero et al., 2002; Adkins, 2013; Siani et al., 2013). Coeval increase in *Gephyrocapsa* mean mass has also been associated with increase in CO₂-rich upwelled waters in the SO (Duchamp-Alphonse et al., 2018). The reason behind these staggered and seemingly disconnected patterns could relate to the ecological competition between diatoms and coccolithophores in the SO. During glacial periods, favoured by higher iron (Fe) availability, diatom productivity outcompetes coccolithophore growth (Martin, 1990; Kohfeld et al., 2005; Martinez-Garcia et al., 2009; Thöle et al., 2019), and call for a weaker CCP. While the availability of Fe dwindled over deglacial periods, reinvigorated upwelling of nutrient-rich waters would have promoted diatom growth during Termination V. In such a scenario, we hypothesize that it is only during peak MIS 11 that silicic acid concentrations became too low, thereby limiting diatom growth (Sarmiento et al., 2004), and thus had the potential to favour coccolith production, enhancing the CCP. Overall, since the upwelled

waters that (re)surface in the AAZ act under the forcing of the prevailing westerlies (Anderson et al., 2009; Lauderdale et al., 2013; Menviel et al., 2018), it is probable that winds migrated poleward between MIS 12 and MIS 11, concomitantly forcing the poleward migration of the entire frontal system (Flores and Sierro, 2007; Kemp et al., 2010; Saavedra-Pellitero et al., 2017a; 2017b). According to the reconstructed SST ($< 8^{\circ}\text{C}$ in mean) as well as the relatively low percentages of *C. leptoporus*, while close to the SAF, it is likely that site MD04-2718 remained in the PFZ. That is only around 390 ka, when SSTs decreased, that westerlies (along with the frontal system) might have shifted back to the North, leading to a decrease in the CCP.




Figure 8 Records of CCP and upwelling strengths in the SO and relation with variation of atmospheric CO₂ concentration. **a** CO₂ (ppm) (Nehrbass-Ahles et al., 2020), **b** Biosphere productivity (F_{bio,t}/F_{bio,PST}) (Brandon et al., 2020), **c** δ¹⁸O of *N. pachyderma* s. from MD04-2718, **d** summer SST reconstructed using *N. pachyderma* s percentages (blue curve) and using Modern Analog Technique method (green curve), **e** δ¹³C of *N. pachyderma* s. from MD04-2718, **f** Δδ¹³C of benthic *Cibicides wuellerstorfi* from ODP Site 1088- ODP Site 1090 (Hodell et al., 2003), **g** *Gephyrocapsa* mean mass (pg), **h** *C. pelagicus* abundance (*10⁶/g sed), **i** *C. leptoporus* abundance (*10⁶/g sed), **j** CaCO₃ (%) derived from Ca_{XRF}, **k** PS75/059-2 *Coccoliths* abundance (10⁹coccoliths/g of sediment, Saavedra-Pellitero et al., 2017b), showed on AICC2012 chronology.

6.3 Increased CCP during TV and MIS 11 and its impact on atmospheric CO₂

The onset of deep-water ventilation following MIS 12 is recorded at ~432 ka, while the CCP starts to increase a few thousand years later, around 426 ka (Fig. 8). This distinct timing suggests that the rapid increase in atmospheric pCO₂ during Termination V was triggered, in part, by the reinvigoration of the SO upwelling early in the deglacial progression. Concomitant with the reinvigoration of SO upwelling, the increase in global biosphere productivity (Brandon et al., 2020) that reflects in part, an increase in vegetation cover and terrestrial carbon storage (Tzedakis et al., 2006; de Vernal and Hillaire-Marcel, 2008; Prokopenko et al., 2009; Melles et al., 2012; Kousis et al., 2018) has probably acted as a sink of atmospheric CO₂, thereby transiently buffering the impact of oceanic CO₂ degassing (Fig. 8). pCO₂ reached maximum concentration (up to ~ 282 ppm) during the second part of MIS 11, as the CCP functioned at maximum strength. Enhanced CCP in the SO contributed to reduce the efficiency of the BCP. Overall, enhanced *Gephyrocapsa* sp. export and burial both in the SO (Saavedra-Pellitero et al., 2017a; 2017b) and at low latitudes (Barker et al., 2006; Rickaby et al., 2007; Alvarez et al., 2010; Amore et al., 2012; Saavedra-Pellitero et al., 2019), would have led to a global increase in carbonate dissolution at the seafloor (Barker et al., 2006; Qin et al., 2018) and to a significant change in the global rain ratio, eventually impacting atmospheric pCO₂ (Sigman et al., 1998; Sigman and Boyle, 2000; Antia et al., 2001; Barker et al., 2006). This hypothesis warrants further investigations to fully quantify changes in the rain ratio and their impact on atmospheric pCO₂. From 398 ka to 381 ka, the gradual, ~35 ppm decrease in atmospheric pCO₂ coincided well with the rapid increase in benthic foraminifera Δδ¹³C in the South Atlantic (Hodell et al., 2003), consistent with enhanced stratification of the SO and increased carbon storage in the ocean interior. The second part of atmospheric CO₂ drawdown during MIS 10 would be supported in part by the reduction in the strength of the CCP combined with a globally more efficient marine BCP.

At a global scale, the most striking pattern of MIS 11 relates to the low eccentricity context that is documented around 400 ka (Laskar et al., 2004). Low eccentricity, through the lengthening of

the growing season, might have triggered an acme of primary producers such as *Gephyrocapsa* sp., at both high and low latitudes (Rickaby et al., 2007). One cannot exclude either that our record reflects the long-term cyclic Noelaerhabdaceae evolutionary pattern recently highlighted for the tropical area over the Pleistocene (Beaufort et al., 2022), with *Gephyrocapsa* blooms during MIS 11 being the consequence of reduced seasonality and phytoplankton niches. In all cases, we propose that, under this particularly low eccentricity context, enhanced terrestrial productivity and carbon storage on land, together with enhanced biological carbonate productivity and CCP strength in the SO may have contributed to shape the 30 ka-long CO₂ plateau characteristic of MIS 11.

7. Conclusion

Our study documents temporal variations in the strength of the CCP based on Southern Ocean sediments and their impact on atmospheric CO₂ concentrations over the period ranging between MIS 12 and MIS 10. First, using micropaleontological and geochemical data, we demonstrate that the sedimentary CaCO₃ signal robustly reflects the production and export of *Gephyrocapsa* sp. and (to a lesser degree) planktonic foraminifera and can therefore be used as a high-resolution proxy of the CCP. Second, using planktic foraminifera, *C. leptoporus* and *C. pelagicus* abundances combined with $\delta^{13}\text{C}$ of *N. pachyderma* s. from the same site, we reconstruct changes in sea surface temperature and nutrient conditions in conjunction with the regional upwelling dynamics, which are closely linked to the latitudinal migration of the westerly winds. Third, the combination of the results obtained in this study together with previous micropaleontological and geochemical data from literature allow us to highlight the sequence of events leading to the deglaciation and to understand the specific atmospheric pCO₂ pattern that characterises this key interval of the past. We show that the abrupt atmospheric CO₂ rise observed during TV is led by the reinvigoration of SO upwelling probably due to a poleward migration of the westerlies and ACC fronts, while the sustained CO₂ plateau observed during MIS 11 is the result of enhanced global primary productivity and increased CCP due to the exceptionally low eccentricity context.

Acknowledgments

This work benefited from funding from the INSU-LEFE-IMAGO program BIOCOD, from the European Research Council under the European Union H2020 Programme (H2020/20192024)/ ERC grant agreement no. 817493 ERC ICORDA as well as from the Institut Pierre Simon Laplace (IPSL). M.B. doctoral scholarship is supported by a public grant overseen by the French National research Agency

(ANR) as part of the « Investissement d'Avenir » program, through the "IDI 2017" project funded by the IDEX Paris-Saclay (grant ANR-11-IDEX-0003-02). S.L.J. acknowledges support from the Swiss National Science Foundation (grants PP00P2_144811 and PP00P2_163003). We thank Sylvain Pont and the Plateau technique de Microscopie Electronique (PtME, MNHN, Paris, France) for their assistance in providing SEM images, which helped for the estimation of the preservation of the carbonate fraction. We also thank Yvonne Hamann for instructions for the XRF core scanner at ETH, Zurich.

Data availability

Data will be made available in the Pangaea database prior to publication.

References

- Abelmann, A., Gersonde, R., Knorr, G., Zhang, X., Chaplignin, B., Maier, E., Esper, O., Friedrichsen, H., Lohmann, G., Meyer, H., Tiedemann, R., 2015. The seasonal sea-ice zone in the glacial Southern Ocean as a carbon sink. *Nat Commun* 6, 8136. <https://doi.org/10.1038/ncomms9136>
- Adkins, J.F., 2013. The role of deep ocean circulation in setting glacial climates. *Paleoceanography* 28, 539–561. <https://doi.org/10.1002/palo.20046>
- Ai, X.E., Studer, A.S., Sigman, D.M., Martínez-García, A., Fripiat, F., Thöle, L.M., Michel, E., Gottschalk, J., Arnold, L., Moretti, S., Schmitt, M., Oleynik, S., Jaccard, S.L., Haug, G.H., 2020. Southern Ocean upwelling, Earth's obliquity, and glacial-interglacial atmospheric CO₂ change. *Science* 370, 1348–1352. <https://doi.org/10.1126/science.abd2115>
- Álvarez, M.C., Flores, J.A., Sierro, F.J., Molina-Cruz, A., 2010. Long-term upwelling evolution in tropical and equatorial Pacific during the last 800 kyr as revealed by coccolithophore assemblages. *Geobios* 8. <https://doi.org/10.1016/j.geobios.2009.07.001>
- Amore, F.O., Flores, J.A., Voelker, A.H.L., Lebreiro, S.M., Palumbo, E., Sierro, F.J., 2012. A Middle Pleistocene Northeast Atlantic coccolithophore record: Paleoclimatology and paleoproductivity aspects. *Marine Micropaleontology* 90–91, 44–59. <https://doi.org/10.1016/j.marmicro.2012.03.006>
- Anderson, R.F., Ali, S., Bradtmiller, L.I., Nielsen, S.H.H., Fleisher, M.Q., Anderson, B.E., Burckle, L.H., 2009. Wind-Driven Upwelling in the Southern Ocean and the Deglacial Rise in Atmospheric CO₂. *Science* 323, 1443–1448. <https://doi.org/10.1126/science.1167441>
- Andrulleit, H., 1997. Coccolithophore fluxes in the Norwegian-Greenland Sea: Seasonality and assemblage alterations. *Marine Micropaleontology* 31, 45–64. [https://doi.org/10.1016/S0377-8398\(96\)00055-2](https://doi.org/10.1016/S0377-8398(96)00055-2)
- Antia, A.N., Koeve, W., Fischer, G., Blanz, T., Schulz-Bull, D., Schölten, J., Neuer, S., Kremling, K., Kuss, J., Peinert, R., Hebbeln, D., Bathmann, U., Conte, M., Fehner, U., Zeitzschel, B., 2001. Basin-wide

641 particulate carbon flux in the Atlantic Ocean: Regional export patterns and potential for
 642 atmospheric CO₂ sequestration. *Global Biogeochem. Cycles* 15, 845–862.
 643 <https://doi.org/10.1029/2000GB001376>

644 Balch, W.M., Bates, N.R., Lam, P.J., Twining, B.S., Rosengard, S.Z., Bowler, B.C., Drapeau, D.T., Garley,
 645 R., Lubelczyk, L.C., Mitchell, C., Rauschenberg, S., 2016. Factors regulating the Great Calcite Belt
 646 in the Southern Ocean and its biogeochemical significance. *Global Biogeochem. Cycles* 30,
 647 1124–1144. <https://doi.org/10.1002/2016GB005414>

648 Balch, W.M., Drapeau, D.T., Bowler, B.C., Lyczkowski, E., Booth, E.S., Alley, D., 2011. The
 649 contribution of coccolithophores to the optical and inorganic carbon budgets during the
 650 Southern Ocean Gas Exchange Experiment: New evidence in support of the “Great Calcite Belt”
 651 hypothesis. *J. Geophys. Res.* 116, C00F06. <https://doi.org/10.1029/2011JC006941>

652 Balestra, B., 2004. Coccolithophorids from the Southeast Greenland Margin (Northern North
 653 Atlantic): production, ecology and the surface sediment record. *Micropaleontology* 50, 23–34.
 654 https://doi.org/10.2113/50.Suppl_1.23

655 Barker, S., Archer, D., Booth, L., Elderfield, H., Henderiks, J., Rickaby, R.E.M., 2006. Globally increased
 656 pelagic carbonate production during the Mid-Brunhes dissolution interval and the CO₂ paradox
 657 of MIS 11. *Quaternary Science Reviews* 25, 3278–3293.
 658 <https://doi.org/10.1016/j.quascirev.2006.07.018>

659 Barker, S., Elderfield, H., 2002. Foraminiferal Calcification Response to Glacial-Interglacial Changes in
 660 Atmospheric CO₂. *Science* 297, 833–836. <https://doi.org/10.1126/science.1072815>

661 Baumann, K.-H., Böckel, B., Frenz, M., 2004. Coccolith contribution to South Atlantic carbonate
 662 sedimentation, in: Thierstein, H.R., Young, J.R. (Eds.), *Coccolithophores*. Springer Berlin
 663 Heidelberg, Berlin, Heidelberg, pp. 367–402. https://doi.org/10.1007/978-3-662-06278-4_14

664 Baumann, K.-H., Čepek, M., Kinkel, H., 1999. Coccolithophores as Indicators of Ocean Water Masses,
 665 Surface-Water Temperature, and Paleoproductivity — Examples from the South Atlantic, in:
 666 Fischer, G., Wefer, G. (Eds.), *Use of Proxies in Paleoceanography*. Springer Berlin Heidelberg,
 667 Berlin, Heidelberg, pp. 117–144. https://doi.org/10.1007/978-3-642-58646-0_4

668 Bazin, L., Landais, A., Lemieux-Dudon, B., Toyé Mahamadou Kele, H., Veres, D., Parrenin, F.,
 669 Martinerie, P., Ritz, C., Capron, E., Lipenkov, V., Loutre, M.-F., Raynaud, D., Vinther, B.,
 670 Svensson, A., Rasmussen, S.O., Severi, M., Blunier, T., Leuenberger, M., Fischer, H., Masson-
 671 Delmotte, V., Chappellaz, J., Wolff, E., 2013. An optimized multi-proxy, multi-site Antarctic ice
 672 and gas orbital chronology (AICC2012): 120–800 ka. *Clim. Past* 9, 1715–1731.
 673 <https://doi.org/10.5194/cp-9-1715-2013>

674 Beaufort, L., 2005. Weight estimates of coccoliths using the optical properties (birefringence) of
 675 calcite. *Micropaleontology* 51, 289–297. <https://doi.org/10.2113/gsmicropal.51.4.289>

676 Beaufort, L., Barbarin, N., Gally, Y., 2014. Optical measurements to determine the thickness of calcite
 677 crystals and the mass of thin carbonate particles such as coccoliths. *Nat Protoc* 9, 633–642.
 678 <https://doi.org/10.1038/nprot.2014.028>

679 Beaufort, L., Bolton, C.T., Sarr, A.-C., Suchéras-Marx, B., Rosenthal, Y., Donnadieu, Y., Barbarin, N.,
 680 Bova, S., Cornuault, P., Gally, Y., 2022. Cyclic evolution of phytoplankton forced by changes in
 681 tropical seasonality. *Nature* 601, 79–84. <https://doi.org/10.1038/s41586-021-04195-7>

682 Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T.F., Fischer, H., Kipfstuhl, S.,
683 Chappellaz, J., 2015. Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before
684 present. *Geophys. Res. Lett.* 42, 542–549. <https://doi.org/10.1002/2014GL061957>

685 Blain, S., Quéguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C.,
686 Brussaard, C., Carlotti, F., Christaki, U., Corbière, A., Durand, I., Ebersbach, F., Fuda, J.-L., Garcia,
687 N., Gerringa, L., Griffiths, B., Guigue, C., Guillerm, C., Jacquet, S., Jeandel, C., Laan, P., Lefèvre,
688 D., Lo Monaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.-H., Picheral, M., Pondaven,
689 P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M., Thuiller, D.,
690 Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L.,
691 Wagener, T., 2007. Effect of natural iron fertilization on carbon sequestration in the Southern
692 Ocean. *Nature* 446, 1070–1074. <https://doi.org/10.1038/nature05700>

693 Boyd, P.W., 2002. Environmental factors controlling phytoplankton processes in the Southern Ocean.
694 *J Phycol* 38, 844–861. <https://doi.org/10.1046/j.1529-8817.2002.t01-1-01203.x>

695 Brandon, M., Landais, A., Duchamp-Alphonse, S., Favre, V., Schmitz, L., Abrial, H., Prié, F., Extier, T.,
696 Blunier, T., 2020. Exceptionally high biosphere productivity at the beginning of Marine Isotopic
697 Stage 11. *Nat Commun* 11, 2112. <https://doi.org/10.1038/s41467-020-15739-2>

698 Broecker, W., Clark, E., 2009. Ratio of coccolith CaCO₃ to foraminifera CaCO₃ in late Holocene deep
699 sea sediments. *Paleoceanography* 24. <https://doi.org/10.1029/2009PA001731>

700 Cebianchi, M., Luciani, V., Lupi, C., Mancin, N., Lirer, F., Pelosi, N., Trattenero, I., Bordiga, M., Hall,
701 I.R., Sprovieri, M., 2012. Pleistocene biogeochemical record in the south-west Pacific Ocean
702 (images site MD97-2114, Chatham Rise). *J. Quaternary Sci.* 27, 519–530.
703 <https://doi.org/10.1002/jqs.2542>

704 Constable, A.J., Melbourne-Thomas, J., Corney, S.P., Arrigo, K.R., Barbraud, C., Barnes, D.K.A., Bindoff,
705 L., Boyd, P.W., Brandt, A., Costa, D.P., Davidson, A.T., Ducklow, H.W., Emmerson, L., Gutt, J.,
706 Hindell, M.A., Hofmann, E.E., Iida, T., Jacob, S., Johnston, N.M., Kokubun, N., Koubbi, P., Lea, M.-
707 A., Makhado, A., Massom, R.A., Meiners, K., Murphy, E.J., Nicol, S., Reid, K., Riddle, M.J.,
708 Rintoul, S.R., Jr, W.O.S., Southwell, C., Stark, J.S., Sumner, M., Takahashi, K.T., Trathan, P.N.,
709 Welsford, D.C., Westwood, K.J., Wienecke, B.C., Wolf, D., 2014. Climate change and Southern
710 Ocean ecosystems I: how changes in physical habitats directly affect marine biota. *Global*
711 *Change Biology* 23. <https://doi.org/10.1111/gcb.12623>

712 de Vernal, A., Hillaire-Marcel, C., 2008. Natural Variability of Greenland Climate, Vegetation, and Ice
713 Volume During the Past Million Years. *Science* 320, 1622–1625.
714 <https://doi.org/10.1126/science.1153929>

715 Duchamp-Alphonse, S., Siani, G., Michel, E., Beaufort, L., Gally, Y., Jaccard, S.L., 2018. Enhanced
716 ocean-atmosphere carbon partitioning via the carbonate counter pump during the last
717 deglacial. *Nat Commun* 9, 2396. <https://doi.org/10.1038/s41467-018-04625-7>

718 Duncan, B., Carter, L., Dunbar, G., Bostock, H., Neil, H., Scott, G., Hayward, B.W., Sabaa, A., 2016.
719 Interglacial/glacial changes in coccolith-rich deposition in the SW Pacific Ocean: An analogue for
720 a warmer world? *Global and Planetary Change* 144, 252–262.
721 <https://doi.org/10.1016/j.gloplacha.2016.08.001>

722 Ferrari, R., Jansen, M.F., Adkins, J.F., Burke, A., Stewart, A.L., Thompson, A.F., 2014. Antarctic sea ice
723 control on ocean circulation in present and glacial climates. *Proceedings of the National*
724 *Academy of Sciences* 111, 8753–8758. <https://doi.org/10.1073/pnas.1323922111>

725 Findlay, C.S., Flores, J.A., 2000. Subtropical Front fluctuations south of Australia (45°09'S, 146°17'E)
726 for the last 130 ka years based on calcareous nannoplankton. *Marine Micropaleontology* 40,
727 403–416. [https://doi.org/10.1016/S0377-8398\(00\)00045-1](https://doi.org/10.1016/S0377-8398(00)00045-1)

728 Findlay, C.S., Giraudeau, J., 2000. Extant calcareous nannoplankton in the Australian Sector of the
729 Southern Ocean (austral summers 1994 and 1995). *Marine Micropaleontology* 40, 417–439.
730 [https://doi.org/10.1016/S0377-8398\(00\)00046-3](https://doi.org/10.1016/S0377-8398(00)00046-3)

731 Flores, J.-A., Gersonde, R., Sierro, F.J., 1999. Pleistocene fluctuations in the Agulhas Current
732 Retroflection based on the calcareous plankton record. *Marine Micropaleontology* 37, 1–22.
733 [https://doi.org/10.1016/S0377-8398\(99\)00012-2](https://doi.org/10.1016/S0377-8398(99)00012-2)

734 Flores, J.-A., Marino, M., Sierro, F.J., Hodell, D.A., Charles, C.D., 2003. Calcareous plankton dissolution
735 pattern and coccolithophore assemblages during the last 600 kyr at ODP Site 1089 (Cape Basin,
736 South Atlantic): paleoceanographic implications. *Palaeogeography, Palaeoclimatology,*
737 *Palaeoecology* 196, 409–426. [https://doi.org/10.1016/S0031-0182\(03\)00467-X](https://doi.org/10.1016/S0031-0182(03)00467-X)

738 Flores, J.-A., Sierro, F.J., 2007. Pronounced mid-Pleistocene southward shift of the Polar Front in the
739 Atlantic sector of the Southern Ocean. *Deep Sea Research Part II: Topical Studies in*
740 *Oceanography* 54, 2432–2442. <https://doi.org/10.1016/j.dsr2.2007.07.026>

741 François, R., Altabet, M.A., Yu, E.-F., Sigman, D.M., Bacon, M.P., Frank, M., Bohrmann, G., Bareille, G.,
742 Labeyrie, L.D., 1997. Contribution of Southern Ocean surface-water stratification to low
743 atmospheric CO₂ concentrations during the last glacial period. *Nature* 389, 929–935.
744 <https://doi.org/10.1038/40073>

745 Freeman, N.M., Lovenduski, N.S., Gent, P.R., 2016. Temporal variability in the Antarctic Polar Front
746 (2002–2014). *J. Geophys. Res. Oceans* 121, 7263–7276. <https://doi.org/10.1002/2016JC012145>

747 Galbraith, E.D., Jaccard, S.L., 2015. Deglacial weakening of the oceanic soft tissue pump: global
748 constraints from sedimentary nitrogen isotopes and oxygenation proxies. *Quaternary Science*
749 *Reviews* 109, 38–48. <https://doi.org/10.1016/j.quascirev.2014.11.012>

750 Gottschalk, J., Battaglia, G., Fischer, H., Frölicher, T.L., Jaccard, S.L., Jeltsch-Thömmes, A., Joos, F.,
751 Köhler, P., Meissner, K.J., Menviel, L., Nehrbass-Ahles, C., Schmitt, J., Schmittner, A., Skinner,
752 L.C., Stocker, T.F., 2019. Mechanisms of millennial-scale atmospheric CO₂ change in numerical
753 model simulations. *Quaternary Science Reviews* 220, 30–74.
754 <https://doi.org/10.1016/j.quascirev.2019.05.013>

755 Gottschalk, J., Michel, E., Thöle, L.M., Studer, A.S., Hasenfratz, A.P., Schmid, N., Butzin, M., Mazaud,
756 A., Martínez-García, A., Szidat, S., Jaccard, S.L., 2020. Glacial heterogeneity in Southern Ocean
757 carbon storage abated by fast South Indian deglacial carbon release. *Nat Commun* 11, 6192.
758 <https://doi.org/10.1038/s41467-020-20034-1>

759 Govin, A., Michel, E., Labeyrie, L., Waelbroeck, C., Dewilde, F., Jansen, E., 2009. Evidence for
760 northward expansion of Antarctic Bottom Water mass in the Southern Ocean during the last
761 glacial inception. *Paleoceanography* 24, n/a–n/a. <https://doi.org/10.1029/2008PA001603>

762 Grelaud, M., Beaufort, L., Cuven, S., Buchet, N., 2009. Glacial to interglacial primary production and El
763 Niño-Southern Oscillation dynamics inferred from coccolithophores of the Santa Barbara Basin.
764 *Paleoceanography* 24, n/a-n/a. <https://doi.org/10.1029/2007PA001578>

765 Gruber, N., Clement, D., Carter, B.R., Feely, R.A., van Heuven, S., Hoppema, M., Ishii, M., Key, R.M.,
766 Kozyr, A., Lauvset, S.K., Lo Monaco, C., Mathis, J.T., Murata, A., Olsen, A., Perez, F.F., Sabine,
767 C.L., Tanhua, T., Wanninkhof, R., 2019. The oceanic sink for anthropogenic CO₂ from 1994 to
768 2007. *Science* 363, 1193–1199. <https://doi.org/10.1126/science.aau5153>

769 Haddam, N.A., Michel, E., Siani, G., Cortese, G., Bostock, H.C., Duprat, J.M., Isguder, G., 2016.
770 Improving past sea surface temperature reconstructions from the Southern Hemisphere oceans
771 using planktonic foraminiferal census data. *Paleoceanography* 31, 822–837.
772 <https://doi.org/10.1002/2016PA002946>

773 Hain, M.P., Sigman, D.M., Haug, G.H., 2010. Carbon dioxide effects of Antarctic stratification, North
774 Atlantic Intermediate Water formation, and subantarctic nutrient drawdown during the last ice
775 age: Diagnosis and synthesis in a geochemical box model. *Global Biogeochem. Cycles* 24,
776 GB4032. <https://doi.org/10.1029/2010GB003790>

777 Hennehan, M.J., Evans, D., Shankle, M., Burke, J.E., Foster, G.L., Anagnostou, E., Chalk, T.B., Stewart,
778 J.A., Alt, C.H.S., Durrant, J., Hull, P.M., 2017. Size-dependent response of foraminiferal
779 calcification to seawater carbonate chemistry. *Biogeosciences* 14, 3287–3308.
780 <https://doi.org/10.5194/bg-14-3287-2017>

781 Hodell, D.A., Venz, K.A., Charles, C.D., Ninnemann, U.S., 2003. Pleistocene vertical carbon isotope and
782 carbonate gradients in the South Atlantic sector of the Southern Ocean. *Geochem. Geophys.*
783 *Geosyst.* 4, 1–19. <https://doi.org/10.1029/2002GC000367>

784 Howard, W.R., Prell, W.L., 1994. Late Quaternary CaCO₃ production and preservation in the Southern
785 Ocean: Implications for oceanic and atmospheric carbon cycling. *Paleoceanography* 9, 453–482.
786 <https://doi.org/10.1029/93PA03524>

787 Husson, L., Pastier, A.-M., Pedoja, K., Elliot, M., Paillard, D., Authemayou, C., Sarr, A.-C., Schmitt, A.,
788 Cahyarini, S.Y., 2018. Reef Carbonate Productivity During Quaternary Sea Level Oscillations.
789 *Geochem. Geophys. Geosyst.* 19, 1148–1164. <https://doi.org/10.1002/2017GC007335>

790 Jaccard, S.L., Galbraith, E.D., Martínez-García, A., Anderson, R.F., 2016. Covariation of deep Southern
791 Ocean oxygenation and atmospheric CO₂ through the last ice age. *Nature* 530, 207–210.
792 <https://doi.org/10.1038/nature16514>

793 Jaccard, S.L., Galbraith, E.D., Sigman, D.M., Haug, G.H., 2010. A pervasive link between Antarctic ice
794 core and subarctic Pacific sediment records over the past 800kyrs. *Quaternary Science Reviews*
795 29, 206–212. <https://doi.org/10.1016/j.quascirev.2009.10.007>

796 Jaccard, S.L., Galbraith, E.D., Sigman, D.M., Haug, G.H., Francois, R., Pedersen, T.F., Dulski, P.,
797 Thierstein, H.R., 2009. Subarctic Pacific evidence for a glacial deepening of the oceanic respired
798 carbon pool. *Earth and Planetary Science Letters* 277, 156–165.
799 <https://doi.org/10.1016/j.epsl.2008.10.017>

800 Jaccard, S.L., Hayes, C.T., Martinez-Garcia, A., Hodell, D.A., Anderson, R.F., Sigman, D.M., Haug, G.H.,
801 2013. Two Modes of Change in Southern Ocean Productivity Over the Past Million Years.
802 *Science* 339, 1419–1423. <https://doi.org/10.1126/science.1227545>

803 Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B.,
804 Nouet, J., Barnola, J.M., Chappellaz, J., Fischer, H., Gallet, J.C., Johnsen, S., Leuenberger, M.,
805 Louergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A.,
806 Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J.P., Stenni, B.,
807 Stocker, T.F., Tison, J.L., Werner, M., Wolff, E.W., 2007. Orbital and Millennial Antarctic Climate
808 Variability over the Past 800,000 Years. *Science* 317, 793–796.
809 <https://doi.org/10.1126/science.1141038>

810 Kemp, A.E.S., Grigorov, I., Pearce, R.B., Naveira Garabato, A.C., 2010. Migration of the Antarctic Polar
811 Front through the mid-Pleistocene transition: evidence and climatic implications. *Quaternary*
812 *Science Reviews* 29, 1993–2009. <https://doi.org/10.1016/j.quascirev.2010.04.027>

813 Kohfeld, K.E., 2005. Role of Marine Biology in Glacial-Interglacial CO₂ Cycles. *Science* 308, 74–78.
814 <https://doi.org/10.1126/science.1105375>

815 Kolla, V., Bé, A.W.H., Biscaye, P.E., 1976. Calcium carbonate distribution in the surface sediments of
816 the Indian Ocean. *J. Geophys. Res.* 81, 2605–2616. <https://doi.org/10.1029/JC081i015p02605>

817 Kousis, I., Koutsodendris, A., Peyron, O., Leicher, N., Francke, A., Wagner, B., Giaccio, B., Knipping, M.,
818 Pross, J., 2018. Centennial-scale vegetation dynamics and climate variability in SE Europe during
819 Marine Isotope Stage 11 based on a pollen record from Lake Ohrid. *Quaternary Science Reviews*
820 190, 20–38. <https://doi.org/10.1016/j.quascirev.2018.04.014>

821 Krumhardt, K.M., Long, M.C., Lindsay, K., Levy, M.N., 2020. Southern Ocean Calcification Controls the
822 Global Distribution of Alkalinity. *Global Biogeochem. Cycles* 34.
823 <https://doi.org/10.1029/2020GB006727>

824 Landschützer, P., Gruber, N., Haumann, F.A., Rödenbeck, C., Bakker, D.C.E., van Heuven, S.,
825 Hoppema, M., Metzl, N., Sweeney, C., Takahashi, T., Tilbrook, B., Wanninkhof, R., 2015. The
826 reinvigoration of the Southern Ocean carbon sink. *Science* 349, 1221–1224.
827 <https://doi.org/10.1126/science.aab2620>

828 Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A.C.M., Levrard, B., 2004. A long-term
829 numerical solution for the insolation quantities of the Earth. *A&A* 428, 261–285.
830 <https://doi.org/10.1051/0004-6361:20041335>

831 Lauderdale, J.M., Garabato, A.C.N., Oliver, K.I.C., Follows, M.J., Williams, R.G., 2013. Wind-driven
832 changes in Southern Ocean residual circulation, ocean carbon reservoirs and atmospheric CO₂.
833 *Clim Dyn* 41, 2145–2164. <https://doi.org/10.1007/s00382-012-1650-3>

834 Le, J., Shackleton, N.J., 1992. Carbonate Dissolution Fluctuations in the Western Equatorial Pacific
835 During the Late Quaternary. *Paleoceanography* 7, 21–42. <https://doi.org/10.1029/91PA02854>

836 Lovenduski, N.S., Gruber, N., Doney, S.C., 2008. Toward a mechanistic understanding of the decadal
837 trends in the Southern Ocean carbon sink. *Global Biogeochem. Cycles* 22, n/a-n/a.
838 <https://doi.org/10.1029/2007GB003139>

839 Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J.,
840 Fischer, H., Kawamura, K., Stocker, T.F., 2008. High-resolution carbon dioxide concentration
841 record 650,000–800,000 years before present. *Nature* 453, 379–382.
842 <https://doi.org/10.1038/nature06949>

843 Malinverno, E., Triantaphyllou, M.V., Dimiza, M.D., 2015. Coccolithophore assemblage distribution
844 along a temperate to polar gradient in the West Pacific sector of the Southern Ocean (January
845 2005). *Micropaleontology* 61, 489–506.

846 Marino, M., Maiorano, P., Lirer, F., 2008. Changes in calcareous nannofossil assemblages during the
847 Mid-Pleistocene Revolution. *Marine Micropaleontology* 69, 70–90.
848 <https://doi.org/10.1016/j.marmicro.2007.11.010>

849 Marino, M., Maiorano, P., Lirer, F., Pelosi, N., 2009. Response of calcareous nannofossil assemblages
850 to paleoenvironmental changes through the mid-Pleistocene revolution at Site 1090 (Southern
851 Ocean). *Palaeogeography, Palaeoclimatology, Palaeoecology* 280, 333–349.
852 <https://doi.org/10.1016/j.palaeo.2009.06.019>

853 Martin, J.H., 1990. Glacial-interglacial CO₂ change: The iron hypothesis. *Paleoceanography* 5, 1–13.

854 Martínez-García, A., Rosell-Melé, A., Geibert, W., Gersonde, R., Masqué, P., Gaspari, V., Barbante, C.,
855 2009. Links between iron supply, marine productivity, sea surface temperature, and CO₂ over
856 the last 1.1 Ma. *Paleoceanography* 24, n/a-n/a. <https://doi.org/10.1029/2008PA001657>

857 Martínez-García, A., Sigman, D.M., Ren, H., Anderson, R.F., Straub, M., Hodell, D.A., Jaccard, S.L.,
858 Eglinton, T.I., Haug, G.H., 2014. Iron Fertilization of the Subantarctic Ocean During the Last Ice
859 Age. *Science* 343, 1347–1350. <https://doi.org/10.1126/science.1246848>

860 Melles, M., Brigham-Grette, J., Minyuk, P.S., Nowaczyk, N.R., Wennrich, V., DeConto, R.M., Anderson,
861 P.M., Andreev, A.A., Coletti, A., Cook, T.L., Haltia-Hovi, E., Kukkonen, M., Lozhkin, A.V., Rosen,
862 P., Tarasov, P., Vogel, H., Wagner, B., 2012. 2.8 Million Years of Arctic Climate Change from Lake
863 El'gygytyn, NE Russia. *Science* 337, 315–320. <https://doi.org/10.1126/science.1222135>

864 Menviel, L., Spence, P., Yu, J., Chamberlain, M.A., Matear, R.J., Meissner, K.J., England, M.H., 2018.
865 Southern Hemisphere westerlies as a driver of the early deglacial atmospheric CO₂ rise. *Nat*
866 *Commun* 9, 2503. <https://doi.org/10.1038/s41467-018-04876-4>

867 Michel, E., Waelbroeck, C., 2017. MD 136/VIGO à bord du R/V Marion Dufresne. La Réunion 3 janvier
868 2004–La Réunion 9 février 2004. Les Rapports de Campagnes à la Mer.

869 Moore, C.M., Mills, M.M., Arrigo, K.R., Berman-Frank, I., Bopp, L., Boyd, P.W., Galbraith, E.D., Geider,
870 R.J., Guieu, C., Jaccard, S.L., Jickells, T.D., La Roche, J., Lenton, T.M., Mahowald, N.M., Marañón,
871 E., Marinov, I., Moore, J.K., Nakatsuka, T., Oschlies, A., Saito, M.A., Thingstad, T.F., Tsuda, A.,
872 Ulloa, O., 2013. Processes and patterns of oceanic nutrient limitation. *Nature Geosci* 6, 701–
873 710. <https://doi.org/10.1038/ngeo1765>

874 Moy, A.D., Howard, W.R., Bray, S.G., Trull, T.W., 2009. Reduced calcification in modern Southern
875 Ocean planktonic foraminifera. *Nature Geosci* 2, 276–280. <https://doi.org/10.1038/ngeo460>

876 Nehrbass-Ahles, C., Shin, J., Schmitt, J., Bereiter, B., Joos, F., Schilt, A., Schmidely, L., Silva, L., Teste,
877 G., Grilli, R., Chappellaz, J., Hodell, D., Fischer, H., Stocker, T.F., 2020. Abrupt CO₂ release to the
878 atmosphere under glacial and early interglacial climate conditions. *Science* 369, 1000–1005.
879 <https://doi.org/10.1126/science.aay8178>

880 Nougier, J., 1970. Contribution à l'étude géologique et géomorphologique des îles Kerguelen (Terres
881 Australes et Antarctiques Françaises). *Com. natl. Fr. Rech. Antarct.* 27, 246P.

882 Orsi, A.H., Whitworth, T., Nowlin, W.D., 1995. On the meridional extent and fronts of the Antarctic
883 Circumpolar Current. *Deep Sea Research Part I: Oceanographic Research Papers* 42, 641–673.
884 [https://doi.org/10.1016/0967-0637\(95\)00021-W](https://doi.org/10.1016/0967-0637(95)00021-W)

885 Paillard, D., Labeyrie, L., Yiou, P., 1996. Macintosh Program performs time-series analysis. *Eos,*
886 *Transactions American Geophysical Union* 77, 379–379. <https://doi.org/10.1029/96EO00259>

887 Park, Y., Durand, I., Kestenare, E., Rougier, G., Zhou, M., d’Ovidio, F., Cotté, C., Lee, J., 2014. Polar
888 Front around the Kerguelen Islands: An up-to-date determination and associated circulation of
889 surface/subsurface waters. *J. Geophys. Res. Oceans* 119, 6575–6592.
890 <https://doi.org/10.1002/2014JC010061>

891 Park, Y. -H., Park, T., Kim, T. -W., Lee, S. -H., Hong, C. -S., Lee, J. -H., Rio, M. -H., Pujol, M. -I.,
892 Ballarotta, M., Durand, I., Provost, C., 2019. Observations of the Antarctic Circumpolar Current
893 Over the Udintsev Fracture Zone, the Narrowest Choke Point in the Southern Ocean. *J.*
894 *Geophys. Res. Oceans* 124, 4511–4528. <https://doi.org/10.1029/2019JC015024>

895 Park, Y.-H., Vivier, F., Roquet, F., Kestenare, E., 2009. Direct observations of the ACC transport across
896 the Kerguelen Plateau. *Geophys. Res. Lett.* 36, L18603. <https://doi.org/10.1029/2009GL039617>

897 Patil, S.M., Mohan, R., Shetye, S.S., Gazi, S., Choudhari, P., Jafar, S., 2020. Interannual changes of
898 austral summer coccolithophore assemblages and southward expanse in the Southern Indian
899 Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* 178, 104765.
900 <https://doi.org/10.1016/j.dsr2.2020.104765>

901 Pichon, J.-J., Labeyrie, L.D., Bareille, G., Labracherie, M., Duprat, J., Jouzel, J., 1992. Surface water
902 temperature changes in the high latitudes of the southern hemisphere over the Last Glacial-
903 Interglacial Cycle. *Paleoceanography* 7, 289–318. <https://doi.org/10.1029/92PA00709>

904 Pollard, R., Tréguer, P., Read, J., 2006. Quantifying nutrient supply to the Southern Ocean. *J.*
905 *Geophys. Res.* 111, C05011. <https://doi.org/10.1029/2005JC003076>

906 Pollard, R.T., Salter, I., Sanders, R.J., Lucas, M.I., Moore, C.M., Mills, R.A., Statham, P.J., Allen, J.T.,
907 Baker, A.R., Bakker, D.C.E., Charette, M.A., Fielding, S., Fones, G.R., French, M., Hickman, A.E.,
908 Holland, R.J., Hughes, J.A., Jickells, T.D., Lampitt, R.S., Morris, P.J., Nédélec, F.H., Nielsdóttir, M.,
909 Planquette, H., Popova, E.E., Poulton, A.J., Read, J.F., Seeyave, S., Smith, T., Stinchcombe, M.,
910 Taylor, S., Thomalla, S., Venables, H.J., Williamson, R., Zubkov, M.V., 2009. Southern Ocean
911 deep-water carbon export enhanced by natural iron fertilization. *Nature* 457, 577–580.
912 <https://doi.org/10.1038/nature07716>

913 Prell, W.L., 1985. Stability of low-latitude sea-surface temperatures: an evaluation of the CLIMAP
914 reconstruction with emphasis on the positive SST anomalies. *U.S. Dep of Energy* 60.

915 Prokopenko, A.A., Bezrukova, E.V., Khursevich, G.K., Solotchina, E.P., Kuzmin, M.I., Tarasov, P.E.,
916 2010. Climate in continental interior Asia during the longest interglacial of the past 500 000
917 years: the new MIS 11 records from Lake Baikal, SE Siberia. *Clim. Past* 18.

918 Qin, B., Li, T., Xiong, Z., Algeo, T.J., Jia, Q., 2018. Deep-Water Carbonate Ion Concentrations in the
919 Western Tropical Pacific Since the Mid-Pleistocene: A Major Perturbation During the Mid-
920 Brunhes. *J. Geophys. Res. Oceans* 123, 6876–6892. <https://doi.org/10.1029/2018JC014084>

921 Quéguiner, B., 2013. Iron fertilization and the structure of planktonic communities in high nutrient
 922 regions of the Southern Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* 90,
 923 43–54. <https://doi.org/10.1016/j.dsr2.2012.07.024>

924 Rae, J.W.B., Burke, A., Robinson, L.F., Adkins, J.F., Chen, T., Cole, C., Greenop, R., Li, T., Littley, E.F.M.,
 925 Nita, D.C., Stewart, J.A., Taylor, B.J., 2018. CO₂ storage and release in the deep Southern Ocean
 926 on millennial to centennial timescales. *Nature* 562, 569–573. [https://doi.org/10.1038/s41586-](https://doi.org/10.1038/s41586-018-0614-0)
 927 018-0614-0

928 Rembauville, M., Meilland, J., Ziveri, P., Schiebel, R., Blain, S., Salter, I., 2016. Planktic foraminifer and
 929 coccolith contribution to carbonate export fluxes over the central Kerguelen Plateau. *Deep Sea*
 930 *Research Part I: Oceanographic Research Papers* 111, 91–101.
 931 <https://doi.org/10.1016/j.dsr.2016.02.017>

932 Rickaby, R.E.M., Bard, E., Sonzogni, C., Rostek, F., Beaufort, L., Barker, S., Rees, G., Schrag, D.P., 2007.
 933 Coccolith chemistry reveals secular variations in the global ocean carbon cycle? *Earth and*
 934 *Planetary Science Letters* 253, 83–95. <https://doi.org/10.1016/j.epsl.2006.10.016>

935 Rigual Hernández, A.S., Trull, T.W., Nodder, S.D., Flores, J.A., Bostock, H., Abrantes, F., Eriksen, R.S.,
 936 Sierro, F.J., Davies, D.M., Ballegeer, A.-M., Fuertes, M.A., Northcote, L.C., 2020. Coccolithophore
 937 biodiversity controls carbonate export in the Southern Ocean. *Biogeosciences* 17, 245–263.
 938 <https://doi.org/10.5194/bg-17-245-2020>

939 Rigual-Hernández, A.S., Trull, T.W., Bray, S.G., Closset, I., Armand, L.K., 2015. Seasonal dynamics in
 940 diatom and particulate export fluxes to the deep sea in the Australian sector of the southern
 941 Antarctic Zone. *Journal of Marine Systems* 142, 62–74.
 942 <https://doi.org/10.1016/j.jmarsys.2014.10.002>

943 Saavedra-Pellitero, M., Baumann, K.-H., Gallagher, S.J., Sagawa, T., Tada, R., 2019. Paleooceanographic
 944 evolution of the Japan Sea over the last 460 kyr – A coccolithophore perspective. *Marine*
 945 *Micropaleontology* 152, 101720. <https://doi.org/10.1016/j.marmicro.2019.01.001>

946 Saavedra-Pellitero, M., Baumann, K.-H., Lamy, F., Köhler, P., 2017a. Coccolithophore variability across
 947 Marine Isotope Stage 11 in the Pacific sector of the Southern Ocean and its potential impact on
 948 the carbon cycle. *Paleoceanography* 32, 864–880. <https://doi.org/10.1002/2017PA003156>

949 Saavedra-Pellitero, M., Baumann, K.-H., Ullermann, J., Lamy, F., 2017b. Marine Isotope Stage 11 in
 950 the Pacific sector of the Southern Ocean; a coccolithophore perspective. *Quaternary Science*
 951 *Reviews* 158, 1–14. <https://doi.org/10.1016/j.quascirev.2016.12.020>

952 Salter, I., Schiebel, R., Ziveri, P., Movellan, A., Lampitt, R., Wolff, G.A., 2014. Carbonate counter pump
 953 stimulated by natural iron fertilization in the Polar Frontal Zone. *Nature Geoscience* 7, 885–889.
 954 <https://doi.org/10.1038/NGEO2285>

955 Sarmiento, J.L., Gruber, N., Brzezinski, M.A., Dunne, J.P., 2004. High-latitude controls of thermocline
 956 nutrients and low latitude biological productivity. *Nature* 427, 56–60.
 957 <https://doi.org/10.1038/nature02127>

958 Schaefer, G., Rodger, J.S., Hayward, B.W., Kennett, J.P., Sabaa, A.T., Scott, G.H., 2005. Planktic
 959 foraminiferal and sea surface temperature record during the last 1 Myr across the Subtropical
 960 Front, Southwest Pacific. *Marine Micropaleontology* 54, 191–212.
 961 <https://doi.org/10.1016/j.marmicro.2004.12.001>

962 Schlitzer, R., 2018. Ocean Data View. <https://odv.awi.de>.

963 Siani, G., Michel, E., De Pol-Holz, R., DeVries, T., Lamy, F., Carel, M., Isguder, G., Dewilde, F.,
 964 Laurantou, A., 2013. Carbon isotope records reveal precise timing of enhanced Southern Ocean
 965 upwelling during the last deglaciation. *Nat Commun* 4, 2758.
 966 <https://doi.org/10.1038/ncomms3758>

967 Sigman, D.M., Boyle, E.A., 2000. Glacial/interglacial variations in atmospheric carbon dioxide. *Nature*
 968 407, 859–869. <https://doi.org/10.1038/35038000>

969 Sigman, D.M., Fripiat, F., Studer, A.S., Kemeny, P.C., Martínez-García, A., Hain, M.P., Ai, X., Wang, X.,
 970 Ren, H., Haug, G.H., 2021. The Southern Ocean during the ice ages: A review of the Antarctic
 971 surface isolation hypothesis, with comparison to the North Pacific. *Quaternary Science Reviews*
 972 254, 106732. <https://doi.org/10.1016/j.quascirev.2020.106732>

973 Sigman, D.M., McCorkle, D.C., Martin, W.R., 1998. The calcite lysocline as a constraint on
 974 glacial/interglacial low-latitude production changes. *Global Biogeochem. Cycles* 12, 409–427.
 975 <https://doi.org/10.1029/98GB01184>

976 Skinner, L.C., Fallon, S., Waelbroeck, C., Michel, E., Barker, S., 2010. Ventilation of the Deep Southern
 977 Ocean and Deglacial CO₂ Rise. *Science* 328, 1147–1151.
 978 <https://doi.org/10.1126/science.1183627>

979 Smith, H.E.K., Poulton, A.J., Garley, R., Hopkins, J., Lubelczyk, L.C., Drapeau, D.T., Rauschenberg, S.,
 980 Twining, B.S., Bates, N.R., Balch, W.M., 2017. The influence of environmental variability on the
 981 biogeography of coccolithophores and diatoms in the Great Calcite Belt. *Biogeosciences* 14,
 982 4905–4925. <https://doi.org/10.5194/bg-14-4905-2017>

983 Sokolov, S., Rintoul, S.R., 2009. Circumpolar structure and distribution of the Antarctic Circumpolar
 984 Current fronts: 1. Mean circumpolar paths. *J. Geophys. Res.* 114, C11018.
 985 <https://doi.org/10.1029/2008JC005108>

986 Sokolov, S., Rintoul, S.R., 2007. On the relationship between fronts of the Antarctic Circumpolar
 987 Current and surface chlorophyll concentrations in the Southern Ocean. *J. Geophys. Res.* 112,
 988 C07030. <https://doi.org/10.1029/2006JC004072>

989 Sowers, T., Bender, M., Labeyrie, L., Martinson, D., Jouzel, J., Raynaud, D., Pichon, J.J., Korotkevich,
 990 Y.S., 1993. A 135,000-year Vostok-SPECMAP common temporal framework. *Paleoceanography*
 991 8, 737–766.

992 Spero, H.J., 2002. The Cause of Carbon Isotope Minimum Events on Glacial Terminations. *Science*
 993 296, 522–525. <https://doi.org/10.1126/science.1069401>

994 Stein, K., Timmermann, A., Kwon, E.Y., Friedrich, T., 2020. Timing and magnitude of Southern Ocean
 995 sea ice/carbon cycle feedbacks. *Proc Natl Acad Sci USA* 117, 4498–4504.
 996 <https://doi.org/10.1073/pnas.1908670117>

997 Stephens, B.B., Keeling, R.F., 2000. The influence of Antarctic sea ice on glacial–interglacial CO₂
 998 variations. *Nature* 404, 171–174.

999 Tagliabue, A., Sallee, J.-B., Bowie, A.R., Levy, M., Swart, S., Boyd, P.W., 2014. Surface-water iron
 1000 supplies in the Southern Ocean sustained by deep winter mixing. *Nature Geoscience* 7, 314–
 1001 320. <https://doi.org/10.1038/ngeo2101>

1002 Therón, R., Paillard, D., Cortijo, E., Flores, J.-A., Vaquero, M., Sierro, F.J., Waelbroeck, C., 2004. Rapid
1003 reconstruction of paleoenvironmental features using a new multiplatform program.
1004 *Micropaleontology* 50, 391–395.

1005 Thöle, L.M., Amsler, H.E., Moretti, S., Auderset, A., Gilgannon, J., Lippold, J., Vogel, H., Crosta, X.,
1006 Mazaud, A., Michel, E., Martínez-García, A., Jaccard, S.L., 2019. Glacial-interglacial dust and
1007 export production records from the Southern Indian Ocean. *Earth and Planetary Science Letters*
1008 525, 115716. <https://doi.org/10.1016/j.epsl.2019.115716>

1009 Toggweiler, J.R., Russell, J.L., Carson, S.R., 2006. Midlatitude westerlies, atmospheric CO₂, and
1010 climate change during the ice ages. *Paleoceanography* 21, n/a-n/a.
1011 <https://doi.org/10.1029/2005PA001154>

1012 Tzedakis, P.C., Hooghiemstra, H., Pälike, H., 2006. The last 1.35 million years at Tenaghi Philippon:
1013 revised chronostratigraphy and long-term vegetation trends. *Quaternary Science Reviews* 25,
1014 3416–3430. <https://doi.org/10.1016/j.quascirev.2006.09.002>

1015 Ullermann, J., 2015. Glacial-interglacial oceanic changes in the central Pacific sector of the Southern
1016 Ocean during the past 500 ka (PhD Thesis). <http://www.suub.uni-bremen.de>.

1017 Vázquez Riveiros, N., Waelbroeck, C., Skinner, L., Roche, D.M., Duplessy, J.-C., Michel, E., 2010.
1018 Response of South Atlantic deep waters to deglacial warming during Terminations V and I. *Earth*
1019 *and Planetary Science Letters* 298, 323–333. <https://doi.org/10.1016/j.epsl.2010.08.003>

1020 Waelbroeck, C., Jouzel, J., Labeyrie, L., Lorius, C., Labracherie, M., Stiévenard, M., Barkov, N.I., 1995.
1021 A comparison of the Vostok ice deuterium record and series from Southern Ocean core MD 88-
1022 770 over the last two glacial-interglacial cycles. *Climate Dynamics* 12, 113–123.

1023 Zondervan, I., 2007. The effects of light, macronutrients, trace metals and CO₂ on the production of
1024 calcium carbonate and organic carbon in coccolithophores—A review. *Deep Sea Research Part*
1025 *II: Topical Studies in Oceanography* 54, 521–537. <https://doi.org/10.1016/j.dsr2.2006.12.004>

1026 Zweng, M., Seidov, D., Boyer, T., Locarnini, M., Garcia, H., Mishonov, A., Baranova, O., Weathers, K.,
1027 Paver, C., Smolyar, I., 2019. *World ocean atlas 2018, volume 2: Salinity*.