



# **The Role of Neutral Hydrogen in Setting the Abundances of Molecular Species in the Milky Way's Diffuse Interstellar Medium. I. Observational Constraints from ALMA and NOEMA**

Daniel R. Rybarczyk, Snežana Stanimirović, Munan Gong, Brian Babler, Claire E. Murray, Maryvonne Gerin, Jan Martin Winters, Gan Luo, T. M. Dame, Lucille Steffes

## **► To cite this version:**

Daniel R. Rybarczyk, Snežana Stanimirović, Munan Gong, Brian Babler, Claire E. Murray, et al.. The Role of Neutral Hydrogen in Setting the Abundances of Molecular Species in the Milky Way's Diffuse Interstellar Medium. I. Observational Constraints from ALMA and NOEMA. *The Astrophysical Journal*, 2022, 928, <10.3847/1538-4357/ac5035>. <insu-03717123>

**HAL Id: insu-03717123**

**<https://insu.hal.science/insu-03717123v1>**

Submitted on 9 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



# The Role of Neutral Hydrogen in Setting the Abundances of Molecular Species in the Milky Way's Diffuse Interstellar Medium. I. Observational Constraints from ALMA and NOEMA

Daniel R. Rybarczyk<sup>1</sup> , Snežana Stanimirović<sup>1</sup> , Munan Gong<sup>2</sup> , Brian Babler<sup>1</sup> , Claire E. Murray<sup>3</sup> , Maryvonne Gerin<sup>4</sup> , Jan Martin Winters<sup>5</sup> , Gan Luo<sup>6,7</sup> , T. M. Dame<sup>8</sup> , and Lucille Steffes<sup>1</sup>

<sup>1</sup> University of Wisconsin–Madison, Department of Astronomy, 475 N Charter St, Madison, WI 53703, USA; [rybarczyk@astro.wisc.edu](mailto:rybarczyk@astro.wisc.edu)

<sup>2</sup> Max-Planck-Institut für Extraterrestrische Physik, Garching by Munich, D-85748, Germany

<sup>3</sup> Department of Physics & Astronomy, Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218, USA

<sup>4</sup> LERMA, Observatoire de Paris, PSL Research University, CNRS, Ecole Normale Supérieure, Sorbonne Université, F-75005 Paris, France

<sup>5</sup> Institut de Radioastronomie Millimétrique (IRAM), 300 rue de la Piscine, F-38406 St. Martin d'Hères, France

<sup>6</sup> School of Astronomy and Space Science, Nanjing University, Nanjing 210093, People's Republic of China

<sup>7</sup> Key Laboratory of Modern Astronomy and Astrophysics (Nanjing University), Ministry of Education, Nanjing 210093, People's Republic of China

<sup>8</sup> Center for Astrophysics | Harvard & Smithsonian, 60 Garden St Cambridge, MA 02138, USA

Received 2021 June 10; revised 2021 September 8; accepted 2021 September 10; published 2022 March 28

## Abstract

We have complemented existing observations of H I absorption with new observations of HCO<sup>+</sup>, C<sub>2</sub>H, HCN, and HNC absorption from the Atacama Large Millimeter/submillimeter Array and the Northern Extended Millimeter Array in the directions of 20 background radio continuum sources with  $4^\circ \leq |b| \leq 81^\circ$  to constrain the atomic gas conditions that are suitable for the formation of diffuse molecular gas. We find that these molecular species form along sightlines where  $A_V \gtrsim 0.25$ , consistent with the threshold for the H I-to-H<sub>2</sub> transition at solar metallicity. Moreover, we find that molecular gas is associated only with structures that have an H I optical depth  $>0.1$ , a spin temperature  $<80$  K, and a turbulent Mach number  $\gtrsim 2$ . We also identify a broad, faint component to the HCO<sup>+</sup> absorption in a majority of sightlines. Compared to the velocities where strong, narrow HCO<sup>+</sup> absorption is observed, the H I at these velocities has a lower cold neutral medium fraction and negligible CO emission. The relative column densities and linewidths of the different molecular species observed here are similar to those observed in previous experiments over a range of Galactic latitudes, suggesting that gas in the solar neighborhood and gas in the Galactic plane are chemically similar. For a select sample of previously observed sightlines, we show that the absorption line profiles of HCO<sup>+</sup>, HCN, HNC, and C<sub>2</sub>H are stable over periods of  $\sim 3$  yr and  $\sim 25$  yr, likely indicating that molecular gas structures in these directions are at least  $\gtrsim 100$  au in size.

*Unified Astronomy Thesaurus concepts:* [Interstellar absorption \(831\)](#); [Interstellar phases \(850\)](#); [Interstellar atomic gas \(833\)](#); [Diffuse molecular clouds \(381\)](#); [Molecule formation \(2076\)](#)

## 1. Introduction

The formation of molecular gas in the diffuse interstellar medium (ISM) marks the first stage of molecular cloud formation and interstellar chemistry. A variety of processes are believed to mediate molecule formation, including, for example, shielding (Krumholz et al. 2009; Sternberg et al. 2014), shock-driven turbulence (Inoue & Inutsuka 2012), and turbulent dissipation (Godard et al. 2009; Lesaffre et al. 2020), but the relative importance of these different effects remains uncertain, from both observational and theoretical perspectives.

Molecular gas forms within the turbulent, multiphase atomic ISM. Since at least the early work of Pikel'Ner (1968) and Field et al. (1969), it has been understood that atomic hydrogen (H I) can exist in a warm, diffuse phase (the warm neutral medium, WNM) and a cold, clumpy phase (the cold neutral medium, CNM). The WNM and CNM have been well characterized theoretically (Wolfire et al. 2003) and observationally in the case of the Milky Way (Heiles & Troland 2003; Murray et al. 2018). However, observations have also established that  $\sim 20\%$  of the H I in the Milky Way is in a

thermally unstable phase (the unstable neutral medium, UNM), with intermediate temperature and density (Murray et al. 2018). Hydrodynamical and magnetohydrodynamical simulations have shown that CNM structures can form out of the diffuse WNM as the result of dynamical processes like shocks (Koyama & Inutsuka 2002; Hennebelle & Audit 2007; Inoue & Inutsuka 2012). Thermal instabilities in shocked gas lead to the condensation of CNM structures with densities several orders of magnitude higher than typical WNM densities. The smallest ( $\lesssim 0.1$  pc) and densest of these structures may represent tiny-scale atomic structure (TSAS; Hennebelle & Audit 2007; Rybarczyk et al. 2020), an overdense, overpressured component of the atomic ISM observed in high-resolution H I absorption measurements (Heiles 1997; Stanimirović & Zweibel 2018, and references therein). Theoretical predictions suggest that molecular hydrogen (H<sub>2</sub>) forms if the H I piled up behind a shock reaches a sufficiently high column density (Inutsuka et al. 2015) and the gas stays cold (Heitsch et al. 2006), although this depends on the metallicity (e.g., Bolatto et al. 2011) and the orientation of the local magnetic field with respect to the shock motion (Inoue & Inutsuka 2012). The H I-to-H<sub>2</sub> transition—where a large fraction of atomic hydrogen is converted to molecular hydrogen—has been observed to occur at a total column density  $N \gtrsim 5 \times 10^{20} \text{ cm}^{-2}$  at solar metallicity (Savage et al. 1977).



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

While it is clear that the turbulent, multiphase ISM is an important ingredient for the formation and survival of molecules, it is still not understood how local properties of H I affect the molecular fraction. Several recent studies have provided hints that the underlying physical properties of the H I, such as the level of turbulence and the presence of thermally unstable H I, play an important role. Observationally, Stanimirović et al. (2014) and Nguyen et al. (2019) have shown that molecular clouds are embedded within atomic gas that has a high CNM fraction relative to random diffuse regions. By modeling the H I-to-H<sub>2</sub> transition in Perseus, Bialy et al. (2015) found that a mixture of CNM and UNM (and perhaps WNM) gas was important in controlling the H I-to-H<sub>2</sub> transition. The widths of the transition regions between warm H I, cold H I, and cold H<sub>2</sub>—and therefore the local properties of the H I-to-H<sub>2</sub> transition—also depend on the level of turbulence (Bialy et al. 2017; Lesaffre et al. 2007), and turbulent mixing between the WNM and CNM can enhance the formation of certain molecular species (Lesaffre et al. 2007; Glover & Clark 2012).

In this work, we investigate the early stages of molecule formation in the diffuse ISM and connect this with the underlying properties of atomic gas using new absorption line observations of HCN, C<sub>2</sub>H, HCO<sup>+</sup>, and HNC in the directions of 20 background radio continuum sources where the 21-SPONGE project (21 cm Spectral Line Observations of Neutral Gas with the Karl G. Jansky Very Large Array; Murray et al. 2015, 2018) previously observed H I in emission and absorption. In Section 2, we present these observations, obtained using the Atacama Large Millimeter/submillimeter Array (ALMA) and the Northern Extended Millimeter Array (NOEMA), along with existing observations in these directions, including observations of H I emission from the Arecibo Observatory and H I absorption from the Very Large Array (VLA) obtained by the 21-SPONGE project (Murray et al. 2015, 2018), maps of interstellar reddening and dust temperature from the Planck satellite (Planck Collaboration et al. 2014a, 2014b), and observations of CO emission from the Dame et al. (2001) survey. 21-SPONGE mainly targeted sources at Galactic latitude  $>10^\circ$ , where H I spectra are simpler and radiative transfer calculations are easier than in the Galactic plane. The sensitivity of our molecular line spectra rivals or exceeds most pre-ALMA surveys (e.g., Lucas & Liszt 1996, 2000; Liszt & Lucas 2001), and our sample represents one of the largest homogeneous samples of Galactic absorption measurements at millimeter wavelengths to date. We outline methods for extracting molecular column densities and decomposing absorption spectra into Gaussian components in Section 3. In Section 4, we test how the observed molecular column densities depend on the line-of-sight gas properties, including the extinction and the CNM and UNM column densities. We also investigate a broad, faint component of the HCO<sup>+</sup> absorption seen in the directions of most background sources. In Section 5, we establish thresholds for the H I optical depth, spin temperature, and turbulent Mach number required for the onset of molecule formation. We then determine the relative abundances and linewidths of the four different molecular species observed in this work, and compare our results with previous works, in Section 6. In Section 7, we investigate the temporal stability of absorption line profiles by comparing our results to previous observations of HCN, C<sub>2</sub>H, HCO<sup>+</sup>, and HNC for select lines of sight. This serves as a

probe for au-scale structure. Finally, we discuss our results in Section 8 and present conclusions in Section 9.

This is the first of two complementary papers. In Rybarczyk et al. (2022; Paper II hereafter), we compare the observational results presented here to predictions from the Gong et al. (2017) photodissociation region (PDR) chemical model and the Gong et al. (2020) ISM simulations.

## 2. Observations

### 2.1. Observations with ALMA (ALMA-SPONGE)

We have observed C<sub>2</sub>H, HCN, HCO<sup>+</sup>, and HNC in absorption with ALMA during observing cycles 6 and 7 (ALMA-SPONGE projects 2018.1.00585.S and 2019.1.01809.S; PI: Stanimirović) in the directions of 20 bright background radio continuum sources previously observed at 21 cm wavelength using the VLA by the 21-SPONGE project. Observations were obtained between 2018 October and 2020 March. The transitions covered by our observations are listed in Table 1. 89 GHz fluxes ranged from 0.03 Jy to 11.8 Jy. The HCO<sup>+</sup> and HNC spectra were obtained at 30.5 kHz frequency channel spacing (0.1 km s<sup>-1</sup> velocity channel spacing), while the HCN and C<sub>2</sub>H spectra were obtained at 61 kHz frequency channel spacing (0.2 km s<sup>-1</sup> velocity channel spacing). In the analysis that follows, all spectra have been smoothed to 0.4 km s<sup>-1</sup> velocity resolution, comparable to that of the 21-SPONGE H I absorption spectra.

With the exception of 3C111 and 3C123, all of the ALMA-SPONGE background radio continuum sources are unresolved and treated as single point sources. 3C111 is a three-component radio galaxy; we have independent observations of all three components, separated by 1'–3', each of which is unresolved. Although the 1420 MHz fluxes vary by less than a factor of 3 across the three components, the 89 GHz fluxes change by over two orders of magnitude. We do not consider the spectra in the direction of 3C111C hereafter, as they lack adequate sensitivity to accurately measure molecular abundances ( $\sigma_\tau \gtrsim 1$ ). The optical depth noise of the C<sub>2</sub>H and HNC spectra are  $\sim 0.3$  for 3C111B, the highest in our sample. As discussed in the next section, we have obtained more sensitive HCO<sup>+</sup> and HCN absorption spectra in the directions of 3C111A and 3C111B with NOEMA. 3C123 is also a multiple-component continuum source. Murray et al. (2018) resolved 3C123 into two components, 3C123A and 3C123B. At ALMA's higher resolution, we resolve 3C123 into four components. Because we make a direct comparison between the 21-SPONGE H I absorption spectra and our molecular absorption spectra, though, we regrid our ALMA observations in the direction of 3C123 using the VLA beam (20''3 × 5''3) from the Murray et al. (2018) observations, before extracting the absorption spectra. As in Murray et al. (2018), we resolve only two components at this lower spatial resolution, which are here referred to as 3C123A and 3C123B to be consistent with their labeling. While the spectra toward 3C123A have a typical optical depth noise of 0.01, the optical depth noise in the 3C123B spectra is  $\sim 0.1$ .

For sources without significant absorption, we extracted the spectra from the continuum-subtracted data cubes output by the default ALMA Science Pipeline at the pixel of peak continuum flux. To calculate upper limits to the optical depth, we added the continuum flux back to the continuum-subtracted spectra (correcting for the spectral index; for a majority of sightlines, the

**Table 1**

The Molecular Line Transitions Covered by the ALMA-SPONGE Spectral Setup

Species	Transition	Frequency GHz	$N/\int\tau dv$ $\text{cm}^{-2}/\text{km s}^{-1}$
C <sub>2</sub> H	$N = 1 - 0 \ J = 3/2 - 1/2 \ F = 2 - 1$	87.3169	$4.34 \times 10^{13}$
C <sub>2</sub> H	$N = 1 - 0 \ J = 3/2 - 1/2 \ F = 1 - 0$	87.3286	
HCN	$J = 1 - 0 \ F = 1 - 1$	88.6304	$1.91 \times 10^{12}$
HCN	$J = 1 - 0 \ F = 2 - 1$	88.6318	
HCN	$J = 1 - 0 \ F = 0 - 1$	88.6339	
HCO <sup>+</sup>	$J = 1 - 0$	89.1890	$1.11 \times 10^{12}$
HNC	$J = 1 - 0$	90.6336	$1.78 \times 10^{12}$

**Note.** HCN and HCO<sup>+</sup> were also covered by the NOEMA-SPONGE spectral setup. Column 1 lists the molecular species. Column 2 lists each transition. Column 3 lists the rest frequency for each transition. Column 4 gives the conversion between the optical depth integral and the column density for each species (see Section 3.1); the values listed in column 4 account for all of the observed transitions for each species and are calculated assuming an excitation temperature equal to the cosmic microwave background (CMB) temperature, 2.725 K.

continuum is relatively flat). For sightlines that showed significant absorption, we reran the ALMA Science Pipeline without continuum subtraction and extracted spectra from the pixel of peak continuum flux to ensure the most reliable estimate of the continuum flux and the optical depths. Except in the cases noted above, at 0.4 km s<sup>-1</sup> velocity resolution, we reach a typical optical depth noise of  $\sim 0.01$  in the ALMA-SPONGE spectra.

Table 2 lists the Galactic coordinates and integrated optical depths in the directions of the 19 ALMA-SPONGE bright background radio continuum sources (excluding 3C111C). 10 sightlines show no molecular absorption at a level of  $3\sigma$ . Six sightlines show strong absorption across all four molecular species. In the direction of 3C111B, we detect HNC absorption, but no C<sub>2</sub>H absorption (HCO<sup>+</sup> and HCN are observed with NOEMA; see the next section). This most likely reflects the poor sensitivity in the direction of 3C111B—the peak C<sub>2</sub>H optical depth in the direction of 3C111A is 0.766, which is only 2.6 times the noise level in the 3C111B spectrum. In the direction of 3C78, we see weak HCO<sup>+</sup> absorption, but no HCN, C<sub>2</sub>H, or HNC absorption. In the direction of J2136, we see weak C<sub>2</sub>H absorption, but no HCN, HCO<sup>+</sup>, or HNC absorption. Although the detections in the directions of 3C123B, 3C78, and J2136 are marginal ( $\sim 3\sigma$ ), they appear significant at lower-velocity resolution. Moreover, the absorption in these directions is aligned in velocity with the H I absorption features. We also see two hyperfine components for J2136, and we know from the 3C123A spectra that molecular gas is present toward 3C123 at the velocities where we detect marginal absorption in the direction of 3C123B. The spatial distribution of all of the ALMA-SPONGE sources is shown in Figure 1, overlaid on a map of CO-integrated emission measured by the Planck Collaboration et al. (2014b).

## 2.2. Observations with NOEMA (NOEMA-SPONGE)

We observed HCO<sup>+</sup> and HCN in absorption at 62.5 kHz channel spacing (0.2 km s<sup>-1</sup> velocity spacing) with NOEMA (NOEMA-SPONGE, projects W19AQ and S20AB) in the directions of 3C111—both A and B components—and BL Lac.

3C111 was observed in 2020 January and BL Lac was observed 2020 October. The measured fluxes were 1.13 Jy for 3C111A, 0.06 Jy for 3C111B, and 2.10 Jy for BL Lac. Both HCO<sup>+</sup> and HCN were placed in high-resolution chunks in the lower sideband, which was tuned to 90 GHz. Standard calibration was carried out using the CLIC and MAPPING software, part of the GILDAS software collection<sup>9</sup> (Pety 2005; Gildas 2013). Because BL Lac and 3C111A were bright, we further performed self-calibration using the 2020 Self Calibration tool in MAPPING. We smooth the final spectra to 0.4 km s<sup>-1</sup> velocity resolution, reaching an optical depth sensitivity of  $\sim 0.002$  for 3C111A and BL Lac and an optical depth sensitivity of  $\sim 0.05$  for 3C111B. The NOEMA-SPONGE sensitivity in the direction of 3C111B is significantly better than the ALMA-SPONGE sensitivity; only the NOEMA-SPONGE HCO<sup>+</sup> and HCN data in this direction are considered hereafter. The HCO<sup>+</sup> and HCN optical depth integrals in these directions are listed in Table 2 (where the NOEMA-SPONGE fluxes are superscripted by a “1”) and their positions are shown in Figure 1.

## 2.3. Additional Data Sets

### 2.3.1. 21-SPONGE

The 21-SPONGE project (Murray et al. 2015, 2018) obtained H I absorption spectra using the VLA and H I emission spectra using the Arecibo Observatory in the directions of 57 bright background radio continuum sources, ranging in latitude from  $|b| = 3^\circ 7'$  to  $|b| = 81^\circ$ . The absorption spectra had an optical depth noise of  $\sim 10^{-3}$  at the 0.4 km s<sup>-1</sup> velocity resolution. Murray et al. (2015, 2018) decomposed the absorption and emission spectra into Gaussian components. They estimated the spin temperatures,  $T_S$ , and H I column densities,  $N(\text{H I})$ , of the H I structures seen in absorption. They also investigated the fraction of gas in the CNM, the WNM, and the UNM phases along each line of sight. We use their absorption spectra and adopt their Gaussian fits in this work. The 21-SPONGE H I absorption spectra are shown in Figure 2.


### 2.3.2. $E(B - V)$ and $T_d$ from Planck

The total hydrogen column density,  $N_{\text{H}} = N(\text{H I}) + 2N(\text{H}_2)$ , can be inferred from the interstellar reddening,  $E(B - V)$ , along the line of sight. We use the dust radiance,  $\mathcal{R}$ , measured by Planck to estimate  $E(B - V)$ .  $\mathcal{R}$  is a more reliable predictor of  $E(B - V)$  than the 353  $\mu\text{m}$  optical depth,  $\tau_{353}$ , at high Galactic latitude (Planck Collaboration et al. 2014a).<sup>10</sup> We extract  $E(B - V)$  at the pixel nearest to each background source using the dust maps Python package (Green 2018). The Planck dust maps have a resolution of  $5'$  and a typical fractional uncertainty of  $\sim 10\%$  in  $E(B - V)$ . The results are listed in Table 2. We also extract the dust temperature,  $T_d$ , derived by the Planck Collaboration et al. (2016) using the generalized needlet internal linear combination method, and a modified blackbody spectral model on Planck temperature maps at 353, 545, 857, and 3000 GHz. We use dust temperature as a proxy for the strength of the interstellar radiation field (ISRF)—the strength of the far-UV (FUV) radiation field scales as  $T_d^{\beta+4}$ , where  $\beta$  is the power-law index characterizing the dust emissivity cross section as a function of frequency (see Equation (7)).<sup>15</sup> of

<sup>9</sup> <https://www.iram.fr/IRAMFR/GILDAS>

<sup>10</sup> We find only negligible differences between the  $E(B - V)$  estimates made from  $\mathcal{R}$  and  $\tau_{353}$  for the sightlines in this work.






**Figure 1.** The distribution of the background radio continuum sources observed in this work plotted over a full-sky map of the CO-integrated intensity from the Planck Collaboration et al. (2014b). Sources with molecular absorption detections at a level  $\geq 3\sigma$  are shown as filled stars; sources with no molecular absorption detections are shown as filled circles. The integrated intensity of the CO emission is shown with a logarithmic color scale, ranging from 1 K km s $^{-1}$  to 35 K km s $^{-1}$ . The different components of the multiple-component continuum sources 3C111 and 3C123 are indistinguishable at this resolution, so are not labeled separately (see Table 2). The dashed lines show Galactic latitude and longitude with 30° spacing.

**Table 2**  
Observed Properties of the Sightlines in This Study

Source	$F_{90}$ Jy	$l/b$ °	$N(H\text{ I})$ $10^{20}\text{ cm}^{-2}$	$E(B - V)$ mag	$\int \tau_{\text{HNC}} dv$ km s $^{-1}$	$\int \tau_{\text{C}_2\text{H}} dv$ km s $^{-1}$	$\int \tau_{\text{HCO}^+} dv$ km s $^{-1}$	$\int \tau_{\text{HNC}} dv$ km s $^{-1}$
3C154	0.390	185.6/4.0	37.12	0.425	$3.325 \pm 0.039$	$1.079 \pm 0.025$	$3.059 \pm 0.056$	$0.819 \pm 0.023$
3C111A	0.769/1.145 <sup>a</sup>	161.7/−8.8	25.96	0.827	$>12.710$	$2.057 \pm 0.033$	$>11.054$	$3.606 \pm 0.129$
3C111B	0.034/0.060 <sup>a</sup>	161.7/−8.8	24.86	0.827	$>11.345$	...	$>6.780$	$>2.028$
BLLac	2.189	92.6/−10.4	18.68	0.350	$4.316 \pm 0.007$	...	$2.546 \pm 0.002$	...
3C138	0.470	187.4/−11.3	20.51	0.263	...	...	...	...
3C123A	0.451	170.6/−11.7	17.33	0.499	$6.029 \pm 0.171$	$1.650 \pm 0.041$	$6.280 \pm 0.234$	$1.100 \pm 0.041$
3C123B	0.270	170.6/−11.7	18.66	0.499	$>4.287$	$0.331 \pm 0.189$	$>4.071$	$1.222 \pm 0.965$
PKS0742	0.304	209.8/16.6	3.37	0.031	...	...	...	...
3C120	2.431	190.4/−27.4	13.33	0.275	$0.031 \pm 0.003$	$0.178 \pm 0.006$	$0.283 \pm 0.006$	$0.006 \pm 0.002$
J2136	2.063	55.5/−35.6	4.34	0.080	...	$0.012 \pm 0.002$	...	...
3C346	0.107	35.3/35.8	5.42	0.082	...	...	...	...
3C454.3	11.838	86.1/−38.2	7.03	0.117	$0.112 \pm 0.004$	$0.106 \pm 0.004$	$0.276 \pm 0.004$	$0.024 \pm 0.002$
3C345	2.489	63.5/40.9	0.75	0.021	...	...	...	...
4C15.05	0.596	147.9/−44.0	4.06	0.057	...	...	...	...
3C78	0.347	174.9/−44.5	10.65	0.162	...	...	$0.043 \pm 0.012$	...
1055 + 018	5.283	251.5/52.8	3.05	0.037	...	...	...	...
3C273	8.668	289.9/64.4	2.16	0.028	...	...	...	...
4C12.50	0.507	347.2/70.2	2.18	0.054	...	...	...	...
3C286	0.671	56.5/80.7	1.21	0.016	...	...	...	...
4C32.44	0.229	67.2/81.0	1.36	0.021	...	...	...	...

**Note.** Column 1: the name of the background radio continuum source. Column 2: the 90 GHz continuum flux. Column 3: the Galactic longitude and latitude of the background source in degrees. Column 4: the total H I column density measured by 21-SPONGE along the line of sight (Murray et al. 2018). Column 5:  $E(B - V)$  inferred from the dust radiance measured by Planck (Planck Collaboration et al. 2014). Columns 6–9: the integrated optical depth for all four species for channels with  $\tau > 3\sigma$ . Lower limits are indicated. As described in Section 2, BL Lac was not observed in C<sub>2</sub>H or HNC absorption.

<sup>a</sup> As described in Section 2, 3C111A and 3C111B were observed by both ALMA (C<sub>2</sub>H, HNC) and NOEMA (HCN, HCO<sup>+</sup>); both fluxes are listed, with the flux for the NOEMA data superscripted by a “1.”



**Figure 2.** The optical depth spectra,  $e^{-\tau}$ , vs. LSR velocity for H I (bottom line, black; Murray et al. 2018),  $\text{HCO}^+$  (second line from bottom, red), HCN (middle line, blue), HNC (second line from top, gray), and  $\text{C}_2\text{H}$  (top line, green). HCN velocities correspond to the  $F = 2 - 1$  transition.  $\text{C}_2\text{H}$  velocities correspond to the  $F = 2 - 1$  transition. The velocity of the  $F = 1 - 0$   $\text{C}_2\text{H}$  transition has been shifted by  $+25 \text{ km s}^{-1}$  for viewing convenience. Gaussian fits to the H I spectra from 21-SPONGE (Murray et al. 2018) and Gaussian fits to the molecular absorption spectra from this work (see Section 3.2) are shown as dashed lines. Absorption spectra that have been amplified are labeled with the multiplication factor.

Lequeux (2005)). To estimate the ISRF, we also extract the corresponding  $\beta$  values at each of our positions.

In several places in this paper, we use values of  $E(B - V)$ ,  $T_d$ , and the ISRF derived from Planck as estimates of the gas properties along the lines of sight. While this provides a uniform approach, because Planck’s resolution is much lower than that of our H I and molecular pencil-beam absorption spectra, we consider these only as rough estimates and not precise measures of the local physical conditions.

#### 2.4. CO Emission Spectra from the CfA 1.2 m Telescope

Dame et al. (2001) obtained CO emission spectra across the entire Galactic plane and select clouds at higher latitudes

using the CfA 1.2 m telescope with an  $8''.4$  beamwidth. For the sources covered by this CO survey, we extract CO emission spectra from the nearest pixel in the Dame et al. (2001) maps, sampled between every other beamwidth ( $0''.25$ ) and every half-beamwidth ( $0''.0625$ ) in the directions of our sources. The brightness temperature noise in these spectra ranges from 0.15 K to 0.30 K. For sources outside the bounds covered by Dame et al. (2001), we use unpublished spectra obtained with the CfA 1.2 m telescope at  $8''.4$  beamwidth and  $0''.25$  sampling. The brightness temperature noise in these unpublished spectra is uniform, 0.18 K. All of the CO emission spectra taken from Dame et al. (2001) have a velocity resolution of  $0.65 \text{ km s}^{-1}$ .




Figure 2. (Continued.)

### 3. Methods

#### 3.1. Deriving Column Densities

For a ground-state transition, the column density,  $N$ , and optical depth integral,  $\int \tau dv$ , for a particular species are related by

$$N = Q(T_{\text{ex}}) \frac{8\pi\nu^3}{c^3} \frac{1}{g_u A_{ul}} [1 - \exp(-h\nu/kT_{\text{ex}})]^{-1} \int \tau dv, \quad (1)$$

where  $\nu$  is the frequency of radiation resulting from the transition from the upper state  $u$  to the lower state  $l$ ,  $g_u$  is the degeneracy of the upper state,  $A_{ul}$  is the Einstein  $A$  coefficient for the transition,  $T_{\text{ex}}$  is the excitation temperature, and  $Q(T_{\text{ex}})$  is the partition function. We calculate  $N/\int \tau dv$  using the constants given in the Cologne Database for Molecular Spectroscopy (Müller et al. 2001; Endres et al. 2016) and the Leiden Atomic and Molecular Database (Schöier et al. 2010). For  $\text{C}_2\text{H}$ , we use  $N/\int \tau dv = 1.6 \times 2.71 \times 10^{13} \text{ cm}^{-2}/\text{km s}^{-1}$ , provided by Lucas & Liszt (2000), where the factor of 1.6 accounts for the fact that we are measuring two of the six satellite  $\text{C}_2\text{H}$  lines. All values of  $N/\int \tau dv$  are calculated assuming an excitation temperature that is equal to the temperature of the CMB, 2.725 K (e.g., Godard et al. 2010; Luo et al. 2020).<sup>11</sup> All of the conversions are listed in Table 1; the  $\text{C}_2\text{H}$  and HCN conversions account for all of the hyperfine transitions listed in Table 1.

For saturated channels where the measured value of  $I(v) \leq 0$ —and therefore  $\tau(v)$  is infinite—we define the optical depth lower limit to be

$$\tau_l(v) = -\ln\left(\frac{2\sigma_I}{I_0}\right), \quad (2)$$

where  $\sigma_I$  is the noise in the spectrum of specific intensity. We then calculate a lower limit to the column density (Equation (1)) using  $\tau(v) = \tau_l(v)$  for the saturated channels. We note that this systematically underestimates the optical depth for spectra with saturated absorption. In the analysis that follows, we therefore consider the optical depths and column densities only for spectra where the absorption is not saturated, unless otherwise noted. The optical depth integrals for saturated absorption spectra in Table 2 should be considered as conservative lower limits.

In Section 4, we estimate the total hydrogen column density,  $N_{\text{H}}$ , from the reddening,  $E(B - V)$ , according to Zhu et al. (2017):

$$N_{\text{H}} = 2.08 \times 10^{21} \times 3.1E(B - V) \text{ cm}^{-2} \text{ mag}^{-1}, \quad (3)$$

where  $3.1E(B - V)$  is here used for the visual extinction,  $A_V$ . This estimate was derived from a compilation of Galactic sources with optical and X-ray observations, and is consistent with other previous estimates (Jenkins & Savage 1974; Bohlin et al. 1978; Krumholz et al. 2009; Rachford et al. 2009a) for sources spanning a wide range of Galactic latitudes and  $N_{\text{H}}$ . Nevertheless, the ratio of  $N_{\text{H}}$  to  $E(B - V)$  is not universal, and

contributes additional uncertainty to the determination of  $N_{\text{H}}$ —see Section 6 for discussion.

#### 3.2. Gaussian Decomposition

We decompose the molecular optical depth spectra into Gaussian functions:

$$\begin{aligned} \tau(v) &= \sum_{i=1}^M \tau_i(v) \\ &= \sum_{i=1}^M \tau_{0,i} e^{-4 \ln 2 (v - v_{0,i})^2 / \Delta v_{0,i}^2}, \end{aligned} \quad (4)$$

where  $\tau_{0,i}$ ,  $v_{0,i}$ , and  $\Delta v_{0,i}$  are the peak optical depth, the central velocity, and the FWHM of the  $i$ th Gaussian component, respectively, and there are  $M$  Gaussian functions fit to the optical depth spectrum. Lucas, Liszt, Gerin, and collaborators have identified dozens of molecular absorption components common across various molecular species, including HCN,  $\text{C}_2\text{H}$ ,  $\text{HCO}^+$ , and HNC. Further, they have found that the central velocities of the absorption components identified at these transitions are nearly identical, and that FWHMs vary across different species by only  $\sim 10\%$ – $30\%$ . Therefore, for each line of sight, we find the best solution to Equation (4) for the  $\text{C}_2\text{H}$ , HCN,  $\text{HCO}^+$ , and HNC optical depth spectra using Python's `scipy.optimize.curve_fit` from sets of initial guesses with the same number of components (modulo an integer factor accounting for the number of hyperfine transitions). The initial guesses for the FWHM and central velocity of each component are the same for all four optical depth spectra. The initial guesses for the central velocities of the different hyperfine  $\text{C}_2\text{H}$  and HCN components are offset according to the known frequency separation of the transitions, and the initial guesses for the optical depths are scaled to the LTE ratios (2:1 and 5:3:1, respectively). For 3C111B and 3C123B, we use the same initial guesses at 3C111A and 3C123A, where the spectra are more sensitive. When fitting, we allow the peak optical depths to vary freely, the FWHMs to vary by a factor of 2 from the initial guess, and the central velocity to vary by  $\pm 1 \text{ km s}^{-1}$  from the initial guess. Only solutions with peak optical depths  $> 3\sigma_\tau$  are classified as detections.<sup>12</sup> For  $\text{C}_2\text{H}$  and HCN, we require only that the strongest transition be detected at a level of  $3\sigma$ .

The Gaussian-fitted components to the absorption spectra are shown in Figure 2. Tables 3, 4, 5, and 6 list the fits for  $\text{C}_2\text{H}$ , HCN,  $\text{HCO}^+$ , and HNC, respectively. Peak optical depths range between 0.008 and 2.5, excluding saturated lines with optical depths  $\gtrsim 3.5$ . FWHMs range between  $0.5 \text{ km s}^{-1}$  and  $3.4 \text{ km s}^{-1}$ . For 10 sightlines, no features are detected (see Table 2). There is one feature identified in  $\text{C}_2\text{H}$  absorption in the direction of J2136 for which there is no corresponding HCN,  $\text{HCO}^+$ , or HNC absorption; and there is one feature identified in  $\text{HCO}^+$  absorption in the direction of 3C78 for which there is no corresponding HCN,  $\text{C}_2\text{H}$ , or HNC absorption. Two features are seen in  $\text{C}_2\text{H}$  and  $\text{HCO}^+$  absorption, but not seen in HNC or HCN absorption. Three absorption features identified in  $\text{C}_2\text{H}$ ,  $\text{HCO}^+$ , and HCN are not detected in HNC.

<sup>11</sup> In Paper II, we investigate this assumption and show that the  $\text{HCO}^+$  ( $1 - 0$ ) excitation temperature begins to rise above the CMB temperature when  $n \gtrsim 300 \text{ cm}^{-3}$ .

<sup>12</sup> HNC lines in the direction of 3C111B are detected at  $< 3\sigma$ , but are included for comparison to 3C111A.



**Table 3**  
Gaussian Fits to the C<sub>2</sub>H Absorption Spectra

Source	$\tau_0$	$\Delta v_0$ km s <sup>-1</sup>	$v_0$ km s <sup>-1</sup>	$\int \tau_0 dv$ km s <sup>-1</sup>
3C111A	0.752 ± 0.036	1.0 ± 0.0	-0.9 ± 0.0	0.78 ± 0.05
3C111A	0.332 ± 0.021	1.5 ± 0.1	-2.5 ± 0.0	0.52 ± 0.05
3C120	0.069 ± 0.001	1.3 ± 0.0	6.6 ± 0.0	0.09 ± 0.01
3C120	0.016 ± 0.002	0.9 ± 0.2	5.1 ± 0.1	0.02 ± 0.01
3C120	0.014 ± 0.001	2.1 ± 0.2	9.3 ± 0.1	0.03 ± 0.01
3C120	0.007 ± 0.002	0.6 ± 0.2	3.7 ± 0.2	0.00 ± 0.01
3C123A	0.372 ± 0.095	1.2 ± 0.5	4.3 ± 0.3	0.49 ± 0.25
3C123A	0.313 ± 0.104	1.4 ± 0.6	5.3 ± 0.3	0.45 ± 0.26
3C123A	0.276 ± 0.070	0.7 ± 0.3	3.1 ± 0.2	0.20 ± 0.11
3C123B	0.584 ± 0.140	0.9 ± 0.6	4.2 ± 0.6	0.59 ± 0.37
3C123B	0.437 ± 0.081	1.1 ± 0.6	5.4 ± 0.8	0.51 ± 0.31
3C123B	0.351 ± 0.072	0.8 ± 0.5	3.0 ± 0.7	0.31 ± 0.18
3C154	0.449 ± 0.050	0.8 ± 0.2	-2.0 ± 0.1	0.38 ± 0.07
3C154	0.221 ± 0.042	1.0 ± 0.3	-0.9 ± 0.1	0.23 ± 0.12
3C154	0.087 ± 0.021	2.3 ± 0.6	-4.0 ± 0.3	0.21 ± 0.10
3C154	0.050 ± 0.015	2.5 ± 1.0	2.7 ± 0.8	0.14 ± 0.06
3C454.3	0.046 ± 0.002	1.2 ± 0.0	-9.2 ± 0.0	0.06 ± 0.01
3C454.3	0.021 ± 0.001	1.2 ± 0.2	-10.5 ± 0.1	0.03 ± 0.01
J2136	0.008 ± 0.001	2.4 ± 0.0	5.3 ± 0.1	0.02 ± 0.01

**Note.** Column 1: source name. Column 2: peak optical depth,  $\tau_0$ . Column 3: FWHM,  $\Delta v_0$ . Column 4: central velocity,  $v_0$ . Column 5: integrated optical depth.

**Table 4**  
Gaussian Fits to the HCN Absorption Spectra

Source	$\tau_0$	$\Delta v_0$ km s <sup>-1</sup>	$v_0$ km s <sup>-1</sup>	$\int \tau_0 dv$ km s <sup>-1</sup>
3C111A	3.639 <sup>†</sup> ± 0.039	1.0 ± 0.0	-0.9 ± 0.0	3.78 <sup>†</sup> ± 0.05
3C111A	1.959 <sup>†</sup> ± 0.009	0.9 ± 0.0	-2.5 ± 0.0	1.87 <sup>†</sup> ± 0.01
3C111B	2.884 <sup>†</sup> ± 1.381	1.0 ± 0.6	-1.0 ± 0.1	3.20 <sup>†</sup> ± 2.49
3C111B	1.678 <sup>†</sup> ± 0.787	1.1 ± 0.6	-2.5 ± 0.1	1.89 <sup>†</sup> ± 1.42
3C120	0.025 ± 0.003	0.7 ± 0.1	6.4 ± 0.0	0.02 ± 0.00
3C120	0.008 ± 0.002	1.2 ± 0.1	5.4 ± 0.2	0.01 ± 0.00
3C123A	2.594 ± 0.826	0.7 ± 0.2	4.4 ± 0.1	1.89 ± 0.78
3C123A	0.926 ± 0.201	1.0 ± 0.3	5.2 ± 0.2	1.01 ± 0.39
3C123A	0.587 ± 0.125	0.9 ± 0.4	3.4 ± 0.2	0.58 ± 0.26
3C123B	2.753 <sup>†</sup> ± 0.948	0.7 ± 0.2	4.2 ± 0.5	1.94 <sup>†</sup> ± 0.82
3C123B	1.265 <sup>†</sup> ± 0.383	0.8 ± 0.4	5.2 ± 0.7	1.12 <sup>†</sup> ± 0.69
3C123B	0.622 <sup>†</sup> ± 0.195	0.7 ± 0.4	3.1 ± 0.7	0.47 <sup>†</sup> ± 0.33
3C154	1.734 ± 0.182	0.7 ± 0.1	-2.0 ± 0.0	1.31 ± 0.17
3C154	0.394 ± 0.046	1.0 ± 0.2	-1.0 ± 0.1	0.41 ± 0.09
3C154	0.056 ± 0.016	2.0 ± 0.8	-4.5 ± 0.7	0.12 ± 0.06
3C154	0.063 ± 0.017	2.3 ± 1.0	2.9 ± 0.7	0.16 ± 0.08
3C454.3	0.068 ± 0.000	1.0 ± 0.0	-9.3 ± 0.0	0.07 ± 0.00
3C454.3	0.010 ± 0.000	0.4 ± 0.0	-10.5 ± 0.0	0.00 ± 0.00
BLLac	0.923 ± 0.053	0.8 ± 0.1	-1.4 ± 0.0	0.83 ± 0.07
BLLac	0.410 ± 0.043	0.9 ± 0.1	-0.6 ± 0.0	0.41 ± 0.07

**Note.** Column 1: source name. Column 2: peak optical depth,  $\tau_0$ . Column 3: FWHM,  $\Delta v_0$ . Column 4: central velocity,  $v_0$ . Column 5: integrated optical depth. Components for sightlines with saturated absorption are superscripted by a dagger (†).

#### 4. Molecular Gas in the Diffuse ISM

The distribution of these background sources across the sky is shown in Figure 1, overlaid on the CO-integrated intensity image from the Planck Collaboration et al. (2014b). We show our ALMA/NOEMA observations of HCN, C<sub>2</sub>H, HCO<sup>+</sup>, and HNC, as well as H I absorption from 21-SPONGE, in Figure 2.

**Table 5**  
Gaussian Fits to the HCO<sup>+</sup> Absorption Spectra

Source	$\tau_0$	$\Delta v_0$ km s <sup>-1</sup>	$v_0$ km s <sup>-1</sup>	$\int \tau_0 dv$ km s <sup>-1</sup>
3C111A	4.172 <sup>†</sup> ± 0.074	1.3 ± 0.0	-0.9 ± 0.0	5.94 <sup>†</sup> ± 0.14
3C111A	3.296 <sup>†</sup> ± 0.043	1.3 ± 0.0	-2.5 ± 0.0	4.72 <sup>†</sup> ± 0.08
3C111B	2.393 <sup>†</sup> ± 0.918	1.4 ± 0.6	-1.0 ± 0.1	3.54 <sup>†</sup> ± 2.12
3C111B	1.749 <sup>†</sup> ± 0.775	1.6 ± 0.8	-2.5 ± 0.1	2.94 <sup>†</sup> ± 1.95
3C120	0.154 ± 0.002	1.3 ± 0.0	6.2 ± 0.0	0.21 ± 0.00
3C120	0.082 ± 0.013	0.4 ± 0.1	5.3 ± 0.0	0.03 ± 0.01
3C120	0.020 ± 0.001	2.4 ± 0.0	8.6 ± 0.1	0.05 ± 0.00
3C120	0.011 ± 0.001	1.2 ± 0.2	3.7 ± 0.1	0.01 ± 0.00
3C123A	3.214 ± 1.100	0.8 ± 0.3	4.3 ± 0.2	2.66 ± 1.49
3C123A	2.123 ± 0.469	1.0 ± 0.3	5.3 ± 0.2	2.31 ± 0.93
3C123A	1.138 ± 0.261	1.2 ± 0.3	3.4 ± 0.2	1.40 ± 0.53
3C123B	3.737 <sup>†</sup> ± 1.137	1.0 ± 0.5	4.4 ± 0.7	3.85 <sup>†</sup> ± 2.36
3C123B	2.110 <sup>†</sup> ± 0.689	0.7 ± 0.3	5.2 ± 0.6	1.57 <sup>†</sup> ± 0.86
3C123B	1.203 <sup>†</sup> ± 0.331	0.6 ± 0.4	3.0 ± 0.5	0.82 <sup>†</sup> ± 0.53
3C154	1.709 ± 0.195	0.7 ± 0.1	-2.2 ± 0.1	1.29 ± 0.16
3C154	0.847 ± 0.078	1.0 ± 0.2	-1.1 ± 0.1	0.92 ± 0.11
3C154	0.139 ± 0.016	2.5 ± 0.3	-4.1 ± 0.3	0.37 ± 0.07
3C154	0.147 ± 0.008	3.1 ± 0.3	2.2 ± 0.2	0.49 ± 0.04
3C454.3	0.212 ± 0.000	1.0 ± 0.0	-9.4 ± 0.0	0.22 ± 0.00
3C454.3	0.055 ± 0.000	0.8 ± 0.0	-10.7 ± 0.0	0.05 ± 0.00
3C78	0.039 ± 0.012	1.3 ± 0.7	-7.9 ± 0.5	0.05 ± 0.03
BLLac	0.898 ± 0.277	1.3 ± 0.4	-1.4 ± 0.0	1.22 ± 0.54
BLLac	0.747 ± 0.018	1.3 ± 0.2	-0.5 ± 0.1	1.07 ± 0.16


**Note.** Column 1: source name. Column 2: peak optical depth,  $\tau_0$ . Column 3: FWHM,  $\Delta v_0$ . Column 4: central velocity,  $v_0$ . Column 5: integrated optical depth. Components for sightlines with saturated absorption are superscripted by a dagger (†).

**Table 6**  
Gaussian Fits to the HNC Absorption Spectra

Source	$\tau_0$	$\Delta v_0$ km s <sup>-1</sup>	$v_0$ km s <sup>-1</sup>	$\int \tau_0 dv$ km s <sup>-1</sup>
3C111A	3.039 ± 0.303	0.9 ± 0.1	-0.9 ± 0.0	2.84 ± 0.33
3C111A	0.675 ± 0.031	1.0 ± 0.0	-2.5 ± 0.0	0.69 ± 0.04
3C111B	1.825 ± 1.734 <sup>†</sup>	0.81 ± 0.55	-1.15 ± 0.38	1.57 ± 1.83 <sup>†</sup>
3C111B	0.493 ± 0.560 <sup>†</sup>	0.82 ± 0.99	-2.71 ± 0.50	0.43 ± 0.49 <sup>†</sup>
3C120	0.018 ± 0.003	0.5 ± 0.1	6.7 ± 0.0	0.01 ± 0.00
3C123A	0.775 ± 0.159	0.8 ± 0.2	4.4 ± 0.1	0.62 ± 0.18
3C123A	0.311 ± 0.089	1.2 ± 0.7	5.1 ± 0.3	0.38 ± 0.24
3C123A	0.183 ± 0.051	0.8 ± 0.4	3.2 ± 0.4	0.16 ± 0.10
3C123B	0.989 ± 0.235	1.1 ± 0.6	4.2 ± 0.6	1.14 ± 0.72
3C123B	0.396 ± 0.094	1.1 ± 0.7	5.2 ± 0.7	0.47 ± 0.30
3C123B	0.214 ± 0.038	0.9 ± 0.5	3.1 ± 0.7	0.21 ± 0.11
3C154	0.825 ± 0.064	0.8 ± 0.1	-2.2 ± 0.0	0.68 ± 0.09
3C154	0.194 ± 0.039	0.7 ± 0.3	-1.2 ± 0.2	0.15 ± 0.07
3C454.3	0.018 ± 0.001	0.9 ± 0.0	-9.4 ± 0.0	0.02 ± 0.00
3C454.3	0.006 ± 0.001	1.4 ± 0.1	-10.0 ± 0.1	0.01 ± 0.00

**Note.** Column 1: source name. Column 2: peak optical depth,  $\tau_0$ . Column 3: FWHM,  $\Delta v_0$ . Column 4: central velocity,  $v_0$ . Column 5: integrated optical depth. Components for sightlines with saturated absorption are superscripted by a dagger (†).

Table 2 summarizes the observational results in these directions, including the H I column density (Murray et al. 2018), reddening (Planck Collaboration et al. 2014a), and molecular optical depth integrals, calculated for channels with an optical depth  $\tau(\nu) \geq 3\sigma_\tau$ . Sources are listed in order of increasing Galactic latitude.



**Figure 3.** Molecular column densities vs. visual extinction ( $A_V$ ; upper left panel), H I CNM column density (upper right panel), H I UNM column density (lower left panel), and the combined column density of the CNM and UNM (lower right panel). The colors indicate the dust temperature derived by the Planck Collaboration et al. (2016). Because the resolution of the Planck extinction maps is  $5'$ , the A and B components of 3C111 and 3C123 have the same  $x$  coordinate in the upper left panel, so are not labeled separately.

For most sources at latitudes  $\lesssim 40^\circ$ , we detect at least one of four transitions. For sources at latitude  $>40^\circ$ , we detect marginal  $\text{HCO}^+$  absorption along only one sightline, and no molecular absorption in any other direction. This is evident in Figure 1, where sources with molecular absorption detections are indicated using stars, while sources with no molecular detections are indicated using circles. There is no significant CO emission observed in the Planck Collaboration et al. (2014a) or Dame et al. (2001) data in the directions of three sightlines with molecular absorption detections—3C120, 3C78, and J2136. These sources have  $|b| = 27^\circ$ – $44^\circ$ . We discuss this further in Section 4.2.

The integrated optical depths—and therefore column densities (Equation (1))—span over two orders of magnitude across our sample for each of the molecular species, indicating that we are probing diverse interstellar environments. In Section 7, we compare the results in Table 2 to previous observations of HCN,  $\text{C}_2\text{H}$ ,  $\text{HCO}^+$ , and HNC in the same directions, taken between 3 and 25 yr ago. All variations are at a level  $\lesssim 3\sigma$ , demonstrating that our observing and processing strategies are sound.

#### 4.1. Thresholds for Molecule Formation

In Figure 3, we compare the observed molecular column densities to the neutral and total gas column densities. The total gas column density is traced via the visual extinction ( $A_V = 3.1E(B - V)$ ; upper left panel). For the H I gas, we investigate the CNM column density (the H I column density occupied by gas with  $T_S < 250$  K; upper right panel), the UNM column density, (the H I column density occupied by gas with

$250 \text{ K} < T_S < 1000$  K; lower left panel), and the sum of the CNM and UNM column densities (lower right panel) for each line of sight. For nondetections, we show  $3\sigma$  upper limits to the molecular column densities assuming an FWHM of three channels or  $1.2 \text{ km s}^{-1}$ . The data points are color-coded, based on the dust temperature provided by Planck, which can be used as a proxy for the ISRF.

The upper left panel of Figure 3 suggests a threshold visual extinction  $A_V \approx 0.25$ , below which no molecular absorption is detected and above which molecular abundances tend to increase with increasing  $A_V$ . A visual extinction  $A_V \gtrsim 0.25$  corresponds to a total hydrogen column density  $N_{\text{H}} \gtrsim 5 \times 10^{20} \text{ cm}^{-2}$  (e.g., Güver & Özel 2009; Zhu et al. 2017), which is roughly the threshold column density for the H I-to- $\text{H}_2$  transition at solar metallicity (e.g., Spitzer et al. 1973; Savage et al. 1977; Krumholz et al. 2009; Gong et al. 2018). Lucas & Liszt (1996) identified a similar threshold from observations of  $\text{HCO}^+$  (as well as OH), and Lucas & Liszt (2000) showed that  $\text{C}_2\text{H}$  was as widespread as  $\text{HCO}^+$ . These results suggest that a similar column density threshold applies in the cases of  $\text{HCO}^+$ , HCN, HNC, and  $\text{C}_2\text{H}$ . Essentially, as soon as conditions are suitable for  $\text{H}_2$  survival, these other species follow quickly. The sightline toward 3C138 is an exception—it has a visual extinction of 0.815, but no detectable molecular absorption. This sightline is explored in greater depth in Section 8.5. The  $A_V$  trend shown in this figure also follows the dust temperature distribution, again highlighting the importance of shielding against the radiation field. The five sources with the highest molecular column densities ( $\gtrsim 10^{12} \text{ cm}^{-2}$  for all four species) have the lowest dust temperature,  $T_d \sim 17$  K, and FUV radiation field  $G' \sim 0.4$ , where  $G'$  is the radiation field strength

relative to the standard Draine (1978) field, calculated using  $T_d$  and  $\beta$  from Planck (see Section 2.3.2). The rest of the sources have molecular column densities  $\lesssim 10^{12} \text{ cm}^{-2}$ ,  $T_d \sim 19 \text{ K}$ , and have  $G' \sim 1$ .

The lower left panel of Figure 3 shows the molecular column densities as a function of the column density of the thermally unstable H I. While this graph lacks a correlation, there is a clear dichotomy of sources: the sources with the highest molecular column densities ( $>10^{12} \text{ cm}^{-2}$  for all species) all have  $N(\text{H I})_{\text{UNM}} \sim 10^{20-21} \text{ cm}^{-2}$ , and the fraction of H I in the UNM is  $\sim 30\%$ – $70\%$  for these sightlines; all other sources have essentially no UNM gas. The only exception is 3C78, which has a high UNM column density and low molecular column densities. This suggests that UNM gas likely plays an important role for molecule formation and survival.

Meanwhile, in the two right panels of Figure 3, we see that there is a threshold of  $\gtrsim 10^{20} \text{ cm}^{-2}$  for the column density of cold H I, below which no molecular absorption is detected (J2136—the only exception—is the lowest-column density source to show molecular absorption, with  $N(\text{H I}) \approx 4 \times 10^{19} \text{ cm}^{-2}$  and very weak  $\text{C}_2\text{H}$  absorption). Again, this threshold is similar to the column density threshold for the H I-to- $\text{H}_2$  transition ( $\sim 5 \times 10^{20} \text{ cm}^{-2}$  Savage et al. 1977). The correlation seen in the lower right plot, where the  $x$ -axis traces the total amount of cold H I, is qualitatively similar to that seen in the upper left plot, and again reflects the importance of shielding.

We do not see any difference here across the different chemical species probed in this study. For example, Godard et al. (2014) suggested that  $\text{HCO}^+$  and  $\text{C}_2\text{H}$  are more sensitive to turbulent dissipation than HCN or HNC. We note, though, that our sample size is modest, and very few structures do not show absorption from all four species.

#### 4.2. A Broad, Faint Component of $\text{HCO}^+$ Absorption

In addition to the narrow, strong absorption features easily identifiable in the spectra shown in Figure 2, we also detect a broad, faint component of  $\text{HCO}^+$  absorption in a majority of sightlines. Typical  $\text{HCO}^+$  optical depths are between 0.01 and 0.1 for the broad component, while narrow absorption lines have typical optical depths  $>0.1$ . In the directions of 3C111A, 3C120, and 3C123A, we detect  $\text{HCO}^+$  absorption across nearly all velocities where 21-SPONGE detected H I absorption (Figure 4). The  $\text{HCO}^+$  absorption in the directions of 3C154 and BL Lac spans  $\gtrsim 10 \text{ km s}^{-1}$ , covering more than half of the velocity range where H I absorption is observed. In the direction of 3C454.3, the  $\text{HCO}^+$  absorption is localized mostly to a narrow range of velocities around  $-10 \text{ km s}^{-1}$ , but there are marginal detections at  $v \approx -30 \text{ km s}^{-1}$  and  $v \approx -15 \text{ km s}^{-1}$ , where H I absorption is also observed. In the direction of 3C78,  $\text{HCO}^+$  absorption is weak, and confined only to a narrow velocity range compared to the H I absorption. Some of these broad features are fit with Gaussians, while some features are only apparent in the residuals (since they have peak optical depths  $\lesssim 0.05$  and widths  $\gtrsim 3 \text{ km s}^{-1}$ ). We do not consider 3C111B or 3C123B here because of the relatively poor optical depth sensitivity in these directions.

##### 4.2.1. Physical Properties of the Broad $\text{HCO}^+$ Absorption

Most  $\text{HCO}^+$  absorption features detected in the literature have narrow linewidths, except in a handful of more sensitive,

recent studies, e.g., Liszt & Lucas (2000) and Liszt & Gerin (2018). To characterize the physical properties of the broad absorption features, we divide the spectra into velocity segments of no  $\text{HCO}^+$  absorption ( $\tau \lesssim 0.01$ ), weak  $\text{HCO}^+$  absorption ( $0.01 \lesssim \tau \lesssim 0.05$ ), and strong  $\text{HCO}^+$  absorption ( $\tau \gtrsim 0.05$ ). These spectral regions are shown in crosshatched gray, diagonally hatched green, and unhatched blue, respectively, in Figure 4.

For each velocity region, we calculate the H I optical depth-weighted mean spin temperature,

$$\langle T_s \rangle = \frac{\int \tau(v) T_B(v) / (1 - e^{-\tau(v)}) dv}{\int \tau(v) dv}. \quad (5)$$

While the definition of  $\langle T_s \rangle$  for selected velocity intervals is subjective, as it depends on where the velocity limits are set, it is useful for comparing atomic and molecular properties within the same velocity intervals (see the discussion below and Liszt & Lucas (2000)).


We also integrate the  $\text{HCO}^+$  optical depth and estimate the H I and  $\text{H}_2$  column densities for each region. We use the isothermal approximation to H I column density,  $N(\text{H I}) = 1.823 \times 10^{18} \text{ cm}^{-2} / (\text{km s}^{-1} \text{ K}) \int \tau(v) T_B / (1 - e^{-\tau(v)}) dv$ . For the  $\text{H}_2$  column density, we assume that  $N(\text{HCO}^+) = 2 \times 10^{-9} N(\text{H}_2)$  for all sightlines (Liszt & Lucas 2000).<sup>13</sup>

Figure 5 shows the integrated  $\text{HCO}^+$  optical depth versus  $\langle T_s \rangle$  in each spectral region. For regions with strong, narrow  $\text{HCO}^+$  absorption, we use blue circles. For regions with broad, weak  $\text{HCO}^+$  absorption, we use green diamonds. Regions with no  $\text{HCO}^+$  absorption are shown as upper limits. Points are sized according to the molecular fraction,  $f_{\text{mol}} = 2N(\text{H}_2)/N_{\text{H}}$ .

Figure 5 shows an anticorrelation between the integrated optical depth of  $\text{HCO}^+$  and the optical depth-weighted mean spin temperature of H I. As shown in numerical simulations by Kim et al. (2014), the CNM fraction  $f_{\text{CNM}} \propto T_c / \langle T_s \rangle$ , where  $T_c$  is the intrinsic CNM temperature. Murray et al. (2015) showed that most 21-SPONGE observations are consistent with  $f_{\text{CNM}} \propto 50 \text{ K} / \langle T_s \rangle$ , although there was a reasonable scatter. The strong  $\text{HCO}^+$  absorption is associated with H I that has  $\langle T_s \rangle < 140 \text{ K}$ , while weak and broad  $\text{HCO}^+$  absorption is primarily associated with  $\langle T_s \rangle$  in the range of 140–400 K (Figure 5). Thus, based on the relationship between  $f_{\text{CNM}}$  and  $\langle T_s \rangle$  established by simulations and observations, the strong  $\text{HCO}^+$  absorption traces gas with a high CNM fraction, while the weak  $\text{HCO}^+$  absorption traces gas with a lower CNM fraction. This is in agreement with the  $\text{H}_2$  fraction shown in the same figure: a high  $\text{H}_2$  fraction is associated with a high CNM fraction and strong  $\text{HCO}^+$  absorption, while lower  $\text{H}_2$  and CNM fractions are associated with weak and broad  $\text{HCO}^+$  absorption.


Our observations suggest that broad  $\text{HCO}^+$  absorption is common and associated with gas that has low CNM and  $\text{H}_2$  fractions. This diffuse molecular gas may originate in the outer layers of molecular structures. Such regions have been proposed as having significant turbulent motions and mixing between the CNM and WNM (Hennebelle & Inutsuka 2006; Valdivia et al. 2016; Lesaffre et al. 2020). Alternatively, this broad  $\text{HCO}^+$  may simply trace lower-density environments.

<sup>13</sup> More recent work (e.g., Liszt et al. 2010) suggests  $N(\text{HCO}^+)/N(\text{H}_2) = 3 \times 10^{-9}$ . We use  $N(\text{HCO}^+)/N(\text{H}_2) = 2 \times 10^{-9}$  to be consistent with Liszt & Lucas (2000), who performed a similar analysis. Using  $3 \times 10^{-9}$  produces results that are qualitatively the same.



**Figure 4.** A comparison of the  $\text{HCO}^+$  optical depth (blue, positive) with HI absorption (red, negative, with Gaussian components shown as dashed lines; Murray et al. 2018). Regions without  $\text{HCO}^+$  absorption are shaded in crosshatched gray. Regions with weak, broad  $\text{HCO}^+$  absorption are shaded in diagonally hatched green. Regions of strong, narrow  $\text{HCO}^+$  absorption are shaded in unhatched blue.






**Figure 5.** The integrated  $\text{HCO}^+$  optical depth vs. the H I optical depth-weighted mean spin temperature in regions with strong  $\text{HCO}^+$  absorption (blue circles), weak  $\text{HCO}^+$  absorption (green diamonds), and no  $\text{HCO}^+$  absorption (gray points, shown as upper limits), as illustrated in Figure 4. The sizes of the plotted points correspond to the molecular fraction,  $f_{\text{mol}} = 2N(\text{H}_2)/N_{\text{H}}$ , and the colors indicate the H I column densities.

Molecular gas is preferentially formed out of the CNM, so the low  $\text{HCO}^+$  column densities may simply be a result of the relatively small quantity of CNM gas at these velocities.

#### 4.2.2. Broad $\text{HCO}^+$ Absorption as a Tracer of the CO-dark Molecular Gas?

It is intriguing to investigate how this broad component of  $\text{HCO}^+$  absorption might contribute to CO-dark molecular gas. Liszt et al. (2018) have shown that  $\text{HCO}^+$  is a CO-dark molecular gas tracer in some environments and that CO line profiles are systematically narrower than  $\text{HCO}^+$  line profiles. In our sample, the CO emission spectra from Dame et al. (2001) do not show emission at velocities where broad, faint  $\text{HCO}^+$  absorption is observed, as shown in Figure 6. The integrated CO intensity is plotted against the integrated  $\text{HCO}^+$  optical depth for the regions outlined in Figure 4, with regions of strong, narrow absorption again shown in blue circles and regions with weak, broad absorption shown in green diamonds.

At a level of  $3\sigma$ , we do not find any CO emission at the velocities where weak, broad  $\text{HCO}^+$  absorption is observed. On the contrary, we detect CO emission in five of the six velocity ranges where strong, narrow  $\text{HCO}^+$  absorption is observed. However, we note that the typical uncertainty in CO brightness temperature is  $\sim 0.15\text{--}0.3$  K per channel, so our sensitivity may not be adequate to detect any potential low-column density CO. Moreover, due to its low excitation temperature, CO may not be detectable in emission (e.g., Lucas & Liszt 1996; Liszt & Lucas 2000). The broad component of  $\text{HCO}^+$  absorption is seen to be CO-dark in at least one experiment with sensitive observations of  $\text{HCO}^+$  and CO in absorption, though—see Figure 1 of Luo et al. (2020). Future ALMA and NOEMA observations of CO absorption in the directions of our sources will be important for determining how much this broad, faint component to the  $\text{HCO}^+$  absorption traces the total budget of the CO-dark molecular gas. Due to the low molecular fraction at these velocities (Figure 5), though, this may trace only a small fraction of the CO-dark molecular gas.



**Figure 6.** The integrated CO intensity ( $W_{\text{CO}} = \int T_b dv$ ) vs. the integrated  $\text{HCO}^+$  optical depth for the velocity ranges with strong, narrow absorption (blue circles) and weak, broad absorption (green diamonds) outlined in Figure 4. The CO spectra are from T. M. Dame et al. (in preparation). Upper limits are indicated with downward arrows.

### 5. Comparison to Atomic Gas Properties

To investigate the connection between atomic gas properties and molecule formation, we compare the Gaussian-fitted components in the H I absorption spectra to those in the molecular absorption spectra (see Figure 2). For each molecular absorption feature, we find the H I absorption feature closest in velocity space and assume that they are associated with the same interstellar absorbing structure.<sup>14</sup> We make exceptions for low-latitude sources, where velocity crowding introduces significant ambiguity. In these cases, if two H I absorption features are within  $1 \text{ km s}^{-1}$ , but one is a broad, weak feature ( $\Delta v > 10 \text{ km s}^{-1}$ ,  $\tau < 0.1$ ), we take the narrower, stronger feature to be the most probable match, even if it is not the closest in velocity space. In cases where multiple molecular absorption features are matched to the same atomic absorption feature, we consider the sum of the column densities of all of the matched molecular components in the following analysis.


In Figure 7, we show the cumulative distribution functions (CDFs) of the spin temperature, optical depth, and turbulent Mach number of the H I structures seen in absorption toward our background sources (all quantities were constrained by 21-SPONGE). The turbulent Mach number—the ratio of the 3D turbulent velocity to the sound speed—is given by

$$M_t = 2.05 \left( \frac{21.866 \Delta v_{H I}^2}{T_s} - 1 \right)^{1/2}, \quad (6)$$

where  $\Delta v_{H I}$  is the FWHM of the Gaussian H I component. The black CDF is for all H I structures, while the blue, orange, green, and red CDFs are for the H I structures that also show  $\text{HCO}^+$ , HCN, HNC, and  $\text{C}_2\text{H}$  absorption, respectively.

From these CDFs, it is immediately clear that H I structures with a molecular component have higher optical depths and lower spin temperatures (so lower kinetic temperatures) than the general population of cold H I structures in the directions of our background sources. In particular, whereas the total population of H I structures exhibits a large range in H I optical depths and spin temperatures, the H I structures with a

<sup>14</sup> Hereafter, “structure” refers to the individual physical entities that correspond to velocity components identified in the 21-SPONGE Gaussian decomposition.




**Figure 7.** CDFs of the H I optical depth (top), the H I spin temperature (middle), and the H I turbulent Mach number (bottom) measured by Murray et al. (2018) in the directions of the background sources in this work. The CDF for all components is shown in black. The CDFs for the H I features associated with HCO<sup>+</sup>, HCN, HNC, and C<sub>2</sub>H absorption are shown in blue, orange, and green, respectively. Error estimates derived from bootstrapping are shown.

molecular component all have optical depths  $>0.1$  and spin temperatures  $\lesssim 80$  K. There is not such a marked difference in the turbulent Mach numbers of the different distributions, but the turbulent Mach numbers do appear higher for structures with a molecular component,  $M_t \gtrsim 2$ . There is no apparent difference between different molecular species, but this analysis does not necessarily include the broad component HCO<sup>+</sup> absorption that is not always fit in our Gaussian decomposition (Section 4.2). Moreover, our sample size is modest, and few structures do not show absorption from all four species.

These results suggest that regions of the CNM with higher optical depths (more CNM increases the production of H<sub>2</sub> and other molecules) and lower temperatures (meaning more shielded against the ISRF) are more conducive to forming molecules. Such regions also tend to have a higher turbulent Mach number. However, in most cases, the 1D turbulent velocities are not systematically higher for features with a molecular component, suggesting that the higher turbulent Mach number is largely a reflection of the lower  $T_S$ .

These results are also summarized in Figure 8, where we compare HCO<sup>+</sup> column densities with H I optical depths and




**Figure 8.** The HCO<sup>+</sup> column density vs. the H I optical depth (top) and spin temperature (bottom) for the Gaussian components identified in the H I and HCO<sup>+</sup> absorption spectra. HCO<sup>+</sup> detections are shown as filled stars. The 3 $\sigma$  upper limits for nondetections are shown as filled circles. All points are colored according to the H I turbulent Mach number,  $M_t$ . Structures for which 21-SPONGE could not determine the spin temperature are not included in these plots.

spin temperatures. Points are colored according to turbulent Mach number. HCO<sup>+</sup> absorption detections are shown as stars, while 3 $\sigma$  upper limits for nondetections are shown as circles. We do not show here features for which 21-SPONGE was unable to determine the H I spin temperature. Again, we see that molecular gas is associated with H I that is colder, optically thicker, and has a higher turbulent Mach number than the mean of the H I cloud sample in this study. For all of the H I structures along these sightlines, the mean values ( $\tau$ ,  $T_S$ ,  $M_t$ ) are (0.2, 171 K, 3.6), while for H I structures with a molecular component, the mean values are (0.7, 42 K, 4.4).

Figure 8 also shows that many components with  $T_S < 80$  K, as well as many components with  $\tau > 0.1$ , do not have detected molecular column densities. This means that low  $T_S$  and high H I optical depth are necessary but not sufficient conditions for molecule production.

Previously, Godard et al. (2010) showed that it is hard to reproduce HCO<sup>+</sup> column densities larger than a few  $\times 10^{12}$  cm<sup>-2</sup> in the diffuse ISM with PDR models. In Figure 8, we see several detections with such high column densities. We also see that these components have H I temperatures in the range 40–80 K (the peak of the CNM distribution was found to be around 50–60 K; Heiles & Troland 2003; Murray et al. 2018), peak H I optical depths  $\sim 1$ , and  $M_t > 3$  (the colors indicate the turbulent Mach numbers). These four detections



**Figure 9.** The same as Figure 8, but the points are colored according to the 1D turbulent velocity,  $v_t$ , of the atomic gas structures.

also have higher 1D turbulent velocities than detections that show only weak molecular absorption,  $v_t \gtrsim 1 \text{ km s}^{-1}$  (see Figure 9, where the points are colored according to the 1D turbulent velocity), suggesting that the higher turbulent Mach numbers are a product of both the colder temperatures and the turbulent velocities. We further discuss these relatively warmer, more turbulent features in Paper II.

## 6. Comparison of Different Molecular Species

A comparison of the different molecular species contains valuable information about formation and destruction processes. We therefore compare the column densities and absorption line properties of HCN,  $\text{C}_2\text{H}$ ,  $\text{HCO}^+$ , and HNC, and place our results in the context of previous studies.

### 6.1. Line-of-sight Abundances

The abundances of the molecular species studied here are determined by the rates of formation and destruction processes, and may depend on different interstellar environments. For example, in cold and dense interstellar clouds, where the gas is largely shielded from external ultraviolet radiation, the HNC/HCN abundance ratio is expected to be close to 1, based on equilibrium chemical models (Aguado et al. 2017). However, in regions illuminated by ultraviolet photons, HNC is photodissociated faster than HCN, resulting in HCN being significantly more abundant than HNC. This could happen in both diffuse interstellar clouds (Liszt & Lucas 2001; Godard et al. 2010) and in photon-dominated regions (Aguado et al.

**Table 7**  
Column Density Ratios for the Different Sightlines Observed in this Work

Source	$N(\text{HNC})/N(\text{HCN})$	$N(\text{C}_2\text{H})/N(\text{HCO}^+)$
3C111A	$0.26 \pm 0.02^\dagger$	$4.54 \pm 0.43^\dagger$
3C120	$0.19 \pm 0.05$	$15.36 \pm 0.60$
3C123A	$0.17 \pm 0.01$	$6.41 \pm 0.29$
3C123B	$0.27 \pm 0.05^\dagger$	$1.99 \pm 0.62^\dagger$
3C154	$0.23 \pm 0.01$	$13.79 \pm 0.58$
3C454.3	$0.20 \pm 0.02$	$9.33 \pm 0.36$

**Note.** Column 1: name of background radio continuum source. Column 2: ratio of HCN column density to HNC column density. Column 3: ratio of  $\text{C}_2\text{H}$  column density to  $\text{HCO}^+$  column density. A dagger ( $^\dagger$ ) indicates that one or both of the measured column densities is a lower limit.

2017). Similarly, considering that both  $\text{HCO}^+$  and  $\text{C}_2\text{H}$  are products of  $\text{CH}_3^+$ , the abundances of these molecules are expected to be correlated (e.g., Godard et al. 2009).

For the four lines of sight where both HNC and HCN are detected (3C120, 3C123A, 3C154, and 3C454.3, excluding lower limits; see Table 7), we find  $\langle N(\text{HNC})/N(\text{HCN}) \rangle = 0.20 \pm 0.02$ . Similarly, for the four lines of sight where both  $\text{C}_2\text{H}$  and  $\text{HCO}^+$  are detected (3C120, 3C123A, 3C154, and 3C454.3, excluding lower limits; see Table 7), we find  $\langle N(\text{C}_2\text{H})/N(\text{HCO}^+) \rangle = 13.3 \pm 6.9$ . This ratio decreases with increasing  $N(\text{HCO}^+)$  (see Section 6.2 for a further discussion of this observation). Our results are consistent with previous measurements from Liszt & Lucas (2001), who found  $\langle N(\text{HNC})/N(\text{HCN}) \rangle = 0.21 \pm 0.05$ , and Lucas & Liszt (2000), who found  $\langle N(\text{C}_2\text{H})/N(\text{HCO}^+) \rangle = 14.5 \pm 6.7$ .


Our results for HCN and HNC are generally not consistent with the predictions of PDR models from the literature. For example, Aguado et al. (2017) used the Meudon PDR code (Le Petit et al. 2006) to model the chemistry of a typical diffuse cloud. They considered a plane-parallel cloud with a total visual extinction of  $A_V = 1$  mag, illuminated at both sides by the ISRF of 1–3 Draine fields, and considered a range of densities of H nuclei of  $10^2$ – $10^4 \text{ cm}^{-3}$ . They found  $N(\text{HNC})/N(\text{HCN}) \sim 0.5$ – $1$ . While these results are only slightly higher than the observed values, they propagate to CN/HCN ratios one or two orders of magnitude above the observed value.

### 6.2. Gaussian Components

#### 6.2.1. Column Density Ratios

We measure the column density of each Gaussian component, where  $\int \tau dv = 1.064 \tau_0 \Delta v_0$  for a Gaussian feature with peak optical depth  $\tau_0$  and FWHM  $\Delta v_0$ . Figure 10 shows the distribution of these column densities. The slope of  $N(\text{C}_2\text{H})/N(\text{HCO}^+)$  becomes shallower for  $N(\text{HCO}^+) \lesssim 10^{12} \text{ cm}^{-2}$ —this explains why  $N(\text{C}_2\text{H})/N(\text{HCO}^+)$  decreases at high  $N(\text{HCO}^+)$ . The HCN column density increases faster than linearly with the  $\text{HCO}^+$  column density. The HNC and HCN column densities scale linearly with each other.

We find  $\langle N(\text{C}_2\text{H})/N(\text{HCO}^+) \rangle = 18.8 \pm 9.5$ ,  $\langle N(\text{HCN})/N(\text{HCO}^+) \rangle = 1.7 \pm 0.9$ , and  $\langle N(\text{HNC})/N(\text{HCN}) \rangle = 0.22 \pm 0.07$  (or, equivalently,  $\langle N(\text{HCN})/N(\text{HNC}) \rangle = 4.9 \pm 1.3$ ). These ratios are consistent with those measured along the total line of sight (Section 6.1), including the large dispersion in  $N(\text{C}_2\text{H})/N(\text{HCO}^+)$  and  $N(\text{HCN})/N(\text{HCO}^+)$ . Previously, Godard et al. (2010) found  $\langle N(\text{HCN})/N(\text{HCO}^+) \rangle = 1.9 \pm 0.9$



**Figure 10.** Comparisons of the  $\text{C}_2\text{H}$  column densities to the  $\text{HCO}^+$  column densities (top), the  $\text{HCN}$  column densities to the  $\text{HCO}^+$  column densities (middle), and the  $\text{HNC}$  column densities to the  $\text{HCN}$  column densities (bottom), for the features identified in our Gaussian decomposition.


and  $\langle N(\text{HCN})/N(\text{HNC}) \rangle = 4.8 \pm 1.3$  at low Galactic latitudes ( $|b| < 1$ ), and Liszt & Lucas (2001) found  $\langle N(\text{HNC})/N(\text{HCN}) \rangle = 4.8 \pm 1.1$  and  $\langle N(\text{HCN})/N(\text{HCO}^+) \rangle = 1.47 \pm 0.86$  at higher Galactic latitudes ( $1.6 < |b| < 38.2$ ). Our results, measured in the directions of sources with latitudes  $4.0 < |b| < 38.2$ , are consistent with both of these previous works, although two of our sources were also included in Liszt & Lucas (2001).<sup>15</sup>

The similarity in line ratios across Galactic latitude seen by Liszt & Lucas (2001), Godard et al. (2010), and this work suggest that gas in the Galactic plane is chemically similar to gas in the solar neighborhood. This trend was noticed by Godard et al. (2010), but the results here verify that this relationship persists to lower column densities (by a factor of a few) than probed by either of the previous experiments.

### 6.2.2. FWHM Ratios

It has long been known that different molecular species in the diffuse ISM do not share the same kinematics. In particular,

<sup>15</sup> We find at most marginal differences between the column densities measured in this work and the column densities measured in previous works for any overlapping sources; see Section 7.



**Figure 11.** A comparison of the FWHMs of the Gaussian-fitted functions (Tables 3, 4, 5, 6).  $\text{HCO}^+$  FWHMs are shown on the  $x$ -axis. The  $y$ -axis displays the FWHMs for  $\text{HCN}$  (black circles),  $\text{HNC}$  (blue triangles), and  $\text{C}_2\text{H}$  (red squares).

differences in the linewidths of absorption/emission features of different species have been suggested as evidence for the formation of these species under different environmental conditions. We compare the FWHMs between the different species observed here in Figure 11. The mean ratio of the  $\text{C}_2\text{H}$  FWHMs to the  $\text{HCO}^+$  FWHMs is roughly 1.2: 1. The mean ratio of the  $\text{HCN}$  FWHMs to the  $\text{HNC}$  FWHMs is approximately 1: 1. The mean ratio of the  $\text{HCN}$  FWHMs to the  $\text{HCO}^+$  FWHMs and the mean ratio of the  $\text{HNC}$  FWHMs to the  $\text{HCO}^+$  FWHMs are both roughly 0.9: 1. Our measurement of the relative narrowness of  $\text{HCN}$  and  $\text{HNC}$  with respect to  $\text{HCO}^+$  is consistent with Godard et al. (2010), whose data set also included measurements from Lucas & Liszt (1996) and Liszt & Lucas (2001). We note, though, that there is considerable scatter in Figure 11, and that these measured ratios have uncertainties of  $\sim 0.3$ , meaning that our estimates for the line ratios are also not statistically significantly different from 1: 1.

Liszt & Lucas (2001) and Godard et al. (2010) previously observed that absorption by CN-bearing molecules is narrower than absorption by  $\text{HCO}^+$  at millimeter wavelengths, while Lambert et al. (1990) showed that CN absorption is systematically narrower than  $\text{CH}^+$  absorption at optical wavelengths. Liszt et al. (2019) also found that CO absorption is narrower than  $\text{HCO}^+$  absorption at millimeter wavelengths. Molecular absorption lines are turbulently broadened, so it has been suggested that  $\text{HCO}^+$  and  $\text{CH}^+$  lines are broader because these species are formed preferentially in turbulent, dynamic environments like turbulent dissipation regions and shocks (Godard et al. 2010; Liszt et al. 2019). Crane et al. (1995) and Pan et al. (2005) suggested that the differences in linewidths indicated that species like  $\text{HCO}^+$  and  $\text{CH}^+$  trace more diffuse regions such as cloud envelopes, while species like CN and HCN trace denser regions or molecular clouds. Liszt & Lucas (2001) also note that the molecular absorption linewidths may be produced by the turbulence of many structures along the line of sight rather than a single “cloud.” We detect  $\text{HCO}^+$  linewidths that are marginally broader than HCN or HNC linewidths, consistent with the idea that  $\text{HCO}^+$  forms in more turbulent environments than CN-bearing species. Nevertheless, a larger data set is needed to place tighter statistical constraints on the relative widths of different species.



**Table 8**  
The Integrated Optical Depths of HCO<sup>+</sup>, HCN, HNC, and C<sub>2</sub>H Measured by Previous Surveys (Column 3) and Measured in This Work (Column 4)

Source	Species	$(\int \tau dv)_1$ 10 <sup>12</sup> cm <sup>-2</sup>	$(\int \tau dv)_2$ 10 <sup>12</sup> cm <sup>-2</sup>	$\Delta t$ yr	$\Delta(\int \tau dv)$ 10 <sup>12</sup> cm <sup>-2</sup>
3C111	HCO <sup>+</sup>	13.32 ± 0.65 <sup>1</sup>	>11.054	26	
	HCN	13.572 ± 0.276 <sup>2</sup>	>12.710	25	
	HNC	3.149 ± 0.025 <sup>2</sup>	3.606 ± 0.129	25	0.457 ± 0.131
	C <sub>2</sub> H	1.913 ± 0.023 <sup>3</sup>	2.057 ± 0.033	25	0.144 ± 0.04
3C273	HCO <sup>+</sup>	nd	nd	24	
3C345	HCO <sup>+</sup>	nd	nd	24	
3C454.3	HCO <sup>+</sup>	0.32 ± 0.02 <sup>1</sup>	0.276 ± 0.004	24	0.044 ± 0.02
	HCN	0.120 ± 0.013 <sup>2</sup>	0.112 ± 0.004	23	0.008 ± 0.014
	HNC	0.046 ± 0.019 <sup>2</sup>	0.024 ± 0.002	23	0.022 ± 0.019
	C <sub>2</sub> H	0.152 ± 0.025 <sup>3</sup>	0.106 ± 0.004	23	0.046 ± 0.025
BL Lac	HCO <sup>+</sup>	2.30 ± 0.07 <sup>1</sup>	2.260 ± 0.004	26	0.04 ± 0.07
	HCN	2.360 ± 0.080 <sup>2</sup>	2.293 ± 0.002	25	0.067 ± 0.080

**Note.** The time elapsed between surveys is listed in column 5, and the change in integrated optical depth is listed in column 6. The superscripts in column 3 indicate references, as follows—1: Lucas & Liszt (1996); 2: Liszt & Lucas (2001); 3: Lucas & Liszt (2000).

## 7. Temporal Stability of Line Profiles

Temporal variations in absorption line profiles against background radio continuum sources have long been used as a probe for small- (au-)scale structure in the ISM (Stanimirović & Zweibel 2018, and references therein). Such variability has been observed in HI (e.g., Crovisier et al. 1985; Greisen & Liszt 1986; Diamond et al. 1989; Faison & Goss 2001; Brogan et al. 2005; Roy et al. 2012) and several molecular lines (Marscher et al. 1993; Moore & Marscher 1995), although the interpretation of the observed variations has been controversial (Thoraval et al. 1996; Heiles 1997; Deshpande 2000). The observed au-scale structure in HI often has densities high enough for the existence of various molecular species. Detecting variability in molecular spectral lines therefore offers an exciting way of probing the internal structure of overdense HI structures.

In Table 8, we compare the integrated optical depths for HCN, C<sub>2</sub>H (considering only the two transitions listed in Table 1), HCO<sup>+</sup>, and HNC measured in this work to those measured in previous surveys (Lucas & Liszt 1996, 2000; Liszt & Lucas 2001).<sup>16</sup> Observations from these previous works were obtained between 1993 and 1997 with different antenna and receiver setups from those used here, and reduced using different software. Our observations were obtained between 2018 and 2020, meaning that we are probing optical depth changes over a span of ~25 yr in Table 8. We find only modest changes in the integrated optical depths, <3.5 $\sigma$  differences in all cases. Two cases with over 3 $\sigma$  significance are both in the direction of 3C111 (HNC and C<sub>2</sub>H) and probe variability on timescales of 25 yr. 3C111 is especially interesting, as Rybarczyk et al. (2020) found significant spatial variations in the HI optical depths between the different components of this source.

Figure 12 further shows the optical depth spectra of HCO<sup>+</sup> in the directions of 3C120 and 3C454.3 measured here and those measured by Luo et al. (2020). Both were obtained with ALMA using similar spectral setups; the ALMA-SPONGE

spectra were obtained in 2018, while the Luo et al. (2020) spectra were obtained in 2015, so these sightlines probe optical depth changes of ~3 yr. The HCO<sup>+</sup> optical depth spectra show no significant change with respect to the Luo et al. (2020) data taken 3 yr earlier—the peak optical depths differ by 0.001–0.002, a <1% difference detected at <1 $\sigma$ .

Our sample of sightlines with multi-epoch measurements is limited to five, two of which are nondetections. For these sightlines, the lack of short-term variations in the absorption profiles suggests that the molecular component of the gas is not highly structured on scales  $\lesssim 100$  au in these directions.<sup>17</sup> This is consistent with previous observations that molecular absorption line profiles are generally stable over time. Liszt & Lucas (2000) found that HCO<sup>+</sup> and OH absorption line profiles were generally stable over 3–5 yr intervals in the directions of multiple background sources. Wiklind & Combes (1997) showed that HCO<sup>+</sup> absorption spectra in the direction of Centaurus A were stable over a roughly 7 yr span, and Han et al. (2017) showed that HCO<sup>+</sup> absorption spectra in the directions of BL Lac and NRAO 150 were stable over ~20 yr. Furthermore, Liszt & Lucas (2000) argued that the modest changes detected in molecular absorption line profiles over time were most likely caused by small-scale changes in chemical abundances rather than by the presence of highly overdense, au-scale structures.


## 8. Discussion

### 8.1. The Atomic Gas Properties Necessary for Molecule Formation

By comparing the Gaussian components identified in the 21-SPONGE HI absorption spectra (Murray et al. 2015, 2018) with the Gaussian components identified in the ALMA-SPONGE and NOEMA-SPONGE molecular absorption spectra, we have established HI optical depth, spin temperature, and turbulent Mach number thresholds for the formation of HCO<sup>+</sup>,

<sup>16</sup> The Lucas & Liszt works do not observe separate components of 3C111, but instead only report one component. We assume that their observations correspond to the A component, as it is the brightest of the three components by over an order of magnitude.

<sup>17</sup> The motion of the Earth around the Sun and the Sun relative to interstellar clouds can be used to estimate the change in the line of sight over time. See Marscher et al. (1993) for discussion. 100 au is an order of magnitude estimate. The precise distance depends on the velocity and distance to each cloud, but 100 au should be appropriate for typical clouds at a distance of order 100 pc /  $\sin|b|$  (Crovisier 1978).



**Figure 12.** The  $\text{HCO}^+$  optical spectra in the directions of 3C120 (top) and 3C454.3 (bottom). The spectra from Luo et al. (2020), obtained by ALMA in 2015, are shown in orange. The spectra from this work, obtained by ALMA in 2018, are shown in blue.

$\text{HNC}$ ,  $\text{HCN}$ , and  $\text{C}_2\text{H}$  (Section 5). We find that molecular gas forms only if the  $\text{HI}$  optical depth of a gas structure exceeds 0.1, the spin temperature—approximately equal to the kinetic temperature—is less than 80 K, and the turbulent Mach number is greater than  $\sim 2$ . In our sample, these conditions appear necessary but not sufficient for the formation of  $\text{HCO}^+$ ,  $\text{C}_2\text{H}$ ,  $\text{HCN}$ , and  $\text{HNC}$ , as several  $\text{HI}$  absorption features satisfy these criteria but do not have a molecular component. We do not find significantly different thresholds for the different species, although our sample size is modest.

The optical depth threshold of 0.1 is similar to that established by Despois & Baudry (1985) for  $\text{CO}$  formation at  $|b| < 2^\circ$ . However, our  $T_s$  threshold ( $T_s \lesssim 80$  K) is significantly lower than theirs ( $T_s \sim 100$ – $350$  K). We note, though, that Despois & Baudry (1985) derived  $T_s$  using an isothermal approximation of channels with  $\text{CO}$  emission, which is known to result in higher spin temperatures relative to Gaussian decomposition (Murray et al. 2015). They also used  $\text{CO}$  emission observations at  $\sim 4'.3$  beamwidth, which probe more gas than the  $\text{HI}$  pencil beam and are less sensitive to low- and even moderate-column density gas than  $\text{CO}$  absorption observations. Observations of 21 cm  $\text{HI}$  and millimeter wavelength molecular absorption in the same direction have been otherwise limited to a very small number of sightlines

( $N = 1$ – $3$ ; Moore & Marscher 1995; Liszt & Lucas 2000, 2004; Liszt & Gerin 2018). For example, Liszt & Lucas (2000) established that strong  $\text{HCO}^+$  absorption was associated with lower  $\langle T_s \rangle$ , 46–103 K, in three directions (see Section 4.2).

The CNM associated with molecule formation in the diffuse ISM is thus systematically colder and more optically thick than the mean of the total cold  $\text{HI}$  cloud population. CNM structures with a molecular component also have higher turbulent Mach numbers than the general  $\text{HI}$  cloud population, which is a reflection of the colder temperature in most cases, as the 1D turbulent velocities are not systematically higher than the general cloud population. It is also interesting to note that the  $\text{HI}$  temperature threshold that we find for molecule formation, 80 K, is near the mean kinetic temperature estimated from FUV observations of  $\text{H}_2$  (55–80 K; Savage et al. 1977; Rachford et al. 2002, 2009b). This temperature agreement is expected if  $\text{HI}$  and  $\text{H}_2$  are cospatial, as is the case for  $\text{H}_2$  forming out of the CNM on the surface of dust grains and being injected back to the CNM. The high turbulent Mach numbers seen here in the direction of diffuse molecular gas are consistent with previous observations of cold  $\text{HI}$  in the direction of Perseus (e.g., Burkhart et al. 2015).

A relationship between the  $\text{HI}$  column density and  $\text{H}_2$  formation has long been known. The  $\text{HI}$ -to- $\text{H}_2$  transition has been extensively studied in theory and using observations of both entire galaxies and individual molecular clouds. The total gas column density threshold for  $\text{H}_2$  formation has been well established (e.g., Savage et al. 1977; Krumholz et al. 2009; Bellomi et al. 2020, and references therein), and many studies have also noticed a threshold for  $\text{CO}$  formation (e.g., Lombardi et al. 2006; Wong et al. 2009; Leroy et al. 2009; Lee et al. 2014; Imara & Burkhart 2016) that is higher (e.g., Lee et al. 2014;  $A_V \sim 1$ – $2$ ) than that of  $\text{H}_2$  ( $A_V \sim 0.25$ ). Observationally established thresholds for the formation of other molecular species have been rare. Our results are consistent with the previous work of Lucas & Liszt (1996) and Lucas & Liszt (2000) for  $\text{HCO}^+$  and  $\text{C}_2\text{H}$ , although our observations are more sensitive than theirs by a factor of 2–3. Both the simulations from Gong et al. (2020; we discuss this more in Paper II) and our ALMA/NOEMA observations show a transition from atomic to molecular gas at  $N(\text{HI}) \sim 5 \times 10^{20} \text{ cm}^{-2}$ . This threshold appears consistent with that of  $\text{H}_2$  and somewhat lower than that of  $\text{CO}$ . This clearly demonstrates the importance of shielding in building molecular abundances.

## 8.2. The Significance of the UNM in Molecule Formation

Murray et al. (2018) quantified the fraction of thermally unstable  $\text{HI}$  along each line of sight. Although we find that the UNM column density is not directly correlated with the total quantity of molecular gas, it is nevertheless clearly important in setting the  $\text{HCO}^+$  column densities. As we have shown in Figure 3, positions with the highest  $\text{C}_2\text{H}$ ,  $\text{HCO}^+$ ,  $\text{HCN}$ , and  $\text{HNC}$  column densities also tend to have a significant quantity of UNM gas,  $10^{20}$ – $10^{21} \text{ cm}^{-2}$ , comprising  $\sim 30\%$ – $70\%$  of all atomic gas along those lines of sight. The UNM plays a role in shielding, but may also reveal the mixing of the CNM and WNM, which is suggested to enhance the abundances of certain molecular species, including  $\text{HCO}^+$  (Valdivia et al. 2016; Lesaffre et al. 2020).

Previous observational and theoretical work has highlighted the role of the multiphase atomic ISM in molecule formation. For example, Bialy et al. (2015) investigated the  $\text{HI}$ -to- $\text{H}_2$

transition in several regions in the Perseus molecular cloud, where Lee et al. (2012) have found a well-defined threshold for  $\text{H}_2$  formation,  $(8\text{--}14) \times 10^{20} \text{ cm}^{-2}$ . By fitting the Sternberg et al. (2014) model for  $\text{H}_2$  formation in equilibrium, and assuming that HI gas is multiphase and in pressure equilibrium, they constrained the volume density of HI in the atomic shielding envelope of Perseus. As the estimated densities were in the range  $\sim 2\text{--}10 \text{ cm}^{-3}$ , they concluded that much of this HI is thermally unstable, possibly in a cooling transition from the WNM to CNM phases. The behavior in Perseus suggests that in addition to CNM, less-dense UNM, and perhaps some diffuse WNM, are important in controlling the HI-to- $\text{H}_2$  transition. Our observations indicate that the UNM is important for enhancing the abundances of species other than  $\text{H}_2$  and CO, as well.

### 8.3. Interpretation of Broad, Faint $\text{HCO}^+$ Absorption

In a majority of sightlines, we find a broad, faint component to the  $\text{HCO}^+$  absorption that spans a large portion of the velocity space where HI absorption is observed (Section 4.2). Our study suggests that this broad  $\text{HCO}^+$  component is likely to be frequently seen in high-sensitivity observations. The HI coincident with this broad  $\text{HCO}^+$  absorption component has higher  $\langle T_S \rangle$  than the HI at velocities where strong, narrow  $\text{HCO}^+$  absorption is observed, indicating that the fraction of HI in the CNM is lower at these velocities (e.g., Kim et al. 2014). If we assume that  $N(\text{HCO}^+)/N(\text{H}_2)$  is constant (Liszt & Lucas 2000; Liszt et al. 2018), then we also find that the fraction of hydrogen in its molecular form is lower for velocities with broad absorption ( $f_{\text{mol}} \lesssim 0.1$ , versus  $f_{\text{mol}} \gtrsim 0.2$  at velocities with a strong, narrow absorption component).

The fact that the broad component of  $\text{HCO}^+$  absorption is associated with gas that has a lower CNM fraction and a lower molecular fraction may indicate that it is probing the outer layers of molecular structures, where the turbulent mixing between the CNM and WNM allows for the formation of  $\text{CH}^+$  and  $\text{HCO}^+$ . This would explain why the broad  $\text{HCO}^+$  absorption spans roughly the same velocity range as the cold HI observed by 21-SPONGE in nearly all sightlines (Figure 4). The relatively broad linewidths of the faint component of  $\text{HCO}^+$  absorption may also suggest formation in warmer, more diffuse interstellar environments (Falgarone et al. 2006). For example, Valdivia et al. (2017) and Lesaffre et al. (2020) showed that  $\text{CH}^+$ —which contributes to the formation of  $\text{HCO}^+$ —is formed efficiently in turbulent dissipation regions, at the edge of molecular clumps, and in slow shocks, where the HI is thermally unstable and the  $\text{H}_2$  formation is out of equilibrium. In the Valdivia et al. (2017) simulations, most of the  $\text{CH}^+$  is produced in regions with  $\text{H}_2$  fractions  $\sim 0.003\text{--}0.3$  and with gas temperatures as high as  $\sim 10^{2-3} \text{ K}$ . Given the role  $\text{CH}^+$  plays in  $\text{HCO}^+$  formation, the broad  $\text{HCO}^+$  may trace the active regions of  $\text{CH}^+$  production.

We see no CO emission at velocities where the broad, faint  $\text{HCO}^+$  absorption is detected, although the resolution and sensitivity of the CO emission observations from Dame et al. (2001) do not match those of our ALMA/NOEMA absorption spectra. Moreover, due to its low excitation temperature, CO may not be detectable in emission (e.g., Lucas & Liszt 1996; Liszt & Lucas 2000). The possibility that the broad  $\text{HCO}^+$  absorption component is CO-dark warrants further investigation, as it may contribute to the explanation of why  $\text{HCO}^+$  is a better tracer of the total gas content than CO in the diffuse

ISM (Liszt et al. 2018). Liszt et al. (2019) have already shown that  $\text{HCO}^+$  linewidths are broader than CO linewidths, which is consistent with the idea that  $\text{HCO}^+$  is enhanced in diffuse, turbulent regions of the ISM relative to CO. Moreover, Figure 1 of Luo et al. (2020) suggests that there is indeed broad, faint  $\text{HCO}^+$  absorption in the direction of 3C454.3—only marginally detected in our less sensitive spectra—that is predominately CO-dark. However, a quantitative comparison of the amount of  $\text{HCO}^+$  occupied by this broad component with the total amount of CO-dark molecular gas has not yet been made.

### 8.4. The Similarity of Gas at Different Latitudes

We show that the relative abundances of different species are similar between this study and several previous surveys, even though these measurements collectively span Galactic latitudes between  $<1^\circ$  and  $\sim 40^\circ$  (Table 7; Liszt & Lucas 2001; Godard et al. 2010). Similarly, we do not find significant differences between the FWHM ratios of different species with those measured in previous experiments over the same range of latitudes (Godard et al. 2010), although these ratios are less tightly constrained.

The similarities in the relative abundances and linewidths of different species suggest that similar chemical pathways operate across the Galaxy, both in the plane and in the solar neighborhood. This is consistent with the interpretation of Godard et al. (2010), but we have added significantly to the total sample of observations of diffuse molecular gas, including gas at column densities lower than those probed by Liszt & Lucas (2001) and Godard et al. (2010) by a factor of a few. Godard et al. (2010) suggested that the lack of dependence on Galactic latitude indicated that the formation mechanism of different species was weakly dependent on the ambient UV field, perhaps pointing to a ubiquitous mechanism like turbulent dissipation. As discussed further in Paper II, it will be important to observe tracers of nonequilibrium processes like  $\text{CH}^+$  or SiO and test PDR predictions to verify this hypothesis.

### 8.5. Is There a Molecular Component to TSAS?

In Section 7, we show that over periods of 3–25 yr,  $\text{HCO}^+$ ,  $\text{C}_2\text{H}$ , HCN, and HNC absorption spectra are generally stable for our small sample of sightlines with multi-epoch absorption observations.

Marscher et al. (1993) and Moore & Marscher (1995) detected temporal variations in the  $\text{H}_2\text{CO}$  optical depth in the direction of 3C111, arguing in favor of dense clouds ( $n \sim 10^6 \text{ cm}^{-3}$ ) in this direction at a scale of  $\sim 10\text{--}100 \text{ au}$ . However, Thoraval et al. (1996) showed that the  $\text{H}_2\text{CO}$  optical depth variations were most probably a result of changes in the relative abundance or excitation of  $\text{H}_2\text{CO}$ , rather than tiny dense structures. We find marginal variations ( $3.5\sigma$ ) in the integrated optical depths of HNC and  $\text{C}_2\text{H}$  in the direction of 3C111 (Section 7), which suggests that there are not extremely dense structures on scales of  $\sim 10 \text{ au}$  in this direction, as suggested by Marscher et al. (1993) and Moore & Marscher (1995), consistent with the interpretation of Thoraval et al. (1996).

No other sources show temporal variations in the  $\text{HCO}^+$ , HCN, HNC, or  $\text{C}_2\text{H}$  optical depths at a level of  $>2.2\sigma$ . BL Lac—in the direction of the Lacerta molecular cloud—shows remarkably stable absorption spectra,  $<1\sigma$  changes over  $\sim 25$



yr (Lucas & Liszt 1996; Liszt & Lucas 2001). Han et al. (2017) previously found no difference in the  $\text{HCO}^+$  absorption spectra obtained with the Korean VLBI Network to the  $\text{HCO}^+$  absorption spectra from Lucas & Liszt (1996) 20 yr earlier in the direction of BL Lac. Similarly, the integrated and peak optical depths measured here in the directions of 3C120 and 3C454.3 show  $<1\sigma$  changes compared to those obtained by Luo et al. (2020) 3 yr earlier.

Spatial variations in the HI optical depths toward the multiple-component sources 3C111 and 3C123 have revealed TSAS at a scale of  $\sim 10^4$ . 3C111 is a particularly intriguing case, as Dickey & Terzian (1978), Greisen & Liszt (1986), Goss et al. (2008), and Rybarczyk et al. (2020) have all measured optical depth variations of  $\sim 0.3$ – $0.4$  between its different components, corresponding to column density changes of  $\gtrsim 10^{20} \text{ cm}^{-2}$  at scales of  $\sim 10^4$  au. Goss et al. (2008) and Rybarczyk et al. (2020) also previously found HI optical depth variations at a level of  $\sim 0.1$  toward the different components of 3C123. Unfortunately, due to a combination of poor sensitivity toward the fainter components and line saturation, we can only put very weak upper limits on the spatial molecular optical depth variations in these directions. Future sensitive observations of unsaturated molecular absorption lines are required to test if these lines vary with the HI on scales of  $10^4$  au.

We note that the HI optical depth in the direction of 3C138 has been observed to vary dramatically, both spatially ( $\sim a$  few  $\times 10$  au) and temporally ( $\lesssim 7$  yr), making it a regular subject of TSAS investigations. Diamond et al. (1989) and Davis et al. (1996) both found spatial variations in the HI optical depth measured across resolved images of 3C138 obtained using very long baseline interferometry techniques. Later, Faison & Goss (2001), Brogan et al. (2005), and Roy et al. (2012) constructed HI optical depth maps against 3C138 using the Very Long Baseline Array across three different epochs. They detected milliarcsecond HI optical depth variations (corresponding to spatial scales of  $\sim 25$  au) in each epoch, as well as significant variations across epochs. These measurements have been attributed to HI structures with densities  $\sim 10^4 \text{ cm}^{-3}$ . Moreover, Deshpande et al. (2000), who were critical of the interpretation of optical depth variations as evidence of dense structures, acknowledged that the optical depth variations in the direction of 3C138 cannot be explained by geometrical effects, and are likely due to true tiny, dense structures. We see no  $\text{HCO}^+$ ,  $\text{C}_2\text{H}$ , HCN, or HNC absorption in the direction of 3C138. The full-sky maps in Pelgrims et al. (2020), made from the Lallement et al. (2019) 3D dust maps, suggest that this line of sight intersects the Local Bubble wall at  $\lesssim 150$  pc, while the HI associated with Taurus lies at a distance of  $\lesssim 140$  pc (Schlafly et al. 2014). Considering that the absorbing gas is located inside the Local Bubble wall (e.g., Lallement et al. 2014), the lack of molecules associated with TSAS in this direction could be caused by the physical conditions associated with the Local Bubble (e.g., Stanimirović et al. 2010).

## 9. Conclusion

We have complemented HI observations (VLA; Murray et al. 2018) with new  $\text{HCO}^+$ , HCN, HNC, and  $\text{C}_2\text{H}$  (ALMA, NOEMA) observations in the directions of 20 background radio continuum sources with  $4^\circ \leq |b| \leq 81^\circ$  to constrain the atomic gas conditions that are suitable for the formation of diffuse molecular gas. In our sample, we find that these

molecular species form along sightlines where  $A_V \gtrsim 0.25$ , corresponding to  $N_{\text{H}} \gtrsim 5 \times 10^{20} \text{ cm}^{-2}$  (e.g., Zhu et al. 2017), consistent with a threshold for the HI-to- $\text{H}_2$  transition at solar metallicity (Savage et al. 1977). This confirms the trend noticed by Lucas & Liszt (1996) and Lucas & Liszt (2000) for  $\text{HCO}^+$  and  $\text{C}_2\text{H}$ , with absorption spectra  $\sim 2$ – $3$  times more sensitive than theirs. Moreover, we find that molecular gas is associated only with structures that have an HI optical depth  $>0.1$ , a spin temperature  $<80$  K, and a turbulent Mach number  $\gtrsim 2$ .

We identify a broad, faint component to the  $\text{HCO}^+$  absorption in a majority of sightlines. In several cases, this component spans nearly the entire range of velocities where HI is seen in absorption. The HI at velocities where only the faint, broad  $\text{HCO}^+$  absorption is observed has systematically higher  $\langle T_s \rangle$  than the HI at velocities where strong, narrow  $\text{HCO}^+$  absorption is observed, indicating that the HI at these velocities has a lower CNM fraction (Kim et al. 2014). We also do not detect CO emission at these velocities, whereas CO emission is detected at nearly all velocities where strong  $\text{HCO}^+$  absorption is observed. The broad  $\text{HCO}^+$  may therefore be a good tracer of the CO-dark molecular gas, and deserves further investigation.

The relative column densities and linewidths of the different molecular species observed here are similar to those observed in previous experiments over a range of Galactic latitudes, suggesting that gas in the solar neighborhood and gas in the Galactic plane are chemically similar. For a select sample of previously observed sightlines, we show that the absorption line profiles of  $\text{HCO}^+$ , HCN, HNC, and  $\text{C}_2\text{H}$  are stable over periods of  $\sim 3$  yr and  $\sim 25$  yr, likely indicating that molecular gas structures in these directions are at least  $\gtrsim 100$  au in size.

In Paper II, we compare the observational results presented here to predictions from the PDR chemical model in Gong et al. (2017) and the ISM simulations in Gong et al. (2020) to investigate in detail the local physical conditions (density, radiation field, and cosmic ray ionization rate) needed to explain the observed column densities.


We would like to thank Ningyu Tang for providing useful supplementary data. We are grateful to Antoine Gusdorf, Benjamin Godard, and Carl Heiles for useful discussions about this work.

Support for this work was provided by the NSF through award SOSP6-023 from the NRAO. S.S. acknowledges the support by the Vilas funding provided by the University of Wisconsin and the John Simon Guggenheim fellowship. This work is based on observations carried out under project numbers W19AQ and S20AB with the IRAM NOEMA Interferometer. IRAM is supported by INSU/CNRS (France), MPG (Germany), and IGN (Spain). This paper makes use of the following ALMA data: ADS/JAO.ALMA#2018.1.00585.S and ADS/JAO.ALMA#2019.1.01809.S. ALMA is a partnership of ESO (representing its member states), NSF (USA), and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO, and NAOJ. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

*Software:* Astropy (Astropy Collaboration et al. 2013, 2018), CASA (McMullin et al. 2007), GILDAS (Pety 2005; Gildas 2013), dustmaps (Green 2018), Scipy (Jones et al. 2001; Virtanen et al. 2020).



## ORCID iDs

Daniel R. Rybarczyk  <https://orcid.org/0000-0003-3351-6831>  
 Snežana Stanimirović  <https://orcid.org/0000-0002-3418-7817>  
 Munan Gong  <https://orcid.org/0000-0003-1613-6263>  
 Brian Babler  <https://orcid.org/0000-0002-6984-5752>  
 Claire E. Murray  <https://orcid.org/0000-0002-7743-8129>  
 Maryvonne Gerin  <https://orcid.org/0000-0002-2418-7952>  
 Jan Martin Winters  <https://orcid.org/0000-0001-6114-9173>  
 Gan Luo  <https://orcid.org/0000-0002-1583-8514>  
 T. M. Dame  <https://orcid.org/0000-0003-0109-2392>  
 Lucille Steffes  <https://orcid.org/0000-0002-9512-5492>

## References

- Aguado, A., Roncero, O., Zanchet, A., Agúndez, M., & Cernicharo, J. 2017, *ApJ*, **838**, 33
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, **156**, 123
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, **558**, A33
- Bellomi, E., Godard, B., Hennebelle, P., et al. 2020, *A&A*, **643**, A36
- Bialy, S., Burkhart, B., & Sternberg, A. 2017, *ApJ*, **843**, 92
- Bialy, S., Sternberg, A., Lee, M.-Y., Le Petit, F., & Roueff, E. 2015, *ApJ*, **809**, 122
- Bohlin, R. C., Savage, B. D., & Drake, J. F. 1978, *ApJ*, **224**, 132
- Bolatto, A. D., Leroy, A. K., Jameson, K., et al. 2011, *ApJ*, **741**, 12
- Brogan, C. L., Zauderer, B. A., Lazio, T. J., et al. 2005, *AJ*, **130**, 698
- Burkhart, B., Lee, M.-Y., Murray, C. E., & Stanimirović, S. 2015, *ApJL*, **811**, L28
- Crane, P., Lambert, D. L., & Sheffer, Y. 1995, *ApJS*, **99**, 107
- Crovisier, J. 1978, *A&A*, **70**, 43
- Crovisier, J., Dickey, J. M., & Kazes, I. 1985, *A&A*, **146**, 223
- Dame, T. M., Hartmann, D., & Thaddeus, P. 2001, *ApJ*, **547**, 792
- Davis, R. J., Diamond, P. J., & Goss, W. M. 1996, *MNRAS*, **283**, 1105
- Deshpande, A. A. 2000, *MNRAS*, **317**, 199
- Deshpande, A. A., Dwarakanath, K. S., & Goss, W. M. 2000, *ApJ*, **543**, 227
- Despois, D., & Baudry, A. 1985, *A&A*, **148**, 83
- Diamond, P. J., Goss, W. M., Romney, J. D., et al. 1989, *ApJ*, **347**, 302
- Dickey, J. M., & Terzian, Y. 1978, *A&A*, **70**, 415
- Draine, B. T. 1978, *ApJS*, **36**, 595
- Endres, C. P., Schlemmer, S., Schilke, P., Stutzki, J., & Müller, H. S. P. 2016, *JMoSp*, **327**, 95
- Faison, M. D., & Goss, W. M. 2001, *AJ*, **121**, 2706
- Falgarone, E., Pineau Des Forêts, G., Hily-Blant, P., & Schilke, P. 2006, *A&A*, **452**, 511
- Field, G. B., Goldsmith, D. W., & Habing, H. J. 1969, *ApJL*, **155**, L149
- Gildas, Team 2013, GILDAS: Grenoble Image and Line Data Analysis Software, Astrophysics Source Code Library, ascl:1305.010
- Glover, S. C. O., & Clark, P. C. 2012, *MNRAS*, **421**, 9
- Godard, B., Falgarone, E., Gerin, M., Hily-Blant, P., & de Luca, M. 2010, *A&A*, **520**, A20
- Godard, B., Falgarone, E., & Pineau Des Forêts, G. 2009, *A&A*, **495**, 847
- Godard, B., Falgarone, E., & Pineau des Forêts, G. 2014, *A&A*, **570**, A27
- Gong, M., Ostriker, E. C., & Kim, C.-G. 2018, *ApJ*, **858**, 16
- Gong, M., Ostriker, E. C., Kim, C.-G., & Kim, J.-G. 2020, *ApJ*, **903**, 142
- Gong, M., Ostriker, E. C., & Wolfire, M. G. 2017, *ApJ*, **843**, 38
- Goss, W. M., Richards, A. M. S., Muxlow, T. W. B., & Thomasson, P. 2008, *MNRAS*, **388**, 165
- Green, G. 2018, *JOSS*, **3**, 695
- Greisen, E. W., & Liszt, H. S. 1986, *ApJ*, **303**, 702
- Güver, T., & Özel, F. 2009, *MNRAS*, **400**, 2050
- Han, J., Yun, Y., & Park, Y.-S. 2017, *JKAS*, **50**, 185
- Heiles, C. 1997, *ApJ*, **481**, 193
- Heiles, C., & Troland, T. H. 2003, *ApJ*, **586**, 1067
- Heitsch, F., Slyz, A. D., Devriendt, J. E. G., Hartmann, L. W., & Burkert, A. 2006, *ApJ*, **648**, 1052
- Hennebelle, P., & Audit, E. 2007, *A&A*, **465**, 431
- Hennebelle, P., & Inutsuka, S.-i. 2006, *ApJ*, **647**, 404
- Imara, N., & Burkhart, B. 2016, *ApJ*, **829**, 102
- Inoue, T., & Inutsuka, S.-i. 2012, *ApJ*, **759**, 35
- Inutsuka, S.-i., Inoue, T., Iwasaki, K., & Hosokawa, T. 2015, *A&A*, **580**, A49
- Jenkins, E. B., & Savage, B. D. 1974, *ApJ*, **187**, 243
- Jones, E., Oliphant, T., & Peterson, P. 2001, SciPy: Open source scientific tools for Python, <https://scipy.org/>
- Kim, C.-G., Ostriker, E. C., & Kim, W.-T. 2014, *ApJ*, **786**, 64
- Koyama, H., & Inutsuka, S.-i. 2002, *ApJL*, **564**, L97
- Krumholz, M. R., McKee, C. F., & Tumlinson, J. 2009, *ApJ*, **693**, 216
- Lallement, R., Babusiaux, C., Vergely, J. L., et al. 2019, *A&A*, **625**, A135
- Lallement, R., Vergely, J. L., Valette, B., et al. 2014, *A&A*, **561**, A91
- Lambert, D. L., Sheffer, Y., & Crane, P. 1990, *ApJL*, **359**, L19
- Le Petit, F., Nehmé, C., Le Bourlot, J., & Roueff, E. 2006, *ApJS*, **164**, 506
- Lee, M.-Y., Stanimirović, S., Douglas, K. A., et al. 2012, *ApJ*, **748**, 75
- Lee, M.-Y., Stanimirović, S., Wolfire, M. G., et al. 2014, *ApJ*, **784**, 80
- Lequeux, J. 2005, *The Interstellar Medium* (Berlin: Springer)
- Leroy, A. K., Bolatto, A., Bot, C., et al. 2009, *ApJ*, **702**, 352
- Lesaffre, P., Gerin, M., & Hennebelle, P. 2007, *A&A*, **469**, 949
- Lesaffre, P., Todorov, P., Levrier, F., et al. 2020, *MNRAS*, **495**, 816
- Liszt, H., & Gerin, M. 2018, *A&A*, **610**, A49
- Liszt, H., Gerin, M., & Grenier, I. 2018, *A&A*, **617**, A54
- Liszt, H., Gerin, M., & Grenier, I. 2019, *A&A*, **627**, A95
- Liszt, H., & Lucas, R. 2000, *A&A*, **355**, 333
- Liszt, H., & Lucas, R. 2001, *A&A*, **370**, 576
- Liszt, H., & Lucas, R. 2004, *A&A*, **428**, 445
- Liszt, H. S., Pety, J., & Lucas, R. 2010, *A&A*, **518**, A45
- Lombardi, M., Alves, J., & Lada, C. J. 2006, *A&A*, **454**, 781
- Lucas, R., & Liszt, H. 1996, *A&A*, **307**, 237
- Lucas, R., & Liszt, H. S. 2000, *A&A*, **358**, 1069
- Luo, G., Li, D., Tang, N., et al. 2020, *ApJL*, **889**, L4
- Marscher, A. P., Moore, E. M., & Bania, T. M. 1993, *ApJL*, **419**, L101
- McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, in ASP Conf. Ser., 376, *Astronomical Data Analysis Software and Systems XVI*, ed. R. A. Shaw, F. Hill, & D. J. Bell (San Francisco, CA: ASP), 127
- Moore, E. M., & Marscher, A. P. 1995, *ApJ*, **452**, 671
- Müller, H. S. P., Thorwirth, S., Roth, D. A., & Winnewisser, G. 2001, *A&A*, **370**, L49
- Murray, C. E., Stanimirović, S., Goss, W. M., et al. 2015, *ApJ*, **804**, 89
- Murray, C. E., Stanimirović, S., Goss, W. M., et al. 2018, *ApJS*, **238**, 14
- Nguyen, H., Dawson, J. R., Lee, M.-Y., et al. 2019, *ApJ*, **880**, 141
- Pan, K., Federman, S. R., Sheffer, Y., & Andersson, B. G. 2005, *ApJ*, **633**, 986
- Pelgrims, V., Ferrière, K., Boulanger, F., Lallement, R., & Montier, L. 2020, *A&A*, **636**, A17
- Pety, J. 2005, in SF2A-2005: Semaine de l'Astrophysique Française, ed. F. Casoli et al. (Les Ulis: EdP-Sciences), 721
- Pikel'Ner, S. B. 1968, *SvA*, **11**, 737
- Planck Collaboration, Abergel, A., Ade, P. A. R., et al. 2014a, *A&A*, **571**, A11
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2014b, *A&A*, **571**, A13
- Planck Collaboration, Aghanim, N., Ashdown, M., et al. 2016, *A&A*, **596**, A109
- Rachford, B. L., Snow, T. P., Destree, J. D., et al. 2009a, *ApJS*, **180**, 125
- Rachford, B. L., Snow, T. P., Destree, J. D., et al. 2009b, *ApJS*, **180**, 125
- Rachford, B. L., Snow, T. P., Tumlinson, J., et al. 2002, *ApJ*, **577**, 221
- Roy, N., Minter, A. H., Goss, W. M., Brogan, C. L., & Lazio, T. J. W. 2012, *ApJ*, **749**, 144
- Rybarczyk, D. R., Gong, M., Stanimirović, S., et al. 2022, *ApJ*, **926**, 190
- Rybarczyk, D. R., Stanimirović, S., Zweibel, E. G., et al. 2020, *ApJ*, **893**, 152
- Savage, B. D., Bohlin, R. C., Drake, J. F., & Budich, W. 1977, *ApJ*, **216**, 291
- Schlaflly, E. F., Green, G., Finkbeiner, D. P., et al. 2014, *ApJ*, **786**, 29
- Schöier, F., van der Tak, F., van Dishoeck, E., & Black, J. 2010, *LAMDA: Leiden Atomic and Molecular Database*, Astrophysics Source Code Library, ascl:1010.077
- Spitzer, L., Drake, J. F., Jenkins, E. B., et al. 1973, *ApJL*, **181**, L116
- Stanimirović, S., Murray, C. E., Lee, M.-Y., Heiles, C., & Miller, J. 2014, *ApJ*, **793**, 132
- Stanimirović, S., Weisberg, J. M., Pei, Z., Tuttle, K., & Green, J. T. 2010, *ApJ*, **720**, 415
- Stanimirović, S., & Zweibel, E. G. 2018, *ARA&A*, **56**, 489
- Sternberg, A., Le Petit, F., Roueff, E., & Le Bourlot, J. 2014, *ApJ*, **790**, 10
- Thoraval, S., Boisse, P., & Stark, R. 1996, *A&A*, **312**, 973
- Valdivia, V., Godard, B., Hennebelle, P., et al. 2017, *A&A*, **600**, A114
- Valdivia, V., Hennebelle, P., Gerin, M., & Lesaffre, P. 2016, *A&A*, **587**, A76
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *NaMe*, **17**, 261
- Wiklund, T., & Combes, F. 1997, *A&A*, **324**, 51
- Wolfire, M. G., McKee, C. F., Hollenbach, D., & Tielens, A. G. G. M. 2003, *ApJ*, **587**, 28
- Wong, T., Hughes, A., Fukui, Y., et al. 2009, *ApJ*, **696**, 370
- Zhu, H., Tian, W., Li, A., & Zhang, M. 2017, *MNRAS*, **471**, 3494