

Molecule formation in dust-poor irradiated jets

I. Stationary disk winds (*Corrigendum*)

B. Tabone^{1,2}, B. Godard^{2,3}, G. Pineau des Forets^{2,4}, S. Cabrit², and E. F. van Dishoeck^{1,5}

¹ Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands
e-mail: tabone@strw.leidenuniv.nl

² LERMA, Observatoire de Paris, PSL Research Univ., CNRS, Sorbonne Univ., 75014 Paris, France

³ Laboratoire de Physique de l'École normale supérieure, ENS, Université PSL, CNRS, Sorbonne Université, Université Paris-Diderot, Sorbonne Paris Cité, 75005 Paris, France

⁴ Université Paris-Saclay, CNRS, Institut d'Astrophysique Spatiale, 91405 Orsay, France

⁵ Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstrasse 1, 85748 Garching, Germany

A&A, 636, A60 (2020), <https://doi.org/10.1051/0004-6361/201937383>

Key words. stars: formation – stars: jets – stars: protostars – ISM: molecules – errata, addenda

Some typographical errors occurred during the production process of the original published article. The lines below Eqs. (14) and (15) should read “where we note as...”. Also, in Appendix C, Eqs. (C.15) and (C.16) were switched and the new Eq. (C.16) is not correct. The sentence that contains the two equations should read:

Defining

$$\eta_X \equiv \frac{n_H}{G_0 n_H^{\text{crit},X}}, \quad (\text{C.15})$$

a fraction

$$\frac{\eta_X}{1 + \eta_X} \quad (\text{C.16})$$

of the XH^+ formed by radiative association is converted in H_2 .