



**HAL**  
open science

## Unexplained Oxygen Variability: New Results on Molecular Oxygen in the Lower Martian Atmosphere from Chemcam and Supercam Passive Sky Observations.

T. H. Mcconnochie, Melissa G. Trainer, M. D. Smith, S. D. Guzewich, H. B. Franz, C. E. Newman, D. Y. Lo, S. Atreya, J. E. Moores, H. M. Sapers, et al.

### ► To cite this version:

T. H. Mcconnochie, Melissa G. Trainer, M. D. Smith, S. D. Guzewich, H. B. Franz, et al.. Unexplained Oxygen Variability: New Results on Molecular Oxygen in the Lower Martian Atmosphere from Chemcam and Supercam Passive Sky Observations.. Seventh International Workshop on the Mars Atmosphere: Modelling and Observations, Jun 2022, Paris, France. pp.id.3403. insu-03746716

**HAL Id: insu-03746716**

**<https://insu.hal.science/insu-03746716v1>**

Submitted on 5 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# UNEXPLAINED OXYGEN VARIABILITY: NEW RESULTS ON MOLECULAR OXYGEN IN THE LOWER MARTIAN ATMOSPHERE FROM CHEMCAM AND SUPERCAM PASSIVE SKY OBSERVATIONS

**T. H. McConnochie**, *Space Science Institute, Boulder CO, USA ([tmccconnochie@spacescience.org](mailto:tmccconnochie@spacescience.org))*, **M. G. Trainer**, **M. D. Smith**, **S. D. Guzewich**, **H. B. Franz**, *NASA Goddard Space Flight Center, Greenbelt, MD, USA*, **C. E. Newman**, *Aeolis Research, Chandler, AZ, USA*, **D. Y. Lo**, **S. Atreya**, *University of Michigan, Ann Arbor, MI, USA*, **J. E. Moores**, **H. M. Sapers**, *Centre for Research in Earth and Space Science, York University, Toronto, Canada*, **M. T. Lemmon**, **M. J. Wolff**, *Space Science Institute, Boulder CO, USA*, **F. Montmessin**, **E. W. Knutsen**, *LATMOS, Guyancourt, France*, **T. Fouchet**, **T. Bertrand**, *LESIA, Paris Observatory, France*, **O. Gasnault**, **J. Lasue**, **O. Forni**, **P. Pilleri**, **S. Maurice**, *IRAP, CNRS, Université de Toulouse, UPS-OMP, Toulouse, France*, **C. Legett IV**, **R. T. Newell**, **D. Venhaus**, **N. Lanza**, *Los Alamos National Laboratory, Los Alamos, NM, USA*, **R. C. Wiens**, *Purdue University, West Lafayette IN, USA*, **M. Hecht**, *MIT Haystack Observatory, Westford MA, USA*, **M.-P. Zorzano**, *Centro de Astrobiología, INTA, Spain*, **A. Khayat**, *NASA Goddard Space Flight Center, Greenbelt, MD, USA* and *CRESST II, Department of Astronomy, University of Maryland, MD, USA*, **Franck Lefèvre**, *LATMOS, CNRS, Sorbonne Université, UVSQ, Paris, France*, **Frank Daerden**, *Royal Belgian Institute for Space Aeronomy, Brussels, Belgium*, **Anna Fedorova**, **Alexander Trokhimovskiy**, *Space Research Institute RAS, Moscow, Russia*

**Background:** MSL's SAM-QMS atmospheric sampling showed that the O<sub>2</sub> volume mixing ratio (VMR) in the near-surface Martian atmosphere has significant seasonal and interannual deviations from the pattern expected and observed for inert non-condensable trace gases (Trainer et al, 2019). Yet photochemical GCM's show that, given known chemistry, O<sub>2</sub> should behave identically to inert trace gases such as argon (Lefevre et al., 2004; Daerden et al., 2019), consistent with photochemical models showing O<sub>2</sub> lifetimes of at least 10 years (Atreya et al., 2006; Lefèvre & Krasnopolsky, 2017).

Trainer et al. (2019) argued that the magnitude of O<sub>2</sub> deviation from inert trace gas behavior is too large to be explained by any atmospheric sources – essentially because CO<sub>2</sub> photolysis is too slow and H<sub>2</sub>O vapor abundance is too small – which suggests that some kind of surface reservoir of oxygen is actively exchanging with the atmosphere on seasonal timescales. The Viking Gas Exchange (GEx) experiment's finding of O<sub>2</sub> release from a soil sample upon humidification (Klein, 1978; Oyama & Berdahl, 1977) supports the idea that such a reservoir might exist. Both superoxides and UV-degraded or ionizing-radiation-degraded chlorates & perchlorates have been proposed (Yen et al. 2000, Quinn et al., 2013, respectively) as the source reservoir for the Viking GEx O<sub>2</sub> release, and both superoxides and chlorates/perchlorates have been proposed to form with an atmospheric source for their O<sub>2</sub> (Yen et al. 2000, Carrier and Kounaves, 2015). Thus, there are proposed mechanisms for both directions of a cyclical exchange of O<sub>2</sub> between the atmosphere and surface. We, therefore, can consider a hypothetical surface-atmosphere “oxygen cycle” as a potential explanation for the unexpected O<sub>2</sub> variability; although it remains unclear whether any of these surface-atmosphere exchange mechanisms can operate with the rates necessary to explain the SAM data. Interestingly, both the uptake and release

mechanisms for O<sub>2</sub> in surface soils, as they have been proposed, involve exposure to UV or higher-energy radiation, but only the release mechanisms are proposed to involve water vapor humidity. This suggests both a relationship between atmospheric O<sub>2</sub> and the water cycle, an idea that Lo et al. (2022) have already begun to explore, and a potentially complicated relationship with UV-blocking aerosol dust which could limit both uptake *and* release of O<sub>2</sub>.

Understanding this apparent surface-atmosphere exchange of oxygen will be important for understanding the formation and likely distribution in space and time of perchlorates and other soil oxidants, which in turn have implications for the preservation of organics on the Martian surface. In addition, having a major active reservoir for O<sub>2</sub> at the surface would potentially complicate the traditional explanation (e. g. Lefèvre & Krasnopolsky, 2017) for the relative abundances of CO and O<sub>2</sub> and their role in recycling CO<sub>2</sub> photolysis products back to CO<sub>2</sub> to stabilize the Martian atmosphere.

**Measurements:** Both ChemCam on the MSL rover (e.g. Wiens et al., 2012) and SuperCam on the Mars2020 rover (e.g. Maurice et al., 2021; Wiens et al., 2021) perform “passive sky” observations following the technique of McConnochie et al. (2018). No laser is used (hence “passive”) and the spectrometer views the sky at two different elevation angles to provide two different path lengths through the absorbing gases. Ratioing the signal from the two different path lengths eliminates instrument response and solar spectrum uncertainties.

The O<sub>2</sub> retrieval technique also follows McConnochie et al. (2018), but in this case fitting the 759 – 767 nm O<sub>2</sub> absorption and updating the molecular line parameters to those from the HITRAN 2016 database. CO<sub>2</sub> absorption lines near 783 nm and 789 nm are used as a reference to correct for the effect of

aerosol scattering uncertainties on line depth. The depth of these O<sub>2</sub> and CO<sub>2</sub> absorption lines is about 0.05% – 0.2% relative to the continuum.

In this spectral region SuperCam uses a transmission spectrometer and an intensifier, while ChemCam uses a reflection grating and no intensifier. Both use nearly-identical CCD detectors. The SuperCam oxygen retrieval precision appears to be limited mainly by fluctuations in intensifier throughput, while for the non-intensified ChemCam it is clear that both systematic and random uncertainties are dominated by detector background. SuperCam and ChemCam thus require very different data processing for passive sky, and with an 8-year head start the ChemCam data processing is relatively mature although some improvements are still planned. SuperCam data processing for filtering intensifier throughput feature is still preliminary and so the results that follow are based solely on the ChemCam passive sky data set.

Both ChemCam and SuperCam passive sky observations are acquired irregularly to minimize conflicts with other rover operations, but we are currently targeting an *average* sampling interval of 14 sols with both instruments. Over the course of the MSL mission, the ChemCam passive sky average sampling interval has been as small as ~7 sols in some periods and as large as ~50 sols in other periods. Since September 2020, we have been adjusting the timing of ChemCam passive sky measurements (without altering the *average* sampling interval) to occur within 1-2 sols of a Trace Gas Orbiter ACS occultation observation (e.g. Korabev et al., 2018) in the vicinity of Gale Crater whenever feasible.

### Results:

*Long-term Averages.* ChemCam O<sub>2</sub> retrievals give 2000 +/- 60 ppm averaged over roughly 180 individual observations spanning about 4.5 Martian years. This is comparable to Trace Gas Orbiter (TGO) ACS occultation results which give ~1900 ppm on average for 15 – 25 km altitudes in the low-to-mid-latitudes (Fedorova et al., EGU General Assembly 2021 presentation materials). MSL’s SAM Quadrupole Mass Spectrometer atmospheric samples show notably less O<sub>2</sub> than ChemCam, giving 1610 ± 90 ppm averaged over Mars Years 31-34. Herschel Space Observatory’s HIFI observed even less O<sub>2</sub> in Mars Year 30 (@ L<sub>s</sub> = 77) reporting a value of only 1400 ± 120 ppm (Hartogh et al., 2010), which was comparable to the previous canonical value of ~1300 ppm from ground based measurements (Barker, 1972; Carleton and Traub, 1972).

Thus, understanding of Martian O<sub>2</sub> abundances has evolved significantly in recent years from a generally-accepted value near 1300 ppm to a new paradigm of average abundances in the 1600 – 2000 ppm range. Furthermore, the difference between the long term average near-surface SAM O<sub>2</sub> (Trainer et al. 2019), and the long term average column measurement from ChemCam, is statistically robust, and


Mars Atmosphere O <sub>2</sub> Measurements		
O <sub>2</sub> VMR	Type and Timing	Reference
2000± 60 ppm	MSL ChemCam, average of MY 31–36	this work
~1900	Trace Gas Orbiter ACS, average of 15-25 km altitudes in low-to-mid-latitudes, MY 3435	Fedorova et al., 2021
1610 ± 90 ppm	MSL SAM, average of MY 31 34	Trainer et al. 2019
1400 ± 120 ppm	Herschel Space Observatory, MY 30, L <sub>s</sub> = 77)	Hartogh et al., 2010
1300 ± 130 ppm	Ground based MY 9, L <sub>s</sub> ~ 300 - 340	Barker, 1972; Carleton and Traub, 1972

TGO-ACS (Fedorova et al., 2021) tends to corroborate ChemCam. Consequently, there is a strong suggestion that O<sub>2</sub> tends to be depleted near the surface at the equator, which is further evidence of O<sub>2</sub> lifetime being much shorter than the > 10 years in existing models and also fits with a hypothesis that release of O<sub>2</sub> from soil is disfavored at the surface near the equator compared to other locations on Mars due to lower humidity.

*Seasonal cycles and interannual changes.* In previous reports on ChemCam O<sub>2</sub> results we had low confidence in the ChemCam evidence of interannual and seasonal variability because the amount of scatter and frequency outliers in the times series was not well explained by our predicted uncertainties. This situation has now changed because the data set has grown sufficiently large and because we have improved the statistics of our smoothed time series by adopting a local regression approach. The local regression finds a smoothed value at a given time by fitting a linear regression to nearby observation points, weighting each observation point by its distance as well as its estimating uncertainty.

In Figure 1 we have suppressed individual observation points in order to show 5 Martian years of the smoothed ChemCam O<sub>2</sub> results with 1-sigma uncertainties represented by the shaded area around each dashed line. Thinner lines with no uncertainty shading are outputs from the LMD (Lefèvre et al., 2004) and GEM (Daerden et al., 2019) photochemical GCM models for comparison. Red points with error bars are SAM O<sub>2</sub> measurements from Trainer et al. (2019), and the green points are Ar from Trainer et al. (2019) to show that the GCM models perform very well for truly inert gases. The season around L<sub>s</sub> = 90 stands out as having clearly significant interannual variability in ChemCam O<sub>2</sub> results. Mars Year 34 also shows notably lower ChemCam O<sub>2</sub> than in other years in the L<sub>s</sub> = 180 season, just before the Mars Year 34 global dust

**Figure 1** O<sub>2</sub> Measurements and models



storm. The very high value in the post-dust-storm period of Mars Year 34 is based on only one ChemCam passive sky observation with relatively poor precision so its large apparent difference from the values in other years is not actually statistically significant, as is apparent from the uncertainty represented by the shading.

One consequence of the large interannual variability in the  $L_s = 90$  season is that the Herschel O<sub>2</sub> measurement, which happened to occur at  $L_s = 77$ , turns out to not be an outlier. Factoring in its uncertainties, it is only slightly smaller than what was observing by ChemCam in MY 35, and, given the wide interannual dispersion, it's entirely plausible that ChemCam might have observed even less in that same season in MY 30 had it been in operation at that time.

#### Discussion:

The large deviation of ChemCam O<sub>2</sub> from the model expectations, and the lack of a strong correlation and different average values between SAM and ChemCam O<sub>2</sub> are further evidence of oxygen chemistry being more active than in current model expectations because they mean that the O<sub>2</sub> chemical lifetimes cannot be much larger than advections or vertical mixing time scales. ChemCam and SuperCam are sensitive only to the bottom 25 km of the atmosphere, with a 75% weight on the bottom 10 km.

A more novel implication comes from the large interannual variability of O<sub>2</sub> in certain seasons. This suggests that the hypothesized oxygen cycle would be quite sensitive to environmental conditions that are themselves quite variable. The primary season of observed O<sub>2</sub> interannual variability happens to coincide with peak water vapor abundance over northern high latitudes, so, speculatively, variability in that peak humidity and its interaction with surface materials or aerosol opacity could drive variable release of O<sub>2</sub> from the soil, which is then rapidly advected into the column over Gale Crater but perhaps less rapidly transported to the near surface atmosphere in Gale as sampled by SAM.

Adding O<sub>2</sub> measurement over Jezero Crater from SuperCam will serve to further constrain such

hypothetical processes. Continuing coordinated observations with TGO-ACS will do so as well. Ultimately, however, the path forward for interpreting O<sub>2</sub> at those multiple sites in terms of processes and preferred locations for surface-atmosphere oxygen exchange will require a GCM that incorporates some representation of the hypothesized surface processes.

#### References:

- Atreya et al. (2006). *Astrobiology*, 6(3), 439–450. doi:10.1089/ast.2006.6.439
- Carrier and Kounaves (2015) *GRL* 42, 3739–3745, doi:10.1002/2015GL064290.
- Daerden et al. (2019). *Icarus*, 326, 197–224, doi: 10.1016/j.icarus.2019.02.030.
- Fedorova et al., 2021, EGU General Assembly EGU21-14406, doi:10.5194/egusphere-egu21-14406.
- Trainer et al., 2019, *JGR* 124. doi:10.1029/2019JE006175
- Hartogh et al. (2010). *Astronomy and Astrophysics*, 521, L49. doi:10.1051/0004-6361/201015160.
- Klein (1978). *Icarus*, 34(3), 666–674. doi :10.1016/0019-1035(78)90053-2
- Korablev et al. (2018) *Space Sci. Rev.* 214, 7 doi: 10.1007/s11214-017-0437-6.
- Lefèvre and Krasnopolsky (2017). In “The Atmosphere and Climate of Mars” (pp. 405–432) doi:10.1017/9781139060172.013
- Lefevre et al. (2004), *JGR* 109, E07004, doi:10.1029/2004JE002268.
- Lo, D. Y. et al., 2022, LPSC #2178.
- Maurice et al. (2021). *Space Sci Rev* 217, 47 doi:10.1007/s11214-021-00807-w.
- McConnochie et al. (2018) *Icarus* 307, 294–326, doi:10.1016/j.icarus.2017.10.043.
- Oyama and Berdahl (1977). *JGR*, 82(28), 4669–4676. doi:10.1029/JS082i028p04669
- Smith et al. (2018). *Icarus* 301, 117–131, doi:10.1016/j.icarus.2017.09.027.
- Wiens et al. (2021). *Space Sci Rev* 217, 4 doi:10.1007/s11214-020-00777-5
- Wiens et al. (2012). *Space Sci. Rev.* 170, 167–227. 10.1007/s11214-012-9902-4.