



HAL
open science

Incidence of H I 21-cm absorption in strong Fe II systems at $0.5 < z < 1.5$

R. Dutta, R. Srianand, N. Gupta, R. Joshi, P. Petitjean, P. Noterdaeme, J. Ge, J. -K. Krogager

► **To cite this version:**

R. Dutta, R. Srianand, N. Gupta, R. Joshi, P. Petitjean, et al.. Incidence of H I 21-cm absorption in strong Fe II systems at $0.5 < z < 1.5$. Monthly Notices of the Royal Astronomical Society, 2017, 465, pp.4249-4264. 10.1093/mnras/stw3040 . insu-03747478

HAL Id: insu-03747478

<https://insu.hal.science/insu-03747478>

Submitted on 8 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Incidence of H I 21-cm absorption in strong Fe II systems at $0.5 < z < 1.5$

R. Dutta,¹★ R. Srianand,¹★ N. Gupta,¹★ R. Joshi,¹ P. Petitjean,² P. Noterdaeme,²
J. Ge³ and J.-K. Krogager^{2,4}

¹Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune 411007, India

²Institut d'Astrophysique de Paris, CNRS and UPMC Paris 6, UMR7095, 98bis boulevard Arago, F-75014 Paris, France

³Astronomy Department, University of Florida, 211 Bryant Space Science Center, PO Box 112055, Gainesville, FL 32611, USA

⁴Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen Ø, Denmark

Accepted 2016 November 21. Received 2016 November 17; in original form 2016 September 29

ABSTRACT

We present the results from our search for H I 21-cm absorption in a sample of 16 strong Fe II systems [$W_r(\text{Mg II } \lambda 2796) \geq 1.0 \text{ \AA}$ and $W_r(\text{Fe II } \lambda 2600)$ or $W_{\text{Fe II}} \geq 1 \text{ \AA}$] at $0.5 < z < 1.5$ using the Giant Metrewave Radio Telescope and the Green Bank Telescope. We report six new H I 21-cm absorption detections from our sample, which have increased the known number of detections in strong Mg II systems at this redshift range by ~ 50 per cent. Combining our measurements with those in the literature, we find that the detection rate of H I 21-cm absorption increases with $W_{\text{Fe II}}$, being four times higher in systems with $W_{\text{Fe II}} \geq 1 \text{ \AA}$ compared to systems with $W_{\text{Fe II}} < 1 \text{ \AA}$. The $N(\text{H I})$ associated with the H I 21-cm absorbers would be $\geq 2 \times 10^{20} \text{ cm}^{-2}$, assuming a spin temperature of $\sim 500 \text{ K}$ (based on H I 21-cm absorption measurements of damped Lyman α systems at this redshift range) and unit covering factor. We find that H I 21-cm absorption arises on an average in systems with stronger metal absorption. We also find that quasars with H I 21-cm absorption detected towards them have systematically higher $E(B - V)$ values than those which do not. Further, by comparing the velocity widths of H I 21-cm absorption lines detected in absorption- and galaxy-selected samples, we find that they show an increasing trend (significant at 3.8σ) with redshift at $z < 3.5$, which could imply that the absorption originates from more massive galaxy haloes at high z . Increasing the number of H I 21-cm absorption detections at these redshifts is important to confirm various trends noted here with higher statistical significance.

Key words: galaxies: ISM – quasars: absorption lines.

1 INTRODUCTION

The resonant absorption lines of Mg II detected towards background quasars have proved to be excellent probes of the gaseous haloes and circumgalactic medium (CGM) of $z \lesssim 2$ galaxies in a luminosity-unbiased way (Lanzetta, Turnshek & Wolfe 1987; Sargent, Steidel & Boksenberg 1988; Petitjean & Bergeron 1990; Bergeron & Boissé 1991; Steidel & Sargent 1992; Srianand & Khare 1994; Steidel 1995; Nestor, Turnshek & Rao 2005; Prochter, Prochaska & Burles 2006; Quider et al. 2011; Zhu & Ménard 2013). While weak Mg II absorbers (i.e. absorbers having rest equivalent width of $\text{Mg II } \lambda 2796$, $W_{\text{Mg II}} < 1.0 \text{ \AA}$) could be tracing co-planar accreting or recycled gas (e.g. Chen et al. 2010b,a; Kacprzak & Churchill 2011; Kacprzak et al. 2011; Lovegrove & Simcoe 2011), strong Mg II absorbers (i.e. absorbers having $W_{\text{Mg II}} \geq 1.0 \text{ \AA}$) are believed to be

tracing galactic winds or outflows (e.g. Bouché et al. 2007; Zibetti et al. 2007; Gauthier, Chen & Tinker 2009; Lundgren et al. 2009; Weiner et al. 2009; Noterdaeme, Srianand & Mohan 2010; Rubin et al. 2010; Ménard et al. 2011; Nestor et al. 2011). Strong Mg II absorbers at $z < 1.65$ have been shown to trace gas with high neutral hydrogen column densities (Rao, Turnshek & Nestor 2006, hereafter R06), like damped Lyman α systems (DLAs; $N(\text{H I}) \geq 2 \times 10^{20} \text{ cm}^{-2}$; see Wolfe, Gawiser & Prochaska 2005, for a review). Further, the incidence of DLAs among Mg II absorbers is found to increase at high z ($2 < z < 6$; Matejek et al. 2013). Murphy et al. (2007) have found a correlation between metallicity and $W_{\text{Mg II}}$ in $z < 2.6$ DLAs, similar to the velocity–metallicity relation seen in $z \geq 1.8$ DLAs (Ledoux et al. 2006; Prochaska et al. 2008a).

The strong Mg II systems in the Mg II absorber–galaxy catalogue of Nielsen et al. (2013) sample a wide range of galaxy impact parameters, i.e. over ~ 10 – 200 kpc . Hence, such strong Mg II absorbers can trace gas in a wide variety of environments like star-forming discs, CGM, galactic winds and outflows. It is likely that $W_{\text{Mg II}}$ is dominated by high column density when the sightlines

* E-mail: rddutta@iucaa.in (RD); anand@iucaa.in (RS); ngupta@iucaa.in (NG)

probe the galactic discs and by velocity spread of the absorbing gas in the other scenarios. If one is interested in studying the cold dense gas around galaxies, other parameters like equivalent width ratios of metal lines are required to select sightlines that probe low impact parameters. R06 have demonstrated that equivalent width ratios of Mg II, Mg I and Fe II absorption can be used to pre-select DLAs more successfully than by just using $W_{\text{Mg II}}$. They detect DLAs with a success rate of ~ 42 per cent by selecting Mg II absorbers with $W_r(\text{Mg II } \lambda 2796)/W_r(\text{Fe II } \lambda 2600) < 2$ and $W_r(\text{Mg I } \lambda 2852) > 0.1 \text{ \AA}$.

Further insights into the origin and physical conditions prevailing in the strong Mg II systems and DLAs can be obtained by studying their associated H I 21-cm absorption. H I 21-cm absorption is an excellent tracer of the cold neutral medium (CNM; $T \sim \text{few } 100 \text{ K}$) of galaxies (Kulkarni & Heiles 1988), and has been observed to arise at small impact parameters ($< 30 \text{ kpc}$) from $z < 0.4$ galaxies (Dutta et al. 2016). There have been numerous searches for H I 21-cm absorption in Mg II systems and DLAs (e.g. Briggs & Wolfe 1983; Lane 2000; Kanekar & Chengalur 2003; Curran et al. 2005; Gupta et al. 2009; Kanekar et al. 2009; Curran et al. 2010; Gupta et al. 2012; Srianand et al. 2012; Kanekar et al. 2014). Gupta et al. (2009, 2012, hereafter G09 and G12, respectively) have shown that the H I 21-cm detection rate in strong Mg II systems can be enhanced with appropriate equivalent width ratio cuts of Mg II, Fe II and Mg I.

In addition, searches of H I 21-cm absorption in samples of Mg II systems and DLAs can be used to trace the redshift evolution of the CNM fraction in galaxies. Spin temperature (T_s) measurements and upper limits, derived using H I 21-cm optical depth and $N(\text{H I})$ measured from DLAs, suggest that most of the gas along these sightlines traces the diffuse warm neutral medium (WNM; $T \sim 10^4 \text{ K}$), and a small fraction of the total $N(\text{H I})$ along these sightlines is associated with the CNM (Srianand et al. 2012; Kanekar et al. 2014). This is also supported by observations of typically low overall molecular fractions of H_2 in $z > 1.8$ DLAs (Petitjean, Srianand & Ledoux 2000; Ledoux, Petitjean & Srianand 2003; Srianand et al. 2005; Noterdaeme et al. 2008), and physical conditions inferred in DLAs using C II* and Si II* fine structure lines (Neeleman, Prochaska & Wolfe 2015). The above studies have indicated that the CNM covering factor in DLAs and Mg II systems may be small (~ 10 – 20 per cent). On the other hand, Heiles & Troland (2003), using H I 21-cm emission and absorption measurements, have found that ~ 40 per cent of the H I gas is in the CNM in the Milky Way (MW). Since the volume filling factor and the physical conditions of the CNM depend on feedback processes related to the in situ star formation (McKee & Ostriker 1977; Wolfire et al. 1995; de Avillez & Breitschwerdt 2004; Gent et al. 2013; Gatto et al. 2015), understanding the redshift evolution of H I 21-cm absorbers can shed light on the star formation history of galaxies as well.

In this work, we wish to explore the connection between Mg II systems, DLAs and cold H I gas by searching for H I 21-cm absorption in a sample of strong Mg II systems that satisfy additional constraints to increase the efficiency of detecting cold gas. Systems showing Si II or C I absorption would have been excellent targets to search for cold gas, since strong Si II absorption is likely to select high-metallicity gas (Prochaska et al. 2008a; Jorgenson, Murphy & Thompson 2013), whereas C I absorption is likely to probe dense molecular regions (Noterdaeme et al. 2011, 2016). However, the strong transitions of Si II $\lambda 1526$ and C I $\lambda 1656$ are not covered simultaneously with the Mg II doublet lines of the same system for a large redshift range. On the other hand, similar to the Mg II doublet, strong Fe II lines, like the $\lambda 2600 \text{ \AA}$ transition, can be observed with ground-based telescopes at $0.4 < z < 2.5$.

Strong Fe II systems can arise from either very high metallicity sub-DLAs or high $N(\text{H I})$ DLAs (Srianand 1996). The H I 21-cm optical depth is directly proportional to $N(\text{H I})$ and inversely to T_s . There are indications for a weak anticorrelation between T_s and the gas phase metallicity (see Kanekar et al. 2014, and references therein). Further, strong absorption from Mg II ($W_{\text{Mg II}} \geq 1 \text{ \AA}$) and Fe II (rest equivalent width of Fe II $\lambda 2600$, $W_{\text{Fe II}} \geq 1 \text{ \AA}$) are usually detected whenever galaxy nebular emission lines are directly detected in the Sloan Digital Sky Survey (SDSS; York et al. 2000) quasar spectra, indicating that impact parameters are $\lesssim 10 \text{ kpc}$ (Noterdaeme et al. 2010, Joshi et al., in preparation). We also notice that ~ 77 per cent of the Mg II systems showing signatures of 2175 \AA bump studied by Jiang et al. (2011) have $W_{\text{Fe II}} \geq 1 \text{ \AA}$. Hence, systems with strong Fe II absorption could provide ideal targets for detecting high metallicity and/or high $N(\text{H I})$ cold gas.

This paper is structured as follows. In Section 2, we describe the motivation behind our sample selection and give details of our sample. In Section 3, we describe our radio observations using the Giant Metrewave Radio Telescope (GMRT) and the Green Bank Telescope (GBT). The results from our search for H I 21-cm absorption are presented in Section 4. The detection rate of H I 21-cm absorption in strong Fe II systems is discussed in Section 5. The connection of H I 21-cm absorption with metal line properties and dust content are discussed in Sections 6 and 7, respectively. Finally, we study the velocity width of the H I 21-cm absorption lines in Section 8. We summarize our results in Section 9. Throughout this work we have adopted a flat Λ cold dark matter cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Omega_M = 0.30$.

2 SAMPLE SELECTION

While Mg II absorption lines at $0.3 < z < 2.3$ can be observed using ground-based optical telescopes, the atmospheric cutoff of light below 3000 \AA restricts ground-based observations of H I Lyman α lines at $z < 1.5$. However, ultraviolet (UV) spectroscopic observations with the *Hubble Space Telescope* (*HST*) make it possible to detect $z < 1.65$ DLAs and sub-DLAs (R06, Rao et al. 2011; Meiring et al. 2011; Battisti et al. 2012; Turnshek et al. 2015; Neeleman et al. 2016). R06 have obtained UV observations of Lyman α absorption in 197 Mg II absorbers with $W_{\text{Mg II}} \geq 0.3 \text{ \AA}$ at $z < 1.65$, using the *HST*. Using their table 1, we plot the cumulative distribution of $N(\text{H I})$ in absorbers with $W_{\text{Mg II}} \geq 1 \text{ \AA}$ in the left-hand panel of Fig. 1. We also show the same for systems with both $W_{\text{Mg II}} \geq 1 \text{ \AA}$ and $W_{\text{Fe II}} \geq 1 \text{ \AA}$, and for those with $W_{\text{Mg II}} \geq 1 \text{ \AA}$ but $W_{\text{Fe II}} < 1 \text{ \AA}$.¹ It can be seen that 32 per cent of the Mg II absorbers with $W_{\text{Mg II}} \geq 1.0 \text{ \AA}$ have $N(\text{H I}) \geq 2 \times 10^{20} \text{ cm}^{-2}$. This fraction becomes 45 per cent when the strong Mg II absorbers also have $W_{\text{Fe II}} \geq 1.0 \text{ \AA}$, i.e. strong Fe II absorbers have a higher (by a factor of ~ 1.4) probability of being DLAs. For comparison only 8 per cent of the absorbers with $W_{\text{Mg II}} \geq 1.0 \text{ \AA}$ and $W_{\text{Fe II}} < 1.0 \text{ \AA}$ are DLAs. We note that the success rate of a $W_{\text{Fe II}} \geq 1.0 \text{ \AA}$ selection in detecting DLAs is consistent with that of a $W_{\text{Mg II}}/W_{\text{Fe II}} < 2$ selection suggested by R06 (see Section 1).

Next, from the SDSS Mg II absorber catalogue of Zhu & Ménard (2013), we identified 741 Mg II absorbers at $2.05 < z < 2.15$ with $W_{\text{Mg II}} \geq 1.0 \text{ \AA}$, which have been searched for Lyman α absorption by Noterdaeme et al. (2012). In the right-hand panel of Fig. 1, we show

¹ For the cumulative distribution of systems with $W_{\text{Mg II}} \geq 1 \text{ \AA}$ and $W_{\text{Fe II}} < 1 \text{ \AA}$, we use the sub-samples 1 and 2 of R06, which were selected purely based on $W_{\text{Mg II}}$ without any constraint on $W_{\text{Fe II}}$ (see R06 for details), to avoid biases.




Figure 1. The cumulative distribution of $\log N(\text{H I})$ in systems with $W_{\text{Mg II}} \geq 1 \text{ \AA}$ (solid), those with both $W_{\text{Mg II}} \geq 1 \text{ \AA}$ and $W_{\text{Fe II}} \geq 1 \text{ \AA}$ (dashed), and those with $W_{\text{Mg II}} \geq 1 \text{ \AA}$ and $W_{\text{Fe II}} < 1 \text{ \AA}$ (dotted). The left-hand panel shows the distribution for systems at $z < 1.65$ with $N(\text{H I})$ measurements from R06 using *HST*. The right-hand panel shows the distribution for systems at $2.05 < z < 2.15$ from Zhu & Ménard (2013) with $N(\text{H I})$ measurements from Noterdaeme et al. (2012) using SDSS. The vertical line in each panel marks $N(\text{H I}) = 2 \times 10^{20} \text{ cm}^{-2}$.

the cumulative distributions of $N(\text{H I})$ in these systems, for the same three equivalent width cuts of Mg II and Fe II as described above. Note that Noterdaeme et al. (2012) provides $N(\text{H I})$ measurements only for systems having $N(\text{H I}) \geq 10^{20} \text{ cm}^{-2}$. Hence, we show the cumulative distributions starting at $N(\text{H I}) = 10^{20} \text{ cm}^{-2}$. Similar to above, we find that an additional cut of $W_{\text{Fe II}} \geq 1 \text{ \AA}$ increases the probability of selecting DLAs by a factor of ~ 1.7 compared to just a $W_{\text{Mg II}} \geq 1 \text{ \AA}$ cut. It can be seen that 57 per cent of the strong Fe II absorbers are DLAs, compared to 34 per cent of the absorbers with just $W_{\text{Mg II}} \geq 1.0 \text{ \AA}$. Further, only 19 per cent of the absorbers with $W_{\text{Mg II}} \geq 1.0 \text{ \AA}$ and $W_{\text{Fe II}} < 1.0 \text{ \AA}$ are DLAs.

From the above discussions it seems that a selection technique based on strong Fe II absorption has higher (by a factor of ~ 1.4 – 1.7) probability of detecting DLAs than that based solely on Mg II absorption strength. Though, as can be seen from Fig. 1, for a limiting rest equivalent width of Fe II, the DLA selection efficiency appears to decrease slightly with decreasing redshift. Similarly, Matejek et al. (2013), using infrared spectra of high- z quasars, have found that the fraction of DLAs in Mg II systems with $W_{\text{Mg II}} \geq 0.3 \text{ \AA}$ increases from ~ 17 per cent at $z < 2$ to ~ 41 per cent at $z \geq 2$. The metallicity in DLAs is known to decline with increasing redshift (e.g. Rafelski et al. 2014). Hence, for a given equivalent width of Mg II or Fe II, the same absorber is likely to arise from a higher $N(\text{H I})$ system at high z compared to at low z . This could be a possible explanation for a larger fraction of the strong Fe II/Mg II absorbers being DLAs at higher redshifts.

We set out to quantify the H I 21-cm absorption detection efficiency in such a strong Fe II-selected sample. Our sample of strong Fe II absorbers is drawn from the Mg II catalogue of Zhu & Ménard (2013) constructed using the SDSS Data Release 12 (DR12; Alam et al. 2015). We selected all the absorbers with $W_{\text{Mg II}} \geq 1.0$ and $W_{\text{Fe II}} \geq 1.0$, over the redshift ranges $0.5 < z < 1.0$ and $1.1 < z < 1.5$. Next, by cross-correlating the positions of the background quasars with radio sources within ~ 1 arcsec radius in the Faint Images of the Radio Sky at Twenty-Centimeters (FIRST; White et al. 1997) survey, we identified 17 and 18 strong Fe II systems at $0.5 < z < 1.0$ and $1.1 < z < 1.5$, respectively, that are in front of quasars with peak flux density at 1.4 GHz greater than 100 mJy. The redshift ranges of $0.5 < z < 1.0$ and $1.1 < z < 1.5$ were chosen to enable observations with the GBT prime-focus PF1 800 MHz receiver and

the GMRT 610 MHz receiver, respectively (see Section 3). The flux density cut of 100 mJy was used in order to achieve 3σ integrated H I 21-cm optical depth sensitivity (for a line width of 10 km s^{-1}) $\lesssim 0.3 \text{ km s}^{-1}$, which is sensitive to detect 100 K gas with $N(\text{H I}) \leq 5 \times 10^{19} \text{ cm}^{-2}$.

We visually inspected the optical spectra and radio images of the quasars to ensure that there are no false identifications. Among the 35 strong Fe II absorbers, 11 are part of Mg II samples that have been searched for H I 21-cm absorption in the literature – 4 in G09, 3 in G12, 3 in Kanekar et al. (2009, hereafter K09) and 1 in Lane (2000, hereafter L00). From the remaining 24 systems, we selected 15 strong Fe II absorbers, whose redshifted H I 21-cm frequencies fall in the radio frequency interference free observing bands of GBT and GMRT. In addition, we included the $z_{\text{abs}} = 1.27384$ Mg II absorber towards J091927.61+014603.0. Note that this system is not in the catalogue of Zhu & Ménard (2013) and the peak flux density of the radio source from FIRST is less than 100 mJy (i.e. 59 mJy). But this system satisfies our criterion of $W_{\text{Mg II}} \geq 1.0$ and $W_{\text{Fe II}} \geq 1.0$, and has been identified as a 2175 \AA absorber candidate by Jiang et al. (2011). Hence, our sample consists of 16 absorbers, of which 7 (at $0.5 < z < 1.0$) were observed with GBT, and 9 (at $1.1 < z < 1.5$) were observed with GMRT. The details of these 16 absorbers are presented in Table 1. The median values of various metal line parameters for our observed sample are: $W_{\text{Mg II}} = 2.23 \text{ \AA}$; $W_{\text{Fe II}} = 1.55 \text{ \AA}$; Mg II doublet ratio, DR = 1.14; $W_r(\text{Mg I } \lambda 2852)/W_{\text{Mg II}}$, R1 = 0.27; $W_{\text{Mg II}}/W_{\text{Fe II}}$, R2 = 1.58.

The 1.4 GHz peak flux densities obtained by the FIRST survey of the radio sources in our sample are listed in Table 2. All the sources except one are compact (deconvolved sizes ≤ 2 arcsec; White et al. 1997) in the FIRST images (resolution ~ 5 arcsec). The radio source J1510+1640 is resolved in the FIRST image, with a deconvolved size of $2.9 \text{ arcsec} \times 1.3 \text{ arcsec}$, and with the peak emission accounting for 86 per cent of the total emission. Low-frequency (≤ 1.4 GHz) sub-arcsecond-scale images are not available for any of the sources in our sample. However, 2.3 GHz Very Long Baseline Array (VLBA) images are available for eight of the sources from the VLBA Calibrator Survey (VCS).² We assume that the covering

² <http://www.vlba.nrao.edu/astro/calib/>

Table 1. Sample of strong Fe II systems.

Quasar name (1)	SDSS name (2)	z_{em} (3)	z_{abs} (4)	$W_r(\text{Mg II } \lambda 2796)$ (\AA) (5)	$W_r(\text{Mg II } \lambda 2803)$ (\AA) (6)	$W_r(\text{Mg I } \lambda 2852)$ (\AA) (7)	$W_r(\text{Fe II } \lambda 2600)$ (\AA) (8)
J0919+0146	J091927.61+014603.0	1.286	1.2738	2.70 ± 0.15	1.99 ± 0.15	0.27 ± 0.13	2.24 ± 0.28
J0921+6215	J092136.22+621552.5	1.447	1.1035	4.11 ± 0.10	3.62 ± 0.13	0.98 ± 0.10	2.86 ± 0.09
J0952+5048	J095227.30+504850.6	1.091	0.9985	2.10 ± 0.04	1.94 ± 0.04	0.40 ± 0.05	1.30 ± 0.04
J1111+3252	J111131.77+325255.8	2.362	1.4753	2.23 ± 0.13	2.35 ± 0.14	0.96 ± 0.10	1.55 ± 0.12
J1111+4507	J111114.81+450735.0	1.486	0.6865	1.83 ± 0.14	1.93 ± 0.14	0.83 ± 0.15	1.73 ± 0.16
J1201+1114	J120140.25+111447.6	2.296	1.4846	2.66 ± 0.14	2.40 ± 0.13	0.31 ± 0.10	1.57 ± 0.12
J1241+6020	J124129.57+602041.3	2.066	1.2379	4.73 ± 0.06	4.48 ± 0.06	1.34 ± 0.07	3.38 ± 0.08
J1245+2232	J124557.77+223205.3	1.478	0.5808	1.64 ± 0.05	1.40 ± 0.05	0.16 ± 0.04	1.14 ± 0.05
J1255+1817	J125531.75+181750.9	1.371	0.7580	2.20 ± 0.18	1.88 ± 0.18	0.71 ± 0.17	1.23 ± 0.18
J1327+4326	J132720.97+432627.9	2.086	0.9542	2.42 ± 0.12	2.09 ± 0.10	0.47 ± 0.16	1.53 ± 0.11
J1342+5110	J134224.31+511012.4	2.598	1.4880	3.18 ± 0.07	2.92 ± 0.07	1.06 ± 0.07	1.74 ± 0.06
J1351+0830	J135116.91+083039.8	1.442	1.4270	2.59 ± 0.05	2.09 ± 0.05	0.85 ± 0.08	1.43 ± 0.07
J1504+2854	J150426.69+285430.5	2.285	1.2208	1.87 ± 0.07	1.58 ± 0.07	0.50 ± 0.08	1.06 ± 0.07
J1510+1640	J151038.63+164010.2	1.827	1.2139	2.22 ± 0.09	1.70 ± 0.08	0.34 ± 0.09	1.29 ± 0.09
J2219+0229	J221930.79+022945.4	2.218	0.9804	2.19 ± 0.12	2.08 ± 0.12	0.64 ± 0.12	1.78 ± 0.13
J2330+1100	J233040.84+110018.6	1.502	0.9488	1.55 ± 0.05	1.32 ± 0.05	0.40 ± 0.06	1.07 ± 0.05

Notes. Column 1: quasar name used in this work. Column 2: SDSS name (J2000) of the quasar. Column 3: redshift of quasar. Column 4: redshift of intervening Fe II system. Columns 5–8: rest frame equivalent widths (in \AA) of the Mg II $\lambda 2796$, Mg II $\lambda 2803$, Mg I $\lambda 2852$ and Fe II $\lambda 2600$ absorption lines, respectively, taken from Zhu & Ménard (2013) (except in the case of J0919+0146, where we measure these ourselves).

Table 2. Details of the radio sources.

Quasar (1)	1.4-GHz peak flux density (mJy beam $^{-1}$) (2)	Morph. (3)	C_f (4)
J0919+0146	59	C	–
J0921+6215	1213	C	0.60
J0952+5048	105	C	0.71
J1111+3252	107	C	–
J1111+4507	239	C	–
J1201+1114	161	C	–
J1241+6020	433	C	0.88
J1245+2232	131	C	–
J1255+1817	366	C	0.36
J1327+4326	589	C	0.91
J1342+5110	144	C	–
J1351+0830	335	C	0.92
J1504+2854	549	C	0.60
J1510+1640	138	R	–
J2219+0229	183	C	–
J2330+1100	1174	C	0.59

Notes. Column 1: quasar name. Column 2: FIRST 1.4-GHz peak flux density in mJy beam $^{-1}$. Column 3: morphology from FIRST image (~ 5 arcsec resolution), where C is for compact and R is for resolved. Column 4: covering factor (assumed to be the core fraction) determined from 2.3 GHz VLBA images (see Section 2 for details).

factor (C_f) of the absorbing gas is equal to the core fraction, i.e. we assume that the gas covers only the core component seen in VLBA images. Hence, we estimate C_f from the ratio of the peak flux density in the VCS image to the total arcsecond-scale flux density (see Table 2). The total arcsecond-scale flux densities at 2.3 GHz have been estimated by interpolating flux densities available from the NASA extragalactic data base³ (Gregory & Condon 1991; Condon et al. 1998).

³ <https://ned.ipac.caltech.edu/>

3 RADIO OBSERVATIONS

The seven systems at $0.5 < z < 1.0$ were observed with the prime-focus PF1 800 MHz receiver at GBT (Proposal ID: 16A–141). We used Versatile GBT Astronomical Spectrometer as the back end with a bandwidth of 11.72 MHz split into 32 768 channels (spectral resolution of ~ 0.1 km s $^{-1}$ per channel, velocity coverage ~ 4000 km s $^{-1}$). The observations were carried out in standard position switching mode, with ~ 2 min spent on-source and same amount of time spent at the off-source position. A dump time of 2 s was used. The data were acquired in two linear polarization products, XX and YY. The nine systems at $1.1 < z < 1.5$ were observed with the 610 MHz receiver at GMRT, using the 2 MHz baseband bandwidth split into 512 channels (spectral resolution ~ 2 km s $^{-1}$ per channel, velocity coverage ~ 1000 km s $^{-1}$). Data were acquired in two polarization products, LL and RR. Standard calibrators were regularly observed during the observations for flux density, bandpass, and phase calibrations. In both the GBT and the GMRT observations, the pointing centre was at the quasar coordinates and the band was centred at the redshifted H I 21-cm frequency.

The GMRT data were reduced using the National Radio Astronomy Observatory (NRAO) Astronomical Image Processing System (AIPS) following standard procedures as described in Gupta et al. (2010). The GBT data were reduced using a pipeline developed using NRAO's GBTIDL package as described in G12. The details of the radio observations of the sources are given in Table 3.

All the nine radio sources observed with GMRT, except J1510+1640, are compact in our GMRT images. Note that the sources, J1111+3252 (deconvolved size ~ 2.2 arcsec \times 0.2 arcsec) and J1351+0830 (deconvolved size ~ 2.0 arcsec \times 0.9 arcsec), are considered as compact since >90 per cent of the total emission is contained in a single Gaussian component. In case of J1510+1640, the synthesized beam of our GMRT image is ~ 6.2 arcsec \times 3.9 arcsec, and the radio emission can be represented by a single Gaussian component with deconvolved size ~ 2.9 arcsec \times 0.5 arcsec, which contains ~ 81 per cent of the total emission, similar to the FIRST image. Note that in all the GMRT data cubes, the H I 21-cm

Table 3. Radio observation log.

Quasar	Date	Time	$\delta\nu$
(1)	(2)	(h)	(km s ⁻¹)
		(3)	(4)
GBT			
J0952+5048	2016 February 18	1.7	0.2
	2016 March 13	0.7	0.2
	2016 May 21	0.7	0.2
J1111+4507	2016 February 24	1.0	0.1
	2016 March 11	0.3	0.1
J1245+2232	2016 February 18	0.8	0.1
J1255+1817	2016 February 18	0.6	0.1
J1327+4326	2016 February 24	0.5	0.1
	2016 March 11	0.2	0.1
	2016 March 13	0.6	0.1
J2219+0229	2016 February 23	0.5	0.1
	2016 March 11	0.4	0.1
J2330+1100	2016 February 23	0.3	0.1
GMRT			
J0919+0146	2015 November 24	6.4	2.0
	2016 June 12	4.2	2.0
J0921+6215	2015 November 25	0.8	1.8
J1111+3252	2015 October 31	5.5	2.1
J1201+1114	2015 December 14	5.1	2.1
J1241+6020	2015 November 22	3.2	1.9
J1342+5110	2015 November 21	5.2	2.1
	2016 April 26	5.4	2.1
J1351+0830	2016 January 22	3.2	2.1
J1504+2854	2015 November 25	3.2	1.9
J1510+1640	2016 January 4	2.9	1.9
	2016 January 5	3.2	1.9

Notes. Column 1: quasar name. Column 2: date of observation. Column 3: time on source in h. Column 4: channel width in km s⁻¹.

absorption spectra were extracted at the location of the continuum peak flux density.

In case of seven radio sources observed with GBT, we obtained the flux density of five of them from the GBT spectrum. For two sources, J1255+1817 and J2219+0229, the flux scale could not be calibrated properly in our GBT observations. For these sources, the flux densities at the redshifted H I 21-cm frequency have been estimated by interpolating between the flux densities from the FIRST and the TEXAS (Douglas et al. 1996) surveys at 1.4 GHz and 365 MHz, respectively. The flux densities at the redshifted H I 21-cm frequency of all the radio sources observed by us are given in Table 4.

4 PARAMETERS DERIVED FROM H I 21-cm ABSORPTION SPECTRA

The results from our search for H I 21-cm absorption in the 16 strong Fe II absorbers are summarized in Table 4. We have detected H I 21-cm absorption in 6 out of these 16 strong Fe II systems. We provide the standard deviation in the optical depth at ~ 2 km s⁻¹ spectral resolution (σ_τ), and 3σ upper limit on the integrated optical depth from spectra smoothed to 10 km s⁻¹ ($\int \tau dv_{10}^{3\sigma}$). In the case of H I 21-cm absorption detections, we provide the peak optical depth (τ_p) and the total integrated optical depth ($\int \tau dv$). From Kanekar et al. (2014), we see that the mean and median T_s in DLAs over $0.5 < z < 1.5$ are 670 and 460 K, respectively. We estimate $N(\text{H I})$ from $\int \tau dv$ of the detections, or 3σ upper limit to it from $\int \tau dv_{10}^{3\sigma}$ in the case of non-detections, assuming a typical T_s of 500 K and $C_f = 1$. The

$N(\text{H I})$ associated with the H I 21-cm absorption will be $\geq 2 \times 10^{20}$ cm⁻² under these assumptions, i.e. these systems will satisfy the definition of being DLAs. The optical depth sensitivity in 8 out of the 10 cases of H I 21-cm non-detection rules out the system being DLA if $T_s = 500$ K and $C_f = 1$. Table 4 also gives the velocity width which contains 90 per cent of the total optical depth (v_{90}), and the velocity offset of the peak H I 21-cm optical depth from the strongest metal component in the SDSS spectrum (v_{off}). The H I 21-cm absorption lines show a wide range of velocity spreads, with $v_{90} \sim 13\text{--}158$ km s⁻¹, and are detected within ± 100 km s⁻¹ of the redshift obtained from metal line absorption in the SDSS spectra.

The H I 21-cm absorption spectra are shown in Fig. 2. We note that tentative absorption features are present towards J1241+6020 and J2330+1100 at the redshift of the metal absorption. However, these features are not present consistently in both the polarizations and are at $\leq 2\sigma$ significance level. Hence, we consider these as non-detections. The Gaussian fits to the H I 21-cm absorption profiles are overplotted in Fig. 2. The number of Gaussian components is determined based on the fit with the minimum χ^2 . The details of the Gaussian fits, i.e. z_{abs} , full width at half-maximum (FWHM) and τ_p of individual Gaussian components, are provided in Table 5. Constraints on the kinetic temperature, T_k , using the FWHM of the absorption lines, i.e.

$$T_k < 21.855 \times \text{FWHM}^2 \text{ K}, \quad (1)$$

and on the $N(\text{H I})$, i.e.

$$N(\text{H I}) = 1.93 \times 10^{18} \tau_p \frac{T_s}{C_f} \text{FWHM cm}^{-2}, \quad (2)$$

are provided in the same table. The FWHMs of the absorption components range from $\sim 5\text{--}131$ km s⁻¹. The large line widths are most likely dominated by mechanisms other than pure thermal broadening, since even the typical temperature ($\sim 10^4$ K) in WNM corresponds to an FWHM of ~ 20 km s⁻¹ (Wolfire et al. 1995; Heiles & Troland 2003). Assuming a typical T_s of 500 K and $C_f = 1$, we find that all the H I 21-cm absorption components will be arising from DLAs.

5 INCIDENCE OF H I 21-cm ABSORPTION IN STRONG Fe II SYSTEMS

We define the detection rate of H I 21-cm absorption (C_{21}) as the fraction of systems showing H I 21-cm detections with $\int \tau dv_{10}^{3\sigma} \leq \tau_0$ and $\int \tau dv \geq \tau_0$, where τ_0 is a 3σ optical depth sensitivity. We use $\tau_0 = 0.3$ km s⁻¹ (corresponding to a sensitivity of $N(\text{H I}) \leq 5 \times 10^{19}$ cm⁻² for $T_s = 100$ K and $C_f = 1$) throughout this work to estimate C_{21} , unless otherwise mentioned.⁴ The quoted errors represent Gaussian 1σ confidence intervals computed using tables of Gehrels (1986) assuming a Poisson distribution. Note that due to lack of uniform measurement of C_f for all the systems, we do not correct the optical depths for C_f in subsequent discussions.

We combine our H I 21-cm measurements with those of strong Mg II systems (i.e. $W_{\text{Mg II}} \geq 1 \text{ \AA}$) at $0.5 < z < 1.5$ from G09 and G12, that satisfy our criterion of being strong Fe II absorbers with $W_{\text{Fe II}} \geq 1.0 \text{ \AA}$. Note that for the absorber at $z_{\text{abs}} = 0.6216$ towards J093035.08+464408.7 from G12, we consider the recent measurement of H I 21-cm absorption (which appeared as a tentative feature

⁴ $\tau_0 = 0.3$ km s⁻¹ is chosen also to enable comparisons with existing H I 21-cm absorption measurements in the literature.

Table 4. Parameters derived from the redshifted H I 21-cm absorption spectra.

Quasar	Peak flux density (mJy beam ⁻¹)	δv (km s ⁻¹)	Spectral rms (mJy beam ⁻¹)	σ_τ	τ_p	$\int \tau dv_{10}^{3\sigma}$ (km s ⁻¹)	$\int \tau dv$ (km s ⁻¹)	$N(\text{H I})$ ($T_s/500$ K) ($1/C_f$) (10^{20} cm ⁻²)	v_{90} (km s ⁻¹)	v_{off} (km s ⁻¹)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
J0919+0146	183	2.0	1.0	0.005	0.02	≤ 0.068	1.61 ± 0.11	15 ± 1	158	-100
J0921+6215	1332	1.8	9.6	0.009	0.05	≤ 0.082	0.40 ± 0.05	4 ± 1	13	13
J0952+5048	155	2.3	2.0	0.013	-	≤ 0.199	-	≤ 1.8	-	-
J1111+3252	239	2.1	3.6	0.015	-	≤ 0.188	-	≤ 1.7	-	-
J1111+4507	184	1.9	2.7	0.015	-	≤ 0.203	-	≤ 1.9	-	-
J1201+1114	109	2.1	3.7	0.034	-	≤ 0.424	-	≤ 3.9	-	-
J1241+6020	259	1.9	2.4	0.009	-	≤ 0.150	-	≤ 1.4	-	-
J1245+2232	180	1.8	2.1	0.012	-	≤ 0.250	-	≤ 2.3	-	-
J1255+1817	856	2.0	6.4	0.007	0.10	≤ 0.095	2.09 ± 0.08	19 ± 1	40	-67
J1327+4326	647	2.3	4.4	0.007	0.02	≤ 0.080	0.36 ± 0.07	3 ± 1	20	-2
J1342+5110	157	2.2	2.7	0.017	0.10	≤ 0.252	3.54 ± 0.23	32 ± 2	56	25
J1351+0830	247	2.1	2.1	0.009	-	≤ 0.129	-	≤ 1.2	-	-
J1504+2854	1082	1.9	2.3	0.002	-	≤ 0.030	-	≤ 0.3	-	-
J1510+1640	257	1.9	0.9	0.003	-	≤ 0.041	-	≤ 0.4	-	-
J2219+0229	200	2.3	3.3	0.016	0.29	≤ 0.300	5.32 ± 0.22	48 ± 2	58	59
J2330+1100	1745	2.2	9.1	0.005	-	≤ 0.066	-	≤ 0.6	-	-

Notes. Column 1: quasar name. Column 2: peak flux density in mJy beam⁻¹ of the local continuum. Column 3: channel width in km s⁻¹. Column 4: spectral rms in mJy beam⁻¹. Column 5: standard deviation of the H I 21-cm optical depth. Column 6: maximum H I 21-cm optical depth in case of H I 21-cm detections. Column 7: 3 σ upper limit on integrated H I 21-cm optical depth from spectra smoothed to 10 km s⁻¹. Column 8: integrated H I 21-cm optical depth in case of H I 21-cm detections. Column 9: $N(\text{H I})$ assuming $T_s = 500$ K and $C_f = 1$, in units of 10^{20} cm⁻² (3 σ upper limit in case of non-detections). Column 10: velocity width that contains 90 per cent of the total optical depth in case of detections. Column 11: velocity offset of the peak H I 21-cm optical depth from the strongest metal component in the SDSS spectrum in case of detections.

Note that the values in Columns 4 and 5 are at the spectral resolution specified in Column 3.

in G12) using deeper observations by Zwaan et al. (2015). The summary of all the strong Fe II absorbers considered here for statistical analyses is given in Table 6. In total, we consider 46 strong Fe II systems out of which 16 show H I 21-cm absorption. We refer to the sample presented in this work (see Table 1) as S1, and the combined sample as S2. Note that sample S2 is homogeneous with regards to analysis of the radio data. Fig. 3 shows the distribution of $W_{\text{Mg II}}$ and $W_{\text{Fe II}}$ in sample S2.

We estimate the detection rate of H I 21-cm absorption in the Fe II samples over $0.5 < z < 1.5$ as, $C_{21}^{\text{Fe II}} = 0.40^{+0.24}_{-0.16}$ ($0.36^{+0.29}_{-0.17}$ for $\tau_0 = 0.2^5$) for S1, and $C_{21}^{\text{Fe II}} = 0.30^{+0.12}_{-0.09}$ ($0.39^{+0.16}_{-0.12}$ for $\tau_0 = 0.2$) for S2. We observe no evolution of $C_{21}^{\text{Fe II}}$ within the uncertainties over the redshift range studied, for both S1 and S2 (see Fig. 4). This is consistent with the constant detection rate of H I 21-cm absorption in strong Mg II systems found by G12. Recall that our sample selection procedure picked three strong Fe II systems that are part of the Mg II sample of K09 and one system that is part of the Mg II sample of L00 (see Section 2). Including these measurements from K09 and L00 in the sample S2 does not change our statistical results within the uncertainties, i.e. $C_{21}^{\text{Fe II}} = 0.27^{+0.11}_{-0.08}$ ($0.39^{+0.15}_{-0.11}$ for $\tau_0 = 0.2$).

To determine the detection efficiency of H I 21-cm absorption of an Fe II equivalent width based selection technique, we plot $C_{21}^{\text{Fe II}}$ as a function of $W_{\text{Fe II}}$ in Fig. 5. For the strong Fe II systems in S2, we plot $C_{21}^{\text{Fe II}}$ in two bins of $W_{\text{Fe II}}$ demarcated at the median value, $W_{\text{Fe II}} = 1.5$ Å. For comparison, we also plot $C_{21}^{\text{Fe II}}$ ($0.08^{+0.18}_{-0.06}$) in strong Mg II systems with $W_{\text{Fe II}} < 1$ Å from G09 and G12. We caution that the strong Mg II systems, that have been searched for H I 21-

cm absorption in the literature, are biased towards high $W_{\text{Fe II}}$, with ~ 65 per cent of them having $W_{\text{Fe II}} \geq 1.0$, compared to 37 per cent of the strong Mg II systems in SDSS at $0.5 < z < 1.5$ from Zhu & Ménard (2013) having $W_{\text{Fe II}} \geq 1.0$. Hence, the number of systems with $W_{\text{Fe II}} < 1$ Å is only one-third of sample S2. We find that the detection efficiency of H I 21-cm absorption in systems with $W_{\text{Fe II}} \geq 1$ Å (sample S2) is a factor of ~ 3.8 times higher than that in systems with $W_{\text{Fe II}} < 1$ Å, albeit not statistically significant due to the large uncertainties. However, the increasing trend of $C_{21}^{\text{Fe II}}$ with $W_{\text{Fe II}}$ is valid over both the redshift ranges, $0.5 < z < 1.0$ and $1.1 < z < 1.5$.

Additionally, we plot in Fig. 5 the cumulative distribution of $C_{21}^{\text{Fe II}}$ as a function of $W_{\text{Fe II}}$, i.e. for $W_{\text{Fe II}} = x$ Å, it gives $C_{21}^{\text{Fe II}}$ in systems with $W_{\text{Fe II}} \geq x$ Å. It can be seen that $C_{21}^{\text{Fe II}}$ increases as we go for samples with higher $W_{\text{Fe II}}$. This increasing trend also holds for different optical depth sensitivities, on correcting the optical depths for C_f , and on considering the additional measurements from K09 and L00. G09 and G12 have found that the detection rate of H I 21-cm absorption can be increased by up to a factor of 2 by imposing constraints on DR, R1 and R2. Here, we have demonstrated that a simple $W_{\text{Fe II}} \geq 1$ Å based selection would work just as well for increasing the probability of detecting H I 21-cm absorption. Additionally, we note that assuming a typical $T_s = 500$ K (see Section 4) and $C_f = 1$, all the 16 H I 21-cm absorption detected in strong Fe II systems would be arising from DLAs. Hence, strong Fe II systems seem to have a high probability of harbouring high $N(\text{H I})$ cold gas.

From the detection rate of H I 21-cm absorption in strong Fe II systems and that of DLAs in strong Fe II systems, we can estimate the detection rate of H I 21-cm absorption in DLAs, i.e. $C_{21}^{\text{DLA}} = C_{21}^{\text{Fe II}} / C_{\text{DLA}}^{\text{Fe II}}$. Using $C_{21}^{\text{Fe II}}$ obtained for S2 (see Fig. 4) and $C_{\text{DLA}}^{\text{Fe II}} = 0.45^{+0.10}_{-0.08}$ obtained for the $z < 1.65$ Mg II sample of R06 (see Section 2), we estimate $C_{21}^{\text{DLA}} = 0.80^{+0.20}_{-0.36}$ at $0.5 < z < 1.0$ and

⁵ Since lower values of τ_0 will select the weaker H I 21-cm absorbers and higher values will select the stronger absorbers, we provide the detection rates for $\tau_0 = 0.2$ for comparison.




Figure 2. H I 21-cm absorption spectra towards the sample of strong Fe II systems, smoothed to $\sim 4 \text{ km s}^{-1}$. In the case of H I 21-cm absorption detections, individual Gaussian components and the resultant fits to the absorption profiles are overplotted as dotted and continuous lines, respectively. The quasar name as given in Table 1 is provided for each spectrum.

Table 5. Details of the Gaussian fits to the H I 21-cm absorption profiles.

Quasar	ID	z_{abs}	FWHM	τ_p	T_k	$N(\text{H I})$
(1)	(2)	(3)	(4)	(5)	(6)	(7)
			(km s^{-1})		(K)	($T_s/500 \text{ K}$) ($1/C_f$)
						(10^{20} cm^{-2})
J0919+0146	1	1.273 08	131 ± 21	0.013 ± 0.004	$<373\ 774$	16 ± 7
J0921+6215	1	1.103 60	9 ± 2	0.04 ± 0.01	<1908	4 ± 2
J1255+1817	1	0.757 61	29 ± 2	0.06 ± 0.01	$<18\ 060$	17 ± 4
	2	0.757 66	5 ± 1	0.05 ± 0.03	<591	3 ± 2
J1327+4326	1	0.954 21	13 ± 7	0.02 ± 0.01	<3600	3 ± 2
	2	1.488 25	39 ± 10	0.06 ± 0.04	$<33\ 987$	22 ± 20
J1342+5110	1	1.488 04	24 ± 5	0.05 ± 0.04	$<12\ 777$	12 ± 10
	2	1.488 25	39 ± 10	0.06 ± 0.04	$<33\ 987$	22 ± 20
J2219+0229	1	0.980 60	13 ± 1	0.10 ± 0.01	<3923	13 ± 3
	2	0.980 75	10 ± 1	0.26 ± 0.02	<2394	26 ± 3
	3	0.980 90	15 ± 2	0.06 ± 0.01	<5023	10 ± 3

Notes. Column 1: quasar name. Column 2: absorption component identification. Column 3: redshift of absorption component. Column 4: FWHM (km s^{-1}) of the Gaussian component. Column 5: peak optical depth of the Gaussian component. Column 6: upper limit on T_k (K), obtained assuming the line width is purely due to thermal motions. Column 7: $N(\text{H I})$ assuming $T_s = 500 \text{ K}$ and $C_f = 1$, in units of 10^{20} cm^{-2} .

Table 6. Summary of strong Fe II absorbers at $0.5 < z < 1.5$ with H I 21-cm absorption measurements considered here

Sample	No. of systems	H I 21-cm detections	Reference
S1	16	6	This work
G09	18	5	Gupta et al. (2009)
G12	12	5 ^a	Gupta et al. (2012)
S2	46	16	S1 + G09 + G12

Note. ^aWith the improved parameters from the H I 21-cm absorption spectrum towards J093035.08+464408.7 reported in Zwaan et al. (2015)




Figure 3. W_{MgII} versus W_{FeII} for the strong Fe II absorbers in our sample are shown as black circles and in the systems from G09 and G12 are shown as red squares. The filled symbols correspond to H I 21-cm detections and the open symbols correspond to non-detections. The typical error bars are plotted in the bottom right corner.




Figure 4. The detection rate of H I 21-cm absorption (for $\tau_0 = 0.3 \text{ km s}^{-1}$) in samples S1 (square) and S2 (diamond) over the two redshift ranges, $0.5 < z < 1.0$ and $1.1 < z < 1.5$ (demarcated by arrows). The dotted and dashed horizontal lines represent the mean C_{21}^{FeII} in samples S1 and S2, respectively, over $0.5 < z < 1.5$ (see Section 5).

$C_{21}^{\text{DLA}} = 0.58^{+0.38}_{-0.25}$ at $1.1 < z < 1.5$. In comparison, the detection rate of H I 21-cm absorption in $z < 1$ DLAs is $0.62^{+0.30}_{-0.21}$ (see Dutta et al. 2016), and that in $2.0 \leq z \leq 3.5$ DLAs is $0.22^{+0.29}_{-0.14}$ (see Srianand et al. 2012). Note that all the detection rates estimates are for $\tau_0 = 0.3 \text{ km s}^{-1}$. Hence, C_{21}^{DLA} appears to be increasing with decreasing redshift, even though the errors are large. This increasing trend of




Figure 5. The detection rate of H I 21-cm absorption (for $\tau_0 = 0.3 \text{ km s}^{-1}$) in three different bins (demarcated by arrows) of W_{FeII} for sample S2. The dotted line is the cumulative distribution of C_{21}^{FeII} , i.e. for $W_{\text{FeII}} = x \text{ \AA}$, it gives C_{21}^{FeII} in systems with $W_{\text{FeII}} \geq x \text{ \AA}$. The C_{21}^{FeII} estimate at $W_{\text{FeII}} < 1 \text{ \AA}$ are using measurements from G09 and G12.

the H I 21-cm absorption detection rate in DLAs with decreasing redshift has also been noted by K09 and G12 (however see also, Curran et al. 2005; Curran & Webb 2006; Curran 2010, 2012). This is similar to the increase in detection rate of H₂ absorption in DLAs/sub-DLAs from ~ 10 per cent at $z \geq 1.8$ (Noterdaeme et al. 2008) to ~ 50 per cent at $z \leq 0.7$ (Muzahid, Srianand & Charlton 2015). In addition, as noted by Dutta et al. (2016), the incidence of H I 21-cm absorption in absorption-selected samples is much higher than that in $z < 0.4$ quasar–galaxy pairs (QGP) that are selected on the basis of galaxies, implying patchy distribution and small sizes (parsec to sub-parsec scale) of cold gas clouds in the extended discs/haloes of galaxies. The increase in incidence of H I 21-cm absorption with decreasing redshift can be explained as a result of the increase in the CNM filling factor in galaxies with time.

6 H I 21-cm ABSORPTION AND METALS

6.1 Metal line properties

The metal line ratios like DR, R1 and R2 are found to be more robust indicators of high $N(\text{H I})$ systems than W_{MgII} alone (R06). Further, G09 and G12 have found that a higher detection rate of H I 21-cm absorption can be obtained in systems limited to restricted ranges of these parameters. Here, we investigate the dependence of H I 21-cm absorption in strong Fe II systems on their metal absorption. For the literature systems, we consider the rest equivalent widths of metal lines as given in G09 and G12. We plot the H I 21-cm absorption strength of the strong Fe II systems in S2 as a function of the metal line equivalent width ratios, DR, R1 and R2 in Fig. 6. We find a positive correlation between $\int \tau dv$ of the H I 21-cm absorption detections and R1: Kendall's rank correlation coefficient, $r_k = 0.38$, with the probability of the correlation arising by chance, $P(r_k) = 0.04$, which is significant at $S(r_k) = 2.0\sigma$ assuming Gaussian statistics; and Spearman rank correlation coefficient, $r_s = 0.49$, $P(r_s) = 0.05$, $S(r_s) = 1.9\sigma$. Since higher R1 values are likely to be arising in systems with high $N(\text{H I})$, this may imply a weak correlation of




Figure 6. H I 21-cm absorption strength of the strong Fe II systems as a function of different metal line equivalent width ratios: (a) DR, (b) R1 and (c) R2. Black circles are for the systems in our sample and red squares are for the systems from G09 and G12. The filled symbols correspond to $\int \tau dv$ of the H I 21-cm detections and the open symbols correspond to $\int \tau dv_{10}^{3\sigma}$ of the non-detections. The typical error bars are plotted in the bottom right corner of each plot.

$\int \tau dv$ with $N(\text{H I})$. However, the correlation becomes less significant (1.4σ) when we include the upper limits ($\int \tau dv_{10}^{3\sigma}$) as censored data points and perform survival analysis using the ‘CENKEN’ function under the ‘NADA’ package in R. We also do not find any correlation of the H I 21-cm absorption strength with $W_{\text{Mg II}}$, $W_{\text{Fe II}}$, DR and R2 (i.e. they are at $\lesssim 1\sigma$ significance). Note that most of the H I 21-cm detections arise in systems with DR ~ 1 , i.e. in the saturated regime.

It is important to keep in mind that the metal line equivalent widths and ratios measured from low-resolution SDSS spectra are averaged over several absorption components. High-resolution optical spectra of Mg II systems show that the metal line absorption usually consists of multiple components, with the number of components increasing with $W_{\text{Mg II}}$ (e.g. Churchill, Vogt & Charlton 2003). Hence, the lack of correlation between the H I 21-cm absorption strength and the metal line parameters estimated from SDSS could be because the H I 21-cm absorption arises from certain specific metal line components (e.g. G09; Rahmani et al. 2012; Srikanth et al. 2012; Dutta et al. 2015). This emphasizes the need for high-resolution optical spectra of the metal absorption lines.

Finally, there are 11 ultrastrong Mg II systems, i.e. those with $W_{\text{Mg II}} \geq 3.0 \text{ \AA}$, in S2. It has been suggested that ultrastrong Mg II absorbers may arise in galactic superwinds (Bond et al. 2001; Nestor et al. 2011). However, their large equivalent widths could also be driven by gas dynamics of the intragroup medium (e.g. Gauthier 2013). Typically ~ 7 per cent of $W_{\text{Mg II}} \geq 1 \text{ \AA}$ absorbers in SDSS satisfy the definition of ultrastrong Mg II absorbers. This fraction becomes ~ 12 per cent when the absorbers also have $W_{\text{Fe II}} \geq 1 \text{ \AA}$. However, ultrastrong Mg II systems constitute ~ 24 per cent of sample S2. H I 21-cm absorption has been detected in four of these. We do not find these absorbers to be different from the rest of the strong Fe II absorbers in terms of their H I 21-cm absorption detection rate, optical depths and velocity widths. While the metal equivalent widths are large in these systems, the velocity widths of the H I 21-cm absorption are in the range of $\sim 12\text{--}60 \text{ km s}^{-1}$.

6.2 Average metal content

To check if there is any difference in the average metal line properties of the systems that show H I 21-cm absorption from those which do not, we stacked the SDSS spectra of the background quasars in sample S2, with and without H I 21-cm absorption separately. The resultant median stacked spectra are shown in panel (a) of Fig. 7,

and the rest equivalent widths of different metal transitions in the stacked spectra are listed in Table 7. We consider the rest wavelength range of 2000–3000 \AA , which is uniformly covered for all the systems. It can be seen that the systems which show H I 21-cm absorption, also show systematically stronger absorption (i.e. larger equivalent widths by $\sim 3\sigma\text{--}4\sigma$) from Mg II, Mg I and Fe II, than those which do not. The absorption lines of Cr II, Mn II and Zn II are also stronger for systems with H I 21-cm detections than for those without, though the differences in equivalent widths are not more than $\sim 2\sigma$ significant. We expect strong Fe II and H I 21-cm absorption to probe sightlines close to galaxies. Nebular emission lines are usually detected in the stacked spectra of strong Mg II absorbers (Noterdaeme et al. 2010; Ménard et al. 2011). However, the number of systems in our stacked spectrum of H I 21-cm absorbers is small, and 3σ upper limits on average luminosity of the nebular emission lines of H β , [O II] and [O III] are 5, 4 and 5, in units of $10^{40} \text{ erg s}^{-1}$, respectively.

Using different absorption lines of Fe II detected in the stacked spectra, we compute the column density (N) and effective Doppler parameter (b) by constructing a single-cloud curve-of-growth (COG) [see panel (b) of Fig. 7]. Note that the spectral resolution of the SDSS spectra does not allow us to measure N and b accurately. However, a single cloud COG can still give an indication of the column density and velocity field. The effective b parameter obtained for Fe II lines in the stacked spectrum of H I 21-cm detections is slightly higher than that for Fe II lines in the stacked spectrum of non-detections. The larger effective b parameter could reflect larger number of absorption components, and hence higher probability of detecting H I 21-cm absorption from a cold gas component. The stacked spectrum of H I 21-cm detections gives two times higher $N(\text{Fe II})$ than that of the non-detections. Even if we consider only the two weak transitions of Fe II $\lambda 2249$ and $\lambda 2260$, which are unlikely to be affected by saturation, we get two times higher $N(\text{Fe II})$ for the stacked spectrum of H I 21-cm detections compared to that of the non-detections. Therefore, it seems that systems which give rise to H I 21-cm absorption are likely to have stronger metal absorption (i.e. higher column density) on average than those which do not. This is also consistent with the increase in detection rate of H I 21-cm absorption with increasing $W_{\text{Fe II}}$ (see Fig. 5).

7 DUST CONTENT OF H I 21-cm ABSORBERS

Dust grains play an important role in the heating and cooling of the neutral gas phases of the ISM (Wolfire et al. 1995), and also in the formation of H₂ molecules in the cold and dense phases (Gould




Figure 7. (a) The median stacked spectrum of SDSS quasars with strong Fe II systems at $0.5 < z < 1.5$ (in rest frame) which are detected in H I 21-cm absorption is shown as the red dashed line, and the same for systems which are not detected in H I 21-cm absorption is shown as the black solid line. The difference of the stacked spectrum of H I 21-cm non-detections from that of the detections is shown as the green line at an arbitrary offset in the y-axis for clarity. The various rest wavelengths of transitions of Mg II, Mg I, Cr II, Mn II, Fe II and Zn II are marked by vertical dotted lines. (b) The COG for Fe II lines detected in the stacked spectrum of H I 21-cm detections is shown as the red dashed line and that for Fe II lines detected in the stacked spectrum of H I 21-cm non-detections is shown as the solid black line. The filled and open circles mark the observed rest-frame equivalent widths and the column densities calculated for Fe II lines in the stacked spectrum of H I 21-cm detections and non-detections, respectively.

Table 7. List of rest equivalent widths of different metal transitions in the stacked spectra discussed in Section 6.2.

Transition	REW (detections) (Å)	REW (non-detections) (Å)
(1)	(2)	(3)
Mg II λ 2796	2.22 ± 0.08	1.88 ± 0.05
Mg II λ 2803	2.08 ± 0.07	1.75 ± 0.05
Mg I λ 2852	0.77 ± 0.07	0.54 ± 0.03
Fe II λ 2600	1.68 ± 0.06	1.40 ± 0.04
Fe II λ 2586	1.23 ± 0.06	0.90 ± 0.04
Fe II λ 2382	1.69 ± 0.07	1.38 ± 0.06
Fe II λ 2374	0.85 ± 0.07	0.59 ± 0.03
Fe II λ 2344	1.06 ± 0.04	1.01 ± 0.02
Fe II λ 2260	0.13 ± 0.04	0.07 ± 0.02
Fe II λ 2249	0.15 ± 0.04	0.09 ± 0.02
Mn II λ 2606	0.14 ± 0.06	0.05 ± 0.04
Mn II λ 2594	0.20 ± 0.06	0.06 ± 0.04
Mn II λ 2576	0.28 ± 0.06	0.07 ± 0.04
Zn II + Cr II λ 2062	0.22 ± 0.06	0.03 ± 0.03
Zn II + Mg I λ 2026	0.25 ± 0.06	0.13 ± 0.02

Notes. Column 1: metal line transition. Columns 2 and 3: rest equivalent widths of the transition in the stacked spectra of quasars with H I 21-cm absorption detections and non-detections, respectively.

& Salpeter 1963; Hollenbach & Salpeter 1971). The large homogeneous data base of SDSS quasar spectra has made it possible to statistically quantify the dust content of absorber populations like Mg II systems, Ca II systems and DLAs through differential reddening measurements (Wild, Hewett & Pettini 2006; York et al. 2006; Ménard et al. 2008; Vladilo, Prochaska & Wolfe 2008; Budzynski & Hewett 2011; Khare et al. 2012; Ménard & Fukugita 2012; Sardane, Turnshek & Rao 2015; Murphy & Bernet 2016). Such studies

have found that Mg II systems rarely show strong reddening due to dust in the spectra of the background quasars [mean $E(B - V) \sim 0.02$]. Jiang et al. (2011) have found only 39 cases that are 2175 Å dust extinction bump candidates among 2951 strong Mg II systems from SDSS with $W_{\text{Mg II}} \geq 1$ Å at $1.0 < z < 1.86$. However, the Mg II line strength is generally found to be correlated with the reddening. For example, Budzynski & Hewett (2011) have found that the dependence of $W_{\text{Mg II}}$ on $E(B - V)$ can be modelled by a power law for $1 \leq W_{\text{Mg II}} (\text{Å}) \leq 5$, and ~ 83 per cent of the high confidence 2175 Å candidates from Jiang et al. (2011) have $W_{\text{Mg II}} \geq 2$ Å and $W_{\text{Fe II}} \geq 1$ Å. Recently, Ledoux et al. (2015) have shown that dust content of C I absorbers at $1.5 < z < 4.5$ selected from SDSS is significant, with mean $E(B - V) = 0.065$ and 30 per cent of them showing the 2175 Å extinction feature. Moreover, C I absorbers provide ideal targets to search for tracers of translucent molecular gas like CO (e.g. Noterdaeme et al. 2011, 2016). Noterdaeme et al. (2008) have found that the presence of H₂ molecular absorption in $z > 1.8$ DLAs is closely related to the dust content and metallicity. H I 21-cm absorption can also provide an efficient way to reveal cold and dusty absorbers (e.g. Srianand et al. 2008). Indeed a possible anticorrelation between the H I 21-cm spin temperature and gas phase metallicity has been reported (see Kanekar et al. 2014, and references therein). Here, we investigate the dust content of strong Fe II absorbers and its connection with H I 21-cm absorption, first in our present sample S1 (Section 7.1) and then in the full sample S2 (Section 7.2).

7.1 Dust content in sample S1

We estimate the quasar reddening, $E(B - V)$, by fitting the quasar spectral energy distribution (SED) using the SDSS composite quasar spectrum (Vanden Berk et al. 2001), reddened by the MW, Small




Figure 8. SED fit to the spectra of (a) J0919+0146, (b) J0921+6215, (c) J1245+2232 and (d) J1327+4326, taking into account the reddening by the intervening absorber. SMC extinction curve gives best fit to all of them. The best-fitting $E(B - V)$ values are provided in each panel. The horizontal regions indicate the spectral range considered in the fitting process. Inset in each panel is the distribution of $E(B - V)$ for a control sample of SDSS quasars (see Section 7.1 for details). The number of quasars, mean, median and standard deviation of $E(B - V)$ in the control sample are also provided. The arrows indicate the $E(B - V)$ values obtained for the respective quasars. The unreddened SDSS quasar composite spectrum (Vanden Berk et al. 2001) is shown as the grey dashed line in panel (a).

Magellanic Cloud (SMC), Large Magellanic Cloud (LMC) or LMC2 supershell extinction curves (Gordon et al. 2003). We follow the same procedure as detailed in the works of Srianand et al. (2008) and Noterdaeme et al. (2009, 2010). In case of multiple epoch spectra available of the same quasar, we fit the SED of all of them and take the fit that gives the minimum χ^2 . We find that the $E(B - V)$ values obtained from fitting multiple epoch spectra of the same quasar can differ by ~ 10 – 20 per cent. This should be taken as the typical systematic error in the $E(B - V)$ values. We note that in the majority of cases, a better fit was obtained for SDSS-DR7 spectra than for SDSS-Baryon Oscillation Spectroscopic Survey (BOSS) spectra of the same quasar. This could be due to flux calibration overestimate of the blue side of the BOSS spectra.

We consider systems with $E(B - V) \geq 0.1$ to be showing signatures of reddening, since below this the distribution of $E(B - V)$ values could be dominated by the quasar SED-induced uncertainty. We find that 4 out of 16 quasars in our sample S1 (J0919+0146,

J0921+6215, J1245+2232 and J1327+4326) have $E(B - V) \geq 0.1$. H I 21-cm absorption has been detected towards three of these quasars (J0919+0146, J0921+6215 and J1327+4326). Note that the system towards J0919+0146 has been identified as a median-confidence (4.8σ) 2175 Å absorber candidate by Jiang et al. (2011). The SED of these four quasars are best fit with SMC extinction law (see Fig. 8). We note that the SED fit of J1245+2232 is not very good, which could be due to the complex intrinsic quasar emission. The errors provided for $E(B - V)$ take into account the uncertainties in the extinction law parameters. For each of these four quasars, we applied the same SED fitting procedure with SMC extinction law to a control sample of SDSS non-Broad Absorption Line quasars within $\Delta z = \pm 0.05 - 0.1$ of z_{cm} and $\Delta r_{\text{mag}} = \pm 0.5 - 1.0$ of r_{mag} of the quasars and having spectra with signal-to-noise ratio ≥ 10 . The details of the control samples are provided in Fig. 8. The standard deviation of the $E(B - V)$ values reflects the typical systematic error in the SED-fitting method due to the dispersion of the unreddened

Table 8. Parameters derived from SED fit to the quasars shown in Fig. 8 and discussed in Section 7.1.

Quasar	$E(B - V)$	$N(\text{H I})$ $\times 1/\kappa 10^{21}$ (cm^{-2})	$(\kappa T_s)/C_f$ (K)	κ
(1)	(2)	(3)	(4)	(5)
J0919+0146	0.20 ± 0.02	7	2500	–
J0921+6215	0.12 ± 0.01	4	6000	>3
J1245+2232	0.31 ± 0.03	11	$\geq 24\,000$	–
J1327+4326	0.11 ± 0.01	4	6000	>2

Notes. Column 1: quasar name. Column 2: $E(B - V)$ obtained from SED fit. SMC extinction curve gives best fit in these cases. Column 3: $N(\text{H I})$ in units of $1/\kappa 10^{21}$ (cm^{-2}) obtained from the relation between $N(\text{H I})$ and A_V in the SMC (Gordon et al. 2003). κ is the dust-to-gas ratio relative to SMC. Column 4: $(\kappa T_s)/C_f$ (K) obtained by comparing Column 3 with $\int \tau dv$ measurements given in Table 4. Column 5: in case of J0921+6215 and J1327+4326 where we have estimates of C_f (see Table 2), we constrain κ by using the upper limit on T_k (see Table 5).

quasar SED. In case of the above four quasars, we find that their reddening is significant at $\sim 2\sigma$ – 4σ level.

In Table 8, we have estimated the $N(\text{H I})$ for the above four systems in terms of the dust-to-gas-ratio relative to SMC (κ), using the observed mean relation between A_V and $N(\text{H I})$ in the SMC (Gordon et al. 2003). Further, comparing this with the $\int \tau dv$ measurement we have estimated $(\kappa T_s)/C_f$. In case of J0921+6215 and J1327+4326, where we have estimates of C_f (see Table 2), we find that the T_s obtained for $\kappa = 1$ is higher by a factor of $\gtrsim 2$ than the upper limit on the gas kinetic temperature from the H I 21-cm line width (see Table 5). Hence, the extinction per hydrogen atom in these systems could be at least a factor of ~ 2 higher than what is observed in the SMC. This could also be the reason for non-detection of H I 21-cm absorption towards J1245+2232. The possibility of the extinction per hydrogen atom being higher in high- z absorbers has been suggested by G12. Further, from fig. 11 of Ledoux et al. (2015), it can be seen that in few of the high- z C I systems, the $E(B - V)$ versus $N(\text{H I})$ relation is consistent with the reddening per hydrogen atom being up to 10 times higher than that seen in the MW and the Magellanic clouds. This could imply a different grain chemistry or small grain

size (i.e. larger total grain surface area) in these high- z absorbers (see e.g. Noterdaeme et al. 2016; Shaw, Rawlins & Srikanth 2016).

7.2 Dust content in sample S2

We estimated $E(B - V)$ for all the strong Mg II systems that have been searched for H I 21-cm absorption (G09, G12). The $E(B - V)$ of the strong Mg II systems is correlated with both $W_{\text{Mg II}}$ and $W_{\text{Fe II}}$ at 2σ significance level. We find that strong Fe II absorbers tend to cause more reddening, with median $E(B - V) = 0.05$ for systems with $W_{\text{Fe II}} \geq 1 \text{ \AA}$, compared to median $E(B - V) = 0.01$ for systems with $W_{\text{Fe II}} < 1 \text{ \AA}$. A two-sided Kolmogorov–Smirnov (KS) test between the $E(B - V)$ distribution of the strong and weak Fe II systems suggests that the maximum deviation between the two cumulative distribution functions is $D_{\text{KS}} = 0.50$ with a probability of $P_{\text{KS}} = 0.01$ (where P_{KS} is the probability of finding this D_{KS} value or lower by chance). Hence, our strong $W_{\text{Fe II}}$ -based selection technique tends to select dusty absorbers. Though we note that among the strong Fe II absorbers there is no correlation of $E(B - V)$ with $W_{\text{Mg II}}$ and $W_{\text{Fe II}}$ (i.e. they are at $\lesssim 1\sigma$ significance). We plot $E(B - V)$ versus $W_{\text{Fe II}}$ for the strong Fe II systems in panel (a) of Fig. 9. In addition, we do not find any redshift evolution of $E(B - V)$.

Next, we check whether among the strong Fe II systems, H I 21-cm absorption leads to more reddening in the quasar spectra. We show the histogram and cumulative distributions of the best-fitting $E(B - V)$ values for the systems with H I 21-cm detections and non-detections in panel (b) of Fig. 9. It can be seen that systems that show H I 21-cm absorption tend to cause more reddening in the quasar spectra. The median $E(B - V)$ for H I 21-cm detections is 0.10, while it is 0.02 for the non-detections. A two-sided KS test between the two distributions gives $D_{\text{KS}} = 0.42$ with $P_{\text{KS}} = 0.09$. The connection between H I 21-cm absorption and dust is further illustrated in panel (c) of Fig. 9, which shows the geometric mean stacked SDSS quasar spectra of the H I 21-cm detections and non-detections. It can be clearly seen that H I 21-cm absorption on average causes more reddening in the quasar spectra. We fit the stacked spectrum of H I 21-cm detections considering the stacked spectrum of H I 21-cm non-detections as a template. The best-fitting extinction curve is SMC, and the differential $E(B - V)$ obtained is




Figure 9. (a) $E(B - V)$ versus $W_{\text{Fe II}}$ for the strong Fe II systems. Filled symbols correspond to H I 21-cm detections and open symbols correspond to non-detections. The typical error bars are plotted in the bottom right corner. (b) The histogram distributions of $E(B - V)$ values for strong Fe II systems that show H I 21-cm absorption (red shaded) and those which do not (black open). The inset shows the cumulative distributions of $E(B - V)$ for systems with and without H I 21-cm absorption as red dashed and black solid lines, respectively. The vertical dashed and solid ticks mark the median $E(B - V)$ for systems with and without H I 21-cm absorption, respectively. (c) The geometric mean stacked spectrum of SDSS quasars with strong Fe II systems at $0.5 < z < 1.5$ (in rest frame) which are detected in H I 21-cm absorption is shown as the red dashed line, and the same for systems which are not detected in H I 21-cm absorption is shown as the black solid line. The spectra have been smoothed by 5 pixels for display purpose. The differential reddening, $\Delta E(B - V)$, is indicated in the figure (see Section 7.2 for details).

0.05. In addition, we find that the H I 21-cm detection rate shows tentative evidence of increasing with reddening, i.e. $C_{21}^{\text{Fe II}} = 0.25_{-0.11}^{+0.17}$ ($0.25_{-0.12}^{+0.20}$ for $\tau_0 = 0.2$) for $E(B - V) < 0.1$ and $C_{21}^{\text{Fe II}} = 0.38_{-0.20}^{+0.36}$ ($0.75_{-0.41}^{+0.73}$ for $\tau_0 = 0.2$) for $E(B - V) \geq 0.1$. However, the increase of $C_{21}^{\text{Fe II}}$ with $E(B - V)$ is not statistically significant due to the large uncertainties, and a larger sample is required to confirm this trend.

Note that in the case of J0919+0146, unlike in the rest, we knew that the quasar showed signatures of reddening, based on the results of Jiang et al. (2011), before searching for H I 21-cm absorption. However, we find that the above results do not change beyond the statistical uncertainties on excluding this system, i.e. this system does not dominate the statistics. For example, on excluding this system, two-sided KS test between the $E(B - V)$ distributions of H I 21-cm detections and non-detections gives $D_{\text{KS}} = 0.39$ with $P_{\text{KS}} = 0.16$, and the stacked spectrum of H I 21-cm detections shows differential $E(B - V)$ of 0.04 with respect to that of the non-detections.

A correlation between $E(B - V)$ and $N(\text{H I})$ is expected based on the observations in the MW and the Magellanic Clouds (Bohlin, Savage & Drake 1978; Gordon et al. 2003; Gudennavar et al. 2012). However, we do not find any significant (1.4σ) correlation of $\int \tau dv$ with $E(B - V)$ for the strong Fe II systems. Finally, we note that 72 per cent of the systems with $E(B - V) \geq 0.1$ have SMC type of dust. This is comparable to 74 per cent of the C I absorbers having SMC type of dust (Ledoux et al. 2015). Two out of the five H I 21-cm detections with $E(B - V) \geq 0.1$ have LMC type of dust and are discussed in detail in Srikanth et al. (2008). The remaining three have SMC type of dust and are discussed in Section 7.1 (see also Fig. 8).

8 VELOCITY WIDTH OF H I 21-cm ABSORPTION

8.1 Redshift evolution

The 16 H I 21-cm absorption lines detected in the strong Fe II systems (i.e. sample S2) exhibit a wide range of velocity widths, i.e. $v_{90} \sim 8\text{--}160 \text{ km s}^{-1}$ (median = 56 km s^{-1}), and FWHM of individual Gaussian components $\sim 3\text{--}130 \text{ km s}^{-1}$ (median = 23 km s^{-1}). We compare the H I 21-cm absorption velocity width in the strong Fe II systems with that in other intervening samples selected based on different criteria – the sample of $z < 0.4$ QGPs (see Dutta et al. 2016, and references therein), and samples of Mg II systems (which do not satisfy our criterion of $W_{\text{Fe II}} \geq 1 \text{ \AA}$ at $0.5 < z < 1.5$) [see G12 and references therein] and DLAs (see Kanekar 2014; Kanekar et al. 2014, and references therein). Note that for this analysis we do not consider proximate absorbers with redshifts within $\sim 3000 \text{ km s}^{-1}$ of the quasar redshift (Møller, Warren & Fynbo 1998; Ellison et al. 2002; Prochaska, Hennawi & Herbert-Fort 2008b), since the absorption could be associated with the quasar and intrinsic H I 21-cm absorption lines tend to be broader (Geréb et al. 2015; Curran et al. 2016). Hence, we do not consider the H I 21-cm absorption towards J0919+0146, which is $\sim 1700 \text{ km s}^{-1}$ blueshifted from the quasar emission redshift. This is the broadest H I 21-cm absorption line among all the Fe II systems (the absorption can be fit with a single Gaussian component of FWHM = 131 km s^{-1}). In addition, we do not consider the very broad (FWHM $\sim 235 \text{ km s}^{-1}$) H I 21-cm absorption line detected towards J162439.09+234512.2 (Curran et al. 2007a), as well as the H I 21-cm absorption lines detected towards gravitational lenses (Chengalur, de Bruyn & Narasimha 1999; Kanekar & Chengalur 2003), where the veloc-

ity widths could be dominated by other factors like the radio structure.

Note that the different samples listed above probe different redshift ranges. When restricting to $z < 1$, we do not find any difference in the v_{90} distributions of the galaxy-selected (or absorption-blind) sample of QGPs and the absorption-selected samples of Mg II/Fe II/DLAs. Similarly, we do not find any difference in the v_{90} distributions of the strong Fe II systems and the other Mg II systems and DLAs, over $0.5 < z < 1.5$. Note that the $\int \tau dv$ distributions in QGPs, DLAs, Mg II and Fe II systems are also similar when we consider common redshift ranges.

The left-hand panel of Fig. 10 shows v_{90} as a function of z_{abs} for the QGPs, the strong Fe II systems and other Mg II systems and DLAs. It can be seen that v_{90} of H I 21-cm absorbers shows an increasing trend with redshift. Test for correlation between v_{90} and z_{abs} for all the H I 21-cm absorbers gives a positive correlation, with $r_k = 0.34$, $P(r_k) = 1.4 \times 10^{-4}$, $S(r_k) = 3.8\sigma$ and $r_s = 0.48$, $P(r_s) = 1.4 \times 10^{-4}$, $S(r_s) = 3.6\sigma$. Note that inclusion of the proximate systems and the systems from Chengalur et al. (1999), Kanekar & Chengalur (2003) and Curran et al. (2007a), leads to the significance of the anticorrelation becoming 3σ . On the other hand, if we consider only the measurements from absorption-selected samples (i.e. excluding the measurements from QGPs), the anticorrelation is significant at 2.5σ .

The increasing trend of v_{90} with redshift indicates that the velocity dispersion of H I gas could be larger in high- z galaxies. From fig. 4 of Ledoux et al. (2006), it can be seen that for a given metallicity, the low-ionization metal lines in DLAs tend to have larger velocity widths at $z > 2.4$. The velocity–metallicity correlation in DLAs is believed to reflect the mass–metallicity relation in galaxies (Tremonti et al. 2004; Møller et al. 2013; Neeleman et al. 2013). Moreover, fig. 3 of Erb et al. (2006) shows that for a given metallicity, galaxies will have larger stellar masses at $z \sim 2$ compared to at $z \sim 0.1$. Hence, this may mean that for a given metallicity, the average velocity width is higher at high z , as it originates from more massive galaxy haloes compared to at low z . Therefore, one possible explanation of the v_{90} versus z correlation is that a typical H I 21-cm absorber may be probed by larger mass haloes at high z .

In addition to v_{90} , the FWHM of individual Gaussian components of the H I 21-cm absorption profiles also show 4σ correlation with z_{abs} . The median FWHM (8 km s^{-1}) of the H I 21-cm absorption lines detected at $z < 0.75$ (which divides the sample into equal halves and corresponds to a lookback time of 6.6 Gyr) gives $T_k \leq 1400 \text{ K}$, while the median FWHM (18 km s^{-1}) at $z \geq 0.75$ gives $T_k \leq 7000 \text{ K}$. The large upper limits on the kinetic temperature indicates that non-thermal motions are most likely to dominate the line widths. The line widths could be driven by turbulent motion, which is closely linked with the supernova rate and hence the star formation rate (Elmegreen & Scalo 2004; Mac Low & Klessen 2004; McKee & Ostriker 2007; Tamburro et al. 2009). Moreover, it is known that the star formation rate on average is higher at higher redshifts (Madau & Dickinson 2014). However, the H I 21-cm velocity width depends not just on the gas structure and kinematics, but on the background radio source structure as well. To gain further insight into the redshift evolution of the H I 21-cm velocity width, we study its dependence on various physical parameters next.

8.2 Dependence on physical parameters

Curran et al. (2007b) have reported a $2.2\sigma\text{--}2.8\sigma$ level correlation between H I 21-cm absorption line widths and $W_{\text{Mg II}}$. However, G09 and K09 do not find similar correlation. Since $W_{\text{Mg II}}$ and $W_{\text{Fe II}}$




Figure 10. (a) The velocity width, v_{90} , of H I 21-cm absorption lines detected in samples of DLAs/Mg II systems (black circles), QGPs (red diamonds) and strong Fe II systems (blue squares). The solid, dashed and dotted vertical (horizontal) ticks mark the median z_{abs} (v_{90}) for the DLAs/Mg II systems, QGPs and Fe II systems, respectively. The measurement towards J0919+0146 from our sample (which is not considered in the analysis; see text for details) is shown as a star. (b) v_{90} of H I 21-cm absorption lines detected at $0.3 < z < 2.3$ as a function of the ratio R2. The symbol convention is same as in panel (a). The typical error bar in R2 is plotted in the bottom right corner.

information cannot be uniformly obtained over the redshift range of H I 21-cm absorption measurements, we focus here on the systems at $0.3 < z < 2.3$. For these systems, we do not find any correlation of v_{90} of the H I 21-cm absorption lines with $W_{\text{Mg II}}$ as well as $W_{\text{Fe II}}$ (i.e. they are at $\lesssim 1\sigma$ significance). The absence of correlation between the H I 21-cm absorption width and $W_{\text{Mg II}}$ or $W_{\text{Fe II}}$ can be explained if the H I 21-cm absorbing gas originates from only one or few of the metal line components, which are not necessarily the strongest. Indeed this has been observed in few cases where high-resolution optical spectra are available (G09; Rahmani et al. 2012; Srianand et al. 2012; Dutta et al. 2015).

Further, we do not find any correlation of v_{90} with DR, R1 and R2 (i.e. they are at $\lesssim 1\sigma$ significance). However, we find that the broadest H I 21-cm absorption lines occur when R2 is ~ 1.2 (see right-hand panel of Fig. 10). If we consider the measurement towards J0919+0146 (shown as a star in Fig. 10), there is a weak 1.5σ anticorrelation between v_{90} and R2 (significance reduces to 1.2σ on excluding it). Assuming that both Mg II and Fe II absorption trace the same velocity field and b parameters in the range 40–60 km s⁻¹ (as indicated by our analysis in Section 6.2), from single-cloud COG we find that values of R2 $\lesssim 1.2$ will occur in the saturated part where $N(\text{Fe II}) > 3 \times 10^{15} \text{ cm}^{-2}$. We can put a lower limit on $N(\text{H I}) > 10^{20} \text{ cm}^{-2}$ assuming solar metallicity and solar abundance of Fe (Asplund et al. 2009). Since Fe is usually prone to depletion on to dust grains (Savage & Sembach 1996; Draine 2003; Jenkins 2009), this is a strict lower limit on $N(\text{H I})$. Hence, such large $N(\text{Fe II})$ are expected from self-shielded regions with high $N(\text{H I})$. The probability of encountering multiple cold H I clouds along such sightlines would be high, which can explain the observed broad H I 21-cm lines. On the other hand, when R2 is higher, $N(\text{H I})$ would be lower and in turn the probability of encountering multiple cold gas clouds along such sightlines would be lower. This can explain the low v_{90} values when R2 ≥ 1.5 . The scatter in the v_{90} values when R2 is close to 1 can be explained if H I 21-cm absorption does not always arise from all the metal line components, as also discussed above.

As noted in Section 8.1, the velocity width of the H I 21-cm absorption lines also depends on the radio structure of the background sources. We note that G12 have not found any correlation of v_{90} with

the linear size of the radio sources measured from VLBA images. However, the three Fe II systems with $v_{90} > 100 \text{ km s}^{-1}$ are towards radio sources with extended morphology at arcsecond-scales (G12). Note that in case of the broad proximate absorber towards J0919+0146, the radio source is compact in our GMRT image. Even when the radio sources are compact at arcsecond-scales, if radio structures are present at parsec-scales, that could lead to broadening of the line (e.g. Srianand et al. 2013, 2015; Biggs et al. 2016). Hence, we emphasize the need for multiwavelength VLBA sub-arcsecond-scale images of all the radio sources towards which H I 21-cm absorption lines are detected in order to characterize their radio structure and address the origin of the velocity widths.

9 SUMMARY

Using strong Mg II systems ($W_{\text{Mg II}} \geq 1 \text{ \AA}$) having measurements of both $N(\text{H I})$ and $W_{\text{Mg II}}$ at $z < 1.65$ and $z \sim 2$, we have found that the probability of having $\log N(\text{H I}) \geq 20.3$, i.e. DLA column densities, can be increased by a factor of ~ 1.4 – 1.7 , by selecting strong Fe II systems with $W_{\text{Fe II}} \geq 1 \text{ \AA}$. Hence, we have searched for H I 21-cm absorption in a sample of strong Fe II systems at $0.5 < z < 1.5$ selected from SDSS-DR12, using GMRT and GBT. We have detected H I 21-cm absorption in six of these systems.

Combining our sample with that of strong Fe II systems from G09 and G12, we have estimated the detection rate of H I 21-cm absorption in strong Fe II systems to be $0.30^{+0.12}_{-0.09}$ for $\tau_0 = 0.3 \text{ km s}^{-1}$. We have found that the detection rate increases with $W_{\text{Fe II}}$, being four times higher in systems with $W_{\text{Fe II}} \geq 1 \text{ \AA}$ compared to in systems with $W_{\text{Fe II}} < 1 \text{ \AA}$. The detection rate of H I 21-cm absorption in strong Fe II systems remains constant within the uncertainties over $0.5 < z < 1.5$. For $T_s = 500 \text{ K}$ (typical of DLAs at this redshift range) and $C_f = 1$, all the H I 21-cm absorption in strong Fe II systems would arise from DLAs. Hence, a $W_{\text{Fe II}}$ -based selection appears to be efficient in detecting high $N(\text{H I})$ cold gas. From the detection rate of H I 21-cm absorption in strong Fe II systems and that of DLAs in strong Fe II systems, we estimate the detection rate of H I 21-cm absorption in DLAs to be $0.67^{+0.31}_{-0.23}$ at $0.5 < z < 1.5$ for $\tau_0 = 0.3 \text{ km s}^{-1}$. This is three times higher than that estimated

in $2 < z < 3.5$ DLAs and may indicate towards an increasing filling factor of cold gas in DLAs with time.

We do not find any significant correlation of the H I 21-cm absorption strength with the metal line properties. However, the metal absorption are systematically stronger in the stacked SDSS spectrum of the systems which show H I 21-cm absorption than in that of the H I 21-cm non-detections. In addition, the H I 21-cm absorbers tend to cause more significant reddening in the spectrum of the background quasars and there is a tendency for the detection rate of H I 21-cm absorbers to be higher towards more reddened quasars. The stacked spectrum of quasars with H I 21-cm absorption detected towards them is more reddened than that of quasars without any H I 21-cm absorption. Hence, the above imply that H I 21-cm absorption is more likely to arise in metal-rich dusty cold gas. Note that highly reddened systems are likely to be missed out in samples selected on the basis of optical/UV spectra. Budzynski & Hewett (2011) have shown that Mg II samples, constructed from flux-limited quasar surveys, suffer 24 per cent and 34 per cent incompleteness for absorbers with $W_{\text{Mg II}} > 1 \text{ \AA}$ and $> 2 \text{ \AA}$, respectively. Hence, upcoming blind H I 21-cm absorption surveys with the Square Kilometre Array pre-cursors could unravel a new population of dusty absorbers towards highly reddened quasars.

By comparing the velocity widths of intervening H I 21-cm absorption lines detected in samples of QGPs, DLAs and Mg II systems at $z < 3.5$, we find evidence for the velocity widths to be increasing with redshift, which is significant at 3.8σ . This could be because a typical H I 21-cm absorber may be originating from a larger mass galaxy halo at high z compared to at low z . However, the H I 21-cm velocity width depends on various factors like kinematics and structure of the absorbing gas as well as structure of the background radio source, and these need to be better understood in order to correctly interpret the redshift evolution. Finally, we emphasize the need for more H I 21-cm absorption detections to confirm different trends noted in this work with higher statistical significance.

ACKNOWLEDGEMENTS

We thank the anonymous referee for useful comments. We thank the staff at GMRT and GBT for their help during the observations. GMRT is run by the National Centre for Radio Astrophysics of the Tata Institute of Fundamental Research. GBT is run by the National Radio Astronomy Observatory (NRAO). The NRAO is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. NG, PN, RS and PPJ acknowledge the support from Indo-French centre for the promotion of Advanced Research (IFCPAR) under Project No. 5504-2. JK acknowledges financial support from the Danish Council for Independent Research (EU-FP7 under the Marie-Curie grant agreement no. 600207) with reference DFF-MOBILEX-5051-00115. This research has made use of the NASA/IPAC Extragalactic Database (NED) that is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the US Department of Energy Office of Science. The SDSS-III web site is <http://www.sdss3.org/>. SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, Carnegie Mellon University, University of

Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington and Yale University.

REFERENCES

- Alam S. et al., 2015, *ApJS*, 219, 12
 Asplund M., Grevesse N., Sauval A. J., Scott P., 2009, *ARA&A*, 47, 481
 Battisti A. J. et al., 2012, *ApJ*, 744, 93
 Bergeron J., Boissé P., 1991, *A&A*, 243, 344
 Biggs A. D., Zwaan M. A., Hatziminaoglou E., Péroux C., Liske J., 2016, *MNRAS*, 462, 2819
 Bohlin R. C., Savage B. D., Drake J. F., 1978, *ApJ*, 224, 132
 Bond N. A., Churchill C. W., Charlton J. C., Vogt S. S., 2001, *ApJ*, 562, 641
 Bouché N., Murphy M. T., Péroux C., Davies R., Eisenhauer F., Förster Schreiber N. M., Tacconi L., 2007, *ApJ*, 669, L5
 Briggs F. H., Wolfe A. M., 1983, *ApJ*, 268, 76
 Budzynski J. M., Hewett P. C., 2011, *MNRAS*, 416, 1871
 Chen H.-W., Wild V., Tinker J. L., Gauthier J.-R., Helsby J. E., Shectman S. A., Thompson I. B., 2010a, *ApJ*, 724, L176
 Chen H.-W., Helsby J. E., Gauthier J.-R., Shectman S. A., Thompson I. B., Tinker J. L., 2010b, *ApJ*, 714, 1521
 Chengalur J. N., de Bruyn A. G., Narasimha D., 1999, *A&A*, 343, L79
 Churchill C. W., Vogt S. S., Charlton J. C., 2003, *AJ*, 125, 98
 Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, *AJ*, 115, 1693
 Curran S. J., 2010, *MNRAS*, 402, 2657
 Curran S. J., 2012, *ApJ*, 748, L18
 Curran S. J., Webb J. K., 2006, *MNRAS*, 371, 356
 Curran S. J., Murphy M. T., Pihlström Y. M., Webb J. K., Purcell C. R., 2005, *MNRAS*, 356, 1509
 Curran S. J., Tzanavaris P., Murphy M. T., Webb J. K., Pihlström Y. M., 2007a, *MNRAS*, 381, L6
 Curran S. J., Tzanavaris P., Pihlström Y. M., Webb J. K., 2007b, *MNRAS*, 382, 1331
 Curran S. J., Tzanavaris P., Darling J. K., Whiting M. T., Webb J. K., Bignell C., Athreya R., Murphy M. T., 2010, *MNRAS*, 402, 35
 Curran S. J., Duchesne S. W., Divoli A., Allison J. R., 2016, *MNRAS*, 462, 4197
 de Avillez M. A., Breitschwerdt D., 2004, *A&A*, 425, 899
 Douglas J. N., Bash F. N., Bozayan F. A., Torrence G. W., Wolfe C., 1996, *AJ*, 111, 1945
 Draine B. T., 2003, *ARA&A*, 41, 241
 Dutta R., Srianand R., Muzahid S., Gupta N., Momjian E., Charlton J., 2015, *MNRAS*, 448, 3718
 Dutta R., Srianand R., Gupta N., Momjian E., Noterdaeme P., Petitjean P., Rahmani H., 2016, *MNRAS*, 465, 588
 Ellison S. L., Yan L., Hook I. M., Pettini M., Wall J. V., Shaver P., 2002, *A&A*, 383, 91
 Elmegreen B. G., Scalo J., 2004, *ARA&A*, 42, 211
 Erb D. K., Shapley A. E., Pettini M., Steidel C. C., Reddy N. A., Adelberger K. L., 2006, *ApJ*, 644, 813
 Gatto A. et al., 2015, *MNRAS*, 449, 1057
 Gauthier J.-R., 2013, *MNRAS*, 432, 1444
 Gauthier J.-R., Chen H.-W., Tinker J. L., 2009, *ApJ*, 702, 50
 Gehrels N., 1986, *ApJ*, 303, 336
 Gent F. A., Shukurov A., Fletcher A., Sarson G. R., Mantere M. J., 2013, *MNRAS*, 432, 1396

- Geréb K., Maccagni F. M., Morganti R., Oosterloo T. A., 2015, *A&A*, 575, A44
- Gordon K. D., Clayton G. C., Misselt K. A., Landolt A. U., Wolff M. J., 2003, *ApJ*, 594, 279
- Gould R. J., Salpeter E. E., 1963, *ApJ*, 138, 393
- Gregory P. C., Condon J. J., 1991, *ApJS*, 75, 1011
- Gudennavar S. B., Bubbly S. G., Preethi K., Murthy J., 2012, *ApJS*, 199, 8
- Gupta N., Srianand R., Petitjean P., Noterdaeme P., Saikia D. J., 2009, *MNRAS*, 398, 201 (G09)
- Gupta N., Srianand R., Bowen D. V., York D. G., Wadadekar Y., 2010, *MNRAS*, 408, 849
- Gupta N., Srianand R., Petitjean P., Bergeron J., Noterdaeme P., Muzahid S., 2012, *A&A*, 544, A21 (G12)
- Heiles C., Troland T. H., 2003, *ApJ*, 586, 1067
- Hollenbach D., Salpeter E. E., 1971, *ApJ*, 163, 155
- Jenkins E. B., 2009, *ApJ*, 700, 1299
- Jiang P., Ge J., Zhou H., Wang J., Wang T., 2011, *ApJ*, 732, 110
- Jorgenson R. A., Murphy M. T., Thompson R., 2013, *MNRAS*, 435, 482
- Kacprzak G. G., Churchill C. W., 2011, *ApJ*, 743, L34
- Kacprzak G. G., Churchill C. W., Evans J. L., Murphy M. T., Steidel C. C., 2011, *MNRAS*, 416, 3118
- Kanekar N., 2014, *ApJ*, 797, L20
- Kanekar N., Chengalur J. N., 2003, *A&A*, 399, 857
- Kanekar N., Prochaska J. X., Ellison S. L., Chengalur J. N., 2009, *MNRAS*, 396, 385 (K09)
- Kanekar N. et al., 2014, *MNRAS*, 438, 2131
- Khare P., vanden Berk D., York D. G., Lundgren B., Kulkarni V. P., 2012, *MNRAS*, 419, 1028
- Kulkarni S. R., Heiles C., 1988, in Vershuur G. L., Kellermann K. I., eds, *Neutral Hydrogen and the Diffuse Interstellar Medium*, 2nd edn. Springer-Verlag, Berlin, p. 95
- Lane W. M., 2000, PhD thesis, Univ. Groningen (L00)
- Lanzetta K. M., Turnshek D. A., Wolfe A. M., 1987, *ApJ*, 322, 739
- Ledoux C., Petitjean P., Srianand R., 2003, *MNRAS*, 346, 209
- Ledoux C., Petitjean P., Fynbo J. P. U., Møller P., Srianand R., 2006, *A&A*, 457, 71
- Ledoux C., Noterdaeme P., Petitjean P., Srianand R., 2015, *A&A*, 580, A8
- Lovegrove E., Simcoe R. A., 2011, *ApJ*, 740, 30
- Lundgren B. F. et al., 2009, *ApJ*, 698, 819
- Mac Low M.-M., Klessen R. S., 2004, *Rev. Mod. Phys.*, 76, 125
- McKee C. F., Ostriker J. P., 1977, *ApJ*, 218, 148
- McKee C. F., Ostriker E. C., 2007, *ARA&A*, 45, 565
- Madau P., Dickinson M., 2014, *ARA&A*, 52, 415
- Matejek M. S., Simcoe R. A., Cooksey K. L., Seyffert E. N., 2013, *ApJ*, 764, 9
- Meiring J. D. et al., 2011, *ApJ*, 732, 35
- Ménard B., Fukugita M., 2012, *ApJ*, 754, 116
- Ménard B., Nestor D., Turnshek D., Quider A., Richards G., Chelouche D., Rao S., 2008, *MNRAS*, 385, 1053
- Ménard B., Wild V., Nestor D., Quider A., Zibetti S., Rao S., Turnshek D., 2011, *MNRAS*, 417, 801
- Møller P., Warren S. J., Fynbo J. U., 1998, *A&A*, 330, 19
- Møller P., Fynbo J. P. U., Ledoux C., Nilsson K. K., 2013, *MNRAS*, 430, 2680
- Murphy M. T., Bernet M. L., 2016, *MNRAS*, 455, 1043
- Murphy M. T., Curran S. J., Webb J. K., Ménager H., Zych B. J., 2007, *MNRAS*, 376, 673
- Muzahid S., Srianand R., Charlton J., 2015, *MNRAS*, 448, 2840
- Neeleman M., Wolfe A. M., Prochaska J. X., Rafelski M., 2013, *ApJ*, 769, 54
- Neeleman M., Prochaska J. X., Wolfe A. M., 2015, *ApJ*, 800, 7
- Neeleman M., Prochaska J. X., Ribaud J., Lehner N., Howk J. C., Rafelski M., Kanekar N., 2016, *ApJ*, 818, 113
- Nestor D. B., Turnshek D. A., Rao S. M., 2005, *ApJ*, 628, 637
- Nestor D. B., Johnson B. D., Wild V., Ménard B., Turnshek D. A., Rao S., Pettini M., 2011, *MNRAS*, 412, 1559
- Nielsen N. M., Churchill C. W., Kacprzak G. G., Murphy M. T., 2013, *ApJ*, 776, 114
- Noterdaeme P., Ledoux C., Petitjean P., Srianand R., 2008, *A&A*, 481, 327
- Noterdaeme P., Ledoux C., Srianand R., Petitjean P., Lopez S., 2009, *A&A*, 503, 765
- Noterdaeme P., Srianand R., Mohan V., 2010, *MNRAS*, 403, 906
- Noterdaeme P., Petitjean P., Srianand R., Ledoux C., López S., 2011, *A&A*, 526, L7
- Noterdaeme P. et al., 2012, *A&A*, 547, L1
- Noterdaeme P. et al., 2016, *A&A*, preprint ([arXiv:1609.01422](https://arxiv.org/abs/1609.01422))
- Petitjean P., Bergeron J., 1990, *A&A*, 231, 309
- Petitjean P., Srianand R., Ledoux C., 2000, *A&A*, 364, L26
- Prochaska J. X., Chen H.-W., Wolfe A. M., Dessauges-Zavadsky M., Bloom J. S., 2008a, *ApJ*, 672, 59
- Prochaska J. X., Hennawi J. F., Herbert-Fort S., 2008b, *ApJ*, 675, 1002
- Prochter G. E., Prochaska J. X., Burles S. M., 2006, *ApJ*, 639, 766
- Quider A. M., Nestor D. B., Turnshek D. A., Rao S. M., Monier E. M., Weyant A. N., Busche J. R., 2011, *AJ*, 141, 137
- Rafelski M., Neeleman M., Fumagalli M., Wolfe A. M., Prochaska J. X., 2014, *ApJ*, 782, L29
- Rahmani H., Srianand R., Gupta N., Petitjean P., Noterdaeme P., Vázquez D. A., 2012, *MNRAS*, 425, 556
- Rao S. M., Turnshek D. A., Nestor D. B., 2006, *ApJ*, 636, 610 (R06)
- Rao S. M., Belfort-Mihalyi M., Turnshek D. A., Monier E. M., Nestor D. B., Quider A., 2011, *MNRAS*, 416, 1215
- Rubin K. H. R., Weiner B. J., Koo D. C., Martin C. L., Prochaska J. X., Coil A. L., Newman J. A., 2010, *ApJ*, 719, 1503
- Sardane G. M., Turnshek D. A., Rao S. M., 2015, *MNRAS*, 452, 3192
- Sargent W. L. W., Steidel C. C., Boksenberg A., 1988, *ApJ*, 334, 22
- Savage B. D., Sembach K. R., 1996, *ARA&A*, 34, 279
- Shaw G., Rawlins K., Srianand R., 2016, *MNRAS*, 459, 3234
- Srianand R., 1996, *ApJ*, 462, 643
- Srianand R., Khare P., 1994, *ApJ*, 428, 82
- Srianand R., Petitjean P., Ledoux C., Ferland G., Shaw G., 2005, *MNRAS*, 362, 549
- Srianand R., Gupta N., Petitjean P., Noterdaeme P., Saikia D. J., 2008, *MNRAS*, 391, L69
- Srianand R., Gupta N., Petitjean P., Noterdaeme P., Ledoux C., Salter C. J., Saikia D. J., 2012, *MNRAS*, 421, 651
- Srianand R., Gupta N., Rahmani H., Momjian E., Petitjean P., Noterdaeme P., 2013, *MNRAS*, 428, 2198
- Srianand R., Gupta N., Momjian E., Vivek M., 2015, *MNRAS*, 451, 917
- Steidel C. C., 1995, in Meylan G., ed., *QSO Absorption Lines*. Springer-Verlag, Berlin, p. 139
- Steidel C. C., Sargent W. L. W., 1992, *ApJS*, 80, 1
- Tamburro D., Rix H.-W., Leroy A. K., Mac Low M.-M., Walter F., Kennicutt R. C., Brinks E., de Blok W. J. G., 2009, *AJ*, 137, 4424
- Tremonti C. A. et al., 2004, *ApJ*, 613, 898
- Turnshek D. A., Monier E. M., Rao S. M., Hamilton T. S., Sardane G. M., Held R., 2015, *MNRAS*, 449, 1536
- Vanden Berk D. E. et al., 2001, *AJ*, 122, 549
- Vladilo G., Prochaska J. X., Wolfe A. M., 2008, *A&A*, 478, 701
- Weiner B. J. et al., 2009, *ApJ*, 692, 187
- White R. L., Becker R. H., Helfand D. J., Gregg M. D., 1997, *ApJ*, 475, 479
- Wild V., Hewett P. C., Pettini M., 2006, *MNRAS*, 367, 211
- Wolfe A. M., Gawiser E., Prochaska J. X., 2005, *ARA&A*, 43, 861
- Wolfe A. M., Hollenbach D., McKee C. F., Tielens A. G. G. M., Bakes E. L. O., 1995, *ApJ*, 443, 152
- York D. G. et al., 2000, *AJ*, 120, 1579
- York D. G. et al., 2006, *MNRAS*, 367, 945
- Zhu G., Ménard B., 2013, *ApJ*, 770, 130
- Zibetti S., Ménard B., Nestor D. B., Quider A. M., Rao S. M., Turnshek D. A., 2007, *ApJ*, 658, 161
- Zwaan M. A., Liske J., Péroux C., Murphy M. T., Bouché N., Curran S. J., Biggs A. D., 2015, *MNRAS*, 453, 1268

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.