



HAL
open science

Structure of differentiated planetesimals: A chondritic fridge on top of a magma ocean

Cyril Sturtz, Angela Limare, Marc Chaussidon, Édouard Kaminski

► To cite this version:

Cyril Sturtz, Angela Limare, Marc Chaussidon, Édouard Kaminski. Structure of differentiated planetesimals: A chondritic fridge on top of a magma ocean. *Icarus*, 2022, 385, pp.115100. <10.1016/j.icarus.2022.115100>. <insu-03748508>

HAL Id: insu-03748508

<https://insu.hal.science/insu-03748508v1>

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Structure of differentiated planetesimals: a chondritic fridge on top of a magma ocean^{*,**}

Cyril Sturtz^{a,*}, Angela Limare^a, Marc Chaussidon^a and Édouard Kaminski^a

^aUniversité Paris Cité, Institut de Physique du Globe de Paris, CNRS, Paris, F-75005, France

ARTICLE INFO

Keywords:
Accretion
Planetesimals
Magma ocean
Meteorites
Erg Chech 002

ABSTRACT

Meteorites are interpreted as relics of early formed planetary bodies, and they provide information about the processes that occurred in the first few Myrs of our solar system. The ages measured for some differentiated meteorites (achondrites), indicate that planetesimals formed a differentiated silicate crust as early as ≈ 3 Myr after the beginning of the solar system. The composition of the recently discovered achondrite Erg Chech 002 (EC002), the oldest andesitic rock known so far, betokens partial melting of a chondritic source taking place as early as ≈ 1 Myr before all other known achondrites. However, thermal models of early accreted planetesimals predict massive melting of the planetesimal during core/mantle differentiation and cannot account for the preservation of a substantial amount of chondritic material. In this paper, we propose a way to interpret petrological and geochemical constraints provided by differentiated meteorites by introducing a refined thermal model of planetesimals formation and evolution. We demonstrate that continuous, protracted accretion of cold undifferentiated material upon a magma ocean over a timescale 2 times larger than the lifetime of the ^{26}Al heat source leads to the preservation of a few km thick chondritic crust. During accretion, the heat released by radioactive decay further induces episodes of partial melting at the base of the crust, which can lead to the formation of andesitic rocks such as EC002. Using the available constraints on the age of EC002 and its cooling rate, the application of our model constraints the terminal radius of its parent body between 70 and 130 km.

1. Introduction

Meteorites are key witnesses of the processes that occurred during the first stages of planetary evolution. In particular, achondritic meteorites sample planetesimals differentiated during the first 5 Myr of the Solar system. Differentiated achondrites are often basaltic and most of them originate from a few parent bodies: the ureilites parent body, the angrites parent body, Vesta 4 - the eucrites parent body (Greenwood, Burbine and Franchi, 2020). Achondrites with andesitic or trachyandesitic compositions are very rare, e.g., GRA 06128 and 06129 (Day, Ash, Liu, Bellucci, III, McDonough, Walker and Taylor, 2009; Shearer, Burger, Neal, Sharp, Spivak-Birndorf, Borg, Fernandes, Papike, Karner, Wadhwa, Gaffney, Shafer, Geissman, Atudorei, Herd, Weiss, King, Crowther and Gilmour, 2010), Almahata Sitta ALM-A (Bischoff, Horstmann, Barrat, Chaussidon, Pack, Herwartz, Ward, Vollmer and Decker, 2014), NWA 11119 (Srinivasan, Dunlap, Agee, Wadhwa, Coleff, Ziegler, Zeigler and McCubbin, 2018) and NWA 11575 (Agee, Habermann and Ziegler, 2018). Recently, a new achondrite, Erg Chech 002 (EC002), has been recognized as the oldest andesitic meteorite known so far (Barrat, Chaussidon, Yamaguchi, Beck, Villeneuve, Byrne, Broadley and Marty, 2021).

Early Solar system relative chronology is anchored to the Ca-Al-rich inclusions (CAIs) from CV3 chondrites which are the oldest components of primitive chondrites according to U-Pb dating (Allègre, Manhès and Göpel, 1995; Bouvier and Wadhwa, 2010; Connelly, Bizzarro, Krot, Nordlund, Wielandt and Ivanova, 2012). The formation of CAIs is generally considered to date the episode of high-temperature condensation, or reprocessing of these condensates, which produced the first Solar system dust in the accretion disk. A crystallization age of 2.255 ± 0.015 Myr after CAIs have been inferred for EC002 from $^{26}\text{Al} - ^{26}\text{Mg}$ chronology (Barrat et al., 2021), i. e. ≈ 1.2 Myr older than all other known achondrites, and contemporaneous with the formation of the metallic cores of the parent bodies of iron

* This study contributes to the IdEx Université de Paris ANR-18-IDEX-0001.

** This work was supported by the Programme National de Planétologie (PNP) of CNRS/INSU, co-funded by CNES.

✉ sturtz@ipgp.fr (C. Sturtz)

ORCID(s): 0000-0002-6329-949 (C. Sturtz); 0000-0001-6755-9843 (A. Limare); 0000-0001-8475-0690 (M. Chaussidon); 0000-0001-6965-8482 (É. Kaminski)

57 meteorites (Kruijer, Touboul, Fischer-Gödde, Bermingham, Walker and Kleine, 2014; Kruijer, Burkhardt, Budde and
58 Kleine, 2017). Petrological and geochemical data on the two major mineral phases in EC002 (pyroxene and plagioclase)
59 indicate that (i) the parent melt of EC002 was produced by 25% partial melting of a source of chondritic composition
60 at a temperature of 1224 ± 20 °C, (ii) the pyroxenes crystallized at 1186 ± 25 °C and re-equilibrated chemically at
61 around 960 °C, (iii) the cooling of the rock or mush (crystal + melt) took place at ≈ 5 °C yr⁻¹ between 1200 °C
62 and 1000 °C, no melt being present anymore at 1000 °C, and (iv) finally the rock was quenched below ≈ 900 °C at
63 a rate $0.1 - 1$ °C day⁻¹ (Barrat et al., 2021). Although the ²⁶Al – ²⁶Mg systematics does not allow to date precisely
64 the episode of melting, a model age for melting between 0.95 Myr and 2.2 Myr (mean 1.4 Myr) after CAIs has been
65 proposed (Barrat et al., 2021). Note that because the ²⁶Al – ²⁶Mg age dates physically the closure temperature of
66 Mg diffusion in EC002 plagioclases, the true crystallization age of EC002 can even be slightly earlier than 2.2 Myr,
67 depending on the cooling rate before the closure of the chemical exchanges. In fact, recent Mg isotopic measurements
68 on mineral separates yielded a ²⁶Al – ²⁶Mg crystallization age of 1.80 Myr. (Fang, Frossard, Boyet, Bouvier, Barrat,
69 Chaussidon and Moynier, 2022).

70 Early differentiation of planetesimals, within the first ≈ 2 Myr of the Solar system, as witnessed by iron meteorites
71 (Kruijer et al., 2014, 2017) and EC002 (Barrat et al., 2021), stems from partial melting caused by the heat released by
72 the radioactive decay of short lived ²⁶Al (and possibly ⁶⁰Fe). Intense radioactive internal heating can indeed induce
73 several phenomena such as partial melting of chondritic materials (Elkins-Tanton, Weiss and Zuber, 2011), solid-
74 state convection (Tkalcic, Golabek and Brenker, 2013; Kaminski, Limare, Kenda and Chaussidon, 2020), iron melt
75 segregation (Neumann, Breuer and Spohn, 2012, 2014), core-mantle differentiation (Sahijpal, Soni and Gupta, 2007),
76 magma ocean episodes (Greenwood, Franchi, Jambon and Buchanan, 2005), and volcanism (Mandler and Elkins-
77 Tanton, 2013; Fu and Elkins-Tanton, 2014). Although the production mechanism of achondritic basaltic or andesitic
78 magmas is still the object of debate, it must rely on the presence of a stable crust, either a primary or a secondary one,
79 at the surface of the planetesimal (Taylor, 1989; Condie, 2016). Primary crusts are formed at the surface of a cooling
80 magma ocean through the flotation of positively buoyant crystals from the magma ocean such as silica (Faure, 2020)
81 or plagioclase that formed the anorthosite crust of the Moon (Wood, Dickey, Marvin and Powell, 1970; Warren, 1985).
82 Partial melting of this primary crust can further lead to the formation of a basaltic secondary crust (Mandler and Elkins-
83 Tanton, 2013). However, the petrology and geochemistry of some differentiated meteorites, such as EC002, indicate
84 that their parent liquids formed directly from chondritic material at shallow depth in planetesimals. The metasomatic
85 minerals found in the CO and CV chondrites (Ganino and Libourel, 2017) could be considered as products resulting
86 from the first step of the thermal heating that could have later induced this partial melting.

87 An additional argument for the presence of a stable chondritic crust in planetesimals is the paleomagnetic record of
88 a magnetic field in the parent body of primitive meteorites which implies both a core – hence a differentiated body –
89 and preserved, undifferentiated, chondritic material. Thermal models of planetesimals available in the literature fail to
90 account for the preservation of a chondritic crust in differentiated planetesimals (Weiss and Elkins-Tanton, 2013). Only
91 few studies specifically considered the link between the thermal evolution of a planetesimal and its accretion history
92 advocating that chondrites and achondrites can originate from the same parent body. For instance, Neumann et al.
93 (2012) considered accretion by radial growth and modeled the differentiation of planetesimals by Darcy flow. They
94 showed that the degree of differentiation varies significantly as a function of the accretion history. Kaminski et al. (2020)
95 introduced a new accretion model for parent bodies of iron meteorites accounting both for planetesimals differentiation
96 and chondritic crust preservation. Their thermal model took into account solid-state convection, but did not consider
97 explicitly the possibility of generating a magma ocean episode. Dodds, Bryson, Neufeld and Harrison (2021) also
98 proposed that crust constituents can be supplied by accretion of undifferentiated material, and considered the possibility
99 of solid-state convection, followed by a magma ocean episode. But their model still predicts metamorphism of the
100 chondritic material in the crust even though they allowed accretion to start relatively late (0.8 Myr) in order to mitigate
101 heating.

102 In the present paper, we introduce and model a refined accretion and differentiation scenario that explains the
103 preservation of a chondritic crust, and produces a thermal record consistent with geochemical and petrological
104 evidences provided by some meteorites. As an illustration of the model predictions, we consider the particular case of
105 the andesite EC002.

Structure of differentiated planetesimals




Figure 1: Accretion curves giving the evolution of the normalized radius of planetesimals (with $R_0/R_f = 0.5$ and $t_{acc} = 0.4$ Myr) for three values of the parameter β : $\beta = 0.5$ (red line), the typical value used in the model, $\beta = 1$ (blue line) corresponding to an exponential law, and $\beta = 1.5$ (purple line). The grey vertical line corresponds to $t = t_{acc} = 0.4$ Myr.

2. Thermal regimes and radial structures of accreting planetesimals

As inferred from petrological data, EC002 was formed by $\approx 25\%$ partial melting of a chondritic source (Barrat et al., 2021). In the following, we establish in which conditions chondritic material can be preserved in a differentiated planetesimal.

2.1. Accretion

Planetesimals are supposed to form by the accretion of cold dust and pebbles in the mid-plane of the accretion disk (Visser and Ormel (2016); Johansen and Lambrechts (2017)). Processes such as turbulence in the disk can produce locally strong concentrations reaching thresholds that trigger quasi-instantaneous accretion of ≈ 100 km size planetesimals (Johansen, Oishi, Low, Klahr, Henning and Youdin, 2007; Cuzzi, Hogan and Shariff, 2008) thus bypassing the so-called meter-size barrier (Weidenschilling, 1977). In such a scenario, most planetesimals should be born big, as also suggested by simulations of the formation of embryos from planetesimals (Morbidelli, Bottke, Nesvorný and Levison, 2009). These accretion thresholds can theoretically be reached at various times and locations in the disk, depending on the availability and concentration of dust and pebbles. Hf – W ages of iron meteorites indicate that their parent bodies accreted over the first 1.5 Myr of the solar system, whereas the accretion of the parent bodies of achondrites and chondrites was probably completed within the first 4 – 5 Myrs (Kleine, Budde, Burkhardt, Kruijjer, Worsham, Morbidelli and Nimmo, 2020). In the inner regions of the accretion disk, within the orbit of Jupiter, primordial dust can be rapidly exhausted, within a Myr or so, because of the barrier to dust migration from the outer disk created by the early formation of Jupiter (Morbidelli, Lambrechts, Jacobson and Bitsch, 2015; Haugbølle, Weber, Wielandt, Benítez-Llambay, Bizzarro, Gressel and Pessah, 2019). However, accretion of planetesimals can continue after 1 Myr since the flux of pebbles is never totally suppressed even for a fully-grown Jupiter.

In the following, we consider that growth of planetesimals can, in some cases, continue after the quasi-instantaneous accretion of a nucleus of several tens of km size (Kaminski et al., 2020). We further choose a Weibull accretion law to parametrize protracted accretion. The radius of the rocky body $R(t)$ grows from R_0 to the terminal radius R_f according to:

$$R(t) = R_f \left[1 - \left(1 - \left[\frac{R_0}{R_f} \right]^3 \right) e^{-(t/t_{acc})^\beta} \right]^{1/3}, \quad (1)$$

with t_{acc} the characteristic accretion timescale and β a constant. Equation (1) will be referred as the “protracted accretion” law hereafter. Three families of accretion scenarios can be defined as a function of the value of β (Figure 1). First, $\beta = 1$ corresponds to a specific case described simply by an exponential law. For $\beta < 1$, the planetesimal accretes material more rapidly than an exponential law at early times ($t < t_{acc}$) and supplied material more slowly at




Figure 2: (a): Influence of R_0/R_f on the accretion timescale $\tau_{99\%}$. (b): Diagram showing the three families of accretion history as a function of the parameters t_{acc} and β . Each solid line represents an iso-value of $\tau_{99\%}$. "Quasi instantaneous accretion" stands for planetesimals whose accretion timescale $\tau_{99\%}$ is smaller than the lifetime of ^{26}Al ($t_{1/2}^{Al}/\ln 2 = 1$ Myr). "Overlong accretion" represents accretion which could take place in debris disk since $\tau_{99\%}$ is greater than the dissipation of the accretion disk (≈ 10 Myr). The red star corresponds to the typical parameters of the accretion law used in the model ($t_{acc} = 0.4$ Myr and $\beta = 0.5$). The orange points correspond to the extreme values of the accretion law used in the sensitivity study, in section 3.5.

135 $t > t_{acc}$, so that the terminal radius is reached later in the history of the planetesimal. For $\beta > 1$, the early accretion
 136 is slower than an exponential law at early time but material is brought more efficiently for $t > t_{acc}$ and the terminal
 137 radius is reached earlier in the planetesimal history.

138 We define the characteristic timescale of this law $\tau_{99\%}$ as the time at which the planetesimal radius reaches 99%
 139 of the terminal radius. After some developments, $\tau_{99\%}$ writes:

$$140 \quad \tau_{99\%} = t_{acc} \left[\ln \left(\frac{1 - (R_0/R_f)^3}{0.03} \right) \right]^{1/\beta} \quad (2)$$

141 The accretion timescale $\tau_{99\%}$ is a function of the three parameters of the Weibull law. In figure 2 (a), the variation of $\tau_{99\%}$
 142 with R_0/R_f shows that for R_0/R_f in the range 0 – 80%, the accretion timescale remains stable. For $R_0/R_f > 80\%$,
 143 the accretion timescale decreases dramatically, and the planetesimal undergoes a quasi-instantaneous accretion. Within
 144 this domain ($0\% < R_0/R_f < 80\%$) only t_{acc} and β have an influence on the accretion history.

145 Moreover, we can add two constraints on $\tau_{99\%}$. First, accretion has to be consistent with the lifetime of
 146 protoplanetary disk. Gas giants like Jupiter may have formed during the first Myr after CAI (Kruijer et al., 2017),
 147 and the dissipation of the solar nebula can range from a few to 10 Myr (Karl E. Haisch, Lada and Lada, 2001). We
 148 thus consider that $\tau_{99\%}$ has to be smaller than 10 Myr. Second, we compare the accretion timescale to the characteristic
 149 timescale of radioactive decay of ^{26}Al . If $\tau_{99\%} \ll t_{1/2}^{Al}/\ln 2$, accretion can be considered instantaneous ($R(t) \approx R_f$)
 150 and the content of radioactive ^{26}Al , hence the heating rate in the planetesimal are maximal. This case happens for
 151 $R_0/R_f \rightarrow 1$ and for small values of t_{acc} . Following these two constrains, we provide an abacus showing three families
 152 of accretion history in Figure 2 (b) : (i) when $\tau_{99\%} < 1$ Myr, accretion is considered quasi instantaneous, (ii) when
 153 $\tau_{99\%} > 10$ Myr, the accretion is too long compared to the lifetime of the disk, (iii) when $\tau_{99\%}$ is between these two
 154 end-members, the accretion is called "protracted".

155 2.2. Composition

156 We consider that the accreted undifferentiated chondritic material is a binary mixture of iron and silicates, with a
 157 mass fraction x_{Fe} and x_{Sil} , respectively. The average density of the undifferentiated material ρ is:

$$158 \quad \frac{1}{\rho} = \frac{x_{\text{Fe}}}{\rho_{\text{Fe}}} + \frac{x_{\text{Sil}}}{\rho_{\text{Sil}}}, \quad (3)$$

159 with ρ_i the density of phase i . We consider that melt crystallization can generate two types of crystals, one that is
 160 negatively buoyant relative to the melt (called H hereafter) and one that is positively buoyant relative to the melt
 161 (called L hereafter). The later produces a flotation crust at the top of the magma ocean, whereas the first induces the
 162 formation of a cumulate at the core-mantle boundary (CMB). The relative proportion of the two families of crystal
 163 is $\phi_{L,0}/\phi_{H,0} = 10/90$. This ratio is based on the proportion of plagioclase crystals relative to olivine and pyroxene
 164 crystals measured in some chondrites (Weisberg, Boesenberg, Kozhushko, Prinz, Clayton and Mayeda, 1995). Crystals
 165 L and H are effective “mineralogical phases” that provide a simplified description of the more complex mineral
 166 assemblages present in chondrites. As a consequence, L and M minerals are characterized by a solidus and a liquidus
 167 temperature. Between the solidus and the liquidus, the solid fraction in the melt, ϕ_{cr} (in vol%), is given by:

$$168 \quad \phi_{cr} = \phi_{L,cr} + \phi_{H,cr}, \quad (4)$$

$$169 \quad \phi_{L,cr} = \phi_{L,0} \frac{T_l^L - \bar{T}}{T_l^L - T_s^L}, \quad (5)$$

$$170 \quad \phi_{H,cr} = \phi_{H,0} \frac{T_l^H - \bar{T}}{T_l^H - T_s^H}, \quad (6)$$

171 where \bar{T} is the volume averaged bulk temperature in the planetesimal, T_l^i and T_s^i are the liquidus and the solidus
 172 temperature of mineral i , respectively, and $\phi_{i,0}$ its initial proportion (in vol%), and $\phi_{i,cr}$ the proportion of crystal i .

173 2.3. Thermal regime: from conduction regime to solid-state convection

174 2.3.1. Conduction regime

175 The bulk energy budget of undifferentiated planetesimal of radius $R(t)$ leads to the following equation (Kaminski
 176 et al., 2020) :

$$177 \quad \rho c_p \frac{d\bar{T}}{dt} = H_0 - \frac{3}{R} Q_s - \rho c_p \frac{3\bar{T}}{R} \frac{dR}{dt}, \quad (7)$$

178 with $c_p = x_{Fe} c_{p,Fe} + x_{Sil} c_{p,Sil}$ the planetesimal average heat capacity, H_0 the total rate of internal heating, and Q_s the
 179 surface heat flux. In equation (7), the dR/dt term corresponds to the effect of accretion of cold material on the thermal
 180 evolution of the planetesimal. Internal heating rates by radioactive elements ^{26}Al and ^{60}Fe are given by:

$$181 \quad H_{0,Fe} = \rho F_{Fe} \left[\frac{^{60}\text{Fe}}{^{56}\text{Fe}} \right]_0 \frac{\ln 2}{t_{1/2}^{Fe}} \mathcal{E}_{Fe} e^{-t \ln 2 / t_{1/2}^{Fe}}, \quad (8)$$

$$182 \quad H_{0,Al} = \rho F_{Al} \left[\frac{^{26}\text{Al}}{^{27}\text{Al}} \right]_0 \frac{\ln 2}{t_{1/2}^{Al}} \mathcal{E}_{Al} e^{-t \ln 2 / t_{1/2}^{Al}}, \quad (9)$$

183 where F_i is the chondritic abundance of i , $[i/j]_0$ is the initial abundance of radioactive and stable isotopes i and j , \mathcal{E}_i
 184 is the decay energy per atom, $t_{1/2}^i$ the half-life.

185 As the planetesimal melts, latent heat is consumed, and likewise, when it crystallizes, latent heat is released. Outside
 186 the solidus to liquidus temperature interval, latent heat is zero. When the temperature falls within the solidus and the
 187 liquidus, latent heat contributes to the energy budget as an additional internal heat source quantified as:

$$188 \quad H_{LH,Fe,0} = -L_{Fe} x_{Fe} \rho \frac{d\bar{T}}{dt} \frac{1}{T_l^{Fe} - T_s^{Fe}}, \quad (10)$$

$$189 \quad H_{LH,Sil,0} = -L_{Sil} x_{Sil} \rho \frac{d\bar{T}}{dt} \frac{1}{T_l^{Sil} - T_s^{Sil}}, \quad (11)$$

with L_i the latent heat of component i (iron or silicate). The total rate of internal heating is thus $H_0 = H_{0,Al} + H_{0,Fe} +$
 $H_{LH,Sil,0} + H_{LH,Fe,0}$.

Our model considers the temperature at the surface of the body imposed by the temperature of the disk, which is
 a reasonably good approximation for planetesimals heated by ^{26}Al (Dodds et al., 2021), and is set to $T_s = 0^\circ\text{C}$. At

Structure of differentiated planetesimals

Parameters	Symbols	Values	Units	Ref.
Iron content (mass fraction)	x_{Fe}	0.18	-	(4)
Silicate content (mass fraction)	$x_{\text{Sil}} = 1 - x_{\text{Fe}}$	0.82	-	(4)
Chondritic abundance (^{60}Fe)	F_{Fe}	$2.14 \cdot 10^{24}$	kg^{-1}	(1)
Chondritic abundance (^{26}Al)	F_{Al}	$2.62 \cdot 10^{23}$	kg^{-1}	(1)
Decay energy per atom (^{60}Fe)	\mathcal{E}_{Fe}	$4.87 \cdot 10^{-13}$	J	(1)
Decay energy per atom (^{26}Al)	\mathcal{E}_{Al}	$4.42 \cdot 10^{-13}$	J	(1)
Initial relative abundance (^{60}Fe)	$[\text{}^{60}\text{Fe}/\text{}^{56}\text{Fe}]_0$	10^{-8}	-	(2)
Initial relative abundance (^{26}Al)	$[\text{}^{26}\text{Al}/\text{}^{27}\text{Al}]_0$	$5 \cdot 10^{-5}$	-	(2)
Half-life (^{60}Fe)	$t_{1/2}^{\text{Fe}}$	2.6	Myr	(1)
Half-life (^{26}Al)	$t_{1/2}^{\text{Al}}$	0.717	Myr	(1)
Iron density	ρ_{Fe}	7800	kg m^{-3}	(4)
Silicate density	ρ_{Sil}	3200	kg m^{-3}	(4)
Crystal/melt drop of density	$\Delta\rho$	100	kg m^{-3}	-
Iron thermal conductivity	λ_{Fe}	50	$\text{W m}^{-1} \text{K}^{-1}$	(4)
Silicate thermal conductivity	λ_{Sil}	3	$\text{W m}^{-1} \text{K}^{-1}$	(4)
Thermal expansion (Fe)	α_{Fe}	$7.7 \cdot 10^{-5}$	K^{-1}	(1)
Thermal expansion (Silicate)	α_{Sil}	$2 \cdot 10^{-5}$	K^{-1}	(1)
Specific heat (Silicate)	$c_{p,\text{Sil}}$	1168	$\text{JK}^{-1} \text{kg}^{-1}$	(3)
Specific heat (Fe)	$c_{p,\text{Fe}}$	622	$\text{JK}^{-1} \text{kg}^{-1}$	(3)
Latent heat (Fe)	L_{Fe}	250	kJ kg^{-1}	(4)
Latent heat (Silicate)	L_{Sil}	500	kJ kg^{-1}	(4)
Liquidus (Fe)	T_l^{Fe}	1615	$^{\circ}\text{C}$	(4)
Solidus (Fe)	T_s^{Fe}	990	$^{\circ}\text{C}$	(4)
Liquidus (Silicate and chondritic mat.)	T_l^{Sil}	1700	$^{\circ}\text{C}$	(4)
Solidus (Silicate)	T_s^{Sil}	1200	$^{\circ}\text{C}$	(4)
Solidus light mineral	T_s^{L}	1200	$^{\circ}\text{C}$	(6)
Solidus heavy mineral	T_s^{H}	1200	$^{\circ}\text{C}$	(6)
Liquidus light mineral	T_l^{L}	1350	$^{\circ}\text{C}$	(6)
Liquidus heavy mineral	T_l^{H}	1700	$^{\circ}\text{C}$	(6)
Activation energy (Sil)	$E_{a,\text{Sil}}$	250	kJ mol^{-1}	(4)
Activation energy (Fe)	$E_{a,\text{Fe}}$	13	kJ mol^{-1}	(4)
Viscosity of iron melt at T_l^{Fe}	η_0	1	Pa s	(3)
Melt fraction factor for silicate	σ_{Sil}	21	-	(4)
Melt fraction factor for iron	σ_{Fe}	4	-	(4)
Suspended crystals radius	r	$3 \cdot 10^{-3}$	m	(5)

Table 1

Parameters used in the thermal model. References are the following : Neumann et al. (2014) (1), Tang and Dauphas (2012) (2), Neumann et al. (2012) (3), Kaminski et al. (2020) (4), Solomatov (2000) (5), adapted and simplified from Righter and Drake (1997) (6).

the early stages of the thermal evolution of the planetesimal, heat is evacuated by conduction and the surface heat flux can be calculated analytically (Carslaw and Jaeger, 1959). For an exponential internal heat generation source given by $H(t) = H_0 \exp(-\lambda t)$ with zero initial and surface temperatures, the surface heat flux is:

$$Q_s(t) = -\frac{\kappa H_0}{\lambda} \left[\sqrt{\frac{\lambda}{\kappa}} \tan^{-1/2} \left(R \sqrt{\frac{\lambda}{\kappa}} \right) - \frac{1}{R} \right] \exp(-\lambda t) - \frac{2RH_0}{\pi} \sum_{n=1}^{+\infty} \frac{\exp\left(-\frac{\kappa n^2 \pi^2 t}{R^2}\right)}{n^2 - \frac{\lambda}{\kappa} \left(\frac{R}{\pi}\right)^2}. \quad (12)$$

190 Here we consider only the heat production by ^{26}Al with $\lambda = \ln 2/t_{1/2}^{\text{Al}}$. Because conduction is not a very efficient heat
191 transfer, the bulk temperature of the planetesimal increases, until new processes occur in the interior of the planetesimal.

192 2.3.2. Onset of solid-state convection

193 As heating proceeds, the planetesimal is bound to melt. Partial melting implies a strong decrease of viscosity and
194 can generate instabilities of pure thermal origin or related to liquid iron percolation that trigger solid-state convection

at planetary scale (Kaminski et al., 2020). Convection is characterized by two dimensionless numbers (Roberts, 1967), the Rayleigh-Roberts number Ra_H^* that accounts for the vigor of convection, and the Prandtl number that quantifies the importance of inertia:

$$Ra_H^* = \frac{\alpha \rho g H^* R^5}{\kappa \eta \lambda}, \quad (13)$$

$$Pr = \frac{\nu}{\kappa}, \quad (14)$$

where α is the thermal expansion coefficient, $g = 4/3\pi\mathcal{G}\rho R$ is the surface gravity with \mathcal{G} the gravitational constant, λ is the thermal conductivity, κ is the thermal diffusivity, η is the dynamic viscosity, $\nu = \eta/\rho$ is the kinematic viscosity, and H^* is the effective rate of internal heating that takes into account secular cooling (Limare, Jaupart, Kaminski, Fourel and Farnetani, 2019; Limare, Kenda, Kaminski, Surducan, Surducan and Neamtu, 2021):

$$H^* = H - \rho c_p \frac{d\bar{T}}{dt}. \quad (15)$$

Convection starts if Ra_H^* is larger than the critical value $Ra_{H,c} = 5758$ (Schubert, Turcotte and Olson, 2001). In the case of a material whose viscosity is strongly temperature dependent, convection will develop under a stagnant lid after a delayed onset time τ_{SL} (Davaille and Jaupart, 1993). In the regime of solid-state convection, the heat budget of the convecting planetesimal is still given by equation (7) but with a surface heat flux now given by the scaling law (Davaille and Jaupart, 1993; Choblet and Sotin, 2000):

$$Q_s = \lambda \left(\frac{\alpha \rho g}{\kappa \eta} \right)^{1/3} \left(\frac{\Delta T_v}{C_T} \right)^{4/3}, \quad (16)$$

$$\Delta T_v = -C_v \eta / \frac{d\eta}{dT}, \quad (17)$$

where ΔT_v is the characteristic viscous temperature scale for convection with a stagnant lid, $C_v = 2.24$ (Davaille and Jaupart, 1993), and C_T a constant that depends on the mechanical boundary condition (here we will use $C_T = 3.31$ following Limare et al. (2021)). The continuity of the heat flux through the stagnant lid leads to the determination of its thickness δ_{SL} :

$$\delta_{SL}(\bar{T}) = \frac{R}{2} \left(1 - \sqrt{1 - \frac{4\lambda(\bar{T} - \Delta T_v - T_s)}{Q_s R}} \right), \quad (18)$$

where T_s is the surface temperature.

The scaling laws introduced above emphasizes the key influence of the viscosity on the dynamics of convection. In planetesimals, this issue is of peculiar importance because of the large variation of viscosity expected during the progressive heating of the material and its partial melting. For the silicate component we use a modified Arrhenius law to account for these dependencies (Scott and Kohlstedt, 2006; Kaminski et al., 2020):

$$\eta(\bar{T}, \phi_{cr}) = \eta_0 f(\phi) \exp \left[\frac{E_{a,Sil}}{R_g} \left(\frac{1}{\bar{T}} - \frac{1}{T_0} \right) \right], \quad (19)$$

$$f(\phi_{cr}) = \exp[-\sigma_{Sil}(1 - \phi_{cr})]. \quad (20)$$

where $f(\phi_{cr})$ encompasses the influence of partial melting on the viscosity, η_0 is the reference viscosity at $T_0 = 1000^\circ\text{C}$, $E_{a,Sil}$ the activation energy for silicates and R_g the ideal gas constant. Because the amount of melt is related to the temperature, the variation of viscosity with temperature is amplified, and the convection may further transition to a magma ocean episode, as explained below.

2.4. Magma ocean episode

2.4.1. Rheological transition: magma ocean and core formation

At a certain degree of melting, the solid matrix becomes unstable and the behavior of the mixture becomes that of a suspension of solid crystals in a liquid magma (Solomatov, 2000). The type of convection changes at rheological

Structure of differentiated planetesimals




Figure 3: (a) Illustration of the stratification inside a planetesimal during a magma ocean episode. (b) Illustration of the iron-silicate differentiation that occurs at the base of the crust. Between $R(t)$ and R_{seg} , the crust remains cold and chondritic. Between R_{seg} and R_{MO} , the crust temperature is high enough to trigger iron-silicate segregation. Iron is drained, and the remaining crust is made of silicate component only. As iron leaves, the lower part of the crust thins.

232 transition $\phi_{RT} = 60\%$ (Elkins-Tanton, 2012; Guazzelli and Pouliquen, 2018), from laminar solid state convection to
 233 liquid-like, turbulent convection. In the so called “magma ocean” regime, the viscosity law (19) is modified so that
 234 $f(\phi)$ is now expressed by a Krieger-Dougherty law (Krieger and Dougherty, 1959):

$$235 \quad f(\phi) = \left(1 - \frac{\phi}{\phi_c}\right)^{-2.5\phi_c}, \quad (21)$$

236 where ϕ stands for the volume fraction of crystals in suspension, whereas ϕ_{cr} is the total amount of crystals in the
 237 mantle, either in suspension or settled. We set ϕ_c at 60.002% to ensure the continuity of the viscosity at $\phi = \phi_{RT}$ and
 238 to reach a viscosity of 14 Pa s for purely liquid magma, consistent with the estimates of Rubie, Melosh, Reid, Liebske
 239 and Righter (2003). The variation of the viscosity as a function of the temperature is given in Appendix 2.4.1 assuming
 240 an equilibrium melting.

241 Beyond the rheological transition, the viscosity is reduced by a few orders of magnitude and opens the way to
 242 metal extraction (Sturtz, Limare, Tait and Kaminski, 2021c). As a matter of fact, before the rheological transition, the
 243 planetesimal is a mush with a low degree of partial melting and iron migration to the center of the planetesimal occurs
 244 by porous flow (Neumann et al., 2012). The timescale for iron to reach the center of the planetesimal is 1 – 10 Myr,
 245 which is long compared to the time required for the onset of the magma ocean (less than 1 Myr, as it will be discussed
 246 later). Hence we assume that no metal-silicate segregation happened before the rheological transition. On the other
 247 hand, once the rheological transition has been reached, there is no more matrix resistance to the iron migration and we
 248 consider that it happens instantaneously (Höink, Schmalzl and Hansen, 2006).

249 A second important consequence of the low viscosity of the convective fluid once the rheological transition has
 250 been reached is the potential development of relative motions between melt and crystals, and ultimately the possibility
 251 of forming layers of segregated crystals: cumulate of thickness δ_H can grow by deposition of the heavy crystals at the
 252 core-mantle boundary (CMB), whereas a layer of light crystals can accumulate below the chondritic material brought
 253 at the surface of the planetesimal, the two contributions forming a crust of total thickness δ_c , as drawn in Figure 3 (a).

254 2.4.2. Thermal state of the bulk magma ocean and the core

255 The magma ocean is treated as a convecting shell. Through the introduction of a geometrical factor for the silicate,
 256 convecting mantle:

$$257 \quad f_c = \frac{R_c}{R_{MO}} = \left(\frac{\rho}{\rho_{Fe}} x_{Fe}\right)^{1/3}, \quad (22)$$

258 where R_c is the core radius and R_{MO} is the distance between the top of the magma ocean and the center of the
 259 planetesimal (figure 3(a)). With parameters summarized in Table 1, $f_c = 0.42$ right after the core differentiation.

260

261 The energy budget for the magma ocean and for the core are:

$$262 \quad \rho_{\text{Sil}} c_{p,\text{Sil}} \frac{dT_{\text{MO}}}{dt} = H_{\text{MO}} + H_{\text{LH,Sil}} + Q_{\text{CMB}} \frac{S_c}{V_m} - Q_s \frac{S_m}{V_m}, \quad (23)$$

$$263 \quad \rho_{\text{Fe}} c_{p,\text{Fe}} \frac{dT_c}{dt} = H_c + H_{\text{LH,Fe}} - Q_{\text{CMB}} \frac{S_c}{V_c}, \quad (24)$$

264 where T_{MO} and T_c are the bulk average temperatures of the magma ocean and of the core, respectively, Q_{CMB}
 265 is the heat flux at the CMB, $S_m = 4\pi R^2 [1 - (\delta_c/R)^2]$ the top surface area of the magma ocean, $V_m =$
 266 $4/3\pi R^3 [(1 - \delta_c/R)^3 - (f_c \{1 - \delta_c/R\} + \delta_H/R)^3]$ the volume of the magma ocean, $S_c = 4\pi f_c^2 R^2 (1 - \delta_c/R)^2$ the
 267 surface area of the core and $V_c = 4/3\pi R^3 f_c^3 (1 - \delta_c)^3$ its volume.

268 As Al is highly lithophile, we assume that all ^{26}Al is fully partitioned into the magma ocean. On the other hand,
 269 we consider that all iron is partitioned into the core, hence that no ^{60}Fe is present in the magma ocean. Accordingly,
 270 the rate of internal heating in each reservoir can be written by modulating (8)-(11) by a concentration factor:

$$271 \quad H_c = H_{0,\text{Fe}} f_c^{-3} \left(1 - \frac{\delta_c}{R}\right)^{-3}, \quad (25)$$

$$272 \quad H_{\text{MO}} = H_{0,\text{Al}} \left[\left(1 - \frac{\delta_c}{R}\right)^3 - \left(f_c \left\{1 - \frac{\delta_c}{R}\right\} + \frac{\delta_H}{R}\right)^3 \right]^{-1}, \quad (26)$$

273 whereas the release of latent heat in the two reservoirs is given by:

$$274 \quad H_{\text{LH,Fe}} = -L_{\text{Fe}} \rho_{\text{Fe}} \frac{dT_c}{dt} \frac{1}{T_l^{\text{Fe}} - T_s^{\text{Fe}}}, \quad (27)$$

$$275 \quad H_{\text{LH,Sil}} = -L_{\text{Sil}} \rho_{\text{Sil}} \frac{dT_{\text{MO}}}{dt} \frac{1}{T_l^{\text{Sil}} - T_s^{\text{Sil}}}. \quad (28)$$

276 The potential presence of a crust at the surface of the magma ocean and at the CMB yields expressions for the heat
 277 fluxes that are more complex than the classical scaling laws used for convective systems. The crust acts like a thermal
 278 insulator and the temperature at the base of the crust $T_{b,c}$ is linked to the convective heat flux through:

$$279 \quad Q_s = \lambda_{\text{Sil}} \left(\frac{\alpha_{\text{Sil}} \rho_{\text{Sil}} g}{\kappa_{\text{Sil}} \eta} \right)^{1/3} \left(\frac{T_{\text{MO}} - T_{b,c}}{C_T} \right)^{4/3}. \quad (29)$$

280 Furthermore, heat transfer occurs by conduction in the crust and follows Fourier's law (Jaupart and Mareschal (2010),
 281 pp.36):

$$282 \quad Q_s = \lambda_{\text{Sil}} \frac{T_{b,c} - T_s}{\delta_c(t)} \frac{R(t)}{R(t) - \delta_c(t)}. \quad (30)$$

283 At the beginning of the magma ocean episode, the crust is only composed of undifferentiated material brought
 284 by accretion. The presence of radioactive elements in the magma ocean later induces high temperatures close to the
 285 liquidus, which in turn lead to partial melting of the base of the crust. As silicates are more refractory than iron, the base
 286 of the crust is hot enough for iron melting to occur. Due to the high density contrast, liquid iron segregates from the
 287 base of the crust, and the crust is composed of a chondritic upper part, and an iron-depleted lower part (Figure 3 (b)).
 288 This consideration has two main consequences. First, when iron is extracted, the crust thickness decreases. Second,
 289 as the crustal thinning occurs at a temperature below the liquidus of silicate, the heat flux increases so the magma
 290 ocean cools at a sub-liquidus temperature. As in the previous section when dealing with the massive iron/silicate
 291 segregation, we consider the temperature at which the iron is supposed to be extracted from the crust as being the
 292 rheological temperature of silicates (1366°C).

293 Sturtz, Kaminski, Limare and Tait (2021a) developed a physical model establishing whether crystals formed during
 294 magma ocean cooling can be entrained by convection or not, based on a dimensionless parameter defined as the ratio

295 between the convective shear stress and the buoyancy of crystals. This dimensionless parameter is a modified Shields
 296 number that combines the convection parameters and the crystals characteristics:

$$297 \quad \zeta = \frac{\kappa\eta}{h_{MO}^2 \Delta\rho g r} Ra_H^{*3/8}, \quad (31)$$

298 with r the crystal radius, $\Delta\rho$ the density difference between the crystals and the melt, and $h_{MO} = R[1 - \delta_c/R -$
 299 $\delta_H/R - f_c(1 - \delta_c/R)]$ the thickness of the magma ocean. If ζ is below a critical value ζ_c crystals are not entrained
 300 by the convection and they can deposit, whereas if $\zeta > \zeta_c$, convective flow can keep crystals in suspension (Shields,
 301 1936; Charru, Mouilleron and Eiff, 2004). Sturtz et al. (2021a) validated this criterion experimentally for the formation
 302 of both cumulates and floating lids, and obtained $\zeta_c = 0.29 \pm 0.17$. In the present case, because the viscosity of the
 303 magma ocean is very low (i.e. liquid like), the Shields number falls well below the critical value.

304 At the CMB, the way heat is transported depends on the magma ocean temperature. In the initial stage of the
 305 magma ocean, its temperature increases by internal heating up to maximum values close to the silicate liquidus. In
 306 these conditions there is no crystal cumulate at the CMB, and we consider that the core is always well mixed and
 307 characterized by an average temperature. On the magma ocean side, a thermal boundary layer forms at the CMB. In
 308 the stagnant lid regime, the TBL thickness at the CMB δ_{CMB} is linked to the stagnant lid thickness δ_{SL} by (Solomatov,
 309 1995; Bryson, Neufeld and Nimmo, 2019; Dodds et al., 2021):

$$310 \quad \delta_{CMB} = \delta_{SL} \frac{\Delta T_v}{T_{MO} - T_s}, \quad (32)$$

311 with $\Delta T_v = -C_v\eta/\partial_T\eta$ if $T_{MO} > T_c$, and $\Delta T_v = -C_v\eta_{Fe}/\partial_T\eta_{Fe}$ if $T_{MO} < T_c$. The CMB heat flux can then be expressed
 312 as a function of the surface heat flux as:

$$313 \quad Q_{CMB} = Q_s \frac{T_{MO} - T_c}{\Delta T_v}. \quad (33)$$

314 As the magma ocean cools down, a cumulate of “heavy silicate crystals” can form at the CMB. As long as the
 315 cumulate thickness is thinner than the TBL thickness at the CMB, heat transfer between the core and the mantle occurs
 316 by convection and equation (33) remains valid. If the cumulate is thicker than the TBL, the conductive heat flux Q_{CMB}
 317 in given by:

$$318 \quad Q_{CMB} = \lambda_{Sil} \frac{T_{b,MO} - T_{CMB}}{\delta_H(t)} \frac{R_c(t) + \delta_H(t)}{R_c(t)}, \quad (34)$$

319 where $T_{b,MO}$ is the temperature at the base of the magma ocean, and T_{CMB} is the temperature at the surface of the core.
 320 If the magma ocean is hotter than the core, we assume that the core acts like a thermostat, and sets $T_{CMB} = T_c$, whereas
 321 $T_{b,MO}$ is obtained from the heat flux using the scaling law:

$$322 \quad Q_{CMB} = \lambda_{Fe} \left(\frac{\alpha_{Sil}\rho_{Sil}g_c}{\kappa_{Sil}\eta} \right)^{1/3} \left(\frac{T_{MO} - T_{b,MO}}{C_T} \right)^{4/3}. \quad (35)$$

323 Similarly, if the the core is hotter than the magma ocean, we assume $T_{b,MO} = T_{MO}$, whereas the temperature T_{CMB} is
 324 obtained from the heat flux at the CMB given by:

$$325 \quad Q_{CMB} = \lambda_{Fe} \left(\frac{\alpha_{Fe}\rho_{Fe}g_c}{\kappa_{Fe}\eta_{Fe}} \right)^{1/3} \left(\frac{T_c - T_{CMB}}{C_T} \right)^{4/3}. \quad (36)$$

326 Within this framework, the thermal evolution of the system is totally determined by (23)-(24). However, two
 327 parameters are missing to solve this set of equations: the thickness of the crust δ_c and the thickness of the heavy
 328 cumulate δ_H .

329 2.4.3. Mechanical evolution of the crust and of the cumulate

330 At the surface, the crust is fed both by the accretion of chondritic material and by the flotation of light crystals
 331 from the magma ocean. The growth of the chondritic part of the crust follows the protracted accretion law (1) while

332 the thickening due to flotation of light crystals follows the law given by Sturtz, Limare, Tait and Kaminski (2021b).
 333 The total growth rate is then given by:

$$334 \quad \frac{d\delta_c}{dt} = \frac{dR}{dt} + c_d v_s \phi_L, \quad (37)$$

335 where ϕ_L is the volume fraction of light crystals in suspension in the magma ocean, v_s is the sedimentation velocity of
 336 these light crystals and c_d is a constant of order unity. Experiments in different types of convective systems have shown
 337 that for diluted suspensions v_s scales with Stokes' velocity (Martin and Nokes, 1988, 1989; Lavorel and Le Bars, 2009;
 338 Sturtz et al., 2021b). To take into account the influence of crystals concentration on the settling velocity v_s , we use the
 339 viscosity described in section 2.4.1, that takes into account the effect of temperature T and the effect of the suspension
 340 on the viscosity:

$$341 \quad v_s = \frac{\Delta\rho g r^2}{\eta(T, \phi)}, \quad (38)$$

342 with $\Delta\rho$ the density difference between the melt and crystals, r the crystal radius, and ϕ the volume fraction of crystals
 343 in suspension.

344 When the MO temperature decreases below the silicate liquidus, heavy crystals nucleate, sink and form a cumulate
 345 at the CMB. The rate of thickening of this layer δ_H is given by:

$$346 \quad \frac{d\delta_H}{dt} = c_d v_s \phi_H, \quad (39)$$

347 with ϕ_H the volume fraction of heavy crystal in suspension in the magma ocean. Note that equations (37) and (39)
 348 are valid only when the magma ocean temperature is below the light and heavy silicate crystals melting temperature,
 349 respectively. If the magma ocean temperature is higher than their liquidus, the layers undergo melting, i.e., are thermally
 350 eroded.

351 **3. Results - what was the size of EC002 parent body?**

352 **3.1. Necessity to take accretion into account**

353 A magma ocean episode is consistent both with an efficient core formation by metal-silicate differentiation and with
 354 the preservation of an undifferentiated crust at the surface. Because EC002 analyses indicate an early accretion, we first
 355 consider a scenario in which its parent body accreted instantaneously at t_0 (Li, Youdin and Simon, 2019). Within the
 356 framework developed above, this scenario simply implies $dR/dt = 0$ in equation (7). We solved numerically equation
 357 (7) using a finite difference scheme. As the conservation of energy is considered for the entire planetesimal, the physical
 358 parameters do not have a radial dependence. They are all volume averaged, for each reservoir, as underlined in equations
 359 23 and 24. As a consequence, we only need a temporal discretization, and we use an order one explicit scheme. In the
 360 parameter space (R, t_0) we track the conditions required to produce a magma ocean episode. As shown in Figure 4 (a),
 361 a magma ocean occurs in a planetesimal of radius R if it accreted early enough. If the accretion takes place too late,
 362 the radioactive source is not sufficient to supply the required energy to reach the rheological transition.

363 Once the magma ocean episode has started (at ≈ 0.45 Myr for a 100 km body, accreted instantaneously at $t_0=0$), a
 364 chondritic crust remains at the surface as part of the stagnant lid with an initial thickness spanning from 0.5 to 1.5 km
 365 depending on the size of the planetesimal. However, as illustrated in Figure 4 (b), this crust is bound to undergo a first
 366 melting at ≈ 0.64 Myr that decreases its thickness. This is due to the fact that the crust insulates thermally the magma
 367 ocean where all ^{26}Al is essentially stored. Note that the crust also contains ^{26}Al , but the effects are negligible on the
 368 global budget. After this first melting episode, the chondritic reservoir that can be preserved at the surface has a very
 369 small thickness. For instance, for a R_f in the range of 50 to 500 km, the chondritic material can only be preserved
 370 in a 100 – 600 m thick crust. Although the crust thickens at the end of the magma ocean episode by addition of light
 371 crystals segregated from the magma ocean, its chondritic part is not thick enough to survive foundering triggered by
 372 impacts (Hevey and Sanders, 2006; Elkins-Tanton et al., 2011). We thus conclude that EC002 cannot be explained by
 373 a scenario of instantaneous accretion and we now consider a scenario of protracted accretion.




Figure 4: Instantaneous accretion. (a): Diagram showing the conditions (radius R and accretion time t_0) of for a planetesimal to differentiate. (b): Evolution of the internal structure evolution of a 100 km radius planetesimal that accreted instantaneously at $t_0 = 0$ Myr.

3.2. Structure evolution during planetesimals accretion

To illustrate the typical thermal history of an accreting planetesimal that undergoes a magma ocean episode, we consider a parent body with a final radius $R_f = 100$ km. To start with, we consider the following parameters of the accretion law, consistent with the “protracted accretion” domain: $R_0/R_f = 0.5$, $t_{acc} = 0.4$ Myr and $\beta = 0.5$. The time evolution of the internal structure, the surface heat flux, the bulk temperature, and the Shields number are shown in Figure 5. During the first 0.7 Myr, heat is dissipated by conduction only. At 0.7 Myr the rheological transition is reached and instantaneous and complete core-mantle differentiation occurs thanks to the low viscosity of the medium. Only a relatively thin crust, relic of the stagnant lid, remains at the surface. As chondritic material keeps being accreted, the thickness of this crust grows, with an increasing insulating effect on the magma ocean. Consequently, the bulk magma ocean temperature becomes larger than the silicate rheological transition temperature which induces the basal melting and iron segregation from the lower crust. Note that this increasing temperature episode also produces the melting to some extent of the first deposited cumulate. As the melting of the chondritic material proceeds, its iron component is incorporated into the core. Protracted accretion keeps bringing new material that globally thickens the crust and contributes to its insulation effect. As ^{26}Al is still active, albeit with a decreasing power, the crust continues to melt and to segregate iron at its base. This feedback effect causes successive melting episodes spanning from 0.7 to 2.3 Myr with several local maxima visible both on the heat flux and bulk temperature. The magma ocean temperature is buffered just above the rheological transition temperature after the first maximum. The value of this absolute maximum can be as high as the silicate liquidus temperature, provided the planetesimal terminal radius is large enough (see Appendix C). After 2.3 Myr, ^{26}Al source has been largely exhausted, and there is no more crust melting. At high temperatures only heavy crystals are formed and sediment, but magma ocean continuous cooling starts to produce light crystals forming a flotation crust beneath the chondritic crust. The rheological transition is reached in the bulk of the silicate mantle at 4 Myr, and the magma ocean freezes. At this point the convection regime transitions back to solid-state convection.

Two key elements should be emphasized in this scenario. First, as the protracted accretion occurs on a longer timescale than the half-life of the radioactive elements, the primordial heat released in the magma ocean is moderate enough to enable the preservation of a km size crust even early in the thermal history. Second, internal heat production induces crustal melting episodes over a few Myr period. The intensity of these phenomena is a function of the amount of primordial heat stored in the magma ocean, and, correspondingly, by the size of the planetesimal. In the following, we discuss how this kind of model can be used to interpret meteoritical data, taking EC002 as an example.

3.3. Lower bound of parent body size: important partial melting episodes in the lower crust

The first constraint we consider deals with the genesis of the parent magma. EC002 is thought to have been formed by 25% partial melting of chondritic material that constitutes the early crust of the parent body. We have found in our model (i) that this type of crust is likely to exist, (ii) that partial melting of the lower part of the crust is usual. To obtain more detailed information on EC002 parent body, we can evaluate the time at which the crustal melting episode occurs as a function of the planetesimal size. Results are displayed in Figure 6 (a). In the case of protracted accretion,




Figure 5: Typical evolution of a 100 km terminal radius planetesimal that begins its protracted accretion at $t = 0$ Myr ($R_0/R_f = 0.5$, $t_{acc} = 0.4$ Myr, $\beta = 0.5$). (a): Evolution of the internal structure of the accreting planetesimal. Magma ocean episode begins at the rheological transition (red vertical dashed line), and yields core/mantle differentiation. The crust is stable and thickens due to accretion supply. Insulation effect of the crust leads to iron extraction at its base between 0.7 to 2.3 Myr. When magma ocean temperature is sub-liquidus, crystals nucleate and deposits form, first a cumulate at CMB, then a floating deposit. Note that the early formed cumulate remelts partially during the crustal thinning episode. We also display the surface heat flux (b), the temperature of the magma ocean and the core (c) and the evolution of the Shields number in the magma ocean (d) during the planetesimal evolution.

408 a magma ocean episode occurs in planetesimals with final radius $R_f > 55$ km. The larger the planetesimal, the earlier
 409 the melting begins and the longer this magma ocean episode lasts. This can be explained by the ratio between the
 410 surface heat losses over the heat generated by the radioactive source that scales with the surface over volume ratio, and
 411 decreases with the planetesimal size. Consequently, the larger the planetesimal, the more heat from radioactive decay
 412 is stored.

413 We assume that the parent magma of EC002 has been formed during this massive partial melting episode of the
 414 crust. As a consequence, the melting period has to encompass the date derived from geochronology. According to
 415 Figure 6 (a), we deduce that the planetesimal final radius has to be larger than 80 km in order to undergo a crustal
 416 melting episode consistent with the time at which EC002 parent magma was produced (≈ 1.8 Myr).

417

418 3.4. Upper bound of the parent body size: EC002 has to be preserved in a “chondritic fridge”

419 A cooling rate of 5 $^{\circ}\text{C yr}^{-1}$ has been inferred for EC002 from petrological data. Such a rate is consistent with
 420 the thermal evolution of a few tens of meters size magmatic reservoirs buried about 1 km below the surface – see
 421 Appendix B. The depth of storage of the magma corresponds to its neutral buoyancy relative to the crustal material.
 422 In the shallowest layers of the planetesimal crust, the density is mainly controlled by the porosity, which itself results
 423 from the brecciation induced by the impacts. It is thus likely that the magma reservoirs form at the base of the regolith.

Structure of differentiated planetesimals




Figure 6: (a) Duration of the crustal melting episode in accreting planetesimal as a function of the planetesimal final radius. (b) Time t_{CF} at which the chondritic material supplied by accretion is preserved as a undifferentiated crust as a function of the planetesimal final radius. (c) Abacus showing the time at which the magma stored in the chondritic crust is preserved as a function of the planetesimal final radius. All simulations are performed with $\beta = 0.5$, $t_{acc} = 0.4$ Myr and $R_0/R_f = 0.5$. In the three sub-panels, the horizontal line gives the time at which the parent magma of EC002 rises in the crust according to petrological data (≈ 1.8 Myr).




Figure 7: Zoom on the upper layers of the planetesimal showing constraints on the time at which a magma produced by 25% of melting in the chondritic crust and that has risen into the crust (dashed purple line) can be preserved from subsequent remelting. Only magma produced after t_{CF} can be preserved from remelting (i.e. their temperature remains below the solidus, solid dashed line). Here, we use the following parameters: $R_f = 100$ km, $R_0/R_f = 0.5$, $t_{acc} = 0.4$ Myr, $\beta = 0.5$.

424 The scenario that models the evolution of the parent magma of EC002 is displayed in Figure 7. The magma parent
 425 forms by 25% partial melting of chondritic material. At 1.8 Myr, the magma is stored at the base of the regolith, at
 426 a depth where it must be preserved from remelting (horizontal dashed line in Figure 7). We assume that this transfer
 427 happened by melt channeling hence is quasi instantaneous (Spiegelman, Kelemen and Aharonov, 2001; Chauveau and
 428 Kaminski, 2008). The depth at which the pluton is preserved from remelting is materialized by the horizontal dashed
 429 line in Figure 7. This depth can be translated into a corresponding time t_{CF} at which the chondritic material supplied
 430 by accretion is preserved as an undifferentiated crust (t_{CF} = “chondritic fridge” time). Figure 6 (b) displays this time
 431 as a function of the planetesimal final radius.

432 We combine the two constraints (magma ascent occurring during a crustal melting episode, and no subsequent
 433 re-melting) to build an abacus relating the size of the parent body to the time at which the magma crystallizes (Figure
 434 6 (c)). This abacus gives the size of the parent body of a meteorite as a function of the date at which the parent magma
 435 crystallizes. For instance, for a crystallization time of 1.8 Myr, as it seems to be the case for EC002, the parent bodies
 436 should have had a final radius between 80 and 130 km.

Structure of differentiated planetesimals




Figure 8: Proposed genesis and thermal history of EC002. (a) Partial melting of the chondritic lower crust generates the parent magma. (b) Magma rises to a shallow storage depth. (c) The parent magma crystallizes quickly and forms a pluton. (d) The pluton is excavated by an impact. Note that the crust thickens continuously during the thermal history, due to accretion of cold material. For sake of simplicity, this figure does not show this effect.

437 The thermal equilibration of the so-formed pluton is quasi-instantaneous (≈ 40 yr) compared to the characteristic
 438 timescale of crustal thermal evolution (several Myr). Then, the magma evolves as a pluton stored in the crust where it
 439 further cools down at a rate of a few $^{\circ}\text{C Myr}^{-1}$. Barrat et al. (2021) proposed that EC002 further experienced a high
 440 cooling rate of $0.1 - 1^{\circ}\text{C/day}$ when it had cooled down below a temperature of 900°C , which would be consistent
 441 with cooling subsequent to an excavation by impact. The genesis and the complete thermal history of EC002 can then
 442 be summarized schematically in Figure 8.

443 3.5. Sensitivity to the accretion law parameters

444 In the previous section we estimated the final radius of the parent body of EC002 between 80 and 130 km, and
 445 we obtained that its chondritic crust was a few km thick. To obtain these results we used a protracted accretion law
 446 based on three parameters (R_0/R_f , t_{acc} and β) that presumably influence the thermal evolution of the planetesimal.
 447 We already highlighted that for R_0/R_f ranging from 0 to 0.8, the accretion history remains largely unchanged. As a
 448 consequence, our results are not affected by R_0/R_f within this range. Parameters β and t_{acc} induce a characteristic
 449 timescale for accretion $\tau_{99\%}$ that (i) has to be consistent with the duration of the dissipation of the protoplanetary disk,
 450 (ii) must be greater than the decay of ^{26}Al . In the following, we explore the sensitivity of β in the range 0.5 and 1.5 for
 451 $t_{acc} = 0.4$ Myr, and we test t_{acc} from 0.07 Myr to 0.87 Myr with $\beta = 0.5$.

452 3.5.1. Influence of β

453 We first consider the influence of β on the onset of the magma ocean episode and of the crustal melting. We fix
 454 the parameters $R_0/R_f = 0.5$ and $t_{acc} = 0.4$ Myr and we present results for three values of β (0.38, 0.5, 1.5 Myr),
 455 resulting from the intersection with the limits of the “protracted accretion” domain in Figure 2(b). Figure 9 shows that
 456 the magma ocean characteristics depends only slightly on β . Planetesimals with final radius $R_f > 55$ km trigger a
 457 magma ocean with an onset time always ranging from 0.5 to 1.5 Myr. The duration of the magma ocean is also robust,
 458 spanning over 2 to 10 Myr. The episode of crustal thinning and melting depends on β . The larger the value of β , the
 459 shorter the duration of crust melting.

460 Figure 10 (a) contains the abacus showing the time at which magma stored in the chondritic crust is preserved
 461 for the three values of β . For relatively small values of β , that correspond to high values of $\tau_{99\%}$, the two zones are

Structure of differentiated planetesimals




Figure 9: Influence of β on the lifetime of the magma ocean (orange) and on the onset and duration of the crustal melting event (red) for planetesimals with terminal radius spanning from 40 to 500 km and $t_{acc} = 0.4$ Myr.




Figure 10: Abacus showing the parent body size required to produce 25% partial melting in the lower crust and later preservation of the produced differentiated rock once it has been stored in the shallow crust. These abacuses are made following the sensitivity study described in the text. (a) Influence of β . (b) Influence of t_{acc} . The horizontal black dashed line gives the age inferred from EC002 (1.8 Myr). The color of the domains are set by the value of $\tau_{99\%}$: purple (10 Myr), red (5 Myr) and grey (1 Myr).

462 almost identical. In this case the accretion is much slower than the decay of radioactive elements and consequently,
 463 after the first crustal melting episode, the chondritic crust is thickened by accretion, reaching a few km thick chondritic
 464 crust (see figure 11 (a), purple and red curves), which will restore the insulation of the magma ocean. Consequently,
 465 heat released by radioactivity is not efficiently evacuated and the magma ocean temperature rises, along with the
 466 lower crust temperature, which, in turn, is able to produce partial melting of the crust. This induces cycles of crustal
 467 thinning and chondritic crust growing, which lengthens the episode of crustal melting. High values of β corresponds to
 468 quasi-instantaneous accretion (grey zone in Figure 10(a)). In this case, when the crust is thinned by basal melting and
 469 iron segregation, accretion is nearly finished, and thus the crust cannot grow though the external supply of chondritic
 470 material, and less than 1 km crust remains at the end of the accretion (see also figure 11 (a), grey curve). In this case,
 471 the first crustal thinning and melting episode increases the heat losses, which rapidly balances the heat released by
 472 radioactive elements.

473 We conclude that β is an important parameter for the history of the accreting planetesimal. Smaller values of β
 474 correspond to parent bodies with a thicker "chondritic fridge" (a few km). The crust melting episode spans the time
 475 range between 1 and 4 Myr. EC002 data are therefore consistent with this family of planetesimals. Higher values of β
 476 correspond to planetesimals that are unable to preserve a substantial undifferentiated crust, as undifferentiated material
 477 is brought early compared to the decay of radioactive elements (in other words they have a small $\tau_{99\%}$).




Figure 11: Thickness of the chondritic crust as a function of planetesimal final radius for different values of β (a) with $t_{acc} = 0.4$ Myr and of t_{acc} (b) with $\beta = 0.5$. $R_0/R_f = 0.5$. The color of the curves are set by the value of $\tau_{99\%}$: purple (10Myr), red (5Myr), blue (2Myr), grey (1Myr).

478 3.5.2. Influence of t_{acc}

479 We also evaluate the influence of t_{acc} on the size of EC002 parent body (Figure 10 (b)). The red zones in Figures
 480 10(a) and (b) are identical and correspond to the values of parameters $\beta = 0.5$ and $t_{acc} = 0.4$ Myr, represented by the
 481 red star in Figure 2 (b). We take the two end-members for $\beta = 0.5$ corresponding to the intersection with the limits of
 482 the “protracted accretion” domain in Figure 2 (b): t_{acc} ranges from 0.07 to 0.87 Myr. Small values of t_{acc} correspond to
 483 an accretion history that is close to an instantaneous one, whereas high values of t_{acc} correspond to a longer accretion
 484 timescale. The parent body of EC002 should have a final radius R_f in the range from 70 to 100 km for $t_{acc} = 0.87$ Myr,
 485 from 80 to 130 km for $t_{acc} = 0.4$ Myr and from 210 to 400 km for $t_{acc} = 0.07$ Myr (Figure 10 (b)). The thickness of
 486 the preserved chondritic crust increases with t_{acc} , and small values of t_{acc} are not consistent with the preservation of a
 487 pluton in an undifferentiated crust (crust thickness < 1km), and hence, cannot explain EC002 formation (Figure 11(b),
 488 grey curve). Figure 11 also displays two curves that correspond to values of β and τ_{acc} from the intersection with
 489 $\tau_{99\%} = 2$ Myr (blue curves). They indicate that a substantial crust, of 2-3 km, can be produced and later preserved only
 490 for larger values of $\tau_{99\%}$. Figures 10 and 11 point out that apparently the most important parameter for the accretion
 491 history is $\tau_{99\%}$, and not the individual β and τ_{acc} producing it.

492 Based on this sensitivity study, the parent body of EC002 radius can be estimated thanks to the Figure 10. If we
 493 exclude cases lying too close to the “quasi instantaneous accretion”, the parent body of EC002 has a final radius R_f
 494 between 70 and 130 km. We highlighted that the characteristics of the preserved chondritic crust and the thermal
 495 history are linked to the comparison between the characteristic accretion timescale $\tau_{99\%}$ and the half-life of ^{26}Al ($t_{1/2}^{Al} = 0.717$ Myr),
 496 which should be a factor of about 3 times larger ($\tau_{99\%} > 2$ Myr).

497 The onset time of crustal melting and the “chondritic fridge” time t_{CF} are the two conditions that set bounds to
 498 the accretion history of a planetesimal consistent with EC002. They both should be lower than the crystallization time
 499 measured for EC002 (1.8 Myr); the first condition ensures the existence of the parent magma and the second one the
 500 preservation of the pluton in the crust. These conditions limit the parameters of the accretion law to the shaded zone
 501 in Figure 12.

502 4. Other geophysical implications

503 4.1. Sustainable “chondritic fridge” at the surface of planetesimals

504 Our model, applied in the previous section to the thermal history of EC002, actually describes general features
 505 in the evolution of planetesimals. In particular, considering a protracted accretion has an important impact on the
 506 planetesimal structure. Early instantaneous accretion induces very intense internal heating, due to the large amount
 507 of ^{26}Al that is stored. As a consequence, the peak bulk temperature is so high that any crust present at the surface is
 508 bound to melt and to become very thin (below 1 km). Protracted accretion will balance internal heating by the addition
 509 of cold material and will thus help to preserve a relatively thick crust that may later undergo melting episodes (Figure

Structure of differentiated planetesimals




Figure 12: Parameters of the accretion law consistent with EC002.

510 11). These phenomena favor the formation of evolved rocks in planetesimals that preserve information from the early
 511 times of their history. For instance, some meteorites bear traces of early dynamos on their parent bodies, as the Kaba
 512 meteorite that recorded a dynamo as early as 10 Myr after CAI (Gattacceca, Weiss and Gounelle, 2016). Dodds et al.
 513 (2021) calculated that planetesimal cores with radius larger than 400 km (i.e., planetesimals with radius higher than
 514 ≈ 918 km) are able to sustain a dynamo at these early times. But the crust in the case of instantaneous accretion remains
 515 only ≈ 1 km thick so that it is unlikely it could register a remanent magnetization. To avoid crustal melting, the authors
 516 made the accretion begin at 0.8 Myr after CAIs, which corresponds roughly to a decay of ^{26}Al to half of its initial
 517 value. In the case of a protracted accretion that begins at $t = 0$ and a planetesimal with $R_f = 920$ km, our model
 518 predicts at 10 Myr a 49 km thick crust including (i) a 15 km thick chondritic upper crust, (ii) a 1 km thick layer and
 519 (iii) a 33 km thick differentiated layer of floating crystals. The undifferentiated part of the crust lies below the Curie
 520 temperature ($\approx 800^\circ\text{C}$) which would provide suitable conditions for the record of the early magnetic activity. This cold
 521 zone is a natural consequence of protracted accretion, in contrast with the runaway accretion scenario used by Dodds
 522 et al. (2021) in which accretion ends early. A more detailed modeling of the possibility to sustain a dynamo in this type
 523 of body is required (Monteux, Jellinek and Johnson, 2011; Bryson et al., 2019; Dodds et al., 2021) in order to provide
 524 a precise information on the time at which the dynamo is triggered.

525 4.2. Geochronology of iron meteorites

526 The Hf – W systematics established for iron meteorites bring some constraints to refine their accretion scenario.
 527 Hf – W age is usually interpreted as the date of instantaneous core formation (Kruijjer et al., 2014). However, if the
 528 contribution of iron from crustal melting occurs over a long period of time, ages inferred from Hf – W systematics
 529 integrate the entire process of differentiation (Neumann, Kruijjer, Breuer and Kleine, 2018). We calculated the
 530 proportion of the core that is supplied by core melting (Figure 13 (a)) and the apparent age of the core inferred from
 531 Hf – W systematics using our model, and we display it as a function of the time of the first differentiation in Figure 13
 532 (b). In order to illustrate this effect we show in these figures only the results obtained for accretion histories characterized
 533 by $\tau_{99\%} = 5$ Myr (red curves) or $\tau_{99\%} = 10$ Myr (purple curves). The proportion of core added during accretion and crust
 534 melting is less important for lower values of the accretion timescale $\tau_{99\%}$ (data not shown). Within the displayed values
 535 (Figure 13 (a)) the volume of core added depends on the values of the group of parameters (β and τ_{acc}), the larger they
 536 are, the bigger the effect. The first differentiation occurs at the rheological transition that happens at 0.5 – 1.5 Myr,
 537 while the apparent age of the core is always younger than the age of the first differentiation, and can be very different,
 538 depending on the values of the accretion parameters we are ready to admit. There is also a dependence of the volume
 539 of core added with the final planetesimal radius R_f which reaches a plateau at large values. The apparent age of the
 540 core does not reach a similar plateau and actually continues to strongly increase with R_f .

541 These considerations can be used to interpret the age dichotomy in the iron meteorites. Earlier Hf – W inferred
 542 ages range from 0.5 to 1.5 Myr for non-carbonaceous (NC) magmatic iron meteorites, and from 2 to 3 Myr for
 543 carbonaceous (CC) magmatic iron meteorites (Kruijjer et al., 2014, 2017). Recent revised Hf – W model ages corrected




Figure 13: (a) Proportion of the core that is brought by iron supply due to progressive melting of the chondritic crust as a function of the planetesimals final radius. (b) Apparent age of the core inferred from Hf – W systematics as a function of the age of the first differentiation that corresponds to the onset of the magma ocean. Black line represent a 1:1 dependence between the two ages. Red curves represent values of $\tau_{99\%} = 5$ Myr and purple curves $\tau_{99\%} = 10$ Myr, full lines $\beta = 0.5$, dashed lines $\beta = 1$ and dashed and dotted lines $\beta = 1.5$.

544 for nucleosynthetic Pt isotope variations are slightly younger (1 – 2 Myr for NC meteorites and 3 – 3.5 Myr for CC
 545 meteorites (Spitzer, Burkhardt, Nimmo and Kleine, 2021)). In the case of NC iron meteorites, these younger ages were
 546 confirmed by Mn – Cr chronometry (0.4 – 2.2 Myr) (Anand, Pape, Wille, Mezger and Hofmann, 2021).

547 Our model can explain the entire age range estimated for NC iron meteorites, for accretion histories characterized
 548 by values of (β and τ_{acc}) rather small, and small to intermediate values of the final planetesimal radius R_f . Ages of
 549 CC iron meteorites can be obtained for large values of R_f and large values of the pair of parameters (β and τ_{acc}) that
 550 result in large values of $\tau_{99\%}$ falling between the red (5 Myr) and purple (10 Myr) lines in Figure 2.

551 Although we can explain the age range encompassed by both NC and CC iron meteorites, we nevertheless
 552 acknowledge that thermal modeling of CC bodies requires a specific study. The substantial presence of water and
 553 other volatiles modifies almost everything in the model, from the heat budget by the intermediate of the latent heat to
 554 other properties susceptible to affect convection such as the viscosity. The present model with the values of parameters
 555 in Table 1, properly describes only the formation of dry, NC bodies.

556 4.3. Degassing despite a “chondritic fridge”?

557 The presence of a “chondritic fridge” on top of planetesimals rises the question of volatile behavior in such systems.
 558 The surface temperature is taken as a constant, which is a reasonably good approximation (less than 10 K variation)
 559 for small radius planetesimals heated by ^{26}Al (Dodds et al., 2021). More rigorously, the surface temperature results
 560 from the balance between the emitted radiative flux, the absorbed solar radiative flux and the heat flux at the surface of
 561 the body (Henke, Gail, Tieloff and Schwarz, 2013). The (quasi)isothermal boundary condition and low temperature
 562 at the surface will determine the preservation of a stagnant lid. A planetary body with a sub-surface magma ocean can
 563 be subject to partial or total crust melting and therefore go through volatile loss and EC002 meteorite can be taken
 564 as a proof of such magmatic activity. The amplitude and timescale of this magmatic activity should depend mainly
 565 on the body radius and the distance to the sun. We cannot rule out a transitory disruption of the crust during the first
 566 few Myr. A transitory disruption of the stagnant lid and the exposure of molten rock at the surface would lead to a
 567 transitory increase of the radiative heat flux which will cool down the surface and restore the isothermal boundary
 568 condition with the aid of the newly accreted, cold material. This transitory episode should be very short with respect
 569 to the duration of the magma ocean, provided planetesimals cannot preserve an atmosphere. This should be the case
 570 for the small radius bodies like the ones considered in the present study. Degassing is hence possible even though the
 571 planetesimal is covered by a “chondritic fridge” as established in a large number of studies (Norris and Wood, 2017;

572 Young, Shahar, Nimmo, Schlichting, Schauble, Tang and Labidi, 2019; Sossi, Klemme, O'Neill, Berndt and Moynier,
573 2019; Collinet and Grove, 2020; Hirschmann, Bergin, Blake, Ciesla and Li, 2021).

574 **5. Conclusion**

575 We highlighted in this paper how geochemical and petrological data obtained from differentiated achondrites can
576 be used to constrain the formation and evolution of their parent bodies. Our results are based on a thermal model that
577 takes into account protracted accretion, core-mantle differentiation, magma ocean episode and crustal thickening by
578 continuous accretion of chondritic material at the surface. This model predicts that the reference internal structure of a
579 planetesimal formed early with a size $\gtrsim 55$ km is a differentiated interior (with a liquid metallic core and a crystallizing
580 magma ocean) covered by an undifferentiated chondritic crust that is stable and will undergo localized basal partial
581 melting. We show how cooling rates provide information about the depth and the size of the reservoir where the parent
582 magma of achondrites crystallized in the chondritic crust. The datation of parent magma formation can then be used
583 to provide some bounds on the parent body size and on the parameters of its accretion laws. This approach is general
584 and could be used to study other parent bodies than that of EC002.

585 **Acknowledgments**

586 This paper is part of Cyril Sturtz's PhD thesis (Université Paris Cité, Institut de Physique du Globe de Paris).
587 The authors would like to thank the editor Doris Breuer for her editorial handling and two anonymous reviewers for
588 their fruitful comments that improve this manuscript. This study contributes to the IdEx Université de Paris ANR-18-
589 IDEX-0001. This work was supported by the Programme National de Planétologie (PNP) of CNRS/INSU, co-funded
590 by CNES.


591 **A. Rheological law and composition as a function of the temperature**


Figure 14: Rheological law (blue) of the planetesimal and liquid fraction (red) in the body as a function of the temperature. We assume an equilibrium melting, so that the liquid content is directly given by (4). The rheological transition occurs for a liquid content of 40% (i.e.: a solid content of 60%).


 592 **B. Cooling model for a magmatic reservoir buried at a certain depth**


Figure 15: Evolution of the size of a magmatic reservoir that undergoes a cooling rate of $5^\circ\text{C}/\text{yr}$ as a function of the depth at which it is stored. The area colored in blue stands for temperature anomalies spanning from 1000 to 1200°C . The forbidden zone represents non-physical solutions that implies reservoirs that are larger than the crust thickness.

The cooling rates measured in some crystals provide constraints on the condition of magma storage in the crust. For a cooling rate of $5^\circ\text{C}\text{yr}^{-1}$, like the one measure in EC002, a simple model for the cooling of a magma reservoir can predict the size and the depth at which magma is stored. We consider a magma body of typical length L at a depth z_0 bearing a thermal anomaly ΔT compared to the surface temperature T_s . During a quasi-static evolution of a period of time dt , the variation of internal energy is $\rho c_p d(\Delta T) LS$, where S is the surface of the system through which heat losses occur. The heat flux can be written $Q_s S dt$, where Q_s is assumed to be conductive (Kaminski and Jaupart, 2000)

and the conservation of energy yields

$$\frac{d\Delta T}{dt} = \frac{\lambda}{\rho c_p} \frac{\Delta T}{z_0 L}, \quad (40)$$

593 where λ is the thermal conductivity of the enclosing rock, ρ is its density and c_p its specific heat. Results of magma
 594 reservoir size as a function of its depth to get the required cooling rate are displayed in Figure 15. To attain 5°C yr^{-1} ,
 595 the magma has to be stored in small reservoir, which means that the reservoir that cools is a few meter size diapir or a
 596 magma lens stored at a few km depth.

597 Furthermore, this approach shows that the shallower the reservoir the larger its size for a given cooling rate. This
 598 calculation predicts also that the crust of the parent body has to be thicker than 100 m in order to sustain a reservoir
 599 whose size is smaller than the crust thickness and at least 300 m in order to have reservoirs that represents 10% of the
 600 crust thickness (Figure 15).

601 C. Maximum temperature reached in the magma ocean




Figure 16: Maximum temperature reached during the crust melting and iron extraction episode as a function of the terminal planetesimal radius R_f .

602 References

- 603 Agee, C., Habermann, M., Ziegler, K., 2018. Northwest africa 11575: Unique ungrouped trachyandesite achondrite, in: Lunar and Planetary Science
 604 Conference, p. 2226.
- 605 Allège, C., Manhès, G., Göpel, C., 1995. The age of the earth. *Geochimica et Cosmochimica Acta* 59, 1445–1456. URL: <https://www.sciencedirect.com/science/article/pii/0016703795000544>, doi:[https://doi.org/10.1016/0016-7037\(95\)00054-4](https://doi.org/10.1016/0016-7037(95)00054-4).
- 607 Anand, A., Pape, J., Wille, M., Mezger, K., Hofmann, B., 2021. Early differentiation of magmatic iron meteorite parent bodies from mn–cr
 608 chronometry. *Geochemical Perspectives Letters* 20.
- 609 Barrat, J.A., Chaussidon, M., Yamaguchi, A., Beck, P., Villeneuve, J., Byrne, D.J., Broadley, M.W., Marty, B., 2021. A 4,565-my-old andesite from
 610 an extinct chondritic protoplanet. *Proceedings of the National Academy of Sciences* 118. URL: [https://www.pnas.org/content/118/11/](https://www.pnas.org/content/118/11/e2026129118)
 611 [e2026129118](https://doi.org/10.1073/pnas.2026129118), doi:[10.1073/pnas.2026129118](https://doi.org/10.1073/pnas.2026129118), arXiv:<https://www.pnas.org/content/118/11/e2026129118.full.pdf>.
- 612 Bischoff, A., Horstmann, M., Barrat, J.A., Chaussidon, M., Pack, A., Herwartz, D., Ward, D., Vollmer, C., Decker, S., 2014. Trachyandesitic
 613 volcanism in the early solar system. *Proceedings of the National Academy of Sciences* 111, 12689–12692. URL: [https://www.pnas.org/](https://www.pnas.org/content/111/35/12689)
 614 [content/111/35/12689](https://doi.org/10.1073/pnas.1404799111), doi:[10.1073/pnas.1404799111](https://doi.org/10.1073/pnas.1404799111), arXiv:<https://www.pnas.org/content/111/35/12689.full.pdf>.
- 615 Bouvier, A., Wadhwa, M., 2010. The age of the solar system redefined by the oldest pb–pb age of a meteoritic inclusion. *Nature Geoscience* 3,
 616 637–641. URL: <https://doi.org/10.1038/ngeo941>, doi:[10.1038/ngeo941](https://doi.org/10.1038/ngeo941).
- 617 Bryson, J.F., Neufeld, J.A., Nimmo, F., 2019. Constraints on asteroid magnetic field evolution and the radii of meteorite parent bodies from
 618 thermal modeling. *Earth and Planetary Science Letters* 521, 68–78. URL: [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S00128221X19303280)
 619 [S00128221X19303280](https://doi.org/10.1016/j.epsl.2019.05.046), doi:<https://doi.org/10.1016/j.epsl.2019.05.046>.
- 620 Carslaw, H.S., Jaeger, J.C., 1959. *Conduction of heat in solids*. Oxford University Press, New York.

- 621 Charru, F., Mouilleron, H., Eiff, O., 2004. Erosion and deposition of particles on a bed sheared by a viscous flow. *Journal of Fluid Mechanics* 519,
622 55–80. doi:10.1017/S0022112004001028.
- 623 Chauveau, B., Kaminski, E., 2008. Porous compaction in transient creep regime and implications for melt,
624 petroleum, and co2 circulation. *Journal of Geophysical Research: Solid Earth* 113. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JB005088>,
625 doi:<https://doi.org/10.1029/2007JB005088>,
626 arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2007JB005088>.
- 627 Choblet, G., Sotin, C., 2000. 3d thermal convection with variable viscosity: can transient cooling be described by a quasi-static scaling law? *Physics*
628 *of the Earth and Planetary Interiors* 119, 321–336.
- 629 Collinet, M., Grove, T.L., 2020. Widespread production of silica- and alkali-rich melts at the onset of planetesimal melting. *Geochimica et*
630 *Cosmochimica Acta* 277, 334–357. URL: <https://www.sciencedirect.com/science/article/pii/S0016703720301666>, doi:<https://doi.org/10.1016/j.gca.2020.03.005>.
- 631
- 632 Condie, K.C., 2016. Chapter 10 - comparative planetary evolution, in: Condie, K.C. (Ed.), *Earth as an Evolving Planetary System*
633 (Third Edition). third edition ed.. Academic Press, pp. 317–367. URL: <https://www.sciencedirect.com/science/article/pii/B9780128036891000109>, doi:<https://doi.org/10.1016/B978-0-12-803689-1.00010-9>.
- 634
- 635 Connolly, J.N., Bizzarro, M., Krot, A.N., Nordlund, Å., Wielandt, D., Ivanova, M.A., 2012. The absolute chronology and thermal processing
636 of solids in the solar protoplanetary disk. *Science* 338, 651–655. URL: <https://science.sciencemag.org/content/338/6107/651>,
637 doi:10.1126/science.1226919, arXiv:<https://science.sciencemag.org/content/338/6107/651.full.pdf>.
- 638 Cuzzi, J., Hogan, R., Shariff, K., 2008. Toward planetesimals: Dense chondrule clumps in the protoplanetary nebula. *The Astrophysical Journal*
639 687, 1432–1447. URL: <https://doi.org/10.1086/591239>, doi:10.1086/591239.
- 640 Davaille, A., Jaupart, C., 1993. Thermal convection in lava lakes. *Geophysical Research Letters* 20, 1827–
641 1830. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/93GL02008>, doi:10.1029/93GL02008,
642 arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/93GL02008>.
- 643 Day, J., Ash, R., Liu, Y., Bellucci, J., III, D., McDonough, W., Walker, R., Taylor, L., 2009. Early formation of evolved asteroidal crust. *Nature* 457,
644 179–182. URL: <https://doi.org/10.1038/nature07651>, doi:10.1038/nature07651.
- 645 Dodds, K.H., Bryson, J.F.J., Neufeld, J.A., Harrison, R.J., 2021. The thermal evolution of planetesimals during accretion and dif-
646 ferentiation: Consequences for dynamo generation by thermally-driven convection. *Journal of Geophysical Research: Planets* 126,
647 e2020JE006704. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JE006704>, doi:<https://doi.org/10.1029/2020JE006704>, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2020JE006704>. e2020JE006704
648 2020JE006704.
- 649
- 650 Elkins-Tanton, L., 2012. Magma oceans in the inner solar system. *Annual Review of Earth and Planetary Sciences* 40,
651 113–139. URL: <https://doi.org/10.1146/annurev-earth-042711-105503>, doi:10.1146/annurev-earth-042711-105503,
652 arXiv:<https://doi.org/10.1146/annurev-earth-042711-105503>.
- 653 Elkins-Tanton, L., Weiss, B., Zuber, M., 2011. Chondrites as samples of differentiated planetesimals. *Earth and Planetary Science Letters* 305,
654 1–10.
- 655 Fang, L., Frossard, P., Boyet, M., Bouvier, A., Barrat, J.A., Chaussidon, M., Moynier, F., 2022. Half-life and initial solar system abundance of
656 ¹⁴⁶Sm determined from the oldest andesitic meteorite. *Proceedings of the National Academy of Sciences* 119, e2120933119.
- 657 Faure, F., 2020. Early silica crust formation in planetesimals by metastable silica-rich liquid immiscibility or cristobalite crystallisation:
658 the possible origin of silica-rich chondrules. *Scientific Reports* 10, 4765. URL: <https://doi.org/10.1038/s41598-020-61806-5>,
659 doi:10.1038/s41598-020-61806-5.
- 660 Fu, R.R., Elkins-Tanton, L.T., 2014. The fate of magmas in planetesimals and the retention of primitive chondritic crusts. *Earth and Planetary*
661 *Science Letters* 390, 128–137. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X1400003X>, doi:<https://doi.org/10.1016/j.epsl.2013.12.047>.
- 662
- 663 Ganino, C., Libourel, G., 2017. Reduced and unstratified crust in cv chondrite parent body. *Nature Communications* 8, 261. URL: <https://doi.org/10.1038/s41467-017-00293-1>, doi:10.1038/s41467-017-00293-1.
- 664
- 665 Gattacceca, J., Weiss, B.P., Gounelle, M., 2016. New constraints on the magnetic history of the cv parent body and the solar nebula from the
666 kaba meteorite. *Earth and Planetary Science Letters* 455, 166–175. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X16304885>, doi:<https://doi.org/10.1016/j.epsl.2016.09.008>.
- 667
- 668 Greenwood, R., Franchi, I., Jambon, A., Buchanan, P., 2005. Widespread magma oceans on asteroidal bodies in the early solar system. *Nature* 435,
669 916–918. doi:10.1038/nature03612.
- 670 Greenwood, R.C., Burbine, T.H., Franchi, I.A., 2020. Linking asteroids and meteorites to the primordial planetesimal population. *Geochimica et*
671 *Cosmochimica Acta* 277, 377–406. URL: <https://www.sciencedirect.com/science/article/pii/S0016703720301058>, doi:<https://doi.org/10.1016/j.gca.2020.02.004>.
- 672
- 673 Guazzelli, E., Pouliquen, O., 2018. Rheology of dense granular suspensions. *Journal of Fluid Mechanics* 852. doi:10.1017/jfm.2018.548.
- 674 Haugbølle, T., Weber, P., Wielandt, D.P., Benítez-Llambay, P., Bizzarro, M., Gressel, O., Pessah, M.E., 2019. Probing the protosolar disk using dust
675 filtering at gaps in the early solar system. *The Astronomical Journal* 158, 55.
- 676 Henke, S., Gail, H.P., Trieloff, M., Schwarz, W., 2013. Thermal evolution model for the h chondrite asteroid-instantaneous formation versus
677 protracted accretion. *Icarus* 226, 212–228. URL: <https://www.sciencedirect.com/science/article/pii/S0019103513002340>,
678 doi:<https://doi.org/10.1016/j.icarus.2013.05.034>.
- 679 Hevey, P., Sanders, I., 2006. A model for planetesimal meltdown by ²⁶Al and its implications for meteorite parent bodies. *Meteoritics*
680 *& Planetary Science* 41, 95–106. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1945-5100.2006.tb00195.x>,
681 doi:10.1111/j.1945-5100.2006.tb00195.x.
- 682 Hirschmann, M.M., Bergin, E.A., Blake, G.A., Ciesla, F.J., Li, J., 2021. Early volatile depletion on planetesimals inferred from c–s systematics of
683 iron meteorite parent bodies. *Proceedings of the National Academy of Sciences* 118.

- 684 Höink, T., Schmalz, J., Hansen, U., 2006. Dynamics of metal-silicate separation in a terrestrial magma ocean. *Geochemistry, Geo-*
685 *physics, Geosystems* 7. URL: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006GC001268>, doi:10.1029/
686 2006GC001268, arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006GC001268>.
- 687 Jaupart, C., Mareschal, J.C., 2010. *Heat Generation and Transport in the Earth*. Cambridge University Press. doi:10.1017/CB09780511781773.
- 688 Johansen, A., Lambrechts, M., 2017. Forming planets vis pebble accretion. *Annu. Rev. Earth Planet. Sci.* 12451, 359–387.
- 689 Johansen, A., Oishi, J., Low, M.M.M., Klahr, H., Henning, T., Youdin, A., 2007. Rapid planetesimal formation in turbulent circumstellar disks.
690 *Nature* 448, 1022–1025. URL: <https://doi.org/10.1038/nature06086>, doi:10.1038/nature06086.
- 691 Kaminski, E., Jaupart, C., 2000. Lithosphere structure beneath the Phanerozoic intracratonic basins of North America. *Earth and Planetary*
692 *Science Letters* 178, 139–149. URL: <https://www.sciencedirect.com/science/article/pii/S0012821X00000674>, doi:[https://doi.org/10.1016/S0012-821X\(00\)00067-4](https://doi.org/10.1016/S0012-821X(00)00067-4).
- 693 Kaminski, E., Limare, A., Kenda, B., Chaussidon, M., 2020. Early accretion of planetesimals unraveled by the thermal evolution of parent bodies
694 of magmatic iron meteorites. *Earth and Planetary Science Letters* 548, 116469. doi:<https://doi.org/10.1016/j.epsl.2020.116469>.
- 695 Karl E. Haisch, J., Lada, E.A., Lada, C.J., 2001. Disk frequencies and lifetimes in young clusters. *The Astrophysical Journal* 553, L153–L156.
696 URL: <https://doi.org/10.1086/320685>, doi:10.1086/320685.
- 697 Kleine, T., Budde, G., Burkhardt, C., Kruijer, T.S., Worsham, E.A., Morbidelli, A., Nimmo, F., 2020. The non-carbonaceous–carbonaceous
698 meteorite dichotomy. *Space Science Reviews* 216, 55. URL: <https://doi.org/10.1007/s11214-020-00675-w>, doi:10.1007/
699 s11214-020-00675-w.
- 700 Krieger, I.M., Dougherty, T.J., 1959. A mechanism for non-newtonian flow in suspensions of rigid spheres. *Transactions of the Society of Rheology*
701 3, 137–152. URL: <https://doi.org/10.1122/1.548848>, doi:10.1122/1.548848.
- 702 Kruijer, T., Touboul, M., Fischer-Gödde, M., Bermingham, K., Walker, R., Kleine, T., 2014. Protracted core formation and rapid accretion of
703 protoplanets. *Science* 344, 1150–1154.
- 704 Kruijer, T.S., Burkhardt, C., Budde, G., Kleine, T., 2017. Age of Jupiter inferred from the distinct genetics and formation times of meteorites.
705 *Proceedings of the National Academy of Sciences* 114, 6712–6716. URL: <https://www.pnas.org/content/114/26/6712>, doi:10.1073/
706 pnas.1704461114, arXiv:<https://www.pnas.org/content/114/26/6712.full.pdf>.
- 707 Lavorel, G., Le Bars, M., 2009. Sedimentation of particles in a vigorously convecting fluid. *Physical Review E, Statistical, nonlinear, and soft matter*
708 *physics* 80, 046324. doi:10.1103/PhysRevE.80.046324.
- 709 Li, R., Youdin, A.N., Simon, J.B., 2019. Demographics of planetesimals formed by the streaming instability. *The Astrophysical Journal* 885, 69.
- 710 Limare, A., Jaupart, C., Kaminski, E., Fourel, L., Farnetani, C., 2019. Convection in an internally heated stratified heterogeneous reservoir. *Journal*
711 *of Fluid Mechanics* 870, 67–105. doi:10.1017/jfm.2019.243.
- 712 Limare, A., Kenda, B., Kaminski, E., Surducun, E., Surducun, V., Neamtu, C., 2021. Transient convection experiments in internally-heated
713 systems. *MethodsX* 8, 101224. URL: <https://www.sciencedirect.com/science/article/pii/S2215016121000170>, doi:<https://doi.org/10.1016/j.mex.2021.101224>.
- 714 Mandler, B., Elkins-Tanton, L., 2013. The origin of eucrites, diogenites, and olivine diogenites: Magma ocean crystallization and shallow magma
715 chamber processes on Vesta. *Meteoritics & Planetary Science* 48, 2333–2349. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/maps.12135>,
716 doi:10.1111/maps.12135, arXiv:<https://onlinelibrary.wiley.com/doi/pdf/10.1111/maps.12135>.
- 717 Martin, D., Nokes, R., 1988. Crystal settling in a vigorously convecting magma chamber. *Nature* 332, 534–536. URL: <https://doi.org/10.1038/332534a0>,
718 doi:10.1038/332534a0.
- 719 Martin, D., Nokes, R., 1989. A fluid-dynamical study of crystal settling in convecting magmas. *Journal of Petrology*
720 30, 1471–1500. URL: <https://doi.org/10.1093/petrology/30.6.1471>, doi:10.1093/petrology/30.6.1471,
721 arXiv:<https://academic.oup.com/petrology/article-pdf/30/6/1471/4314225/30-6-1471.pdf>.
- 722 Montoux, J., Jellinek, A., Johnson, C., 2011. Why might planets and moons have early dynamos? *Earth and Planetary Science Letters* 310, 349–359.
- 723 Morbidelli, A., Bottke, W., Nesvorný, D., Levison, H., 2009. Asteroids were born big. *Icarus* 204, 558–573. URL: <https://www.sciencedirect.com/science/article/pii/S0019103509003029>, doi:<https://doi.org/10.1016/j.icarus.2009.07.011>.
- 724 Morbidelli, A., Lambrechts, M., Jacobson, S., Bitsch, B., 2015. The great dichotomy of the solar system: Small terrestrial embryos and massive
725 giant planet cores. *Icarus* 258, 418–429.
- 726 Neumann, W., Breuer, D., Spohn, T., 2012. Differentiation and core formation in accreting planetesimals. *A&A* 543, A141. URL: <https://doi.org/10.1051/0004-6361/201219157>,
727 doi:10.1051/0004-6361/201219157.
- 728 Neumann, W., Breuer, D., Spohn, T., 2014. Differentiation of Vesta: Implications for a shallow magma ocean. *Earth and Planetary Science Letters*
729 395, 267–280.
- 730 Neumann, W., Kruijer, T., Breuer, D., Kleine, T., 2018. Multistage core formation in planetesimals revealed by numerical modeling and hf-w
731 chronometry of iron meteorites. *Journal of Geophysical Research: Planets* 123, 421–444.
- 732 Norris, C.A., Wood, B.J., 2017. Earth’s volatile contents established by melting and vaporization. *Nature* 549, 507–510.
- 733 Righter, K., Drake, M., 1997. A magma ocean on Vesta: Core formation and petrogenesis of eucrites and diogenites. *Meteoritics & Planetary*
734 *Science* 32, 929–944. doi:10.1111/j.1945-5100.1997.tb01582.x.
- 735 Roberts, P., 1967. Convection in horizontal layers with internal heat generation. theory. *Journal of Fluid Mechanics* 30, 33–49. doi:10.1017/
736 S0022112067001284.
- 737 Rubie, D., Melosh, H., Reid, J., Liebske, C., Righter, K., 2003. Mechanisms of metal-silicate equilibration in the terrestrial magma ocean. *Earth*
738 *and Planetary Science Letters* 205, 239–255.
- 739 Sahijpal, S., Soni, P., Gupta, G., 2007. Numerical simulations of the differentiation of accreting planetesimals with ²⁶Al and ⁶⁰Fe as the heat sources.
740 *Meteoritics & Planetary Science* 42, 1529–1548.
- 741 Schubert, G., Turcotte, D.L., Olson, P., 2001. *Mantle convection in the Earth and planets*. Cambridge University Press, Cambridge.
- 742 Scott, T., Kohlstedt, D.L., 2006. The effect of large melt fraction on the deformation behavior of peridotite. *Earth Planet. Sci. Lett.* 216, 177–187.

- 746 Shearer, C., Burger, P., Neal, C., Sharp, Z., Spivak-Birndorf, L., Borg, L., Fernandes, V., Papike, J., Karner, J., Wadhwa, M., Gaffney, A., Shafer,
747 J., Geissman, J., Atudorei, N.V., Herd, C., Weiss, B., King, P., Crowther, S., Gilmour, J., 2010. Non-basaltic asteroidal magmatism during the
748 earliest stages of solar system evolution: A view from antarctic achondrites graves nunatak 06128 and 06129. *Geochimica et Cosmochimica*
749 *Acta* 74, 1172–1199. URL: <https://www.sciencedirect.com/science/article/pii/S0016703709006681>, doi:[https://doi.org/](https://doi.org/10.1016/j.gca.2009.10.029)
750 [10.1016/j.gca.2009.10.029](https://doi.org/10.1016/j.gca.2009.10.029).
- 751 Shields, A., 1936. Anwendung der aehnlichkeitsmechanik und der turbulenzforschung auf die geschiebewegung, mitteilung preussischen
752 versuchsanstalt wasserbau schiffbau [application of similarity mechanics and turbulence research on shear flow]. Preussischen Versuchsanstalt
753 für Wasserbau und Schiffbau, Berlin 26.
- 754 Solomatov, V., 2000. Fluid dynamics of a terrestrial magma ocean. *Origin of the Earth and Moon* 1.
- 755 Solomatov, V.S., 1995. Scaling of temperature- and stress-dependent viscosity convection. *Physics of Fluids* 7, 266–274. URL: <https://doi.org/10.1063/1.868624>, doi:[10.1063/1.868624](https://doi.org/10.1063/1.868624), arXiv:<https://doi.org/10.1063/1.868624>.
- 756 Sossi, P.A., Klemme, S., O'Neill, H.S.C., Berndt, J., Moynier, F., 2019. Evaporation of moderately volatile elements from silicate melts: experiments
757 and theory. *Geochimica et Cosmochimica Acta* 260, 204–231.
- 758 Spiegelman, M., Kelemen, P., Aharonov, E., 2001. Causes and consequences of flow organization during melt transport: The reaction
759 infiltration instability in compactible media. *Journal of Geophysical Research: Solid Earth* 106, 2061–2077. URL: [https://](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2000JB900240)
760 agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2000JB900240, doi:<https://doi.org/10.1029/2000JB900240>,
761 arXiv:<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2000JB900240>.
- 762 Spitzer, F., Burkhardt, C., Nimmo, F., Kleine, T., 2021. Nucleosynthetic pt isotope anomalies and the hf-w chronology of core formation in inner and
763 outer solar system planetesimals. *Earth and Planetary Science Letters* 576, 117211. URL: [https://www.sciencedirect.com/science/](https://www.sciencedirect.com/science/article/pii/S0012821X21004660)
764 [article/pii/S0012821X21004660](https://www.sciencedirect.com/science/article/pii/S0012821X21004660), doi:<https://doi.org/10.1016/j.epsl.2021.117211>.
- 765 Srinivasan, P., Dunlap, D., Agee, C., Wadhwa, M., Coleff, D., Ziegler, K., Zeigler, R., McCubbin, F., 2018. Silica-rich volcanism in the
766 early solar system dated at 4.565 ga. *Nature Communications* 9, 3036. URL: <https://doi.org/10.1038/s41467-018-05501-0>,
767 doi:[10.1038/s41467-018-05501-0](https://doi.org/10.1038/s41467-018-05501-0).
- 768 Sturtz, C., Kaminski, E., Limare, A., Tait, S., 2021a. The fate of particles in a volumetrically heated convective fluid at high prandtl number. *Journal*
769 *of Fluid Mechanics* 929, A28. doi:[10.1017/jfm.2021.862](https://doi.org/10.1017/jfm.2021.862).
- 770 Sturtz, C., Limare, A., Tait, S., Kaminski, É., 2021b. Birth and decline of magma oceans. part 1: erosion and deposition of crystal layers in evolving
771 magmatic reservoirs. arXiv:2108.00910.
- 772 Sturtz, C., Limare, A., Tait, S., Kaminski, É., 2021c. Birth and decline of magma oceans. part 2: wobbling thermal history of early accreted
773 planetesimals. arXiv:2108.00995.
- 774 Tang, H., Dauphas, N., 2012. Abundance, distribution, and origin of 60fe in the solar protoplanetary disk. *Earth and Planetary Science Letters*
775 359-360, 248–263.
- 776 Taylor, S.R., 1989. Growth of planetary crusts. *Tectonophysics* 161, 147–156. URL: [https://www.sciencedirect.com/science/article/](https://www.sciencedirect.com/science/article/pii/0040195189901510)
777 [pii/0040195189901510](https://www.sciencedirect.com/science/article/pii/0040195189901510), doi:[https://doi.org/10.1016/0040-1951\(89\)90151-0](https://doi.org/10.1016/0040-1951(89)90151-0). growth of the continental crust.
- 778 Tkalcec, B., Golabek, G., Brenker, F., 2013. Solid-state plastic deformation in the dynamic interior of a differentiated asteroid. *Nature Geoscience*
779 6, 93–97.
- 780 Visser, R.G., Ormel, C.W., 2016. On the growth of pebble-accreting planetesimals. *Astron. Astrophys.* 586, A66, 1–11.
- 781 Warren, P., 1985. The magma ocean concept and lunar evolution. *Ann. Rev. Earth Planet. Sci.* 13, 201–240.
- 782 Weidenschilling, S., 1977. Aerodynamics of solid bodies in the solar nebula. *Monthly Notices of the Royal Astronomical Society* 180, 57–70.
783 doi:[10.1093/mnras/180.2.57](https://doi.org/10.1093/mnras/180.2.57).
- 784 Weisberg, M.K., Boesenberg, J.S., Kozhushko, G., Prinz, M., Clayton, R.N., Mayeda, T.K., 1995. EH3 and EL3 Chondrites: A Petrologic-Oxygen
785 Isotopic Study, in: *Lunar and Planetary Science Conference*, p. 1481.
- 786 Weiss, B., Elkins-Tanton, L., 2013. Differentiated planetesimals and the parent bodies of chondrites. *Annual Review of Earth and Planetary Sciences*
787 41, 529–560. URL: <https://doi.org/10.1146/annurev-earth-040610-133520>, doi:[10.1146/annurev-earth-040610-133520](https://doi.org/10.1146/annurev-earth-040610-133520),
788 arXiv:<https://doi.org/10.1146/annurev-earth-040610-133520>.
- 789 Wood, J., Dickey, J., Marvin, U., Powell, B., 1970. Lunar anorthosites and a geophysical model of the moon, in: *Proceedings of the Apollo 11 Lunar*
790 *Science Conference*, pp. 965–988.
- 791 Young, E., Shahar, A., Nimmo, F., Schlichting, H., Schauble, E., Tang, H., Labidi, J., 2019. Near-equilibrium isotope fractionation during
792 planetesimal evaporation. *Icarus* 323, 1–15.
- 793

1 Highlights

2 **Structure of differentiated planetesimals: a chondritic fridge on top of a magma ocean**

3 Cyril Sturtz, Angela Limare, Marc Chaussidon, Édouard Kaminski

- 4 • We study the formation and the thermal evolution of planetesimals that accreted undifferentiated material during
5 the first few Myr following the CAI formation.
- 6 • The protracted accretion of cold chondritic material results in the preservation of a few to tens of km thick
7 undifferentiated crust on top of a magma ocean.
- 8 • Basal crustal melts rise in the crust where they stall and crystallize to produce andesitic rocks such as Erg Chech
9 002.