



**HAL**  
open science

## 1.5 Martian Years of Monitoring of the Martian Water Ice Clouds with TGO/ACS-MIR

Aurélien Stcherbinine, Franck Montmessin, Mathieu Vincendon, Michael J. Wolff, Margaux Vals, Christopher S. Edwards, A. Trokhimovskiy, Oleg Korablev, Anna Fedorova, M. Luginin, et al.

► **To cite this version:**

Aurélien Stcherbinine, Franck Montmessin, Mathieu Vincendon, Michael J. Wolff, Margaux Vals, et al.. 1.5 Martian Years of Monitoring of the Martian Water Ice Clouds with TGO/ACS-MIR. Seventh International Workshop on the Mars Atmosphere: Modelling and Observations, Jun 2022, Paris, France. pp.3204. insu-03751710

**HAL Id: insu-03751710**

**<https://insu.hal.science/insu-03751710v1>**

Submitted on 15 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


# 1.5 MARTIAN YEARS OF MONITORING OF THE MARTIAN WATER ICE CLOUDS WITH TGO/ACS-MIR.

**A. Stcherbinine**, *Dept. of Astronomy and Planetary Science, Northern Arizona University, Flagstaff, AZ, USA / LATMOS/IPSL, UVSQ, Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France (aurelien.stcherbinine@nau.edu)*, **F. Montmessin**, *LATMOS/IPSL, UVSQ, Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France*, **M. Vincendon**, *Institut d'Astrophysique Spatiale, Université Paris-Saclay, CNRS, Orsay, France*, **M. J. Wolff**, *Space Science Institute, Boulder, CO, USA*, **M. Vals**, *LATMOS/IPSL, UVSQ, Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France*, **C. S. Edwards**, *Dept. of Astronomy and Planetary Science, Northern Arizona University, Flagstaff, AZ, USA*, **A. Trokhimovskiy**, **O. Korablev**, **A. Fedorova**, **M. Luginin**, *Space Research Institute (IKI), Moscow, Russia*, **G. Lacombe**, **L. Baggio**, **A. Braude**, *LATMOS/IPSL, UVSQ, Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France*.


## Introduction

Water ice clouds play an important role in the Martian water cycle and climate. They are a major actor in the inter-hemispheric water exchange and affect the radiative structure of the atmosphere, as their crystals absorb and scatter the incoming Solar radiation [1, 3, and references contained within]. Thus, monitoring the spatial and temporal evolution of the Martian water ice clouds, along with their physical properties (crystal effective radius, opacity) is of importance to improve our understanding and modeling of the current Martian climate.

The Atmospheric Chemistry Suite (ACS) MIR channel onboard the ESA-Roscosmos Trace Gas Orbiter (TGO) [2] probes the Martian atmosphere in the 2.3–4.2  $\mu\text{m}$  spectral range using the Solar Occultation technique since April 2018. This observing geometry provided detailed vertical profiles of the atmospheric transmission, which allows us to detect and derive the physical properties of the Martian water ice clouds as a function of the altitude for more than one full Martian Year (MY) now.



**Figure 1** – Spatial (latitude) and temporal ( $L_s$  and local time) distribution of the 452 ACS-MIR position 12 observations used in this study. The orange region corresponds to the period of the 2018/MY34 Global Dust Storm.



**Figure 2** – Example of a vertical profile showing the water ice particle size  $r_{\text{eff}}$  (left) and the vertical variation of the extinction coefficient  $k_{\text{ext}}$  at 3.2  $\mu\text{m}$  from a single ACS-MIR observation. The light blue areas represent the altitudes where water ice clouds have been identified, and the greys (and blues) regions indicated all the observed altitudes below the haze top.

## Data & Methods

In this study, we use data acquired using the secondary grating position which covers the 3.1–3.4  $\mu\text{m}$  spectral range (referred to as "position 12") [2] between April 2018 and August 2021. After filtering the data to remove observations in which there was no full spectral coverage ("partial frames"), the dataset analyzed consists of 452 observations ranging from  $L_s = 163^\circ$  (MY 34) to  $L_s = 83^\circ$  (MY 36). The spatial and temporal distribution of these observations is shown in Figure 1. This temporal coverage notably includes the MY 34 Global

Dust Storm (GDS) period and allows us to compare these observations with the ones acquired at a similar epoch during MY 35 (without GDS).

Based on the methodology described in [5], we extract the global continuum transmission spectrum for each observed altitude in ACS-MIR profiles and we convert them into extinction spectra through an onion-peeling vertical inversion algorithm. Then, we compare each ACS-MIR spectrum with theoretical modeled spectra for various sizes of water ice and dust particles. This allows us to simultaneously derive the presence of water ice crystals in each atmospheric layer, along with their effective radii. Figure 2 shows an example of the extinction profile and the water ice crystal sizes obtained from an ACS-MIR observation.

### Latitudinal variations


To isolate the latitudinal variations from the seasonal ones, Figure 3 presents the ACS-MIR cloud profiles acquired between  $L_s = 140^\circ$  and  $L_s = 230^\circ$  (MY 35). We observe that the mean altitude of the water ice clouds follows a bell-shaped distribution as a function of latitude: moving from  $\sim 10$ – $40$  km in the polar regions to  $\sim 40$ – $80$  km around the equator.

We note that regardless of the latitude, there is still a vertical gradient of the particles sizes within the cloud: the size of the ice crystals decreases in the upper layers of the clouds. Larger crystals ( $r_{\text{eff}} \geq 1.5 \mu\text{m}$ ) thus occupy the lower layers of the clouds and are observed up to 55 km around the equator, while the smallest particles ( $r_{\text{eff}} \leq 0.1 \mu\text{m}$ ) can reach 85 km for the same profiles. Clouds are usually founded at the top of the profiles, thus capping the lower layers composed of other types of aerosols (either dust, large water ice crystals, or a mix of them) that may be present in the Martian atmosphere [4].

Another noteworthy point is that this bell-shaped latitudinal distribution noted for the general distribution of the clouds is also observed if we only consider a fixed size of ice crystals. Thus, there is no bijective relation between the altitude and the size of the crystals, other parameters such as the latitude have to be taken into account. This is coherent with the decrease of the scale height of the atmosphere in the higher altitudes.

### Seasonal variations

As noted above, the latitude across the planet affects the vertical distribution of the water ice clouds within the atmosphere. Thus, to decorrelate the seasonal variations from the latitudinal trends, Figure 4 presents the evolution of the vertical clouds profiles from mid-MY 35 to the beginning of MY 36 for the equatorial latitudes (between  $45^\circ\text{S}$  and  $45^\circ\text{N}$ ).




**Figure 3** – Vertical profiles of water ice cloud crystal sizes in the Martian atmosphere as observed by ACS-MIR between  $L_s = 140^\circ$  and  $L_s = 230^\circ$  (MY 35) as a function of the latitude of observation.

We observe that the maximal altitude of the clouds goes from 60 km around  $L_s = 150^\circ$  to 85 km around the perihelion ( $L_s = 270^\circ$ ) and then starts decreasing after  $L_s = 300^\circ$  to reach 50 km at  $L_s = 50^\circ$  (MY 36). Similarly to the latitudinal effect, we also observe that these seasonal variations of the cloud’s altitude affect all the ice particles, regardless of their size. If the proportion of small crystals increases for the high-altitude clouds, at first order the entire size vertical distribution is shifted towards higher altitudes when the average altitude of the clouds increases. I.e., for a fixed value of  $r_{\text{eff}}$  water ice particles are observed at higher altitudes around the perihelion than at the aphelion.

### Conclusion

To conclude, we present here the results of the monitoring of the distribution and characteristics of the Martian water ice clouds with ACS-MIR since the beginning of TGO’s science phase in April 2018. This new dataset allows us to derive the vertical profiles of the clouds’ extinction and crystals sizes over all latitudes for 1.5 MY now, including the MY 34 GDS followed by one classic year.

For a given  $L_s$  range, we observe that clouds are detected about 40 km higher in the equatorial regions than under polar latitudes. And, regarding the equatorial latitudes, we also observe a seasonal evolution of the main cloud’s altitude, with water ice crystals detected about 30 km higher around the perihelion than at the aphelion. We also note that water ice crystals larger than  $1.5 \mu\text{m}$  can be observed up to 50 km under equatorial latitudes around the aphelion during a classic (i.e., GDS-free) year.



**Figure 4** – Vertical profiles of water ice clouds crystals sizes in the Martian atmosphere as observed by ACS-MIR between  $L_s = 140^\circ$  (MY 35) and  $L_s = 90^\circ$  (MY 36) over the equatorial latitudes.

### Acknowledgments

ExoMars is a space mission of ESA and Roscosmos. The ACS experiment is led by IKI Space Research Institute in Moscow. The project acknowledges funding by Roscosmos and CNES. Science operations of ACS are funded by Roscosmos and ESA. Science support in IKI is funded by Federal agency of science organization (FANO). Raw ACS data are available on the ESA PSA at <https://archives.esac.esa.int/psa/#!Table%20View/ACS=instrument>.

### References

- [1] Clancy, R. T., Montmessin, F., Benson, J., et al. 2017, in *The Atmosphere and Climate of Mars*, ed. F. Forget, M. D. Smith, R. T. Clancy, R. W. Zurek, & R. M. Haberle, Cambridge Planetary Science, Cambridge: Cambridge University Press, 76–105. doi:10.1017/9781139060172.005
- [2] Korabev, O., Montmessin, F., Trokhimovskiy, A., et al. 2018. *The Atmospheric Chemistry Suite (ACS) of Three Spectrometers for the ExoMars 2016 Trace Gas Orbiter*, *Space Science Reviews*, 214, 7. doi:10.1007/s11214-017-0437-6
- [3] Montmessin, F., Smith, M. D., Langevin, Y., Mellon, M. T., & Fedorova, A. 2017, in *The Atmosphere and Climate of Mars*, ed. F. Forget, M. D. Smith, R. T. Clancy, R. W. Zurek, & R. M. Haberle, Cambridge Planetary Science, Cambridge: Cambridge University Press, 338–373. doi:10.1017/9781139060172.011
- [4] Smith, M. D., Wolff, M. J., Clancy, R. T., Kleinböhl, A., & Murchie, S. L. 2013. Vertical Distribution of Dust and Water Ice Aerosols from CRISM Limb-Geometry Observations: CRISM LIMB AEROSOLS, *Journal of Geophysical Research: Planets*, 118, 321. doi:10.1002/jgre.20047
- [5] Stcherbinine, A., Vincendon, M., Montmessin, F., et al. 2020. Martian Water Ice Clouds During the 2018 Global Dust Storm as Observed by the ACS-MIR Channel Onboard the Trace Gas Orbiter, *Journal of Geophysical Research: Planets*, 125, e2019JE006300. doi:10.1029/2019JE006300