

Vertebrate paleobiodiversity of the Early Cretaceous (Berriasian) Angeac-Charente Lagerstätte (southwestern France): implications for continental faunal turnover at the J/K boundary

Ronan Allain, Romain Vullo, Lee Rozada, Jérémy Anquetin, Renaud Bourgeais, Jean Goedert, Maxime Lasseron, Jeremy Martin, Adán Pérez-García, Claire Peyre de Fabrègues, et al.

▶ To cite this version:

Ronan Allain, Romain Vullo, Lee Rozada, Jérémy Anquetin, Renaud Bourgeais, et al.. Vertebrate paleobiodiversity of the Early Cretaceous (Berriasian) Angeac-Charente Lagerstätte (southwestern France): implications for continental faunal turnover at the J/K boundary. Geodiversitas, 2022, 44 (25), pp.683-752. 10.5252/geodiversitas2022v44a25. insu-03752198v2

HAL Id: insu-03752198 https://insu.hal.science/insu-03752198v2

Submitted on 16 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

geodiversitas 2022-44-25

Vertebrate paleobiodiversity of the Early Cretaceous (Berriasian) Angeac-Charente Lagerstätte (southwestern France): implications for continental faunal turnover at the J/K boundary

> Ronan ALLAIN, Romain VULLO, Lee ROZADA, Jérémy ANQUETIN, Renaud BOURGEAIS, Jean GOEDERT, Maxime LASSERON, Jerémy E. MARTIN, dán PÉREZ-GARCÍA, Claire PEYRE DE FABRÈGUES, Rafael ROYO-TORRES, Dominique AUGIER, Gilles BAILLY, Lilian CAZES, Yohan DESPRES, Auréliane GAILLIÈGUE, Bernard GOMEZ, Florent GOUSSARD, Thierry LENGLET, Renaud VACANT, MAZAN & Jean-François TOURNEPICHE

> > PUBLICATIONS SCIENTIFIQUES

art. 44 (25) — Published on 21 July 2022 www.geodiversitas.com DIRECTEUR DE LA PUBLICATION / PUBLICATION DIRECTOR : Bruno David, Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / EDITOR-IN-CHIEF: Didier Merle

ASSISTANT DE RÉDACTION / ASSISTANT EDITOR : Emmanuel Côtez (geodiv@mnhn.fr)

MISE EN PAGE / PAGE LAYOUT: Emmanuel Côtez

COMITÉ SCIENTIFIQUE / SCIENTIFIC BOARD : Christine Argot (Muséum national d'Histoire naturelle, Paris) Beatrix Azanza (Museo Nacional de Ciencias Naturales, Madrid) Raymond L. Bernor (Howard University, Washington DC) Henning Blom (Uppsala University) Jean Broutin (Sorbonne Université, Paris, retraité) Gaël Clément (Muséum national d'Histoire naturelle, Paris) Ted Daeschler (Academy of Natural Sciences, Philadelphie) Bruno David (Muséum national d'Histoire naturelle, Paris) Gregory D. Edgecombe (The Natural History Museum, Londres) Ursula Göhlich (Natural History Museum Vienna) Jin Meng (American Museum of Natural History, New York) Brigitte Meyer-Berthaud (CIRAD, Montpellier) Zhu Min (Chinese Academy of Sciences, Pékin) Isabelle Rouget (Muséum national d'Histoire naturelle, Paris) Sevket Sen (Muséum national d'Histoire naturelle, Paris, retraité) Stanislav Štamberg (Museum of Eastern Bohemia, Hradec Králové) Paul Taylor (The Natural History Museum, Londres, retraité)

COUVERTURE / COVER: Réalisée à partir des Figures de l'article/Made from the Figures of the article.

Geodiversitas est indexé dans / Geodiversitas is indexed in:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Physical, Chemical, and Earth Sciences®
- Scopus®

Geodiversitas est distribué en version électronique par / Geodiversitas is distributed electronically by: – BioOne® (http://www.bioone.org)

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Geodiversitas* sont référencés par / Articles and nomenclatural novelties published in Geodiversitas are referenced by: – ZooBank® (http://zoobank.org)

Geodiversitas est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris Geodiversitas is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish: Adansonia, Zoosystema, Anthropozoologica, European Journal of Taxonomy, Naturae, Cryptogamie* sous-sections *Algologie, Bryologie, Mycologie, Comptes Rendus Palevol* Diffusion – Publications scientifiques Muséum national d'Histoire naturelle CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France) Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40 diff.pub@mnhn.fr / http://sciencepress.mnhn.fr

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2022 ISSN (imprimé / print): 1280-9659/ ISSN (électronique / electronic): 1638-9395

Vertebrate paleobiodiversity of the Early Cretaceous (Berriasian) Angeac-Charente Lagerstätte (southwestern France): implications for continental faunal turnover at the J/K boundary

Ronan ALLAIN

Muséum national d'Histoire naturelle, Centre de Recherche en Paléontologie – Paris (CR2P), CNRS/MNHN/Sorbonne Université, CP 38, 57 rue Cuvier, F-75231 Paris cedex 05 (France) ronan.allain@mnhn.fr (corresponding author)

Romain VULLO

Univ Rennes, Géosciences, CNRS UMR 6118, Campus de Beaulieu, bât. 15, 263 avenue du Général Leclerc, F-35042 Rennes cedex (France)

Lee ROZADA

Muséum national d'Histoire naturelle, Centre de Recherche en Paléontologie – Paris (CR2P), CNRS/MNHN/Sorbonne Université, CP 38, 57 rue Cuvier, F-75231 Paris cedex 05 (France)

Jérémy ANQUETIN

Jurassica Museum, Route de Fontenais 21, 2900 Porrentruy, Switzerland Département des Géosciences, Université de Fribourg, Chemin du Musée 6, 1700 Fribourg (Switzerland)

Renaud BOURGEAIS

Résidence du Clos des Ormes, F-91820 Vayres-sur-Essonne (France)

Jean GOEDERT

Université de Bordeaux, De la préhistoire à l'actuel : culture, environnement et anthropologie, UMR 5199, Bâtiment B8, Allée Geoffroy Saint Hilaire, CS 50023 F-33615 Pessac cedex (France)

Maxime LASSERON

Muséum national d'Histoire naturelle, Centre de Recherche en Paléontologie – Paris (CR2P), CNRS/MNHN/Sorbonne Université, CP 38, 57 rue Cuvier, F-75231 Paris cedex 05 (France)

Jeremy E. MARTIN

Univ Lyon, ENSL, Univ Lyon 1, CNRS, LGL-TPE, F-69007 Lyon (France)

Adán PÉREZ-GARCÍA

Grupo de Biología Evolutiva, Facultad de Ciencias, UNED, Paseo Senda del Rey 9, 28040 Madrid (Spain)

Claire PEYRE DE FABRÈGUES

Centre for Vertebrate Evolutionary Biology, Institute of Palaeontology, Yunnan University, Kunming, 650091 (China)

Rafael ROYO-TORRES

Fundación Conjunto Paleontológico de Teruel-Dinópolis, 44002 Teruel (Spain)

Dominique AUGIER Gilles BAILLY

683

Musée d'Angoulême, 1 rue Friedland, F-16000 Angoulême (France)

Lilian CAZES Yohan DESPRES

Muséum national d'Histoire naturelle, Centre de Recherche en Paléontologie – Paris (CR2P), CNRS/MNHN/Sorbonne Université, CP 38, 57 rue Cuvier, F-75231 Paris cedex 05 (France)

Auréliane GAILLIÈGUE

Institut des Sciences du Calcul et des Données (ISCD/MNHN/LJLL) 4, place Jussieu – F-75005 Paris (France)

Bernard GOMEZ

CNRS-UMR 5276 Terre, Planètes, Environnement, Université Lyon 1, 2, rue Raphaël Dubois, F-69622 Villeurbanne, France

Florent GOUSSARD

Muséum national d'Histoire naturelle, Centre de Recherche en Paléontologie – Paris (CR2P), CNRS/MNHN/Sorbonne Université, CP 38, 57 rue Cuvier, F-75231 Paris cedex 05 (France)

Thierry LENGLET

76 rue de la République, F-17150 Mirambeau (France)

Renaud VACANT

Muséum national d'Histoire naturelle, Centre de Recherche en Paléontologie – Paris (CR2P), CNRS/MNHN/Sorbonne Université, CP 38, 57 rue Cuvier, F-75231 Paris cedex 05 (France)

MAZAN

Rue du Bourg, F-16290 Champmillon (France)

Jean-François TOURNEPICHE

Musée d'Angoulême, 1 rue Friedland, F-16000 Angoulême (France)

Submitted on 13 November 2020 | accepted on 29 May 2021 | published on 21 July 2022

urn:lsid:zoobank.org:pub:EA12DCB7-A5BE-4763-B805-25087EBD726D

Allain R., Vullo R., Rozada L., Anquetin J., Bourgeais R., Goedert J., Lasseron M., Martin J. E., Pérez-García A., Peyre De Fabrègues C., Royo-Torres R., Augier D., Bailly G., Cazes L., Despres Y., Gailliègue A., Gomez B., Goussard F., Lenglet T., Vacant R., Mazan & Tournepiche J.-F. 2022. – Vertebrate paleobiodiversity of the Early Cretaceous (Berriasian) Angeac-Charente Lagerstätte (southwestern France): implications for continental faunal turnover at the J/K boundary. *Geodiversitas* 44 (25): 683-752. https://doi.org/10.5252/geodiversitas2022v44a25. http://geodiversitas.com/44/25

ABSTRACT

This contribution describes the continental micro- and macrovertebrate fauna of Angeac-Charente (Berriasian, Early Cretaceous). The rich and diversified fauna includes at least 38 different vertebrate taxa from all major clades, and is represented by more than 50 000 specimens. The Angeac-Charente locality includes the most diverse earliest Cretaceous mixed continental bonebed and the only Lager-stätte known to date in the World, and it provides a good picture of a Purbeckian paleocommunity. It includes remarkable taxa such as a new ornithomimosaur, a large turiasaur, an helochelydrid turtle and numerous mammals. The vertebrate fauna of Angeac-Charente has beyond all a Purberckian character. Many exclusively European genera and species belong to families with an essentially Laurasian paleogeographic distribution. Some taxa nevertheless suggest dispersal events between Africa and Europe at the Jurassic/Cretaceous transition. The successive Charentese faunas of Chassiron (Tithonian), Cherves-de-Cognac (Berriasian) and Angeac-Charente improve our poor knowledge of the evolution of continental vertebrate faunas at the Jurassic/Cretaceous transition. Rather than reflecting an important faunal turnover between the Tithonian and the Berriasian, they record environmental changes related to the sea–level regression that characterizes the end of the Jurassic.

KEY WORDS Vertebrata, Dinosauria, Mammalia, Amphibia, Reptilia, Early Cretaceous, Jurassic/Cretaceous boundary, Berriasian, France, Lagerstätte, Faunal list.

RÉSUMÉ

Paléobiodiversité des vertébrés du Lagerstätte du Crétacé inférieur (Berriasien) d'Angeac-Charente (sud-ouest de la France), implications pour le renouvellement faunique continental à la limite J/K.

La faune de vertébrés continentaux, représentée à la fois par des macro- et des microrestes, du Berriasien d'Angeac-Charente est décrite ici. Cette faune riche et diversifiée comprend au moins 38 taxons différents de vertébrés provenant de tous les grands clades et est constituée de plus de 50000 spécimens. Le gisement d'Angeac-Charente est le plus diversifié des bonebeds mixtes et le seul Lagerstätte continental du Crétacé connu à ce jour dans le monde. Il donne une bonne image de la paléocommunauté purbeckienne. Cette dernière comprend des taxons remarquables tels qu'un nouvel ornithomimosaure, un grand turiasaure, une tortue hélochelydride et de nombreux mammifères. La faune d'Angeac-Charente présente avant tout un cachet purbeckien. De nombreux genres et espèces exclusivement européens appartiennent à des familles dont la répartition paléogéographique est essentiellement laurasiatique. Certains taxons suggèrent néanmoins des événements de dispersion entre l'Afrique et l'Europe à la transition Jurassique/Crétacé. Les faunes charentaises des gisements de Chassiron (Tithonien), de Cherves-de-Cognac (Berriasien) et d'Angeac-Charente améliorent nos connaissances sur l'évolution des faunes continentales à la transition Jurassique-Crétacé. Plutôt que de refléter un important renouvellement faunistique entre le Tithonien et le Berriasien, elles enregistrent les changements environnementaux liés à la régression globale qui caractérise la fin du Jurassique.

MOTS CLÉS Vertebrata, Dinosauria, Mammalia, Amphibia, Reptilia, Crétacé inférieur, limite Jurassique-Crétacé, Berriasien, France, Lagerstätte, liste faunique.

INTRODUCTION

The Purbeckian facies the type section of which is found in the Isle of Purbeck, Dorset, UK, was initially recognized by the French naturalist Alexandre Brongniart (1829). This term applied to the sediments accumulated in very shallow environments, often of the lagoon or lake type, which spanned the transition between the Late Jurassic and the earliest Cretaceous in southern England. These deposits are related to the significant global sea-level regression that characterizes the Jurassic/Cretaceous (J/K) boundary (Hallam 2001). The Purbeckian beds in southern England have also long been known for their fossils (Owen 1853, 1871; Milner & Batten 2002). Since that time, several "Purbeckian" sections and fossil localities have been reported from outside England in western Europe (e.g. Strasser 1986; De Cisneros & Vera 1993; Arp & Mennerich 2008). In the past two decades, three new productive "Purbeckian" fossil localities have been discovered in Charente and Charente-Maritime departments in western France: Chassiron, Cherves-de-Cognac and Angeac-Charente (Fig. 1).

Chassiron, the westernmost and oldest of these localities is located at the extreme northern point of Oléron Island (Fig. 1). It has yielded abundant and diverse fossil remains of plants and animals from an early Tithonian littoral ecosystem that comprises a mix of terrestrial, freshwater and marine taxa, including at least 31 vertebrate taxa (Schnyder *et al.* 2012; Vullo *et al.* 2014). The gypsum quarry of Champblanc is located near Cherves-de-Cognac, 5 km north of Cognac. It records Tithonian and Berriasian deposits from a coastal lagoon to a continental lake (Colin *et al.* 2004; El Albani *et al.* 2004). The alternation of levels of gypsum, claystones, marlstones and limestones has yielded microremains of a diverse vertebrate assemblage, as well as numerous macroremains of fish, turtles and crocodyliforms (Pouech 2008; Pouech et al. 2015; Louchart & Pouech 2017). The easternmost and youngest of these localities, Angeac-Charente, located equidistant from Cognac to the west and Angoulême to the east, yielded its first fossils in 2008 (Allain et al. 2017). The preliminary results, based on the material collected during the first two excavation campaigns in 2010 and 2011, have made it possible to highlight the taxonomic diversity of the site (Néraudeau et al. 2012). Further studies, based on the same material, have clarified the age of the deposit showing it to be late Berriasian (Benoit et al. 2017; Polette et al. 2018), and emphasized the originality of this continental fauna (Allain et al. 2014). Taphonomic and sedimentological studies conducted during the first ten years of excavations have shown that the Angeac-Charente locality represents a "snapshot" of an Early Cretaceous continental swampy ecosystem, and that the richness, diversity and preservation of the fossils qualify the site as a fossil Lagerstätte (Rozada et al. 2014; Gônet et al. 2019; Rozada 2019; Rozada et al. 2021).

The main aims of this study are to describe in outline and figure the complete vertebrate fauna of Angeac-Charente, including the microfauna collected over the past ten years, and to examine the diversity and paleoenvironmental characteristics of this fossil assemblage. Our goal is also to compare this fauna with those known, during the restricted interval that spans the Tithonian and the Berriasian in western France and England and, to a lesser extent, in the Iberian Peninsula and Northern Africa. In the context of the poorly known J/K continental faunal transition (Tennant *et al.* 2017; Campos-Soto *et al.* 2019; Lasseron *et al.* 2020), the latter comparisons are useful to test whether continental vertebrate assemblages record any extinction events, at least at a continental scale.



Fig. 1. – Geographical distribution of the three continental fossil localities documenting the Late Jurassic to Early Cretaceous transition in Western France: Chassiron (Tithonian, Department of Charente-Maritime), Cherves-de-Cognac (Berriasian, Department of Charente) and Angeac-Charente (Berriasian, Department of Charente).

MATERIAL AND METHODS

INSTITUTIONAL ABBREVIATIONS

- ANG Angeac-Charente Collection, Musée d'Angoulême, Angoulême, France;
- DORCM Dorset County Museum, Dorchester, United Kingdom; FMNHN Field Museum of Natural History, Chicago, United
- FMNHN Field Museum of Natural History, Chicago, United States.

FIELDWORK

The first dinosaur bones were discovered at Angeac-Charente as early as 2008, in a quarry operated by the Audoin & Fils Company. By exploiting the sands and gravels left during the Pleistocene by the Charente River, which now flows 2 km northeast of the site, the quarry worker Jean-Pierre Paillot discovered a vertebra, a metatarsal and some fragments of large sauropod bones (Néraudeau *et al.* 2012; Allain *et al.* 2017). Following the discovery of new bones by quarrymen in January and February 2010, a first excavation campaign involving about 15 people from the Musée d'Angoulême, the Rennes University and the Muséum national d'Histoire naturelle was carried out between late August and early September 2010 (Fig. 2A). The excavation area of 20 m² yielded 599 fossils of vertebrates, invertebrates and plants, including a 2.2 m long sauropod femur (Néraudeau et al. 2012: fig. 11). On the strength of these results, from the following year, and thanks to the support of the local authorities and the Audoin Company, larger scale excavations were set up. Thus, every year for one month since summer 2011, a team of around forty people has been manually excavating the fossiliferous clays of Angeac-Charente (Fig. 2). The paleontological excavations are divided into two main plots, a northwestern plot called CG, and a southwestern plot called R (Fig. 3; Rozada et al. 2021). "CG" means "Conseil Général" and refers to the fact that this part of the site is today the property of the Department of Charente, while "R" stands for "Rodet", the family name of the owner of this part of the quarry. By the end of the 2019 campaign, more than 800 m² had been excavated on an average thickness of 1 m. More than 7500 macrofossils were collected, identified, mapped using theodolite and total station, and fully prepared. To these can be added around 67000 unidentified specimens (i.e. bone fragments), 3350 coprolites and tens of thousands of microremains (Rozada et al. 2021).



Fig. 2. – Views of the Angeac-Charente gravel quarry: **A**, CG1 plot at the end of the 2010 excavation campaign; **B**, CG1 and CG3 plots at the beginning of the 2013 excavation campaign; **C**, CG4, CG6, CG7 plots (on the left), and CG9 plot (on the right) during the 2018 excavation campaign.

Microremains

Nearly all the microremains presented in this study were collected in 2014 from within a large lens in Unit 3 clay of R1 plot (Fig. 3), composed of sand, soft white calcareous clasts, wood fragments and vertebrate remains, including a nearcomplete turtle shell (Gônet *et al.* 2019; Rozada *et al.* 2021). Several hundred kilograms of sediment from other plots have also been sampled but have yet to be fully sorted. All sediment samples were dried before treating by screen-washing. In the field, the dissociation of clay was only performed with water. Screen-washing was made through four successive sieves of 4, 1.2, 0.8 and 0.5 mm mesh. In the laboratory of the Centre de Recherche en Paléontologie – Paris (CR2P), at the Muséum national d'Histoire naturelle, the lithoclastic and bioclastic sand sieved following this process was reconcentrated using formic acid for carbonate dissolution and sodium dithionite to remove iron-bearing mineral phases. The size fractions were then separated using sieves of 2, 1 and 0.5 mm mesh



Fig. 3. — Spatial distribution of identified vertebrate bodyfossil remains of Angeac-Charente recovered from 2013 to 2018: **A**, aerial view of the Angeac-Charente site with the location of the manual (R1-3 and CG1-8) excavation plots and some remarkable fossils (drone picture ©D. Abit, July 2017); **B**, diagram of the Angeac-Charente locality with the position of vertebrate body fossil remains according to taxon type. Each point represents the position of one specimen (3-D SIG data measured from 2013 to 2018 projected with ArcGIS Desktop 10.3.1.). SIG data from the CG1 plot (2010-1012) are lacking (in grey) (See Rozada *et al.* 2021).

and sorted under stereomicroscope by two of us (G. Bailly and R. Vacant). The vertebrate microremains were counted using the 'Count Image Elements' script, specifically developed by A. Gailliègue (pers. comm., April 2017) and based on the principle of picture segmentation (see Lasseron *et al.* 2020). SEM photomicrographs were made at the CR2P and at Rennes University.

REPOSITORY

All fossil specimens collected in Angeac-Charente were officially donated to the Musée d'Angoulême (Charente, France) by the owners of the different plots of land that made up the fossiliferous part of the quarry: the Audoin & Fils Company, Mrs Rodet and the Charente department. These fossils are deposited in the collections of the Musée d'Angoulême, under collection numbers with the following syntax: *ANGyy-nnnn, ANGyy-Rnnnn* or *ANG M-nnnn*, where "ANG" is an abbreviation of Angeac-Charente, "M" means microremain, "R" means Rodet, "yy" year of discovery and "nnnn" the specimen number.

SEDIMENTOLOGICAL FRAMEWORK AND TAPHONOMY

DEPOSITIONAL ENVIRONMENT

A detailed account of the general taphonomy and sedimentology of the Angeac-Charente Lagerstätte has already been published (Rozada *et al.* 2021). The site represents a "snapshot" into a Lower Cretaceous swampy environment and the sedimentological assemblage consists mainly of clay deposits. We provide here a synthetic sedimentary log of the Angeac-Charente bonebed (Fig. 4). The sandy clay (Unit 2), clay of decantation (Unit 3), and sand/conglomerate (Unit 4) deposits contain the bonebed and have the same paleontological content, but they correspond to different energy of deposition episodes (Fig. 4):

- Deposition Event 1 (DE 1): The system begins with the deposition of a high energy sandy clay (Unit 2), which erodes the underlying green clay (Unit 1) and brings into it many plant and vertebrate fossils.

- Deposition Event 2 (DE 2): The clay of decantation (Unit 3) is deposited over a longer time interval and under a stagnant water column. There is no discordance or break in sedimentation between Units 2 and 3. A few mudcracks may indicate occasional, localized and temporary exposure of the clay. This low energy deposit is occasionally interrupted by streams of variable energy, depositing lenses rich in coarser grains and fossils. Unit 3 has yielded most of the Angeac-Charente fossils.

– Deposition Event 3 (DE 3): A more or less clayey and silty sandbank that sometimes grades laterally into sandstone and conglomerate (Unit 4) was deposited at a higher energy level. It probably corresponds to a discrete flooding event from a nearby river. During the 2019 field campaign, interbedded lamination of silt and sand that may indicate deposits left by a river channel have been observed in this unit.

In parallel with these three depositional events, synsedimentary deformational structures formed in water-saturated sediments can be observed such as convolute lamination in clay, folded figures in sands and rebalance loads. These numerous sedimentological structures, allied with taphonomic "frozen scenes", suggest a coeval and continuous deposition of all three units (Rozada et al. 2021). The homogeneous composition of the clay mineral assemblage of the whole lithostratigraphic section (Néraudeau et al. 2012) and the Rare Earth Elements and Yttrium profiles measured in sediment thoughout the lithological section and on dinosaur fossil biogenic apatite samples (Rozada et al. 2021) also confirm that all the sediments and vertebrate remains were deposited under similar, coeval, poorly oxygenated burial and diagenetic conditions. These mineralogical and geochemical analyses also indicate that the Angeac-Charente organisms fossilized near their living habitat and/or place of death in a local depositional environment and demonstrate a limited time averaging of the fossil concentration.

Thus, the site represents a "snapshot" into an Early Cretaceous swampy ecosystem, very likely a floodplain connected to a nearby river. The Angeac-Charente bonebed accumulated on a poorly oxygenated freshwater floodplain dominated by cheirolepidiacean vegetation in a tropical to subtropical climate (Néraudeau *et al.* 2012; Polette *et al.* 2018; Rozada *et al.* 2021). The fossil richness, diversity and preservation permit designation of the site as a fossil Lagerstätte (Rozada *et al.* 2021).

SPATIAL DISTRIBUTION OF ANGEAC-CHARENTE VERTEBRATE MACROREMAINS

Rozada *et al.* (2021) provide the first data concerning the spatial distribution of vertebrate remains based on identified and unidentified macroremains collected between the 2010 and 2017 field campaigns. Herein, we supplement these data by mapping the position of identified vertebrate remains recovered between the 2013-2018 field campaigns (Fig. 3B).

Turtle remains are mainly represented by isolated, complete or fragmented shell plates and a few bones, and are quite homogeneously distributed throughout the whole site (Fig. 3B). The collected remains of *Pleurosternon bullockii* belong at least to twelve individuals of different sizes. The remains are generally well-preserved, with no evidence of weathering or erosion, although they are frequently fragmented. A very well-preserved and nearly complete turtle shell of *Pleurostenon bullockii* was collected in 2014, from a lens in R1 (Gônet *et al.* 2019; Fig. 3A). At several loci of the site, connected plates of *P. bullockii* were also collected. In 2017, numerous plates belonging to a nearly complete specimen have been found in loose articulation in the R3 plot (Fig. 3A). All these observations suggest that pleurosternid turtles were autochthonous.

Crocodyliforms are known from numerous isolated teeth (mostly shed due to continuous tooth replacement), osteoderms, and other bones including numerous skull bone fragments, randomly distributed throughout the whole site. Most of these remains belong to the generalist genus *Goniopholis* (see below). In 2018, numerous *Goniopholis* bones (skull, mandibles, vertebrae, limb, girdle bones and osteoderms) belonging



Fig. 4. – Angeac-Charente synthetic log based on the lithostratigraphic sections of R3, CG3 and CG9, of Rozada *et al.* (2021), and the main corresponding depositional events. The thickness is measured above and below the Unit 1/Unit 2 limit, which marks the base of the bonebed. Abbreviations: **DE**, Depositional Event; **U**, Unit.



FIG. 5. — Scanning electron micrographs of hybodont shark teeth from Angeac-Charente: **A**, **B**, lateral tooth of *Parvodus celsucuspus* Rees, Cuny, Pouech & Mazin, 2013 (ANG M-107) in labial (**A**) and occlusal (**B**) views; **C**, anterior tooth of *Parvodus celsucuspus* (ANG M-17) in labial view; **D**, dermal denticle (ANG M-109) in lateral view; **E**, **F**, dermal denticle (ANG M-108) in lateral (**E**) and apical (**F**) views. Scale bar: C, 1 mm; A, B, D-F, 400 µm.

to a single individual were found in loose articulation in the CG9 plot (Fig. 3A). Otherwise, the remains of Pholidosauridae, Bernissartiidae and Atoposauridae are rarer and mainly recovered from sieved microremains, and are therefore under represented on the map (Fig. 3B).

Ornithomimosaur remains are concentrated in the CG1-9 plots (85% of the ornithomimosaur bones) and become progressively rarer toward the SE and in the NW part of the site, in the CG9 plot (Fig. 3). This spatial distribution supports a mass mortality event involving a herd (Rozada *et al.* 2021).

So far, no postcranial remains of carnivorous dinosaurs have been discovered at Angeac-Charente. However, nearly 150 isolated teeth have been collected. They are randomly distributed and most of them have not been mapped.

Sauropod remains, represented by numerous teeth and bones from all parts of the skeleton, come from at least seven individuals with no apparent connection to each other. They are abundant in most areas, but become rarer toward the NW and the SE. Although they seem mainly concentrated in R1 plot, this is a false impression as bones are heavily fragmented, and each fragment has been mapped at some point. The R2 and CG10 plots have yielded the well-preserved hindquarters of a single individual, including the last dorsal vertebra, the sacrum, the two ischia, a few chevrons and ribs, and a complete femur (Figs 3A; 27).

Stegosaurs are represented by a few bones, mostly vertebrae and ribs and scarce appendicular and skull bones. Although they are present in all plots of the site, they are particularly concentrated in CG4. In this area, the bones show a homogeneous state of preservation, and probably belong to a single individual, which died near the site. Other ornithischian remains (camptosaurid ornithopods, hypsilophodontids, heterodontosaurids and ankylosaurs) are rare.

SYSTEMATIC PALEONTOLOGY

VERTEBRATA Lamarck, 1801 PISCES Linnaeus, 1758 CHONDRICHTHYES Huxley, 1880 ELASMOBRANCHII Bonaparte, 1838 HYBODONTIFORMES Patterson, 1966 Family LONCHIDIIDAE Herman, 1977 Genus *Parvodus* Rees & Underwood, 2002

Parvodus celsucuspus Rees, Cuny, Pouech & Mazin, 2013 (Fig. 5)

DESCRIPTION

This species is characterized by high-cusped anterior teeth, and is well represented at Angeac-Charente. The material includes numerous isolated teeth and dermal denticles (Fig. 5), as well as a few incomplete cephalic and dorsal fin spines (Néraudeau *et al.* 2012), that are identical to those of Cherves-de-Cognac (Rees *et al.* 2013).

OSTEICHTHYES Huxley, 1880 ACTINOPTERYGII Klein, 1885 PYCNODONTIFORMES Berg, 1937 Family MESTURIDAE Nursall, 1996 Genus *Micropycnodon* Hibbard & Graffham, 1945

cf. *Micropycnodon* sp. (Fig. 6A-C)

Description

Teeth attributed to Mesturidae, sub-circular to oval in occlusal view, are characterized by the presence of several



Fig. 6. — Actinopterygian remains from Angeac-Charente: **A**, vomerine dentition of cf. *Micropycnodon* sp. (ANG M-69) in occlusal view; **B**, **C**, detailed view of an isolated vomerine tooth of cf. *Micropycnodon* sp. (ANG M-28) in (**B**) anterior and (**C**) occlusal view; **D**, **E**, isolated tooth of Pycnodontidae indet. (ANG M-50) in occlusal (**D**) and lateral (**E**) views; **F**, isolated tooth of Pycnodontidae indet. (ANG M-59) in lateral view. Scale bars: A, 2.5 mm; B, C, F, 500 µm; D, E, 1 mm.

small, sharp tubercles, arranged in rows or surrounding a shallow central depression (Fig. 6A-C). A nearly complete, relatively broad vomerine dentition shows six tooth rows (Fig. 6A). Crown morphology and tooth arrangement are similar to those of *Micropycnodon* (Dunkle & Hibbard 1946; Cronin & Shimada 2019). *Micropycnodon* is known from the Late Cretaceous of North America, but similar genera (possibly subjective junior synonyms) known from the Early Jurassic of Germany (*Grimmenodon*; Stumpf *et al.* 2017) and the Early Cretaceous of Texas (*Texasensis* pro *Callodus*; Thurmond 1974; Özdikmen 2009) suggest a long temporal range for this lineage. The form in Angeac-Charente also occurs in the Purbeckian facies of Cherves-de-Cognac, and was described as "pycnodontiform tooth morphotype 10" by Pouech *et al.* (2015: fig. 8e-f).

Family PYCNODONTIDAE Agassiz, 1835

Pycnodontidae indet. (Fig. 6D-F)

Description

A second pycnodontiform taxon is represented by beanshaped to suboval teeth with a transverse groove or depression



Fig. 7. – Actinopterygian remains from Angeac-Charente: **A**, **B**, isolated tooth of Ginglymodi indet. (ANG M-30) in anterior/posterior (**A**) and occlusal (**B**) views; **C**, Ginglymodi indet. scale (ANG M-68); **D**, Ginglymodi indet. scale (ANG M-67); **E**, upper jaw fragment of Amiiformes indet. (ANG M-62) in lingual view; **F**, left dentary of Amiiformes indet. (ANG M-117) in labial view; **G**, **H**, fragmentary left dentary of Amiiformes (ANG 15-4062) in lingual (**G**) and labial (**H**) views. Scale bar represents: A, B, 1 mm; C, 4 mm; D, 2 mm; E, 2.5 mm; F, 10 mm; G, H, 5 mm.

and one or two lateral tubercles (Fig. 6D-F). This taxon is also present in the Purbeckian facies of Cherves-de-Cognac, and has been described as "tooth morphotype 7" by Pouech et al. (2015: fig. 8b). A very similar and likely congeneric form has also been described from the Wealden facies of southern England as Coelodus sp. (Sweetman et al. 2014) then assigned to Ocloedus sp. (Sweetman 2016). However, the English material includes nearly complete vomerine dentitions with three tooth rows (Sweetman et al. 2014: fig. 10e), indicating that this unnamed species does not belong to either Coelodus or Ocloedus, which both have five vomerine tooth rows (Poyato-Ariza & Wenz 2002). Numerous ventral keel scales showing contacting spines of anteroposteriorly increasing size, can be assigned to Pycnodontidae (see Poyato-Ariza & Wenz 2002; Sweetman et al. 2014).

HOLOSTEI Müller, 1844 (*sensu* Grande 2010) GINGLYMODI Cope, 1872 (*sensu* López-Arbarello & Sferco 2018)

Ginglymodi indet. (Fig. 7A-D)

DESCRIPTION

Material from Angeac-Charente mainly includes isolated teeth and ganoid scales (Fig. 7A-D). Most teeth show a typical sub-hemispheric, unornamented crown. A small tip is sometimes developed at the apex of the crown. Scales are relatively thick and rhomboid in shape. An anteroventral process can be present in addition to the anterodorsal process. The posterior margin is not serrated. Based on the available material, one or two ginglymodian taxa may be present, with possibly a callipurbeckiid (Semionotiformes) and/or a lepidotid (Lepisosteiformes) (*sensu* López-Arbarello & Sferco 2018).



Fig. 8. — Actinopterygian remains from the Berriasian of Angeac-Charente: **A**, isolated tooth of Amiiformes indet. (ANG M-39); **B**, **C**, isolated tooth of Amiiformes indet. (ANG M-110); **D**, dentulous vomerine element of Amiiformes indet. (ANG M-56); **E-G**, isolated tooth of Ionoscopiformes indet. (ANG M-111); **H**, jaw fragment of Actinopterygii indet. (ANG M-38) in lingual view; **I**, vertebra of Actinopterygii indet. (ANG M-069). Scale bar: A-C, E-H, 500 µm; D, 250 µm; I, 2 mm.

HALECOMORPHI Cope, 1872 AMIIFORMES Hay, 1929 (*sensu* Grande & Bemis 1998)

> Amiiformes indet. (Figs 7E-H; 8A-D)

DESCRIPTION

Amiiform fish are represented in the assemblage by isolated teeth, jaw remains (Fig. 7E-H) and vertebrae. Most of the labiolingually compressed teeth show a typical triangular crown apex and bear two well-developed carinae (Fig. 8A-C). Such teeth were previously referred to *Caturus* sp. (Sweetman *et al.* 2014; Vullo *et al.* 2014; Pouech *et al.* 2015), although this dental morphology is also present in more derived members of Amiiformes (i.e., Amiidae). Some tooth associations are interpreted as vomerine dentitions (Fig. 8D). These elements bear teeth of various sizes, subconical and devoid of carinae, as those found in many Amiiformes (Grande & Bemis 1998). It is worth noting that this tooth morphology corresponds

to that usually assigned to the aspidorhynchid *Belonostomus* (e.g., Sweetman *et al.* 2014; Pouech *et al.* 2015). However, the rounded morphology of the vomerine tooth plates from Angeac-Charente and the fact that they are paired elements preclude assignment to *Belonostomus* (in which the vomer is unpaired); therefore, all isolated small teeth with a conical crown devoid of carinae are here referred to Amiiformes indet.

IONOSCOPIFORMES sensu Grande & Bemis (1998)

> Ionoscopiformes indet. (Fig. 8E-G)

DESCRIPTION

Small, curved teeth with a pointed apex bearing two short, blunt carinae are assigned to an indeterminate ionoscopiform (Fig. 8E-G). Small ganoid scales characterized by a finely serrated posterior margin may also belong to this group. Such a material is similar to that described from the Purbeckian beds of Chassiron (Vullo *et al.* 2014). As some recent studies suggest that Ionoscopiformes is not a monophyletic clade (Ebert 2018; López-Arbarello & Sferco 2018), the material from Angeac-Charente is assigned to Ionoscopiformes *sensu* Grande & Bemis (1998).

AMPHIBIA Linnaeus, 1758 LISSAMPHIBIA Haeckel, 1866 ALLOCAUDATA Fox & Naylor, 1982 Family Albanerpetontidae Fox & Naylor, 1982

> Albanerpetontidae indet. (Fig. 9)

DESCRIPTION

Albanerpetontids are represented in Angeac-Charente by numerous and diverse bones, including dentaries, premaxillae, maxillae, vertebrae and forelimb bones (Fig. 9A-K). All the material was recovered from screen washing residues. It is always disarticulated, and almost always fragmentary. Several diagnostic characters allow their assignment to the Albanerpetontidae, including: 1) intertonguing symphyseal joint between dentaries, in a mortise-and-tenon style (Fox & Naylor 1982; Milner 1988; McGowan & Evans 1995; McGowan 1996, 2002; Gardner 1999b, 2000; Gardner et al. 2003; Sweetman & Gardner 2013; Matsumoto & Evans 2018). In the Angeac-Charente specimens, there are two symphyseal prongs (Fig. 9A, B); 2) pleurodont, chisel-like and regularly arranged non-pedicellate teeth, labiolingually compressed and bearing three faint, mesiodistally aligned cuspules (Fig. 9A-G; Fox & Naylor 1982; Gardner 1999a, b, 2000; McGowan & Evans 1995; Sweetman & Gardner 2013; Matsumoto & Evans 2018); and 3) foramina arranged in line on external face of dentary (Fig. 9F; Fox & Naylor 1982; Gardner 2000).

In Angeac-Charente albanerpetontids, the maxilla is unornamented labially, except for scattered external nutritive foramina that are characteristic of the group (Fig. 9D; Fox & Naylor 1982; Gardner 2000). In this respect, they differ from Albanerpeton inexpectatum from the Miocene of France (Gardner 1999a). As in other albanerpetontid for which trunk vertebrae have been described (Estes & Hoffstetter 1976; McGowan 1996, 2002; McGowan & Ensom 1997; Sweetman & Gardner 2013; Matsumoto & Evans 2018), those from the Angeac-Charente material (Fig. 9I-K) are amphicoelous, hourglass-shaped and bear a short unicipital transverse process. The centrum is narrowly constricted at its center (Fig. 9I, K) and the cotyles are circular in outline and have thickened rims (Fig. 9J). As in other albanerpetontid trunk vertebrae described elsewhere, the notochordal canal is anteroposteriorly continuous (Fig. 9J; Sweetman & Gardner 2013) and thus the vertebrae are fully notochordal.

Angeac-Charente specimens also show numerous features that have been described in other albanerpetontids and allow the distinction from other lissamphibians and lizards. The premaxilla (Fig. 9D, E) and maxilla (Fig. 9C) have a deep *pars dentalis* and the dentary (Fig. 9A, B, F, G) has a tall dental parapet, allowing the attachment of highly pleurodont teeth (Gardner 2000). Upper jaws are also characterized by a prominent, shelf-like *pars palatinum* lingually (Fig. 9E; Gardner 2000). The maxilla has a low, posteriorly tapered *pars facialis* (Fig. 9C; Gardner 1999a, 2000). On the dentary, the Meckelian canal is closed anteriorly, and the subdental shelf is low, narrow and gutter-like anteriorly (Fig. 9A; Gardner 1999a, b, 2000). As in other known albanerpetontid species, the humeral condyle is spherical, fully ossified and larger than the adjacent radial epicondyle (Fig. 9H; Sweetman & Gardner 2013). Above this humeral ball is a triangular and well-defined cubital fossa, at the proximal extremity of which a small foramen can be seen (Fig. 9H), as also reported for *Albanerpeton inexpectatum* (Estes & Hoffstetter 1976) and *Wesserpeton evansae* (Sweetman & Gardner 2013).

The albanerpetontids from Angeac-Charente differ from Anoualerpeton and Albanerpeton nexuosum, but resemble other Albanerpeton species, Celtedens and Wesserpeton in having dentaries and maxillae with relatively straight occlusal margins, and teeth weakly heterodont in size (Sweetman & Gardner 2013). Unfortunately, the diagnostic characters necessary for identification at the generic level are lacking. Neither the maxilla nor the dentary is diagnostic for Albanerpeton and Celtedens (Gardner 2000), as their diagnoses are based on frontal characters (Gardner 1999a, b, 2000; McGowan & Evans 1995; McGowan 2002). We did not find any frontal bones in the Angeac-Charente material, and so we cannot attribute the albanerpetontid material to these genera with certainty. Within Albanerpeton, the premaxilla is the most taxonomically informative bone for species (Gardner 1999b, 2000), but the specimens found to date at Angeac-Charente are too incomplete to be relevant.

Thus, considering the characters mentioned above, we can attribute the Angeac-Charente specimens to the family Albanerpetontidae, but so far the material is too incomplete to allow a further attribution at the generic and specific level.

ANURA Fischer von Waldheim, 1813

Anura indet. (Fig. 10)

DESCRIPTION

Despite a swampy paleoenvironment that might be suitable for their occurrence and preservation, anurans are relatively scarce among the microvertebrate material from Angeac-Charente. In Angeac-Charente, anurans are represented by isolated, fragmentary bones, such as fused zeugopods (Fig. 10A, B) and urostyles (Fig. 10C, D)

Among fused zeugopods, some are sufficiently well preserved to allow their identification. For instance, the general shape and length (even if the proximal part is missing) of specimen ANG M-63 (Fig. 10A, B), as well as several other features, allow its identification as a tibiofibula (Thomas 1996): presence of a tibiofibular groove, deeper on the anterior face, between the fused zeugopod, and presence of a tibial crest on the anterior face (Fig. 10A). Among well-known Early Cretaceous anuran families are the Alytidae (= "Discoglossidae"), but within this



Fig. 9. — Albanerpetontid remains from Angeac-Charente: A, B, left dentary of Albanerpetontidae indet. (ANG M-101) in lingual (A) and dorsal (B) views; C, premaxillary of Albanerpetontidae indet. (ANG M-102) in ventrolingual view; D, E, left maxillary of Albanerpetontidae indet. (ANG M-51) in labial (D) and lingual (E) views; F, G, dentary fragment of Albanerpetontidae indet. (ANG M-35) in labial view showing foramina discussed in the text (F) and lingual (G) view; H, distal end of a left humerus of Albanerpetontidae indet. (ANG M-36) in ventral view; I-K, dorsal vertebra of Albanerpetontidae indet. (ANG M-103) in dorsal (I), anterior (J) and lateral (K) views. Scale bars: A-E, H-K, 400 µm; F, G, 1 mm.



FIG. 10. – Anuran remains from Angeac-Charente: A, B, right tibiofibular of Anura indet. (ANG M-63) in anterior (A) and posterior (B) views; C, D, urostyle of Anura indet. (ANG M- 64) in dorsal (C) and ventral (D) views. Scale bar represents: A, B, 5 mm; C, D, 2 mm.

family, the tibia and fibula remain unfused (Roček 2000), unlike the Angeac-Charente specimen.

The specimen ANG M-64 (Fig. 10C, D) bears two broad and flattened articular facets anteriorly, and a small crest arises from its dorsal face. Four weakly marked ridges can be seen alongside this small crest, two on each side (Fig. 9C). This morphology identifies it as a urostyle. To date, incompleteness of the material limits comparisons with other Mesozoic anurans.

CAUDATA? Scopoli, 1777

Caudata? indet. (Fig. 11)

DESCRIPTION

Among the microvertebrates from Angeac-Charente, a specimen could be referred to Caudata. Indeed, the vertebra ANG M-71 (Fig. 11) is anteroposteriorly elongated, with a broad, almost circular cotyle (Fig. 11A) and with well-defined preand postzygapophyses. The posterior part of the centrum is abraded, but the vertebra is clearly procoelous. Two small subcentral foramina are present on the ventral face (Fig. 11F). The general shape, presence of transverse processes extending posterolaterally, and broad vertebral cotyle are reminiscent of Caudata (Rage et al. 1993). In Caudata, the trunk vertebrae are, however, commonly opisthocoelous or amphicoelous and the procoelous condition is rare (see Estes 1981; Rage et al. 1993; Alloul et al. 2018 for examples of procoelous caudates). On the contrary, the procoelous condition is common within Squamata, to which this vertebra could be alternatively referred. Moreover, anterior basapophyses, which are present in many caudate groups (Estes 1981), are not discernable, and the presence or absence of a notochordal pit, which is usually observed on caudate vertebrae (Alloul et al. 2018), cannot be inferred because of the abraded condyle. However, the left transverse process (= rib-bearer), although broken, has an expanded head (Fig. 11C, E, F), indicating that it may be bilobed as in salamanders, and a ridge extending between the transverse process and the condyle seems to be present, as in Caudata (Alloul *et al.* 2018), but the preservation is too poor to reach a conclusion. Thus, this vertebra is tentively referred to Caudata?, although an assignment to Squamata cannot be excluded.

TESTUDINATA Klein, 1760

PRELIMINARY STATEMENT

Turtles are well represented in Angeac-Charente, notably by numerous isolated shell plates and bones of the axial and appendicular skeletons, and more rarely by articulated shell material (see Table 1; Fig. 35A). Only a few pieces of skulls, including a maxillary and a mandible, have been recovered. Néraudeau *et al.* (2012) have previously reported the presence of three distinct turtle taxa in Angeac-Charente: a pleurosternid, a solemydid (now helochelydrid), and a third taxon characterized by shell bones without surface ornamentation (Figs 12; 13). Since then, additional and more complete material confirms these preliminary conclusions (see below). The third taxon is now identified as a thalassochelydian.

> PERICHELYDIA Joyce, 2017 Family HELOCHELYDRIDAE Nopcsa, 1928 (sensu Joyce et al. 2016)

> > Helochelydridae indet. (Fig. 12A, B)

DESCRIPTION

In Angeac-Charente, helochelydrids are mostly represented by incomplete isolated shell elements covered by typical



FIG. 11. – Vertebra of Caudata indet. from Angeac-Charente: A-F, (ANG M-71) in anterior (A), posterior (B), left lateral (C), right lateral (D), dorsal (E) and ventral (F) views. Scale bar: 2 mm.



Fig. 12. — Helochelydrid and thalassochelydian turtle remains from Angeac-Charente: **A**, peripheral plate of Helochelydridae indet. in dorsal view; **B**, pleural plate of Helochelydridae indet. in dorsal view; **C**, **D**, right hyoplastron of *Hylaeochelys belli*? (Mantell, 1844) (ANG18-5915) in ventral (**C**) and dorsal (**D**) views; **E**, left humerus of *Hylaeochelys*? *belli*? (ANG18-5937) in ventral view; **F**, right humerus (ANG18-5938) in dorsal view; ; **G**, **H**, Peripheral of *Hylaeochelys*? *belli*? (ANG 18-5993) in dorsal (**G**) and ventral (**H**) views. Scale bar: A, 1 cm; E-H, 2 cm; B, 3 cm; C, D, 4 cm.

tubercles, pustules and crests (Fig. 12A, B) that diagnose this clade (Lapparent de Broin & Murelaga 1999; Joyce 2017). Probably based on the previous stratigraphical claim by Néraudeau et al. (2012) that Angeac-Charente was Hauterivian-Barremian in age, Joyce (2017) provisionally referred this material to Helochelydra nopcsai, a species otherwise diagnosed by a shell surface ornamentation consisting of distinct, easily dislocated tubercles. Our observations reveal, however, that the helochelydrid material from Angeac-Charente usually displays distinct tubercles that do not easily dislocate. This ornamentation is more similar to the morphotypes of "Helochelydra" anglica and "Helochelydra" bakewelli (Joyce 2017), respectively from the Purbeck Group (Berriasian) of Dorset and the Wealden (Valanginian) of Sussex, United Kingdom (Joyce et al. 2011). That being said, the systematics of Early Cretaceous helochelydrids remains highly confused. For that reason, the helochelydrid material from Angeac-Charente is herein conservatively referred to Helochelydridae indet.

The posterior part of a large carapace collected in 2018 and 2020 provides new information on the morphology of the helochelydrid from Angeac-Charente. However, the material is not fully prepared yet and will be described in more detail elsewhere. A raised midline keel on the posterior neurals is clearly observable, which is a character shared with other helochelydrids (Lydekker 1889; Lapparent de Broin & Murelaga 1999; Milner 2004; Vullo et al. 2010; Joyce et al. 2014; Joyce 2017; Pérez-García et al. 2020). The general morphology resembles that of other species in which this region is sufficiently known, such as "Helochelydra" anglica, Naomichelys speciosa, Aragochersis lignitesta, and Solemys vermiculata (Lydekker 1889; Lapparent de Broin & Murelaga 1999; Milner 2004; Joyce et al. 2014; Joyce 2017; Pérez-García et al. 2020). Neurals V and VI are elongate and hexagonal. Neurals VII and VIII are probably fused. They are wide and hexagonal anteriorly and posteriorly, and narrower on the midline. There are three suprapygals, which are reminiscent of the condition in FMNH PR273, a nearly complete specimen referred to the North American species Naomichelys speciosa. However, the number of suprapygals is known to be rather variable in basal turtles. The morphology of the suprapygals and pygal differs from other known helochelydrids, but detailed comparisons are needed in order to assess the systematic value of these characters. The shell surface sculpturing is less prominent on the neurals and becomes more pronounced on the distal part of the costals and on the peripherals, suprapygals, and pygal.

Helochelydrids are considered by numerous authors to be terrestrial forms (Lapparent de Broin & Murelaga 1999; Joyce *et al.* 2011; Scheyer *et al.* 2015). They are known to have limbs and neck covered with osteoderms (Barrett *et al.* 2002; Joyce *et al.* 2014; Scheyer *et al.* 2015). Such osteoderms have also been found at Angeac-Charente. Known cranial material suggests a durophagous diet, although it is still unclear exactly which hard-shelled food elements these turtles fed on (Joyce *et al.* 2011, 2014). PARACRYPTODIRA Gaffney, 1975 (sensu Lyson & Joyce 2011) Family PLEUROSTERNIDAE Cope, 1868 (sensu Lyson & Joyce 2011) Genus Pleurosternon Owen, 1853

Pleurosternon bullockii (Owen, 1842) (Fig. 13)

Description

Pleurosternids are the most abundant turtles in Angeac-Charente representing nearly 85% of the turtle material (Table 1). Most of the material consists of isolated shell plates characterized by an external surface covered by regular pits with fine linear striations perpendicular to the plate margins. This character combination is only shared with the Portuguese Late Jurassic (Kimmeridgian) Selenemys lusitanica and the British and French Late Jurassic-Early Cretaceous (Tithonian to Berriasian) Pleurosternon bullockii (Pérez-García & Ortega 2011). An almost complete shell collected in 2014 confirmed the attribution to the species Pleurosternon bullockii (Fig. 13; Gônet et al. 2019). The carapace is oval in outline and depressed with no nuchal emargination (Fig. 13A). The nuchal is trapezoidal, wider posteriorly than anteriorly. The cervical scute is absent. The vertebral scutes are wide and cover about half of the costals. Unlike British specimens, suprapygal 2 is a wide hexagonal element that presents several morphotypes in Angeac-Charente. The plastron has a pair of mesoplastra and a wide anal notch posteriorly (Fig. 13B). The large entoplastron is triangular anteriorly and rounded posteriorly. The posterior plastral lobe is longer than the anterior one. The near-complete shell and the remaining pleurosternid material in general will be described in detail elsewhere. The pleurosternid from Angeac-Charente differs from Selenemys lusitanica in characters including: nuchal width less than twice its maximum length; anterior nuchal edge constituting the anterior carapace rim; posterior margin of the first peripheral two times shorter than the anterior one; overlap of the first vertebral on the first pair of peripherals; first vertebral wider than the nuchal; first pair of marginals wider than long; and development of an anal notch.

Pleurosternon bullockii is known by numerous specimens from the Berriasian Purbeck Group of Dorset (United Kingdom), including complete shells, carapaces, plastra, elements of the appendicular skeleton, and one skull (Evans & Kemp 1975; Milner 2004; Sterli *et al.* 2010; Evers *et al.* 2020), but it is also identified in the Tithonian record of Dorset and the French locality of Wimille (Boulogne-sur-Mer) (Guerrero & Pérez-García 2020).

Pleurosternids are freshwater turtles, of which most of the remains have been collected from fluvial and lacustrine sedimentary deposits (Joyce & Anquetin 2019). This hypothesis is confirmed by the depressed shell shape, appendicular bone morphology, and feeding specializations. *Pleurosternon bullockii* presents an elongate skull with low labial margins which suggests a gape and suction feeding (Pritchard 1984; Foth *et al.* 2017; Joyce & Anquetin 2019). The abundance of pleurosternid material and the presence of different growth stages suggest that the Angeac-Charente paleoecosystem probably corresponds to the preferred habitat of this turtle.



Fig. 13. — Pleurosternid turtle from the Berriasian of Angeac-Charente: **A**, carapace of *Pleurosternon bullockii* (Owen, 1842) (ANG14 R-454) in dorsal view; **B**, plastron and right peripheral plates 1 and 2 of *Pleurosternon bullockii* (ANG14 R-474 & ANG15 R-1082) in ventral view. Scale bar: 10 cm.

THALASSOCHELYDIA Anquetin, Püntener & Joyce, 2017 Family indet. Genus *Hylaeochelys* Lydekker, 1889

Hylaeochelys belli? (Mantell, 1844) (Fig. 12C-H)

DESCRIPTION

The third turtle taxon is represented by many isolated plates and one incomplete specimen discovered in 2018, consisting of parts of the plastron, some peripherals and both humeri (Fig. 12C-H). The shell bone surface is nearly smooth with shallow grooves, which clearly distinguishes this taxon from the two others previously described. The shell bones are relatively thin. The vertebral scutes are very wide and cover more than half of the costals laterally. The neurals are narrow and elongated, especially in the mid-posterior part of the carapace. The plastron bears a central plastral fontanelle. The mesoplastron is absent. The humerus shape is unremarkable and similar to that recently described in Late Jurassic thalassochelydians (e.g., Püntener et al. 2014, 2017). Based on the available characters, this form is possibly closely related to the Early Cretaceous (Purbeck and Wealden of the United Kingdom) thalassochelydian Hylaeochelys belli (Lydekker 1889; Hirayama *et al.* 2000; Pérez-García 2012; Pérez-García & Ortega 2014; Anquetin & André 2020).

Hylaeochelys belli is generally considered to be a freshwater turtle based on the sedimentary contexts of the localities in which it occurs (Milner *et al.* 2012; Pérez-García & Ortega 2014; Anquetin & André 2020). Most of the Angeac-Charente material represents juveniles or sub-adults, which may suggest that the swampy environment was a rookery for this turtle.

> LEPIDOSAUROMORPHA Benton, 1983 LEPIDOSAURIA Haeckel, 1866 RHYNCHOCEPHALIA Günther, 1867 SPHENODONTIA Williston, 1925 Family indet. Genus *Opisthias* Gilmore, 1909

> > cf. Opisthias (Fig. 14)

DESCRIPTION

ANG M-120 (Fig. 14) is a posterior dentary fragment bearing two preserved acrodont teeth in addition to seven broken teeth, there are six anteriorly and one posteriorly. Tooth crowns are



Fig. 14. — Rhynchocephalian right dentary fragment from Angeac-Charente: A-C, cf. *Opisthias* sp. (ANG M-120) in labial (A), lingual (B) and dorsal (C) views. Scale bar: 2.5 mm.

sub-pyramidal in shape, angulous anteriorly, more rounded posteriorly, and slightly inclined anteriorly throughout the dentition (Fig. 14A). As suggested by tooth bases, teeth are gradually increasing in size posteriorly. In occlusal view, the tooth row is straight (Fig. 14C). The coronoid process is broken at its base and the mandibular ramus is broken ventrally at the level of the Meckelian groove (Fig. 14B).

ANG M-120 is here tentatively referred to cf. Opisthias because of the tooth morphology and gradual heterodonty pattern which is similar to that observed in previously described Opisthias dentaries (Gilmore 1910; Evans & Fraser 1992). In lateral view, the dentary of Homoeosaurus shows wider tooth bases and crown apices that are less acute and not anteriorly inclined (Cocude-Michel 1963). Tingitana from the Tihonian-Berriasian of Morocco, has dentary teeth bearing a strong, inwardly concave, anterolingual crest between a large posterior cone and a small anterolingual one (Evans & Sigogneau-Russell 1997). It is worth noting that a sphenodontian from Cherves-de-Cognac, represented by a partial skeleton, was referred to Homoeosaurus by Buffetaut et al. 1989. A revision of this material, housed in a private collection, including a detailed comparison with the Angeac-Charente specimen would be useful to assess sphenodontian diversity in western Europe.

> SQUAMATA Oppel, 1811 SCINCOMORPHA Camp, 1923 Family PARAMACELLODIDAE Estes, 1983 Genus *Paramacellodus* Hoffstetter, 1967

> > Paramacellodus sp. (Fig. 15A-C)

Description

ANG M-20 (Fig. 15A-C) is a jaw fragment bearing two closely spaced teeth, with the anterior tooth slightly larger

than the posterior. Curvature of the teeth suggests that this fragment might come from a premaxilla or the anterior part of a dentary. Tooth implantation is pleurodont. Both tooth crowns are gently recurved and twisted so that the lingual face is slightly inclined posteriorly. The anterior marginal zone has a shoulder or angulus mesialis, following the terminology of Richter (1994). A similar shoulder, that would represent the *angulus distalis*, is absent in the distal marginal zone. The labial face is cambered and smooth whereas the lingual face is ornamented with longitudinal ridges, which are more numerous in the anterior tooth. The sharp and pointed tooth apex shows two small cusps (i.e., cuspis labialis and *cuspis lingualis*) closely adpressed and linked by a short carina (i.e., carina intercuspidalis). The ornamented central zone of the lingual face (i.e., *pars furcata*), delineated by long striae dominantes, is moderately wide. In lingual view, several foramina are present between the columnar roots of the teeth (Fig. 15A). The tooth crown shape (which is twisted with an acute apex and shows an angulus mesialis) and ornamentation combined with the presence of *cuspis labialis* and *cuspis* lingualis united by a short carina intercuspidalis is consistent with referral of ANG M-20 to Paramacellodus (Broschinski & Sigogneau-Russell 1996; Evans & Searle 2002). The other closely related genus Parasaurillus has more pointed crowns, no angulus mesialis, and longer lingual ridges (Evans & Searle 2002). Three species of *Paramacellodus* are known to date: P. oweni from the Late Jurassic of U.S.A and the Berriasian of England, P. marocensis from the Tithonian and/or Berriasian of Morocco, and P. sinuosus from the Barremian of Spain (Hoffstetter 1967; Richter 1994; Broschinski & Sigogneau-Russell 1996; Evans & Chure 1998; Evans & Searle 2002). A fourth species from the late Early Cretaceous (Aptian-Albian) of the United States, characterized by a welldeveloped cuspule on the anterior carina, was tentatively referred to Paramacellodus (cf. P. keebleri; Nydam & Cifelli 2002). Other occurrences tentatively referred to Paramacellodus have been reported from the Bathonian-Barremian of Europe, North America, Asia and Africa (Evans & Searle 2002). The Angeac-Charente form might belong to P. oweni or P. marocensis; however, pending the discovery of additional material, we refer it to Paramacellodus sp.

Numerous osteoderms (Fig. 15D-F) have been collected from Angeac-Charente. These elongated plate-like elements are subrectangular and show more or less numerous pits visible on their external surface. A low, longitudinal ridge can be present, either in the midline or displaced laterally. The unornamented imbrication shelf is variably developed. These isolated osteoderms are morphologically consistent with scincomorph osteoderms (e.g., Krause *et al.* 2003; Nydam *et al.* 2013), including those referred to paramacellodids (e.g., Hoffstetter 1967; Richter 1994; Broschinski & Sigogneau-Russell 1996; Nydam & Cifelli 2002; Sweetman & Evans 2011).

> ARCHOSAUROMORPHA von Huene, 1946 CHORISTODERA Cope, 1876



Fig. 15. — Scincomorph lizard remains from Angeac-Charente: A-C, fragment of left dentary of cf. *Paramacellodus* sp. (ANG M-20) in lingual (A), labial (B) and distal (C) views; D-F, osteoderms of Scincomorpha indet., ANG M-46 (D), ANG M-49 (E), ANG M-58 (F) in dorsal view. Scale bar: 500 µm.

Family CTENIOGENYIDAE Seiffert, 1975 Genus *Cteniogenys* Gilmore, 1928

> cf. *Cteniogenys* (Fig. 16A-E)

DESCRIPTION

A characteristic robust vertebra (Fig. 16A-C), as well as a smaller vertebra that probably belongs to a juvenile individual (Fig. 16D, E), have been discovered among the microremains. Both centra are amphicoelous, with an opened neurocentral suture. Dorsally, a distinct longitudinal ridge is laterally bordered by longitudinal grooves on the floor of the neural canal (Fig. 16A). In ventral view, the lower half of the cen-

trum is pinched into a keel (Fig. 16D). The margins of the cotyle are thick (Fig. 16E). All these characters are reminiscent of choristoderan reptiles (Evans 1991; Averianov *et al.* 2006; Vullo *et al.* 2014; Haddoumi *et al.* 2016; Lasseron *et al.* 2020). ANG M-20 is nearly identical to the vertebra of *Cteniogenys* figured by Evans & Milner (1994: fig.18.5). Hence, it is here tentatively assigned to this genus, although this identification needs to be substantiated with additional material. These freshwater or amphibious diapsids are well known in Laurasia (Matsumoto *et al.* 2013), but have also been reported from the Jurassic-Cretaceous transition of North Africa (Lasseron *et al.* 2020). In the Lower Cretaceous of Asia, the diversity and abundance of choristoderans is particularly high in the faunal assemblages, in which crocodyliforms are



Fig. 16. – Choristoder remains from Angeac-Charente: A-C, anterior dorsal vertebra of cf. *Cteniogenys* sp. (ANG M-20) in dorsal (A), ventral (B) and posterior (C) views; D, E, juvenile dorsal vertebral centrum (ANG M-47) in ventral (D) and oblique anterior (E) views. Scale bar: A-C, 1 mm; D, E, 200 µm.

scarce (Matsumoto *et al.* 2015; Skutschas & Vitenko 2017). Conversely, the abundance and diversity of crocodyliforms at Angeac-Charente may explain the scaricity of choristoderans.

> ARCHOSAURIA Cope, 1870 CROCODYLOMORPHA Hay, 1930 MESOEUCROCODYLIA Whetstone & Wybrow, 1983 NEOSUCHIA Benton and Clark, 1988 Family Atoposauridae Gervais, 1871 Genus *Theriosuchus* Owen, 1879

> > *Theriosuchus* sp. (Fig. 17A-D)

DESCRIPTION

Small isolated teeth are commonly recovered from residues obtained from bulk screening. Many of them (Fig. 17A-D) are leaf-shaped, labiolingually compressed, low-crowned and pseudoziphodont, a characteristic of the posterior dentition in the atoposaurid genus *Theriosuchus* (Schwarz & Salisbury 2005; Lauprasert *et al.* 2011; Martin *et al.* 2014b). Members of the genus *Theriosuchus* are small crocodyliforms with total body lengths less than 1m (Schwarz & Salisbury 2005). Besides teeth, two skull bones might belong to this genus: a right jugal (ANG 16-5179) and a fragmentary frontal (ANG 14-3692). The jugal is about 1 cm long and its ornamented lateral surface is thin and projects ventrally. The frontal preserves the interorbital area and displays an acute median keel on its dorsal surface. A finely sculpted osteoderm, longer than wide and possessing a shallow median keel (Fig. 19D), matches the morphology of dorsal osteoderms of the tail region in *Theriosuchus pusillus* (Owen 1879; Clark 1986).

COELOGNATHOSUCHIA

Martin, Lauprasert, Buffetaut, Liard & Suteethorn, 2014 NEOSUCHIA Benton & Clark, 1988 Family GONIOPHOLIDIDAE Cope, 1875 Genus *Goniopholis* Owen, 1841

> Goniopholis sp. (Fig. 18)

DESCRIPTION

In addition to fragmentary cranial remains and numerous isolated teeth, a nearly complete skull and mandibular elements



Fig. 17. — Crocodyliform teeth from Angeac-Charente: A-D, broadened teeth of *Theriosuchus* sp., ANG M-13 (A), ANG M-07 (B), ANG M-19 (C), ANG M-12 (D); E, tooth of Bernissartiidae indet. (ANG M-11). Scale bar: 1 mm.

(ANG1 8-5914, 5920, 5921 and 5925) of a single individual have been excavated and prepared in 2018 (Fig. 18). The skull is broken transversally in front of the orbits and the rostrum is split longitudinally along the nasals and premaxillae. The associated mandible preserves both dentaries. In comparison with other goniopholidids from Europe, this skull is notable for its exquisite preservation, being minimally compacted dorsoventrally.

Based on recent works (Salisbury et al. 1999; Schwarz 2002; Andrade et al. 2011; Buscalioni et al. 2013; Puértolas-Pascual et al. 2015; Martin et al. 2016a), several diagnostic characters used in combination permit assignment of this specimen to Goniopholididae, including: 1) a mesorostrine condition with a rostrum to skull ratio c. 0.65; 2) constricted outline of the upper jaw at the level of the premaxillary-maxillary suture; 3) double-caniniform maxillary teeth 4+5; 4) confluent dentary alveoli 3+4; 5) orbits smaller than supratemporal fenestrae and slightly smaller than frontal width at interorbital level; 6) skull table and dorsal surface of quadratojugal and jugal densely covered by large circular cupules; 7) internal choanae anteriorly bound by palatines; 8) semi-arched and widely expanded premaxillae with fifth alveolus more laterally placed than the rest of the premaxillary tooth row; and 9) absence of contact between ectopterygoid and posterior margin of maxillary tooth row.

The Angeac-Charente specimen (ANG18-5914, 5920, 5921 and 5925) also exhibits several characters that permit provisional assignment to the genus *Goniopholis*. Among these,

antorbital and interorbital ornamentation is informative. The presence of periorbital crests on the prefrontal and lacrimal are present in Goniopholis (Andrade & Hornung 2011) but absent in Anteophthalmosuchus and Hulkepholis (Salisbury & Naish 2011; Buscalioni et al. 2013; Martin et al. 2016a). The interorbital ridge is restricted to the frontal in Hulkepholis (Salisbury & Naish 2011; Buscalioni et al. 2013). In the Angeac-Charente specimen, the presence of both antorbital and interorbital crests is also a character shared with Goniopholis. A short postorbital spine is present in the Angeac-Charente specimen, a condition similar to other specimens of the genus Goniopholis (Salisbury et al. 1999; Schwarz 2002; Andrade et al. 2011) but unlike Anteophthalmosuchus in which this spine is longer, almost contacting the anterior margin of the orbits (Salisbury & Naish 2011; Martin et al. 2016a). The Angeac-Charente specimen is also similar in its maxillary tooth count (20) to Goniopholis kiplingi (Andrade et al. 2011) and with Anteophthalmosuchus, which possesses 19 alveoli (Martin et al. 2016a). The maxillary alveolar count has not been established for Goniopholis simus (Salisbury et al. 1999). However, this count is less than that of the goniopholidid Hulkepholis willetti, in which at least 24 maxillary alveoli are present (Salisbury & Naish 2011). Obvious differences in rostrum proportions are observed with Vectisuchus and Hulkepholis, which possess long and narrow rostra (Buffetaut & Hutt 1980; Salisbury & Naish 2011; Buscalioni et al. 2013). As in Anteophthalmosuchus and Goniopholis, the palpebral in the specimen from Angeac-Charente is small and not large and triangular as in



Fig. 18. — Goniopholidid crocodyliform Angeac-Charente: A, D, left dentary of *Goniopholis* sp. (ANG18-5925) in dorsal (A) and ventral (D) views; B, C, skull of *Goniopholis* sp. (ANG18-5914, ANG18-5920, ANG18-5921) in dorsal (B) and ventral (C) views. Scale bar: 10 cm.

Hulkepholis (Salisbury & Naish 2011; Buscalioni *et al.* 2013; Martin *et al.* 2016a). Comparison with *Nannosuchus* will not be discussed here as this taxon has been viewed either as a juvenile of *Goniopholis* (Salisbury 2002) or as a valid taxon (Andrade *et al.* 2011). Therefore, considering the characters mentioned above, we tentatively refer the Angeac-Charente specimens to *Goniopholis* sp.

A detailed description of the Angeac-Charente material will be provided elsewhere. In the meantime, a study of the numerous goniopholid specimens from Cherves-de-Cognac and assigned to *Goniopholis* (Raslan-Loubatié 2007) will provide a useful basis for discussing morphological variability within a goniopholidid population. Preliminary observations concerning the Angeac-Charente specimen do not reveal any obvious morphological differences between the goniopholidids at both localities but further study is required. Family PHOLIDOSAURIDAE von Zittel & Eastman, 1902 Genus *Pholidosaurus* Meyer, 1841

> Pholidosaurus sp. (Fig. 19E-J)

Description

The genus *Pholidosaurus* is attested to at Angeac-Charente by the presence of numerous isolated teeth (Fig. 19E-J). These are characteristic being slender, curved and bearing numerous apicobasal ridges that converge near the apex. Identical teeth are known from the Tithonian of Chassiron (Vullo *et al.* 2014), and in *Pholidosaurus purbeckensis* recently described from the nearby locality of Cherves-de-Cognac (Martin *et al.* 2016b). A right anterior portion of a dentary (ANG14-2959) displaying an extensive splenial symphysis can also be assigned to *Pholidosaurus*.



FIG. 19. — Crocodyliform remains from Angeac-Charente: **A**, **B**, tooth of Bernissartiidae indet. (ANG10-268) in lingual/labial (**A**) and occlusal (**B**) views; **C**, tooth of Bernissartiidae indet. (ANG10-76) in lingual/labial view; **D**, osteoderm of Atoposauridae? (ANG 10-167) in dorsal view; **E**, **F**, tooth of *Pholidosaurus* sp. (ANG11-960) in labial (**E**) and mesial/distal (**F**) views; **G**, **H**, tooth of *Pholidosaurus* sp. (ANG11-883) in labial (**G**) and mesial/distal (**H**) views **I**, **J**, tooth of *Pholidosaurus* sp. (ANG10-369) in labial (**I**) and mesial/distal (**J**) views. Scale bar represents: A-C, 2 mm; D-J, 8 mm.

Family BERNISSARTIIDAE Dollo, 1883

Bernissartiidae indet. (Figs 17E; 19A-C)

DESCRIPTION

Several mesiodistally elongate, bulbous and low-crowned isolated teeth have been recovered at Angeac-Charente (Figs 17E; 19A-C) and their morphology is reminiscent of the tribodont condition described for the small neosuchian *Bernissartia* (Buffetaut & Ford 1979; Martin *et al.* 2020). An incomplete left jugal (ANG15R-1205) is inflated on its lateral surface and might also belong to a bernissartiid. A small osteoderm (ANG16-4907) also resemble the dorsal row of osteoderms figured by Buffetaut (1975) in *Bernissartia fagesii*. Here, the specimen is slightly wider than long although it is not as rectangular as in *Bernissartia fagesii*. A double-keel runs on its dorsal surface and an anterolateral process is present. The recently described bernissartiid *Koumpiodontosuchus aprosdokiti* (Sweetman *et al.* 2015) also shares tribodont teeth with *Bernissartia fagesii* and, for this reason, we cannot yet ascertain a generic or specific attribution for the Angeac-Charente material.

PTEROSAURIA Kaup, 1834 Superfamily PTERODACTYLOIDEA Plieninger, 1901

Pterodactyloidea indet. A (Fig. 20A-D)

Description

This indeterminate pterodactyloid taxon is represented by small, labiolingually compressed teeth (Fig. 20A-D). The crown is relatively low, triangular and broad-based in labial and lingual views (Fig. 20A-C). The basal part of the labial and lingual faces is devoid of enamel, except along the mesial and distal carinae.



Fig. 20. — Pterosaur teeth from Angeac-Charente: **A**, Pterodactyloidea indet. A (ANG M-24) in labial view; **B**, Pterodactyloidea indet. A (ANG M-10) in lingual view; **C**, **D**, Pterodactyloidea indet. A, (ANG M-42) in lingual (**C**) and mesial (**D**) views; **E**, **F**, Pterodactyloidea indet. B, (ANG M-43) in lingual (**E**) and mesial (**F**) views; **G**, **H**, Pterodactyloidea indet. B, (ANG M-112) in lingual (**G**) and mesial (**H**) views; **I**, Pterodactyloidea indet. B, (ANG M-40) in lingual view; **J**, **K**, Pterosauria indet. (ANG M-41) in labial (**J**) and distal (**K**) views; **L**, **M**, Pterosauria indet. (ANG M-113) in lingual (**L**) and mesial (**M**) views. Scale bar: 500 µm.

The crown base is slightly inflated and the recurved apex is more or less slender, so that in labial view the carinae are convex and concave in the basal half and distal half of the crown, respectively. A weak enamel ornamentation consisting of faint vertical folds is present in some teeth. This taxon, characterized by a gradual heterodonty, corresponds to the morphotypes 5-7 reported from the Purbeckian beds of Chassiron (Vullo *et al.* 2014: fig. 17eg) and to the isolated tooth MPZ2011/46 described from the Valanginian-Hauterivian of Pochancalo 1 in northeastern Spain (Gasca *et al.* 2012: fig. 3k). Since this tooth morphology is present in some archaeopterodactyloids (e.g., *Pterodactylus, Germanodactylus*) and istiodactyliforms (e.g., *Haopterus, Longchengopterus, Mimodactylus*) (Lü *et al.* 2008; Wang *et al.* 2008; Kellner *et al.* 2019), we refer this taxon to Pterodactyloidea indet. A.

Pterodactyloidea indet. B (Fig. 20E-I)

DESCRIPTION

A second indeterminate pterodactyloid taxon is present at Angeac-Charente (Fig. 20E-I). The teeth mainly differ from those of taxon A by their slenderer crown and narrower base in labial view. In cross section, the lingual face is more convex than the labial face. The enamel distribution is similar to that observed in taxon A. The enamel surface shows some short, irregular vertical folds. This taxon also occurs in the Berriasian of Cherves-de-Cognac (Pouech 2008: fig. 38, pl. 23b).

Pterosauria indet. (Fig. 20J-M)

DESCRIPTION

A third morphological group of pterosaurian teeth corresponds to slender, recurved crowns that are characterized by a subcircular cross section and poorly distinct carinae (Fig. 20J-M). The enamel is smooth whereas the basal part of the crown devoid of enamel (i.e., basal portion of labial and lingual faces) shows an irregular rugose texture. Teeth of this morphology can be observed in some non-pterodactyloid taxa (e.g., the Late Jurassic rhamphorhynchid genus *Bellubrunnus*; Hone *et al.* 2012) and archaeopterodactyloids (e.g., ctenochasmatids; Perea *et al.* 2018); therefore, this third morphotype cannot be identified beyond Pterosauria indet.

DINOSAURIA Owen, 1842

ORNITHISCHIA Seeley, 1888

DESCRIPTION

With the exception of thyreophorans, the ornithischian dinosaurs of Angeac-Charente are mainly represented by isolated teeth. Most of these teeth show evidence of pre-burial transport and are either broken or eroded. However, the few remaining well-preserved teeth indicate a great diversity of ornithischians, represented by at least five families. Family HETERODONTOSAURIDAE Kuhn, 1966 Genus *Echinodon* Owen, 1861

Echinodon sp. (Fig. 21A-D)

Description Premaxillary teeth

Two well-preserved, isolated heterodontosaurid premaxillary teeth (ANG15-R672 & ANG14-3368) were recovered in Angeac-Charente (Fig. 21A-D). Only the crown, possibly from a shed tooth, is preserved in the first one (Fig. 21A, B), the other also shows a part of the root (Fig. 21C, D). Both crowns are very similar in being swollen and recurved folidont (Hendrickx et al. 2015a). They are rather short with a mesiodistal basal length of 6 mm and 7.5 mm and a preserved crown height of 8 and 9 mm, respectively. The teeth resemble those of Echinodon from the Purbeck Group of England described by Norman & Barrett (2002) and Sereno (2012). The crown is slightly concave lingually and gently convex labially, with an elliptical cross-section at mid-height. The main axis of the crown is recurved distally, so that its apex is slightly distal to the center of the crown base. The apex is blunt and bears a wear facet lingually. In lingual and labial views, the mesial border of the crown is convex with a bulge at its base, whereas the distal border is concave. As in Echinodon (Sereno 2012), and in contrast to many ornithischians, the carinae of the premaxillary teeth do not bear denticles (Galton 2009). The marked lingual wear facet, presumably from occlusion with the predentary bill, is apicobasally oriented (Sereno 2012). The crown enamel ends at the same level on every surface. The enamel has a relatively smooth texture, but exhibits small striations.

At the cervix, a slightly pronounced constriction separates the crown from the base of the root. The latter being incomplete in one specimen and lacking in the other, we cannot assess its length relatively to the length of the crown. The base of the root is large. The root labiolingual width is stable along the preserved section, practically equivalent to the crown mesiodistal basal length, and slightly inferior to the maximum width of the crown. The root is labiolingually narrower than mesiodistally wide. In lingual and labial views, it has subparallel and slightly convex borders. At the fracture point, the root has an oval cross-section with a large pulp cavity.

Family HYPSILOPHODONTIDAE Dollo, 1882

Hypsilophodontidae indet. (Fig. 21E, F)

DESCRIPTION

Premaxillary tooth

A well-preserved isolated hypsilophodontid crown (ANG M-119) only lacks its apex (Fig. 21E, F). In general shape, it is very similar to the aforementioned heterodontosaurid premaxillary teeth, but has a less bulbous morphology, bears denticles and lacks a wear facet on the lingual surface of the



Fig. 21. — Ornithischian teeth from Angeac-Charente: **A**, **B**, Heterodontosaurid premaxillary tooth of *Echinodon* sp. (ANG15-R671) in lingual (**A**) and labial (**B**) views; **C**, **D**, Heterodontosaurid premaxillary tooth of *Echinodon* sp. (ANG14-3368) in lingual (**C**) and labial (**D**) views; **E**, **F**, hypsilophodontid premaxillary tooth (ANG M-119) in lingual (**E**) and labial (**F**) views; **G**, **H**, hypsilophodontid dentary tooth (ANG15-R927) in labial (**G**) and lingual (**H**) views; **I-L**, hypsilophodontid maxillary tooth (ANG 10-153) in lingual (**J**), labial (**K**) and mesial (**L**) views. Scale bars: A-D, I-L, 5 mm; E-H, 3 mm.

crown. Based on its shape, it is identified as a premaxillary tooth, the maxillary and dentary teeth being more quadrangular in shape (Norman et al. 2004: fig. 18.3). The crown is recurved folidont (Hendrickx et al. 2015a) and short, with a mesiodistal basal length of 3 mm and a preserved crown height of 4 mm. It appears quite similar to the premaxillary teeth of *H. foxii* figured by Galton (2009: fig. 2G, L-P). The crown is compressed labiolingually and has a slightly convex surface on both sides. At mid-height, the cross-section is elliptical. The main axis of the crown is recurved, so that in lingual and labial views, the apex appears distal to the midpoint of the base of the crown. Both the mesial and distal margins are mesially and distally expanded respectively at their base. However, the mesial margin is convex, whereas the distal margin is concave for most of its length. The carinae extend from the apex along two-thirds of the crown and bear a series of fine, bulbous denticles. The mesial carina is damaged, but all the denticles are visible on the distal border. The density of the denticles is approximately 6 per mm. The crown surfaces have a relatively smooth texture. The crown enamel does not extend further basally on one particular side as for dentary teeth (see below). There is a visible constriction forming a cervix, at the base of the root.

Maxillary tooth

One well-preserved and complete isolated hypsilophodontid maxillary tooth (ANG10-153) (Fig. 21I-L) has been recovered from Angeac-Charente. It is 27 mm long with the root being twice as long as the crown. The crown is sub-quadrangular and short, with a mesiodistal basal length of 6 mm and a crown height of 10 mm. It looks similar to the maxillary crown of H. foxii (Galton 1974: fig. 14a, b). The crown is expanded mesiodistally, compressed labiolingually, and has a sub-oval cross section at mid height. The lingual surface is relatively flat, whereas the labial surface bears a longitudinal concavity in the central area of the crown and is convex mesiodistally. In lingual and labial views, the main axis of the crown is straight, but the apex is displaced distally relatively to the midpoint of the crown base. The rounded apex is however not distally bent, but strictly ventrally oriented. The mesial and distal margins are subparallel, even if the mesial margin is slightly convex, whereas the distal one is straight with a small angle at mid-height. The carinae are located on the upper portions of the mesial and distal edges, and on the apical border of the crown. The tooth carinae are covered by small denticles throughout their length, in contrast to the H. foxii tooth figured by Galton (1974: fig. 14a, b), in which the apical border is worn and, therefore lacks denticles. The mesial carina is damaged, but probably exhibited denticles. The distal and apical carinae exhibit large and regular denticles, at a frequency of approximately 1 per mm. They are rounded in shape, apicobasally oriented and their size slightly increases towards the apex. The denticles on the apical margin are smoothly worn and, consequently, an extended single and oblique wear facet is visible on the dorsal part of the lingual surface. There is no wear on the labial surface, but small ridges extend ventrally to denticles, towards the crown base.

The longest visible is almost 6 mm long and is directly mesial to the apex. The enamel texture is relatively smooth, except towards the base of the crown where it is more irregular. The crown enamel is clearly more basally extended on the labial side than on the lingual side of the tooth. Consequently, the base of the crown appears swollen on the labial surface and forms an incipient cingulum, as in most basal euornithopods (Norman *et al.* 2004).

At the level of the cervix, there is a constriction in labial and lingual views. It is not particularly pronounced, in which respect it differs from the condition observed in *H. foxii* (Galton 1974: fig. 14a, b). The root is long, tubular and narrower than the crown. Its mesiodistal width decreases towards the apex, whereas its labiolingual thickness is more or less equivalent on most of the root length and only decreases slightly towards the apex.

Dentary tooth

An isolated hypsilophodontid dentary crown, only lacking the apex, has also been recovered at Angeac-Charente, (ANG R-927; Fig. 21G-H). The crown is straight folidont (Hendrickx et al. 2015a) and short, with a mesiodistal basal length of 4 mm and a preserved crown height of 5 mm. At mid-length, it has a semicircular cross-section. In lingual and labial views it is diamond-shaped, and the proportions and shape are quite similar to the dentary teeth of *H. foxii* (Galton 1974: fig. 15; 2009: fig. 3). The crown is compressed labiolingually, with a convex labial surface and a slightly concave lingual surface. The main axis of the crown is straight, and the apex was most likely directed strictly dorsally. The mesial and distal borders are strongly convex, forming an angle of approximately 100° at mid-height of the crown. The carinae extend from there towards the apex. They bear a series of large denticles. The denticles are regular and semicircular, and have an apicobasal orientation. Ventral to each denticle, a blunt ridge extends towards the crown base on the lingual surface. It seems that the same ridges are also present on the labial surface, but this cannot be assessed with certainty because the tooth is eroded. On the lingual surface, the longest and most pronounced ridge is positioned below the apex. On the labial surface, three subvertical and subparallel ridges are visible on the central area of the crown. The enamel texture is irregular. The crown enamel extends further basally on the labial side than on the lingual side. A marked constriction is visible at the base of the crown, as in the dentary teeth of *H*. foxii (Norman et al. 2004: fig. 18.3E).

> IGUANODONTIA Dollo, 1888 Family Camptosauridae Marsh, 1885

> > Camptosauridae indet. (Fig. 22)

Description

Several diagnostic features allow the referral of the material described below to Camptosauridae, including: 1) lozenge-



FIG. 22. — Camptosaur remains from Angeac-Charente: A, dentary tooth of Camptosauridae indet. (ANG11-1120) in lingual view; B-E, left femur of Camptosauridae indet. (ANG14-R563) in anterior (B), medial (C), posterior (D) and lateral (E) views. Scale bar: A-D, 5 cm; E, 1 cm.

shaped teeth with prominent vertical keel more strongly developed on maxillary teeth than on dentary teeth, several secondary vertical ridges on sculptured surface of crown with anterior and posterior ridges bordering the non-denticulate half of maxillary tooth crown, posterior ridge on equivalent part of dentary teeth (Galton & Powell 1980); and 2) femur robust, with a dorsally arched shaft, a midshaft positioned and pendant fourth trochanter, and with a dorsally open, troughlike anterior intercondylar groove (Norman & Barrett 2002).

Dentary tooth

One extremely well-preserved, isolated camptosaurid tooth (ANG11-1120) exhibits an almost complete crown and a root broken at approximately mid-length (Fig. 22E). The crown is straight folidont (Hendrickx et al. 2015a), moderately elongated, and relatively robust. The mesiodistal basal length is 7 mm and the preserved crown height is 17 mm. It appears very similar to dentary teeth from the Kimmeridgian of England described by Galton & Powell (1980) and to the Owenodon sp. material figured by Galton (2009). The crown is labiolingually compressed, flat lingually, and strongly convex labially, with a D-shaped cross-section at mid-height. In lingual view, the main axis of the crown is straight, with a crown apex that was most likely pointing strictly dorsally. Both the mesial and the distal margins are symmetrical, with a marked angle at mid-height, conferring to the crown a diamond-shaped aspect in lingual and labial views. On the distal border, the lower edge of the crown is flexed mesiolingually, forming a pseudo-cingulid, as seen in Camptosaurus prestwichii (Norman & Barrett 2002: fig. 5) and Camptosaurus medius (Galton 2009: fig. 10D). The carinae extend from the apex to half of the length of the crown, until the mesial and distal angulations. They appear shorter than in some *Camptosaurus* specimens, in which the carinae extend along almost two-thirds of the edges (Norman & Barrett 2002: fig. 5A; Galton 2009: fig. 10D). Both carinae are straight and oblique. Denticles are visible on the entire length of the preserved carinae. The mesial carina is damaged, but it apparently exhibits the same density of denticles as the distal carina with 3 denticles per mm. The denticles are fine, slightly mesially and distally hooked, and apicobasally oriented. They present a marked increase in size, the largest being the most apical. Ventral to the denticles, small ridges extend from their base towards the crown base. Most are approximately 1 mm in length but the longest is almost 5 mm in length and extends from the third well-developed denticle. The longitudinally concave lingual crown surface bears two strongly developed central and apicobasally oriented ridges separated by a flute. In contrast to the condition observed in some dentary teeth of the holotype of C. prestwichii (Norman & Barrett 2002: fig. 5A; Galton 2009: fig. 9K) and C. medius (Galton 2009: fig. 10D), but as seen in Owenodon sp. (Galton 2009: fig. 18J), the ridges are parallel throughout their length. The crown enamel extends more basally in the mesiolingual part

of the crown than in the distolingual part. The enamel has a smooth texture and its surface bears transverse undulations contacting both carinae. They are more marked on the basal half of the crown.

A pronounced constriction occurs at the base of the crown forming a cervix. Breakage of the root renders it impossible to assess whether or not it was longer than the crown. It is, however, clearly narrower. The width of the root is the same throughout the preserved portion, and it is roughly equivalent to the mesiodistal basal length measured at the level of the cervix. The root is labiolingually narrow and exhibits subparallel mesial and distal borders. On its lingual surface, a shallow concavity is visible. At the fracture point, it has a suboval cross section.

Femur

One subcomplete left femur is part of the material from Angeac-Charente referred to Camptosauridae. It is well-preserved, but both ends are crushed and eroded, and a small part of the distal end is missing (Fig. 22B-E). The preserved length of the femur is 265 mm. The diaphysis is relatively stout. In anterior and posterior views, the femur is straight (Fig. 22B, D). In lateral and medial views, the femur is curved with convex anterior and concave posterior margins.

The femoral head is positioned in the same plane relative to the transverse axis of the distal condyles. The femoral head is discrete, unlike that of *C. dispar* (Galton & Powell 1980: fig.2F) and oval in anterior and posterior views. In those views, it is inclined at approximately 40° with respect to the main axis of the bone. The neck is large and it merges obliquely with the femoral shaft. In some *Camptosaurus* specimens, the neck looks more constricted (Norman & Barrett 2002: fig. 6; Galton 2009: fig. 5T).

The lesser trochanter is located on the anterior surface of the femur, on the proximal extremity. It is eroded, but it still appears as a strong and high protrusion extending proximodistally in the anterolateral angle of the femur. Its proximal extremity is lower than the proximal margin of the femoral head, but it still appears more strongly developed than in most described *Camptosaurus* species (Galton & Powell 1980; Norman & Barrett 2002; Carpenter & Wilson 2008; Galton 2009). There is no deep cleft visible between the lesser trochanter and the greater trochanter, in contrast to the condition observed in *C. aphanoecetes* (Carpenter & Wilson 2008). The greater trochanter is difficult to discern, probably as a result of erosion.

The femur diaphysis has a quadrangular cross section. It is as robust as in other camptosaurids. Under the fourth trochanter, the diaphysis is slightly narrower transversely than anteroposteriorly. On the posterior surface of the diaphysis, the pendent fourth trochanter is located on the medial edge of the bone, just above midshaft (Fig. 22C-E). It is strongly developed, as in most species of *Camptosaurus* (Carpenter & Wilson 2008; Galton 2009), and it is blade-shaped in lateral view (Fig. 22E). In posterior view, the base of the trochanter is straight (Fig. 22D). The proximal margin of the trochanter is elongated and gradually rises from the diaphysis, at an angle of approximately 115° relatively to the main axis of the shaft. The distal margin of the fourth trochanter is shorter and steeper.

The distal extremity of the femur is incomplete. A longitudinal bulge is visible above the distal condyles, but the condyles themselves are not preserved. On the anterior surface of the femur, a very shallow intercondylar groove is present. On the posterior surface, the popliteal fossa is visible.

> THYREOPHORA Nopcsa, 1915 ANKYLOSAURIA Osborn, 1923

> > Ankylosauria indet. (Fig. 23A-C)

DESCRIPTION

The ankylosaurs are only represented in Angeac-Charente by a single tooth (ANG15-3980) and an osteoderm (ANG18-6585). The tooth (Fig. 23A, B) was collected in 2015 from the unit 3 of the CG3 plot. The osteoderm (Fig. 23C) was found in 2018, in the CG9 plot, at the base of the bonebed (unit 2 of Rozada *et al.* 2021).

ANG15-3980 consists of a nearly complete, but worn tooth crown, with only the basal part of the root attached. The crown height is 9 mm, and the maximum width at the crown base is 8.5 mm. The crown is folidont, labiolingually compressed, and slightly recurved posteriorly. The labial and lingual crown surfaces are smooth and swollen around the base. The base of the crown is raised on the labial side (Fig. 23B), while there is a distinct cingulum at the base of the lingual side (Fig. 23A). A broad primary ridge extends vertically from the swollen base on both sides of the crown to form the apex of the tooth. The mesial and distal carinae bear 6 denticles, but there is no evidence of fluting as in stegosaurs and many Late Cretaceous ankylosaurs. There is a slight constriction of the root just below the crown. ANG15-3980 is nearly identical to NHMUK R2940, which is an isolated tooth from the Purbeck Group of Lulworth Cove, Dorset, referred to an ankylosaur (Galton 1980, 1983). It is also similar to ankylosaur teeth reported from the Valanginian and Barremian of southern England (Blows & Honeysett 2014).

Based on its shape and its size, ANG18-6585 is clearly distinct from other osteoderms found in Angeac-Charente that belong either to turtles, crocodyliformes or scincomorphs (see above). It has a pentagonal shape, with a maximum length of *c*. 10 cm and a thickness of *c*. 15 mm. ANG18-6585 possesses a marked keel on its exterior surface (Fig. 23C), while its base is nearly flat. The exterior surface of the osteoderm is marked with numerous grooves that give it a spongy texture. Such a morphology and structure have been reported in nodosaurid ankylosaurs (Blows 2001), and is very similar to the large ossicles observed on the sacral shield of *Polacanthus* from the Barremian of the Isle of Wight (Hulke 1887; Blows 1987, Pereda-Suberbiola 1994). Although the material is very limited, the discovery of ankylosaur remains at Angeac-Charente is significant because these animals are very rare in the European fossil record in Purbeckian facies. Except for the Lulworth tooth (Galton 1983) and a cervico-pectoral lateral spine associated with a distal humerus from the Early Valanginian of Gronau in Germany (Sachs & Hornung 2013), there are no other known ankylosaur remains in Europe around the Jurassic/ Cretaceous boundary.

> STEGOSAURIA Marsh, 1877 Family Stegosauridae Marsh, 1880 Subfamily Dacentrurinae Mateus, Maidment & Christiansen, 2009 Genus *Dacentrurus* Lucas, 1902

> > Dacentrurus sp. (Fig. 23D-J)

DESCRIPTION

Stegosaurs are represented at Angeac-Charente by 84 identified skeletal elements, most of which are vertebrae and ribs. Other remains include a few appendicular (phalanges) and skull (squamosal and braincase elements) bones, and one tooth (Fig. 23D-J). Except for the tooth, which was collected following washing and screening, all the other elements are from the same area (Fig. 3[CG4 to CG7 plots]). They very likely belong to the same individual because none of the identified bones represent the same skeletal element. Moreover, the relative proportion of the bones is also consistent with their belonging to a single individual, as suggested by the reconstructed vertebral series (Fig. 23F-J).

The single collected dentary tooth crown (Fig. 23D, E) has the same proportions as the stegosaur tooth from Chervesde-Cognac described by Billon-Bruyat et al. (2010). It has a preserved height of 4.2 mm and a maximum mesiodistal width basally of 4 mm. The root is broken at its base. The presence of a sharp demarcation, either swelling or cingulum, almost at the same level on each crown surface together with the mesiodistal symmetry of the tooth lead us to refer this tooth to a stegosaurian (Billon-Bruyat et al. 2010). Nevertheless, the morphology of the crown is quite distinct from that of other known stegosaur teeth. As in Huayangosaurus, the base of the crown is sharply differentiated from the root by a prominent swelling, but there is no true ring-shaped cingulum as in other stegosaurs (Sereno & Dong 1992). In labial view, a single rounded median protuberance is present, but there are no well-defined ridges. A broad, 45° angled wear surface is present above the bulbous crown base on the labial side and has truncated a part of the dentary crown, so that the denticles are no longer visible (Fig. 23D). Vertical coarse ridges, semicircular in cross section and divided by narrow grooves are visible in lingual view.

A dozen cervical vertebrae, including the axis as well as four dorsal and two caudal vertebrae have been collected (Fig. 23F-J). The neural arches are often broken, probably due to trampling (Rozada et al. 2021), and numerous neural spines or transverse processes are found isolated. The axial skeleton shows some features permitting clarification of the phylogenetic position of the Angeac-Charente stegosaur. The transverse processes of dorsal vertebrae project at a high angle to the horizontal as in all Thyreophoroidea (Fig. 23I), and the prezygapophyses are fused on some dorsal vertebrae as in most Eurypoda. The dorsal centra are wider than long (ANG 15-3937, centrum witdh 126 mm; centrum length 95 mm). This is a feature diagnostic of *Dacentrurus* according to Galton (1985) and Maidment et al. (2008), and of Dacentrurinae (Dacentrurus + Miragaia) (Mateus et al. 2009; Costa & Mateus 2019). Cervical ribs are fused to para- and diapophyses in at least two cervical vertebrae. This feature is also diagnostic of Dacentrurinae (Mateus et al. 2009; Costa & Mateus 2019).

A taxonomic revision of the Late Jurassic to Early Cretaceous European stegosaurs is beyond the scope of this paper. However, if we acknowledge the existence of several species of stegosaurs closely related to Dacentrurus (Costa & Mateus 2019), we concur with Cobos et al. (2010) that, given the available diagnostic characters, and despite the discovery of a new specimen of Miragaia (Costa & Mateus 2019), Dacentrurus and Miragaia cannot be clearly differentiated at a generic level, at this time. Moreover, we do not understand the taxonomical logic of creating, on the one hand the new European genus, Miragaia, and the new sub-family Dacentrurinae, while the original diagnostic characters of *Dacentrurus*, such as dorsal vertebral centra that are wider transversely than they are long anteroposteriorly (see Galton 1985; Maidment et al. 2008), largely encompass Miragaia, and on the other hand, synonymizing a third American genus Alcovasaurus with Miragaia (Costa & Mateus 2019). Pending the complete description and revision of the Late Jurassic to Early Cretaceous European stegosaur material and a phylogenetic analysis of the specimens concerned, we still prefer to synonymize Miragaia with Dacentrurus (Cobos et al. 2010), which is not incompatible with the results of phylogenetic analyses (Raven & Maidment 2017). Thus, based on Costa & Mateus (2019), the Angeac-Charente stegosaur may be more closely related to Dacentrurus *longicollum* (comb. nov.), because the cervical transverse processes of a posterior cervical vertebra project ventral to the ventral margin of the prezygapophyses. Nevertheless, it is very difficult to assess the condition of this feature in *D. armatus*, given that only one half of a neural arch is preserved in the type species (Owen 1875; Galton 1985). In any case, the Angeac-Charente taxon is closely related to the English, Spanish and Portuguese stegosaurs, and it is the youngest occurrence of *Dacentrurus*. In addition to bony remains, about 100 stegosaur natural track casts in a range of sizes have been identified and documented on a single bedding plane in the CG3 plot, and assign to the ichnotaxon Deltapodus (Rozada 2019; Rozada et al. 2021).



Fig. 23. — Thyreophoran remains from Angeac-Charente: **A**, **B**, ankylosaur maxillary tooth (ANG15-3980) in lingual (**A**) and labial (**B**) views; **C**, ankylosaur osteoderm (ANG18-6585) in dorsal view; **D**, **E**, dentary tooth of *Dacentrurus* sp. (ANG M-14) in labial (**D**) and lingual (**E**) views; **F**, axis of *Dacentrurus* sp. (ANG18-6203) in ventral view; **G**, anterior cervical vertebra of *Dacentrurus* sp. (ANG12-1878) in ventral view; **H**, reconstructed cervical series of *Dacentrurus* sp. (ANG16-6748, ANG16-4660, ANG12-1749, ANG14-3202, ANG14-2912, ANG14-3094) in ventral view; **I**, **J**, dorsal vertebra of *Dacentrurus* sp. (ANG18-6548) in anterior (**I**) and right lateral (**J**) views. Scale bars: A, B, 5 mm; C, D, 2.5 mm; E, 2.5 cm; F-J, 5 cm.


Fig. 24. — Sauropod teeth from Angeac-Charente: A-D, posterior left maxillary tooth of Turiasauria indet. (ANG11-837) in labial (A), mesial (B), lingual (C) and distal (D) views; E-H, left maxillary tooth of Turiasauria indet. (ANG13-2330) in labial (I), mesial (J), lingual (K) and distal (L) views; M-P, left premaxillary tooth of Turiasauria indet. (ANG13-2330) in labial (I), mesial (J), lingual (K) and distal (L) views; M-P, left premaxillary tooth of Turiasauria indet. (ANG14-R289) in labial (M), mesial (N), lingual (O) and distal (P) views; Q-T, dentary tooth of Turiasauria indet. (ANG14-3495) in labial (Q), mesial (R), lingual (S) and distal (T) views; U-X, left dentary tooth of Turiasauria indet. (ANG14- R435) in labial (U), mesial (V), lingual (W) and distal (X) views; Y-AA, maxillary tooth of Macronaria indet. (ANG R-1732) in labial (Y), distal (Z) and lingual (AA) views. Scale bar: A-X, 2 cm; Y-AA, 1 cm.



Fig. 25. — Sauropod embryonic or hatchling teeth from Angeac-Charente: A, tooth of Macronaria indet. (ANGM-18) in lingual view; B, C, tooth of Turiasauria indet. (ANGM-118) in labial (B) and lingual (C) views. Scale bar: 500 µm.

SAURISCHIA Seeley, 1888 SAUROPODA Marsh, 1878 EUSAUROPODA Upchurch, 1995 TURIASAURIA Royo-Torres, Cobos & Alcalá, 2006

> Turiasauria indet. (Figs 24A-X; 25B, C; 26; 27)

DESCRIPTION

Sauropod remains are especially abundant in the Early Cretaceous of Angeac-Charente. The locality has yielded many teeth (N = 146), bones (N = 784), and track casts of this group of dinosaurs (Néraudeau et al. 2012; Rozada et al. 2021). All parts of the skeleton are represented including the braincase, some skull bones, teeth, cervical, dorsal and caudal vertebrae, chevrons, pelvic girdle and all the limb bones (Figs 24-27). Based on the number of femurs and their size, as well as the teeth, there are at least seven different individuals preserved in the site. With the exception of two teeth (see below), all this material belongs to a single taxon. All remaining teeth are reminiscent of the Turiasauria clade (Allain et al. 2013, 2017). We can classify them based on a small number of diagnostic characters. Teeth are heart-shaped in labial and lingual views, with an asymmetric shape induced by a concave distal margin towards the apex (Royo-Torres et al. 2006, 2017; Royo-Torres & Upchurch 2012; Mocho et al. 2016). This feature has been observed in most of the sauropod teeth that have been collected from Angeac-Charente (Figs 24; 25B, C). A second character permits referral of these teeth to Turiasauria. When the root is well preserved, several long longitudinal grooves are visible in Turiasaurus, Losillasaurus (Royo-Torres et al. 2021) and Moabosaurus (Britt et al. 2017 and RRT personal observation). These grooves are also present in the Angeac-Charente taxon (Fig. 24I-P, U-X) and may be diagnostic for Turiasauria (Royo-Torres et al. 2021). Moreover, the teeth of Angeac-Charente show a range of crown morphotypes and this variability of forms has also been described in turiasaur teeth from Portugal (Mocho *et al.* 2016) and in *Mierasaurus* (Royo-Torres *et al.* 2017) and *Losillasaurus* (Royo-Torres *et al.* 2021). Teeth, in private collections, identical in every way to those of Angeac-Charente, are also present in the Berriasian of Cherves-de-Cognac (RA, TL pers. obs.).

The caudal vertebrae are also useful in determining the systematic position of the Angeac-Charente sauropod (Fig. 26). The anterior caudal vertebrae are procoelous with a slightly convex posterior articulation (Fig. 26A-I) whereas the middle become amphicoelous or amphyplatyan (Fig. 26J-L). The presence of a convex posterior articulation on sauropod caudal vertebrae was acquired several times during sauropod evolution (Wilson 2002; Upchurch et al. 2004; D'Emic 2012; Mannion et al. 2017, 2019) and can be seen in diplodocids, titanosaurs and mamenchisaurids. The procoelous condition was also acquired in Turiasauria, as described for the Late Jurassic Turiasaurus and Losillasaurus (Casanovas et al. 2001; Royo-Torres et al. 2006, 2021). It has also been reported in the posterior series of Early Cretaceous Mierasaurus and Moabosaurus (Royo-Torres et al. 2017; Britt et al. 2017). This feature is considered to be synapomorphic for Turiasauria in some phylogenetic analyses (Carballido & Sander 2014). The neural arch of anterior caudal vertebrae is restricted to the anterior half of the centrum. This character is shared with Turiasaurus, Losillasaurus, Moabosaurus, Mierasaurus, Cetiosaurus and the Titanosauriformes (Upchurch et al. 2004; D'Emic 2012; Britt et al. 2017; Royo-Torres et al. 2017). The presence in the Angeac-Charente taxon of caudal vertebrae with short lateral processes ('caudal ribs') that do not extend beyond the posterior end of the centrum suggests affinities with Titanosauriformes (Mannion et al. 2019; Royo-Torres et al. 2021).

Two additional possible synapomorphic characters for Turiasauria seen in specimens from Angeac-Charente include slightly opisthocoelous posterior dorsal centra, as well as a high neural arch below the postzygapophyses of the posterior dorsal vertebrae (Carballido & Sander 2014). MACRONARIA Wilson & Sereno, 1998

Macronaria indet. (Figs 24Y-AA; 25A)

DESCRIPTION

In addition to the turiasaur, a second sauropod taxon may be present at Angeac-Charente site. It is only represented by a single abraded tooth and a tooth recovered from microremains (Figs 24Y-AA; 25A). They are spatulate and characterized by straight and subparallel distal and mesial edges at the base of the crown, and by the presence of a convex labial and concave lingual surface. Based on these features, these teeth are assigned to a macronarian sauropod probably close to *Camarasaurus* (Wilson 2002; Upchurch *et al.* 2004; Mocho *et al.* 2017).

Sauropod track casts have also been recorded at Angeac-Charente. Thay are represented by casts of pes and manus footprints (Rozada *et al.* 2021). In 2018, a sauropod footprint cast was observed above and in contact with an *in-situ* broken sauropod radius. It represents a spectacular "instantaneous" preservation of the action of a sauropod pes or manus crushing a sauropod long bone, and inducing bone modifications (breakage, displacement and reorientation) and sediment deformations (Rozada *et al.* 2021). The footprints are identified as Sauropoda indet. because of the general circular morphology of the pes, the characteristic tubular metacarpal arrangement of the manus and also the huge size of the prints (Carrano & Wilson 2001; Wilson 2005).

> THEROPODA Marsh, 1881 TETANURAE Gauthier, 1986 Family MEGALOSAURIDAE? Fitzinger, 1843

> > Megalosauridae? indet. (Fig. 28K-N)

DESCRIPTION

All the large (> 3 cm) blade-like theropod teeth found at Angeac-Charente are tentatively referred to a single taxon, although two tooth morphotypes are present. The first morphotype probably corresponds to mesial teeth (Fig. 28K-M), the crown of which are slender and more elongated than in lateral teeth (Fig. 28N). The crown height ratio (Hendrickx et al. 2015a) ranges between 1.91 (ANG M-121) for a lateral tooth and 2.43 for a mesial one (ANG 17-5650). Apart from that, the teeth have the same characters. They are strongly compressed labiolingually. The mesial margin is convex and the distal margin is only slightly concave. Both the distal and mesial carinae are denticulate, but the latter only occurs on the apical half to one third of the crown. The carinae are centrally positioned on both the mesial and distal margins of crowns. There are 12 (ANG 17-5650) to 18 (ANG M-121) denticles per 5 mm along the mesial carinae, and 12 to 15 along the distal carinae at two thirds of the crown height. The denticles are longer mesiodistally than they are basoapically high, and they have a horizontal subrectangular outline. Interdental sulci are present. The enamel is transversely wrinkled and both transverse and marginal undulations are present (Hendrickx *et al.* 2015a).

Teeth of large carnivorous dinosaurs, morphologically very close to those of Angeac-Charente, have been reported from the Tithonian of Chassiron and from the Purbeckian of England. The first have been referred with caution to Megalosauridae (Vullo *et al.* 2014), while the latter have been alternately referred to Megalosauridae or Allosauroidea (see Milner 2002). All dental characters described above have been recognized in Megalosauridae (Hendrickx *et al.* 2015b), and we tentatively refer the large dinosaur teeth of Angeac-Charente to this group, but an assignement to another basal tetanuran clade cannot be definitively excluded.

COELUROSAURIA Huene, 1914 ORNITHOMIMOSAURIA Barsbold, 1976

Ornithomimosauria indet.

Description

Ornithomimosaurs are by far the most commonly represented vertebrates in Angeac-Charente, with more than 3800 macroremains collected (Figs 29-30), accounting for more than 50% of the identified vertebrate material (Rozada et al. 2021). The minimum number of individuals (MNI) is approximately 70 based on the distal end of left tibiae. Ornithomimosaur remains are mainly concentrated in the CG1 and CG3 loci, in which they represent 70 % of all the ornithomimosaur remains identified. Such a concentration and high number of individuals are congruent with a mass mortality event of an ornithomimosaur herd (Allain et al. 2011, 2014; Néraudeau et al. 2012; Rozada et al. 2021). However, no articulated skeletons have been observed due to the intense trampling (dinoturbation) affecting this area (Rozada et al. 2021). The only articulated remains of ornithomimosaurs found so far come from the northwestern part of the quarry (CG9 plot) and they include the zeugopod and the autopod of the forelimb of a single individual, as well as the the zeugopod and autopod of the hindlimb of another single individual.

Except for the most fragile elements such as the maxillary and palate bones, which have probably suffered from trampling and have not yet been identified, the skeleton of the Angeac-Charente ornithomimosaur is virtually complete (Fig. 31). A complete description of the entire skeleton of this new taxon is beyond the scope of this study. Nevertheless, it seems important to highlight here key anatomical features of the Angeac-Charente ornithomimosaur: first, because this clade was hitherto unknown in Europe at the beginning of the Cretaceous (Allain et al. 2014); secondly because it may be the oldest known ornithomimosaur to date (Choiniere et al. 2012; Cerroni et al. 2019); thirdly, because it shows very close anatomical similarities to Limusaurus, which is a Late Jurassic Chinese theropod that is not considered a member of the Ornithomimosauria, but a ceratosaurian (Xu et al. 2009). These similarities include a very large external mandibular fenestra and short forelimbs with



Fig. 26. — Sauropod caudal vertebrae from Angeac-Charente: A-C, anterior caudal vertebra of Turiasauria indet. (ANG15-R679) in left lateral (A), posterior (B) and dorsal (C) views; D-F, anterior caudal vertebra of Turiasauria indet. (ANG15-R698) in right lateral (D), posterior (E) and dorsal (F) views; G-I, anterior caudal vertebra of Turiasauria indet. (ANG15-R692) in right lateral (G), posterior (H) and dorsal (I) views; J-L, middle caudal vertebra of Turiasauria indet. (ANG15-R652) in right lateral (J), posterior (K) and dorsal (L) views. Scale bar: 10 cm.



FIG. 27. — Femur of Turiasauria indet. (ANG19-7000) from the Berriasian of Angeac-Charente discovered during the 2019 campaign. The length of the femur is 2 m. © L. Bocat.

manual digit reduction (RA pers. obs.). Relationships between ceratosaurians and ornithomimosaurs have long been confusing (e.g. Marsh 1895; Janensch 1925; Galton 1982; Holtz 1994; Rauhut 2003). Some taxa, including *Elaphrosaurus*, *Deltadromeus*, *Limusaurus*, *Nqwebasaurus* and probably the Angeac-Charente taxon do not have a clearly established phylogenetic position, and their anatomy may also reflect unexpected and unrecognized relationships between ceratosaurians and ornithomimosaurs. Pending a comparative and detailed phylogenetic study, we provide herein some anatomical features that clearly indicate the ornithomimosaurian affinity of Angeac-Charente material.

Besides the features already mentioned by Allain et al. (2014), we mainly used the anatomical characters discussed in the recent reappraisal of the phylogenetic position of Afromimus by Cerroni et al. (2019). The edentulous and downturned dentary (Fig. 29A) is an ornithomimosaurian synapomorphy convergently acquired by numerous other coelurosaurian groups (Zanno & Makovicky 2010). It is worth noting that outside coelurosaurs only Limusaurus displays a toothless skull and mandible in mature individuals (Wang et al. 2017). The pedal unguals of the Angeac-Charente theropod have a weak longitudinal curvature and exhibit the reduction of the flexor tubercle to a ventral platform seen in ornithomimosaurs, but also in abelisauroids (Fig. 29B, C; Cerroni et al. 2019: fig. 7). Nevertheless, they are more reminiscent of ornithomimosaurs, being slender, and having a triangular cross-section and a single ventral groove (Longrich 2008), whereas pedal unguals of Afromimus and Masiakasaurus are shorter and possess a dorsal vascular groove.

The centrum of the middle and distal caudal vertebrae is long and low (Fig. 29D-L). The anterior and posterior articular surfaces are slightly wider than tall, with a reniform contour (Fig. 29H, L). A broad and shallow sulcus is present on the ventral surface, and it is laterally delimited by two prominent ridges (Fig. 29E, J). All these features are present in ornithomimosaurs (Osmolska *et al.* 1972; Longrich 2008) but also in *Elaphrosaurus* (Rauhut & Carrano 2016). As in all ornithomimosaurs, the robust and tongue-shaped prezygapophyses of the Angeac-Charente taxon are elongated anteroposteriorly, up to three-quarters the length of the centrum. They are horizontally directed (Fig. 29E, K) and do not diverge laterally from the sagittal plane (Fig. 29D, I). Conversely, the zygapophyses of ceratosaurs are slender, shorter and directed anterodorsally (Carrano *et al.* 2002; O'Connor 2007, Cerroni *et al.* 2019).

The tibia of the Angeac-Charente ornithomimosaur has already been described in detail (Allain *et al.* 2014). Here, we figure new material to highlight the features that best differentiate it from a ceratosaur tibia (Fig. 30A-D). The proximal end of the tibia is markedly different from that of *Ceratosaurus, Masiakasaurus, Carnotaurus, Majungasaurus, Afromimus* and *Elaphrosaurus* having a fibular crest clearly separated from the proximal articular surface (Fig. 30A-C), as in tetanuran theropods and thus all the ornithomimosaurs. The elliptical scar present on the posterior surface of the proximal end of the tibia of some ceratosaurs is not visible in the Angeac-Charente taxon (Cerroni *et al.* 2019). As in all ornithomimosaurs, the anterior surface of the distal end of the tibia of the Angeac-Charente taxon bears a tall and transversely expanded flat articular surface for the ascending process of the astragalus (Fig. 30D). There is no medial buttress to accommodate the ascending process as in many basal tetanurans and ceratosaurs, including *Berberosaurus*, *Masiakasaurus*, *Majungasaurus* and *Ceratosaurus*.

The medial face of the fibula bears a deep and proximodistally elongate elliptical fossa for the insertion of *musculus popliteus*. This fossa opens medially and is anteriorly and posteriorly bounded by sharp rims (Fig. 30E). Such a condition is only known in coelurosaurs and *Elaphrosaurus*, and markedly differs from the condition seen in coelophysoids and ceratosaurs, in which the fossa is covered anterodorsally by the tibial crest and thus opens posteriorly (Rauhut 2003; Allain *et al.* 2007; Rauhut & Carrano 2016; Cerroni *et al.* 2019).

In common with the tibia and fibula, the astragalus has a morphology typical of the coelurosaurs and very different from that of the ceratosaurs (Fig. 30F-H). In contrast to Ceratosaurus, Elaphrosaurus, Masiakasaurus, and abelisaurids, the astragalus is fused neither to the calcaneum nor the tibia or fibula (Fig. 30H). The height of the blade-like ascending process of the astragalus is more than twice the height of astragalar body and the process arises from the complete breadth of the astragalar body (Fig. 30F-G). In contrast, all ceratosaurs exhibit a low and narrow ascending process. In addition, the fibular facet on the astragalus is strongly reduced on the lateral side of the ascending process of the astragalus (Fig. 30H). In contrast, the distal end of the fibula of numerous abelisauroids including Berberosaurus, Masiakasaurus, Afromimus and Majungasaurus is transversely expanded and the flared distal end partially overlaps the ascending process of astragalus, the fibular facet of which is large. As previously stated (Néraudeau et al. 2012, Allain et al. 2013, 2014), all surveyed anatomical features agree with assignment of the Angeac-Charente theropod to Ornithomimosauria.

Cerroni *et al.* (2019) have recently questioned the ornithomimosaurian phylogenetic affinities of the Early Cretaceous African *Nqwebasaurus* (Choiniere *et al.* 2012). If confirmed, it would imply that the Charentais taxon is the oldest known ornithomimosaur, based on the Berriasian age of the Lägerstatte of Angeac-Charente (Benoit *et al.* 2017; Polette *et al.* 2018). Moreover, ornithomimosaurs would then have an exclusively Laurasian distribution. Nevertheless, based on first hand examination of fossil specimens by one of us (R.A.), the phylogenetic affinities of *Limusaurus* and *Deltadromeus* are far from certain. More detailed descriptions regarding their anatomy are required to draw conclusions regarding the origin and evolution of ornithomimosaurs.

Superfamily TYRANNOSAUROIDEA Osborn, 1905

Tyrannosauroidea indet. (Fig. 28H-J)

Description

There are at least eight teeth with a very characteristic morphology. They are D-shaped in cross-section with the mesial carina lingually displaced (Fig. 28H-J). The mesial margin is



Fig. 28. — Theropod teeth from Angeac-Charente: A-C, Archaeopterygid tooth (ANG M-09) in lingual (A) and labial (B, C) views; D, Archaeopterygid tooth (ANG M-08) in lingual view; E, F, cf. *Nuthetes* sp. (ANG M-45) in lingual (E) and distal (F) views; G, cf. *Nuthetes* sp. (ANG M-61) in lingual view; H, tooth of Tyranno-sauroidea indet. (ANG17-5342) in lingual view; I, J, tooth of Tyrannosauroidea indet. (ANG M-73) in lingual (I) and distal (J) views; K, Megalosauridae? indet. (ANG17 R-1748) in lingual view; J, L, M, Megalosauridae? indet. (ANG17-5650) in labial (L) and (M) lingual views; N, Megalosauridae? indet. (ANG M-121) in lingual view. Scale bar: A, B, 1 mm; C-F, 400 µm; G, 5 mm; H-N, 1 cm.



Fig. 29. — Ornithomimosaur remains from Angeac-Charente: **A**, right dentary (ANG11-776) in lateral view; **B**, pedal ungual (ANG11-1335) in lateral view; **C**, pedal ungual (ANG11-898) in lateral view; **D**-**H**, middle caudal vertebra (ANG12-1622) in dorsal (**D**), ventral (**E**), right lateral (**F**), anterior (**G**) and posterior (**H**) views; **I-L**, distal caudal vertebra (ANG14-3084) in dorsal (**I**), ventral (**J**), right lateral (**K**) and posterior (**L**) views. Scale bar: 2 cm.

only slightly convex. The distal carina is denticulate along its entire length, while the mesial carina is only denticulate on its apical quarter. The denticles extend over the apex. There are 12-13 denticles per 5 mm on mesial and distal carinae. Two longitudinal grooves are present on the lingual side of ANG M-73, below the denticulate mesial carina (Fig. 28I). When preserved, the enamel texture is irregular (Hendrickx *et al.* 2015a).

The morphology of these teeth and in particular the lingual offset of the mesial carina are typical of tyrannosauroids (Holtz 2004). They are very similar to that described in the Bathonian *Proceratosaurus bradleyi* and the Barremian *Eotyrannus* (Hutt *et al.* 2001; Rauhut *et al.* 2010). This discovery confirms the presence of Tyrannosauroidea in Europe at the beginning of the Early Cretaceous.

Family DROMAEOSAURIDAE Colbert & Russell, 1969 Genus Nuthetes Owen, 1854

cf. *Nuthetes* sp. (Fig. 28E-G)

DESCRIPTION

Several isolated dromaeosaurid teeth, including the two reported here (ANG M-61; ANG M-45), were recovered following sampling and screening-washing of a gravel lens, rich in lignite and fossil bone remains (Fig. 28E-G). These teeth lack a serrated mesial carina, as in Tsaagan mangas (Norell et al. 2006) and some teeth of Nuthetes destructor (Milner 2002). ANG M-61 is a complete tooth crown, and is more rounded. It is strongly recurved and its distal carina runs along half the length of the lingual face of the crown (Fig. 28G). There are 32 denticles per 5 mm on distal carina. This tooth is identified as a mesial tooth. ANG M-45 is the apical part of a second tooth (Fig. 28E-F). Its crown is more labiolingually compressed, and this tooth is identified as a lateral tooth. The distal carina bears numerous distinct denticles. In both teeth, the enamel surface is devoid of ornamentation.

Teeth identical in every way to those of Angeac-Charente have also been reported from the Berriasian of Cherves-de-Cognac (Pouech 2008) and from the Tithonian of Chassiron (Vullo *et al.* 2014). All these teeth from Charente are similar to those of the dromaeosaurid *Nuthetes destructor* from the Purbeck Group of southern England (Milner 2002), and are tentatively referred to this genus.

> AVES Linnaeus, 1758 Family Archaeopterygidae Huxley, 1872

> > Archaeopterygidae indet. (Fig. 28A-D)

DESCRIPTION

Archaeopterygid birds are represented in Angeac-Charente by at least five teeth. One of these teeth is complete (ANG M-09, Fig. 28A-C), whereas the others are broken at the base of the crown (Fig. 28D). The total height of the complete tooth is 3.2 mm. The crown height is only 1.2 mm and its basal length is 0.63 mm. The crown is strongly compressed labiolingually. There are no obvious enamel ornamentation and serrations are totally absent on the slight carinae. The apical quarter of the tooth is strongly distally recurved. Both the mesial and distal edges of the crown are sigmoid. By comparison with teeth of other theropods, we consider that the most convex side of the tooth corresponds to the labial surface (Fig. 28B-D).

The specimens from Angeac-Charente are nearly identical to the single tooth collected at Cherves-de-Cognac, which in turn was assigned to an indeterminate archaeopterygid, based on the above characteristics (Louchart & Pouech 2017). Angeac-Charente material is the youngest temporal occurrence of this extinct European family of early birds.

> SYNAPSIDA Osborn, 1903 MAMMALIA Linnaeus, 1758

> > Mammalia indet. (Fig. 32C, D)

Description

A tooth from the mammal material of Angeac-Charente, ANG M-34 (Fig. 32C, D), preserves a high, main central cusp, one accessory cusp and one root. Given the strong development of the accessory cusp and the inclination of the main cusp apex, this side of the tooth, and thus the only preserved root, is considered to be distal. The section of the tooth breakage shows that a second, mesial root was present. A symmetrical and similar accessory cusp was maybe present on the mesial side. Such morphology is reminiscent to that of the posterior premolariforms of the Middle Jurassic eutriconodont Amphilestes and other "Amphilestidae" (Kielan-Jaworowska et al. 2004). However, without the second half of the tooth, it is difficult to reach a conclusion, as it could also correspond to other, more derived mammals. Without any more diagnostic characters, this specimen is thus cautiously identified as Mammalia indet.

> Family THEREUODONTIDAE Sigogneau-Russell & Ensom, 1998 Genus *Thereuodon* Sigogneau-Russell, 1989

> > *Thereuodon* cf. *taraktes* (Fig. 32G-I)

DESCRIPTION

An isolated tooth crown (Fig. 32G-I) is assigned to the genus *Thereuodon* of the monogeneric family Thereuodontidae, on the basis of several characters (Sigogneau-Russell 1989; Sigogneau-Russell & Ensom 1998; Kielan-Jaworowska *et al.* 2004): it is labio-lingually narrow, with a great development of the parastylar lobe, forming almost a basin (Fig. 32I); a small vertical concavity can be seen on the anterior face of the paracone; the stylocone is well-developed (Fig. 32G); cusp D occurs just behind the stylocone, although it is heavily abraded on this specimen (Fig. 32G); a vertical ridge, the mediocrista, links the paracone to the cusp D, although it is weakly pronounced (Fig. 32I). Thus, three transversely narrow basins



Fig. 30. — Ornithomimosaur remains from Angeac-Charente: A-C, proximal end of right tibia (ANG12-1893) in posterior (A), anterior (B) and lateral (C) views; D, distal end of right tibia (ANG10-56) in anterior view; E, proximal end of right fibula (ANG11-696) in medial view; F-H, Left astragalus and calcaneum (ANG12-1803) in anterior (F), posterior (G) and dorsal (H) views. Scale bar: A-E, 4 cm; F-H, 2 cm.

are present in occlusal view. Moreover, the paracone is high compared to the labial wall and labially but not posteriorly recurved (Fig. 32G, I). This crown is well-preserved, lacking only the metastyle.

Two species are known for the genus *Thereuodon*: the type-species *T. dahmanii* (Sigogneau-Russell 1989) from the Jurassic/Cretaceous transition of the Ksar Metlili Formation, Morocco, and *T. taraktes* (Sigogneau-Russell & Ensom 1998), from the Early Cretaceous (Berriasian) Purbeck Group, England. In the Angeac-Charente specimen,

the great protrusion of the parastylar cusp and the weakly marked mediacrista are reminiscent of *T. taraktes*, but the well-developed metacone is close to what is observed in *T. dahmanii*. The stylocone is closer to cusp D and less sharp than in *T. dahmanii*, but this conformation is similar to that of *T. taraktes*. Finally, as in *T. taraktes* but unlike *T. dahmanii*, there is no cuspule "c". In conclusion, we tentatively assign this specimen to *Thereuodon* cf. *taraktes*.

Sigogneau-Russell (1989) and Sigogneau-Russell & Ensom (1998) considered the teeth they referred to the



Fig. 31. — Reconstruction of the Angeac Ornithomimosaur, based on 3D surface scans of 232 bones. All the bones of the 3D reconstruction were scaled on the basis of a 40 cm long femur.

two species of Thereuodon as "symmetrodontan" permanent molars. However, the upper deciduous premolars of Nanolestes drescherae, a stem-Zatheria from the Late Jurassic of Guimarota, Portugal (Martin 1999, 2002), show some similarities with the teeth of *Thereuodon*. Thus, the DP?3-5 of N. drescherae share with the teeth of Thereuodon an obtuse-angled trigone, a low and recurved paracone, and a trigone basin divided by a ridge connecting stylocone and paracone. These characters are diagnostic of Holotheria, a clade that include the last common ancestor to Kuehneotherium and Theria, and all of its descendants (Hopson 1994; Wible et al. 1995). Martin (2002) thus reinterpreted the teeth of Thereuodon as holotherian, probably zatherian upper deciduous premolars. Moreover, Bonaparte (1990) and Sigogneau-Russell & Ensom (1998) observed a gross resemblance between the teeth of Thereuodon and the upper cheek teeth of Barberenia, from the Late Cretaceous of Argentina. These were subsequently identified by Martin (1999) as upper deciduous premolars, probably belonging to the "dryolestoidean" Brandonia from the same formation, an opinion then followed by Bonaparte (2002). Thus, it seems that Thereuodon should be considered as a stem-Zatheria (Martin 2002). However, from a nomenclatural point of view, the

fact that *Thereuodon* is known only by milk teeth does not invalidate the genus, which is why we use it here to precise the identification of this specimen.

EUTRICONODONTA Kermack, Mussett & Rigney, 1973 (Fig. 32A, B, E, F)

Description

Some mammalian teeth show a distinctive morphology comprising three main cusps placed serially in anteroposterior alignment on a transversely compressed crown (Fig. 32A, B, E, F). This cusp arrangement and general morphology is typical of eutriconodontans (Kielan-Jaworowska *et al.* 2004).

Orientation of these isolated teeth follows the criteria of Godefroit & Battail (1997). The most convex side of the crown is considered to be the labial face, and the most concave the lingual face. Accessory cusps are usually more numerous or better developed on the distal side, and thus the side bearing the most developed and/or the most differentiated accessory cusps is considered to be distal. The distal inclination of the cusps, when present, also helps to distinguish the mesial and distal faces of the teeth.



Fig. 32. — Mammal teeth from Angeac-Charente: **A**, **B**, premolariform or molariform tooth of *Gobiconodon*? sp. (ANG M-21) in occlusal (**A**) and labial (**B**) views; **C**, **D**, premolariform tooth of Mammalia indet. (ANG M-34) in lingual (**C**) and labial (**D**) views; **E**, **F**, left lower molar of *Triconodon* sp. (ANG M-02) in lingual (**E**) and occlusal (**F**) views; **G**-**I**, left upper molar of *Thereuodon* cf. *taraktes* (ANG M-23) in labial (**G**), lingual (**H**) and occlusal (**I**) views. Scale bar: 500 µm.

Family GOBICONODONTIDAE Chow & Rich, 1984 Genus *Gobiconodon*? Trofimov, 1978

Gobiconodon? sp. (Fig. 32A, B)

DESCRIPTION

Within the eutriconodontan material from Angeac-Charente, the specimen ANG M-21 (Fig. 32A, B) consists of an isolated tooth crown, the main central cusp a being surrounded by two unequal cusps b and c (see figures 4.7 and 7.2 by Crompton & Jenkins [1968] and Kielan-Jaworowska et al. [2004] respectively for an explanation of the cusp-numbering system). The root is missing but the tooth was probably uniradiculate. The crown is labiolingually compressed and the lingual face appears almost planar in occlusal view, while the labial face is convex at its base (Fig. 32A). There is no trace of either a labial or lingual cingulid. Cusp a dominates the crown and is triangular in lateral profile (Fig. 32B). Its mesial surface is rounded, but its distal edge shows a well-defined crest (Fig. 32A). An oval wear facet can be observed on the labiodistal surface of cusp a (Fig. 32A). Cups b is smaller than cusp c; it is also placed higher relatively to cusp a and less separated from it than cusp c. The apex of cusp c appears to be slightly flexed labially. Both cusps b and c bear small labially oriented wear facets. The presence of a probably single root indicates a tooth from the anterior dentition, but the well-developed accessory cusps show that it is a distal premolar and not a more mesial tooth. Among Laurasian Early Cretaceous mammals, single-rooted distal premolars are only seen within the Gobiconodontidae family (Sweetman 2006). Moreover, the specimen ANG M-21 closely resembles the gobiconodontid distal premolar (possibly right p4) described by Sweetman (2006) in the Early Cretaceous Wessex Formation of the Isle of Wight, southern England, and tentatively referred to Gobiconodon. Considering this, and the general morphology of ANG M-21 (tricuspid tooth with mesiodistally aligned cusps; dominant cusp a and small but well-developed accessory cusps; labial inflation of the crown; lack of cingulid), the specimen is identified as a left distal premolar, probably a p3 (as the accessory cusps are less developed than in the possible p4 described by Sweetman [2006]), of a gobiconodontid mammal, and is tentatively referred to Gobiconodon.

Family TRICONODONTIDAE Marsh, 1887 Genus *Triconodon*? Owen, 1859

Triconodon? sp. (Fig. 32E, F)

DESCRIPTION

The specimen ANG M-02 (Fig. 32E, F) consists of an almost complete tooth, only lacking the mesial root.

The crown is labiolingually compressed and bears three mesiodistally aligned cusps. A main central cusp a is largely dominant. It is triangular in lateral profile and surrounded by two smaller accessory cusps: cusp c (distal) reaches half of the height of cusp a, and cusp b (mesial) is about three times smaller than cusp c. Cusp a is slightly flexed lingually. Cusps a and c bear distolingually oriented wear facets. Both labial and lingual cingulid are present. Thus, considering these characters, ANG M-02 can be identified as an eutriconodontan left lower premolariform. It is somewhat similar to the p3 of *Triconodon* from the Berriasian of Britain (Jäger *et al.* 2021), so the specimen is tentatively referred to this genus.

ALLOTHERIA Mars, 1880

MULTITUBERCULATA Cope, 1884 (Fig. 33)

PRELIMINARY STATEMENT

Multituberculates are the most commonly represented mammals in Angeac-Charente with nearly fifteen isolated teeth already collected by screening-washing. We follow here the classification of Mesozoic allotherian mammals provided by Kielan-Jaworowska et al. (2004). All teeth are provisionally referred to the Pinherodontidae (Hahn & Hahn 1999), except for one P4 (Fig. 33I, J) that is very similar to the holotype of Sunnyodon notleyi (Kielan-Jaworowska & Ensom 1992), and one ?P5 that has only two rows of cusps (Fig. 33K-L), whereas there are three in Pinherodontidae (Hahn & Hahn 1999; Kielan-Jaworowska et al. 2004) and is thus identified as Multituberculata indet. A systematic revision of European Late Jurassic to Early Cretaceous Multituberculates is beyond the scope of this paper, but it is worthy to note that, for the moment, based only on their morphology, no tooth really suggests the presence of more than two taxa of multituberculates in Angeac-Charente.

Family PINHEIRODONTIDAE Hahn & Hahn, 1999

Pinheirodontidae indet. (Fig. 33A-H, M-P)

Description

Most of the multituberculate teeth discovered in Angeac-Charente can be referred to the family Pinheirodontidae, but it has not been possible to refer them to any existing genera. The material figured here includes a left p4 (ANG M-72), a left P1 (ANG M-03), a right P2 (ANG M-06), a left P3 (ANG M-22), a left ?m2 (ANG M-105) and a left M2 (ANG M-32) (Fig. 33A-H, M-P).

The most mesial part of the blade-like p4 (ANG M-72) is missing (Fig. 33A). It possesses at least six serrations, and no basal labial cusp, a diagnostic feature of *Pinheirodon* (Hahn &



Fig. 33. — Multituberculate mammal teeth from Angeac-Charente: **A**, **B**, left p4 of Pinheirodontidae indet. (ANG M-72) in lingual (**A**) and labial (**B**) views; **C**, **D**, left P1 of Pinheirodontidae indet. (ANG M-03) in occlusal (**C**) and distolingual (**D**) views; **E**, **F**, right P2 of Pinheirodontidae indet. (ANG M-06) in occlusal (**E**) and lingual (**F**) views; **G**, **H**, left P3 of Pinheirodontidae indet. (ANG M-02) in occlusal (**C**) and distolingual (**D**) views; **E**, **F**, right P2 of Pinheirodontidae indet. (ANG M-06) in occlusal (**E**) and lingual (**F**) views; **G**, **H**, left P3 of Pinheirodontidae indet. (ANG M-22) in occlusal (**G**) and lingual (**H**) views; **I**, **J**, left ?P4 of *Sunnyodon* sp. (ANG M-04) in occlusal (**I**) and labial (**J**) views; **K**, **L**, left ?P5 of Multituberculata indet. (ANG M-106) in mesio-occlusal (**K**) and occluso-labial (**L**) views; **M**, **N**, left ?m2 of Pinheirodontidae indet. (ANG M-105) in occlusal (**M**) and labial (**N**) views; **O**, **P**, right M2 of Pinheirodontidae indet. (ANG M-32) in occlusal (**O**) and lingual (**P**) views. Scale bar: A, B, 1 mm; C, D, 750 µm; E-P, 500 µm.

Hahn 1999; Kielan-Jaworowska *et al.* 2004). Nevertheless, the Angeac-Charente specimen has a straight, vertical, distal margin (Fig. 33B), while the latter is strongly convex just above the root in *Pinheirodon*.

The ?m2 (ANG M-105) has a well-developed central basin. Labial cusps are missing and there are only two lingual cusps as in *Bernardodon* (Fig. 33N), whereas there are three in *Pinheirodon* (Hahn & Hahn 1999). Nevertheless, the tooth shape is considerably longer than wide, and precludes assigning it to *Bernardodon* (Fig. 33M).

The M2 (ANG M-32) shows a prominent anterolingual shelf (Fig. 33O), like that observed in the molars of *Bernarodon* and *Pinheirodon* (Kielan-Jaworowska *et al.* 2004). As in the latter two genera there are three lingual cusps, but ANG M-32 differs from other pinheirodontids in lacking the labial cusps (Fig. 33P; Hahn & Hahn 1999).

The anterior upper premolars P1 (ANG M-03), P2 (ANG M-06) and P3 (ANG M-22) have 3-4 ribbed cusps arranged in two rows like in other "plagiaulacidans" (Fig. 33C-H; Kielan-Jaworowska *et al.* 2004). They are not very diagnostic and only tentatively assigned here to Pinheirodontidae, based on their similarities with the pinheirodontid teeth described by Hahn & Hahn (1999: fig. 58).

Family ?PAULCHOFFATIIDAE Hahn, 1969 Genus *Sunnyodon* Kielan-Jaworowska & Ensom, 1992

> Sunnyodon sp. (Fig. 33I, J)

DESCRIPTION

ANG M-04 is roughly oval in shape and has only two rows of cusps, with four main lingual cusps, and two main labial cusps and one posterior labial cuspule (Fig. 33I, J). It also possesses an incipient distal lingual ridge, with a small cuspule. All these features led us to assign this tooth to the genus Sunnyodon, a monospecific genus, S. notleyi, based on a single ?P5 from the Berriasian of the Purbeck Group in England (Kielan-Jaworowska & Ensom, 1992). The Angeac-Charente tooth differs from the latter by the labial cusps that are more anteriorly located, whereas they are arranged symmetrically at the middle of the tooth length in S. notleyi, and by the absence of the anterior labial cuspule (Fig. 33I). This difference could be explained by the fact that these teeth belong to two different loci and/or two different species. This tooth is considered here to be a ?P4 and it is referred to Sunnyodon sp., although the validity of this genus can be debated. An upper posterior premolar of Sunnyodon has also been reported from the Berriasian strata of the Rabekke Formation on the island of Bornholm, Denmark (Lindgren et al. 2004).

TRECHNOTHERIA McKenna, 1975

Family SPALACOTHERIIDAE Marsh, 1887 Genus *Spalacotherium* Owen, 1854

Spalacotherium evansae Ensom & Sigogneau-Russell, 2000 (Fig. 34A, B)

DESCRIPTION

Three teeth of Spalacotheriidae have been recognized including two molars, which have the characteristic "symmetrodont" pattern with acute angulation of the principal cusps, seen in other spalacotheriids (Fig. 34A; Kielan-Jaworowska *et al.* 2004).

The best preserved of the molars ANG M-26 lacks the posterior root and the hypoconulid (cusp d) (Fig. 34A, B). The protoconid, the metaconid and the paraconid are welldeveloped. The talonid was probably much reduced. Mesially, a single cusp e is lingually placed on the cingulid (Fig. 34B), and allows the identification of this tooth as a left lower molar. ANG M-26 is very similar, if not identical, to a tooth referred to Spalacotherium evansae (DORC GS 360, Ensom & Sigogneau-Russell 2000). Its smaller size compared to that of the molars of other species of Spalacotherium, the incompleteness of the labial cingulid associated with the equivalent height of the paraconid and metaconid, are diagnostic features (Ensom & Sigogneau-Russell 2000), that allow us to assign this specimen to Spalacotherium evansae. The latter species has been decribed from various localities of the Purbeck Group, including the Berriasian Lulworth Formation at Sunnydown Farm, U.K.

> CLADOTHERIA McKenna, 1975 Superfamily DRYOLESTOIDEA Butler, 1939 Family DRYOLESTIDAE Marsh, 1879

> > Dryolestidae indet. (Fig. 34C-F)

DESCRIPTION

Dryolestidae are represented by only two fragmentary molars lacking the roots and the talonid (Fig. 34C-F). Based on the configuration of the trigonid and the lack of a cuspid in median position on the labial side of the molars, those teeth are considered to be left lower molars (Kielan-Jaworowska et al. 2004). Both lower molars are shorter mesiodistally than wide labiolingually, but the most posterior molar (ANG M-01; Fig. 34E-F) is strongly shortened and widened compared to the anterior molar (ANG M-05; Fig. 34C, D). This feature is diagnostic of Dryolestidae (Kielan-Jaworowska et al. 2004). The trigonid is well-developed in both lower molars. The paraconid of the posterior molar ANG M-01 is broken at its base, but it is labiolingually more elongated than the metaconid. Both are separated by a narrow incision (Fig. 34E). The metaconid is enlarged and almost as high as the protoconid (Fig. 34F). Based on these features, the dryolestid molars from Angeac-Charente seem more closely related to either Guimarotodus from the Kimmeridgian of Guimarota in Portugal (Martin 1999) or Phascolestes from the Berriasian of Dorset in England (Owen 1871; Kielan-Jaworowska et al. 2004) than any other Dryolestidae. However, additional material is required to allow a more accurate attribution.



FIG. 34. — Trechnotherian mammal teeth from the Berriasian of Angeac-Charente: **A**, **B**, left lower molar of *Spalacotherium evansae* Ensom & Sigogneau-Russell, 2000 (ANG M-26) in occlusal (**A**) and lingual (**B**) views; **C**, **D**, left lower molar (m1 or ?m2) of Dryolestidae indet. (ANG M-05) in occlusal (**C**) and linguodistal (**D**) views; **E**, **F**, left lower molar (m6 or ?m7) of Dryolestidae indet. (ANG M-01) in occlusal (**E**) and mesial (**F**) views; **G-I**, left lower molar (m3?) of *Peramus* sp. (ANG M-25) in occlusal (**I**) views. Scale bars: 400 µm.

ZATHERIA McKenna, 1975 Family Peramuridae Kretzoi, 1946

Peramus sp. (Fig. 34G-I)

DESCRIPTION Peramuridae are represented by at least two lower molars. Only a left lower molar (ANG M-25), the talonid and the roots of which are missing, is figured herein (Fig. 34G-I). The talonid is present on the other molar, and it clearly displays a hypoconulid separated from the hypoconid, a diagnostic feature of Peramuridae (Davis 2012). The main cusp of the trigonid, i.e. the protoconid, is slightly inclined posteriorly (Fig. 34H). The paraconid is mesiodistally narrower and apicobasally shorter than the metaconid. It is located higher and in a more labial position. The metaconid is visible in labial view, whereas it is hidden by the protoconid in the Moroccan genus *Minimus* (Sigogneau-Russell 1999). Both the anterolabial (cusp e) and the anterolateral (cusp f) marginal cuspules are preserved (Fig. 34G). The latter is much more developed than the former. They define a strongly developed hypoconulid embrasure. All these features are more reminiscent of the lower molars (m2 and m3) of *Peramus tenuirostris* and *P. dubius* from the Berriasian of Dorset, England (Clemens & Mills 1971), than to any other peramuran taxon (Sigogneau-Russell 1999). However, the scarcity of material prevents any comparison based on measurements, and does not allow a reliable specific identification (Davis 2012). Thus, we assign this tooth to *Peramus* sp.

DISCUSSION

ORIGINALITY OF THE VERTEBRATE FAUNA AND PALEOENVIRONMENTAL IMPLICATIONS *Fossil and taxonomic diversities*

The continental vertebrate fauna from Angeac-Charente is both very diverse and abundant. It includes wide size range of bones and teeth from 0.5 mm to 2.2 m, and includes three types of ichnofossil: coprolites, track casts, and traces on bone surfaces, including trampling and tooth marks (Rozada et al. 2021). But above all, what makes this fossil locality so exceptional is the great taxonomic diversity. With at least forty different vertebrate taxa (see the Appendix 1, Systematic paleontology of the complete faunal list and Table 1 below), Angeac-Charente is the most diverse earliest Cretaceous mixed continental bonebed known to date (see Eberth et al. 2007). It has been demonstrated that the Angeac-Charente taphocoenosis provides a good picture of the local paleocommunity, because the site represents a 'snapshot' of a Lower Cretaceous ecosystem (Rozada et al. 2021), but also because both macro- and microremains have been collected (Carrano et al. 2016). This paleocommunity also reflects the Purbeckian paleometacommunity as a whole, as all major clades known for the Jurassic/Cretaceous transition in Europe are represented including rare fossils such as birds and mammals (Table 1). Recognizing this diversity was only possible as a result of the time and space scales of fieldwork, the laboratory preparation of all macroremains from one year to the next, and the continuous sorting of microfossils.

Abundance

Besides the diversity, the abundance of fossils is also remarkable. Relative abundances of large vertebrates (*i.e.* macroremains) have been calculated based on the material collected between 2010 and 2017 (Fig. 35A). For microremains, we used the material collected from the gravel lens of the R1 plot (Fig. 3), which was rich in lignite and bones including the complete shell of *Pleurosternon bullockii* (Fig. 13; Gônet *et al.* 2019), and the material collected at the base of the Unit 3, of the R3 plot (Fig. 35B, C). The overall paleoenvironmental interpretation and significance of these relative abundances is presented below in more detail, as well as the paleoecological insights they provide based on their relationships with spatial distribution and taphonomic data. The ornithomimosaur is the most abundant taxon throughout. However, its spatial distribution over the site is clearly heterogeneous, as 85% of the ornithomimosaur remains are concentrated in the CG1-8 plots (Fig. 3). This high local abundance of exquisitely-preserved bones supports the hypothesys of a mass mortality event of an ornithomimosaur herd (Rozada *et al.* 2021). While their carcasses may have been transported over a short distance after a mass drowning (see Subalusky *et al.* 2017), their complete skeletons were deposited *in situ* before being scattered by trampling (Rozada *et al.* 2021).

Goniopholidid crocodyliforms, pleurosternid turtles and sauropod dinosaurs are both abundant and homogeneously distributed (Fig. 3). It suggests that these taxa were autochthonous, which is supported by the occurrence of 1) plates of *Pleurosternon bullockii* found in close association in R3 and belonging to a nearly complete specimen, 2) a disarticulated *Goniopholis* sp. skeleton found in CG9, and 3) numerous sauropod track casts found at several loci of the site, as well as the hindquarters of a single individual.

Atoposaurid, bernissartiid and pholidosaurid crocodyliforms are mostly represented by teeth. Atoposaurids and bernissartiids mostly occur as microremains, and they are abundant. They are homogeneously distributed, and were likely part of the autochthonous fauna.

Helochelydrid and thalassochelydian turtles and stegosaurs are only known from a small amount of material. Their distribution appears heterogeneous, probably due to their scarcity. Nevertheless, the three taxa are represented by several associated elements belonging to single individuals. Because of the presence of track casts of the *Deltapodus* type, the stegosaur is considered to be an autochthonous taxon, whereas the two turtles may have been parautochthonous.

Isolated megalosaurid teeth are common and have a homogeneous spatial distribution. Most of the teeth are broken at their base (Fig. 28G-N), suggesting that these breakages occurred during feeding in the Angeac-Charente wetland. However, the absence of carnivorous theropod bones in the assemblage indicates that these dinosaurs probably only fed opportunistically and/or seasonaly in this area.

Pterosaurs, ankylosaurs, heterodontosaurids, camptosaurids, hypsilophodontids, dromaeosaurids, tyrannosauroids, sphenodontians and mammals are only represented by isolated teeth or a single bone (e.g. camptosaurid and ankylosaur). They are much rarer, heterogeneously distributed in the field, and often abraded. This suggests short-distance transport and thus parautochthonous origin for these taxa.

As abundant anuran remains are known in some chaotic debris flows (e.g., Plant Debris Beds of the Barremian Wessex Formation, Isle of Wight, southern England; Sweetman & Insole 2010), their low abundance at Angeac-Charente cannot be explained by the fragility of their bones, but rather seems to reflect genuine scarcity. There may have been a more important sorting upstream of the deposit environment, or a local hydrodynamism difference, impacting the conservation of the anuran remains and/or the presence of anurans at the time of deposition.



FIG. 35. – **A**, Relative abundance of Angeac-Charente large vertebrate taxa based on identified macroremains collected from 2010 to 2017. White numbers indicate the Minimum Number of Individuals (**MNI**) (after Rozada *et al.* 2021); **B**, Relative abundance of Angeac-Charente taxa, based on microremains collected in 2017, by water screen-washing (diameter of mesh = 0.8 mm), at the base of the unit 3, of the R3 plot.

Paleoenvironment

Sedimentology, mineralogogy, and geochemistry as well as non-vertebrate paleontology indicate a continental freshwater wetland depositional environment for the Angeac-Charente bonebed (Rozada *et al.* 2021). The sedimentological assemblage is dominated by clay in suspension deposited from standing TABLE 1. — List of vertebrate taxa from the Jurassic-Cretaceous transition of Charente with their relative abundances in each locality, habitat, stratigraphic range and Mesozoic geographical distribution. The black, grey and white squares represent the preferred habitats of each taxon. Abbreviations : **A**, abundant; **C**, common; **CZ**, costal zone; **FW**, freshwater; **J**, Middle Jurassic; **LC**, Lower Cretaceous; **LJ**, Lower Jurassic; **LP**, Lower Permian; **LT**, littoral; **MT**, Middle Triassic; **NA**, North America; **PR**, paralic; **R**, rare; **SA**, South America; **TR**, terrestrial; **UC**, Upper Cretaceous; **UCa**, Upper Carboniferous; **UJ**, Upper Jurassic; **UP**, Upper Permian; **UT**, Upper Triassic.

		Locality									
		uo	~								
		Sir	ves	ac		Env	viron	ment		Mesozoic	Mesozoic
		has	her	nge	Co	ont.	Li	tt.	Mar.	stratigraphic	geographical
Таха		ΰ	Ö	Ā	TR	FW	PR	LT	CZ	range	distribution
Chondrichthyes Hybodontiformes	Planohybodus Parvodus celsucuspus	R A	А	С		×	•	××		Bathonian MJ/Barremian LC Tithonian UJ/Berriasian LC	Europe/SA/NA Europe
Rajiformes	Belemnobatis		R	-				•	×	Bathonian MJ/Aptian LC	Europe
Osteichthyes Huxley, 1880 Ginglymodi	indet.	А	А	А		×	•			Kimmeridgian UJ/Aptian LC	Europe/Asia/Africa/
lonoscopiformes	indet.	С		R		×	•	×		Anisian MT/Cenomanian UC	Europe/Asia/Africa/
Amiiformes	indet.	R	R	С		×	•	×		Ladinian MT/Maastrichtian UC	Europe/Asia/NA/
Mesturidae	Micropycnodon		R	С		×	•	×		Norian UT/Maastrichtian UC	Europe/Asia/Africa/ NA/SA
Pycnodontidae	indet.	R	R	С		×	•	×		Kimmeridgian UJ/Maastrichtian UC	Europe/Asia/Africa/ NA/SA
Aspidorhynchidae	Belonostomus		R				•	×		Tithonian UJ/Maastrichtian UC	Europe/Africa/NA/
Ichthyodectidae	Thrissops	С	R				•	×		Kimmeridgian UJ/Tithonian UJ	Europe
Lissamphibia Haeckel,1866 Allocaudata	Albanerpetontidae	С	R	С	×	•				Bathonian MJ/Maastrichtian UC	Europe/Asia/Africa/
Caudata	indet.	С		R	×	•				Bathonian MJ/Maastrichtian UC	Europe/Asia/Africa/
Anura	indet.	R		R	×	•				Hettangian LJ/Maastrichtian UC	Europe/Asia/Africa/ NA/SA
Reptilia Laurenti,1768 Testudinata Pleurosternidae Plesiochelyidae	Pleurosternon bullockii Jurassichelon	C A	С	A	×	•	• ×	×	×	Tithonian UJ/Berriasian LC Kimmeridgian UJ/Tithonian UJ	Europe Europe
Helochelydridae Lepidosauria	Hylaeochelys? indet.		R	R R	×	•				Tithonian ÜJ/Valanginian LC Tithonian UJ/Campanian UC	Europe Europe/NA
Sphenodontia Scincomorpha Choristodera	Opisthias Paramacellodus			R R		•				Tithonian UJ/Berriasian LC Tithonian UJ/Barremian LC	Europe/NA Europe/Africa
Crocodylomorpha	Cteniogenys?	R		R		•				Bathonian MJ/Aptian LC	Europe/Asia/NA
Teleosauridae Atoposauridae Bernissartiidae Goniopholididae Pholidosauridae	Steneosaurus Theriosuchus Goniopholis Pholidosaurus	R C C C C C	A A C C	C C A C	•	• • •	× × ×	•	×	Toarcian LJ/Berriasian LC Bathonian MJ/Cenomanian LC Berriasian LC/Aptian LC Kimmeridgian UJ/Berriasian LC Tithonian UJ/Berriasian LC	Europe/Africa/Asia Europe/Asia/Africa Europe Europe Europe
Theropoda	Megalosauridae	R	R	С	•					Bajocian MJ/Berriasian LC	Europe/Asia/Africa/
Dromaeosauridae	Nuthetes Ornithomimosauria Tyrannosauroidea	R	R	R A R	•	-				Tithonian UJ/Berriasian LC Berriasian LC/Maastrichtian UC Bathonian MJ/Maastrichtian UC	Europe Europe/Asia/NA Europe/Asia/NA
Sauropoda	Turiasauria Macronaria indet.	R	R	с R	•					Kimmeridgian UJ/Aptian LC Bajocian MJ/Maastrichtian UC	Europe/Africa/NA Europe/Asia/Africa/ NA/SA
Ornithischia Heterodonto- sauridae	Camptosauridae Echinodon	R	R	R R	•					Kimmeridgian UJ/Berriasian LC Berriasian LC	Europe/NA Europe
Stegosauria	Hypsilophodontidae <i>Dacentrurus</i> Ankylosauria	R	R	R R R	•					Bathonian MJ/Maastrichtian UC Kimmeridgian UJ/Berriasian LC Pliensbachian LJ/Maastrichtian UC	Europe/NA Europe Europe/NA/Asia

TABLE 1. - Continuation.

		Locality									
Таха		Chassiron	Cherves	Angeac	Co	Env ont. FW	riron Li PR	men tt. LT	t Mar. CZ	Mesozoic stratigraphic range	Mesozoic geographical distribution
Pterosauria	Rhamphorhynchidae Ctenochasmatidae Pterodactyloidea A	R R R		A	•					Hettangian LJ/ Berriasian LC Oxfordian UJ/Aptian LC Bathonian MJ/Maastrichtian UC	Europe/Asia/Africa Europe/Asie/SA Europe/Asia/Africa/ NA/SA
	Pterodactyloidea B		R	С	•					Bathonian MJ/Maastrichtian UC	Europe/Asia/Africa/ NA/SA
Mammalia Linnaeus, 1758 Thereuodontidae	Thereuodon		R	R	•					Tithonian U.I/Berriasian I C	Europe/Africa
Eutriconodonta Gobiconodontidae	Gobiconodon		R	R	•					Bathonian MJ/Cenomanian UC	Europe/Asia/Africa/
Triconodontidae Multituberculata	Triconodon		R	R	•					Berriasian LC	Europe
Chalacathariidaa	Pinheirodontidae indet. Sunnyodon	R	R	R R	•					Oxfordian UJ/Barremian LC Berriasian LC	Europe Europe
Cladotheria	Spalacotherium evansae		R	R	•					Berriasian LC	Europe
	Dryolestidae indet. <i>Peramus</i>	R	R R	R R	•					Bathonian MJ/Campanian UC Tithonian UJ/Berriasian LC	Europe/Africa/NA Europe/Africa

water, locally interrupted by higher energy deposits (Néraudeau et al. 2012; Rozada et al. 2021). PAAS (Post-Archean Australian Shale)-normalized REY (Rare Earth Element and Yttrium) patterns of sediments and fossil biogenic apatites from the bonebed also indicate that Angeac-Charente sediments and fossils were deposited in a strictly freshwater environment (Rozada et al. 2021). Angeac-Charente plant microfossils are dominated by spores of liverworts (marchantiophytes) and lycopods (lycophytes), with a good diversity of fern spores (monilophytes) and gymnosperm pollen grains (Polette et al. 2018). Palynological assemblages are, however, strongly biased by huge differences in the quantity of produced sporomorphs, and there are mostly, if not always, mixocoenoses that include a large regional component. In strong contrast, both plant mesofossils and megafossils show a large dominance of conifers (Néraudeau et al. 2012; Daviero-Gomez & Gomez in Allain et al. 2017). Mesofossils mostly consist of wood preserved as jet (vitrain) and charcoal (fusain), out of which some belong to the genus Agathoxylon Hartig. Apart from wood, there are abundant twigs and isolated leaves of Brachyphyllum Brongn., which bear well-preserved cuticles, three-dimensionally preserved cones and seeds, and much rarer fern stipes and frond fragments (Daviero-Gomez & Gomez 2017, unpublished data). Dealing with megafossils, a number of conifer trunks observed in the excavation field remain attached to their rooting systems and branches, and at least one is more than twelve meters in length (Rozada et al. 2021: fig. 6B). Overall, the autochthonous and parautochthonous fossil deposit displays a picture of a local plant community dominated by paucispecific, cheirolepidiacean conifer forest. The fossil algae *Ovoidites parvus* and *Lecaniella* sp. (Néraudeau *et al.* 2012), as well as the large dominance of the fossil charophyte family Clavatoraceae compared to the Porocharaceae (Benoit *et al.* 2017) support a long lasting, fully freshwater, wetland environment. Finally, the abundance of freshwater invertebrates such as viviparid gastropods, and unionoid bivalves preserved in life-position in clay, and ostracods (e.g. *Cypridea* sp.) (Néraudeau *et al.* 2012) are also in agreement with such an environment.

The Angeac-Charente vertebrate fauna is in perfect agreement with and corroborates this sedimentological and paleobotanical-based environmental interpretation (Table 1). Of the about 40 identified vertebrate taxa that compose this fauna, 23 taxa (57.5%) are elsewhere only found in terrestrial environments (helochelydrid turtles, atoposaurid crocodyliforms, dinosaurs, pterosaurs, lepidosaurs and mammals), and 11 (27.5%) in freshwater environments (lissamphibians, pleurosternids, choristoders, bernissartiids, goniopholidids and pholidosaurids). The six remaining taxa (15%) are euryhaline and can live in both freshwater and brackish water. This is the case of the hybodontid shark Parvodus as well as all osteichthyan taxa. About 70% of the identified taxa and 98% of the anatomically unidentifiable macroremains collected from Angeac-Charente are derived from terrestrial taxa (Rozada et al. 2021). Conversely, microremains are dominated by freshwater taxa (Fig. 35B). Given the overall size of the latter, especially the osteichtyans, this suggests that the water was shallow, which is consistent with the occurrence of dinosaurs tracks and trampling marks, as well as the local record of some mudcracks.

It is worth noting that most of the isolated Parvodus teeth found in Angeac-Charente lack a root, which contrasts with what is observed at Cherves-de-Cognac (Rees et al. 2013). Taphonomically, this may be explained by the resorption of the root during tooth replacement (Underwood & Cumba 2010). These rootless teeth therefore reflect a loss during the animal's lifetime, in agreement with a wetland connected with a watercourse. It should also be noted that the turtle Hylaeochelys belli was recently referred to Thalassochelydia, a group otherwise exclusively composed of coastal marine forms (Anguetin & André 2020). However, this species is always found in sediments of freshwater origin (middle and upper Purbeck and Wealden facies of England) and is apparently the only known freshwater thalassochelydian even if the calcium isotopic composition of its biogenic apatite may indicate this taxon spent time in more saline water and was probably euryhaline.

In summary, the composition of the vertebrate fauna, that comprises a mixture of freshwater and terrestrial taxa, supports the environment of a continental, exclusively freshwater wetland without marine inputs (Fig. 36). Our interpretation is consistent with recent sedimentological, ichnological and geochemical data (Rozada et al. 2021), and challenges the previously held assumption of at least occasional, marine water inputs (Néraudeau et al. 2012; Benoit et al. 2017; Polette et al. 2018). The variety of habitats inhabited by the Angeac-Charente vertebrate fauna reflects the complex mosaic of micro-habitats, necessary to sustain a wide range taxa. These include terrestrial, amphibious and aquatic animals in a generally wetland environment. Similar conclusions were drawn concerning the late Barremian freshwater - terrestrial fossil assemblage and environment of Las Hoyas, La Huérguina Formation, Cuenca province, Spain (Buscalioni & Poyato-Ariza 2016).

BIOCHRONOLOGICAL IMPLICATIONS

Age of the Angeac-Charente Lagerstätte based on the vertebrate fauna

Several taxa present in the Angeac-Charente assemblage cannot be identified at low taxonomic levels or are persistent for geologically long periods of time. These taxa do not provide useful information concerning the age of the vertebrate-bearing deposits (e.g. osteichthyians, and pterosaurs). In contrast, many taxa characterized by relatively short stratigraphical ranges can be used as biostratigraphic markers, and we discussed them below.

Fishes. Among the ichthyofauna, the hybodont shark *Parvodus celsucuspus* is an endemic species that is so far restricted to the Tithonian-Berriasian of the Charentes region (Rees *et al.* 2013; Vullo *et al.* 2014). Its presence clearly indicates a similar age for the Angeac-Charente assemblage.

Turtles. The earliest helochelydrid remains have been reported from the Tithonian of England and France, but they become more common during the Early Cretaceous

(Joyce 2017; Pérez-García et al. 2020). The group is only known in Europe and North America and probably went extinct at the end of the Cretaceous (Nopcsa 1928; Lapparent de Broin & Murelaga 1996, 1999; Hirayama et al. 2000; Joyce et al. 2011; Joyce 2017). Earliest Cretaceous, European species are poorly known due to an unsatisfactory fossil record (Joyce 2017; Pérez-García et al. 2020). As mentioned above, the shell surface sculpturing observed in the Angeac-Charente helochelydrid material is closer to the ornamentation of 'Helochelydra' anglica from the Berriasian of Dorset and 'Helochelydra' bakewelli from the Valanginian of Sussex, than to that of Helochelydra nopcsai from the Barremian of the Isle of Wight in England, and Spain. It also differs to that of Helochelys danubina from the Cenomanian of Germany, and from Plastremys lata from the Albian-Cenomanian of England. It is therefore consistent with either a Berriasian or Valanginian age for the Angeac-Charente locality.

In Europe, pleurosternids are known by several Kimmeridgian-Albian species (Lydekker 1889; Milner 2004; Pérez-García & Ortega 2011; Pérez-García 2015; Pérez-García *et al.* 2015a; Joyce & Anquetin 2019; Guerrero & Pérez-García 2020). In North America, they are limited to the Late Jurassic (Gaffney 1979; Joyce & Anquetin 2019). The Angeac-Charente pleurosternid material is clearly referable to the species *Pleurosternon bullockii*, which is known from the Tithonian of Boulogne-sur-Mer (France), and from the Tithonian-Berriasian Purbeck Group of Dorset, England (Joyce & Anquetin 2019; Guerrero & Pérez-García 2020).

Hylaeochelys is known from the Purbeck (Tithonian-Berriasian) and Wealden (Berriasian-Valanginian) of the UK by the type species *Hylaeochelys belli* (Lydekker 1889; Hirayama *et al.* 2000; Milner 2004; Pérez-García 2012) and in the Tithonian of Portugal by *Hylaeochelys kappa* (Pérez-García & Ortega 2014). The material from Angeac-Charente is more consistent with a tentative referral to *Hylaeochelys belli*, although this awaits confirmation. This suggests that the locality is either Berriasian or Valanginian in age.

In brief, the turtle assemblage from Angeac-Charente is similar to that of the predominantly Berriasian Purbeck Group of England and it supports a Berriasian age for this locality.

Lepidosaurs. *Opisthias rarus* is a sphenodontian reptile species from the Late Jurassic Morrison Formation of western North America (Gilmore 1910). An indeterminate species is also known from the latest Tithonian and Berriasian beds of the English Purbeck Group Lulworth Formation (Evans & Fraser 1992; Evans & Searle 2002).

Four species of *Paramacellodus* are known to date, all from Early Cretaceous localities: *P. oweni* and *P. marocensis* from the Berriasian of England and Morocco, respectively, *P. sinuosus* from the Barremian of Spain, and cf. *P. keebleri* from the Aptian-Albian of the United States (Hoffstetter 1967; Richter 1994; Broschinski & Sigogneau-Russell 1996; Evans & Searle 2002; Nydam & Cifelli 2002). In addition, a few indeterminate species tentatively referred to *Paramacellodus* have been reported from various Middle and Upper Jurassic localities (see Evans & Searle 2002). The morphological features of the scarce, fragmentary material from Angeac-Charente suggest that it represents a form close to either *P. oweni* or *P. marocensis*. Although more material is needed to confirm our preliminary identifications, the lepidosaurs from Angeac-Charente tend to indicate a Berriasian age.

Crocodyliforms. Atoposaurids are diverse from the Jurassic to the Late Cretaceous in Europe (e.g. Clark 1986; Schwarz & Salisbury 2005; Martin et al. 2014a; Tennant & Mannion 2014; Schwarz et al. 2017), but also in Asia (Lauprasert et al. 2011) and North America (Foster 2018), and they include several genera beyond the genus Theriosuchus. Atoposaurids can be preserved as complete skeletons on slabs or, more often, as isolated disarticulated specimens, which makes determination of their taxonomic identity problematic. For this reason, they have a biostratigraphic potential but this has not been defined yet. It is noteworthy that the locality of Cherves-de-Cognac yielded two complete skeletons of Theriosuchus, which will be compared in a future work with the relatively complete type material of Theriosuchus pusillus from the Purbeck Group of England (Clark 1986).

In Europe, the oldest goniopholidids have been reported from the Late Jurassic. Goniopholis baryglyphaeus is from the Kimmeridgian of Guimarota, Portugal (Schwarz 2002) and more fragmentary specimens have been reported from the Late Jurassic near Boulogne-sur-Mer (Buffetaut 1986). The genus Goniopholis is a common component of semi-aquatic faunas of the earliest Lower Cretaceous being well-known from the Berriasian Purbeck Group of England (Owen 1879; Salisbury et al. 1999; Andrade et al. 2011) and from the Obernkirchen Sandstone of Germany (Koken 1887; Salisbury et al. 1999). The genus Nannosuchus has also been reported from the Purbeck Group (Owen 1879; Salisbury et al. 1999; Andrade et al. 2011). Other European goniopholidids (Hulkepholis, Ant eophthalmosuchus, Vectisuchus) are all younger in age with most of the European record consisting of specimens from the Barremian-Aptian interval (Buffetaut & Hutt 1980; Salisbury & Naish 2011; Martin et al. 2016b) or from the Albian (Buscalioni et al. 2013; Puértolas-Pascual et al. 2015). Although a Late Jurassic age cannot be excluded, the present recognition of *Goniopholis* sp. supports an early Early Cretaceous age for the locality of Angeac-Charente, in agreement with the Berriasian age proposed for the close-by locality of Cherves-de-Cognac (Colin et al. 2004).

The genus *Pholidosaurus* is restricted to the latest Jurassic and earliest Cretaceous of Europe, where two species have been named: *P. schaumburgensis* von Meyer, 1841 from the Obernkirchen Sandstone of Germany and *P. purbeckensis* (Mansell-Pleydell 1888) from the Purbeck Group of England (Salisbury *et al.* 1999). The occurrence of *Pholidosaurus* sp. in the Tithonian of Chassiron and of *P. purbeckensis* from the Berriasian nearby locality of Cherves-de-Cognac (Martin *et al.* 2016b) indicates that Angeac-Charente is Tithonian-Berriasian in age.

Bernissartia fagesii Dollo, 1883 has been described from the Barremian-Aptian of Belgium. The species was subsequently reported from the Wealden of the Isle of Wight (Buffetaut & Ford 1979), but it has recently been demonstrated that this material pertains to the bernissartiid Koumpiodontosuchus (Sweetman et al. 2015). B. fagesii has also been reported from the Berriasian-Aptian of Galve in Spain (Buscalioni & Sanz 1990). For a review of the distribution of bernisartiids see Martin et al. 2020. The Spanish specimen precise age is uncertain but a Berriasian age cannot be discarded. Bernissartia cf. fagesii has also been reported from the nearby locality of Cherves-de-Cognac (Pouech 2008), which could lend support for a Berriasian age for Angeac-Charente. Nevertheless, given current knowledge concerning bernissartiid stratigraphic distribution, their occurence in Angeac-Charente can only indicate an Early Cretaceous age without further refinement.

Dinosaurs. Among theropods, two taxa (*Nuthetes* and Archaeopterygidae) are exclusively known from the Tithonian-Berriasian of western Europe (Milner 2002; Pouech 2008; Vullo *et al.* 2014; Louchart & Pouech 2017; Rauhut *et al.* 2018). This strongly supports a latest Jurassic-earliest Cretaceous age for the Angeac-Charente assemblage.

The only stratigraphically short-lived ornithischian taxon is the genus *Echinodon*, which is known exclusively from the Berriasian (see Sereno 2012). This corroborates the age provided for the two above-mentioned theropod taxa.

Mammals. Isolated mammalian teeth are often highly diagnostic and recognized genera and species are usually stratigraphycally short-lived taxa, thus providing useful biostratigraphical information. This is the case for the four taxa *Sunnyodon, Thereuodon, Spalacotherium evansae* and *Peramus*, which are restricted to the Berriasian stage (Kielan-Jaworowska & Ensom 1992; Sigogneau-Russell & Ensom 1998; Ensom & Sigogneau-Russell 2000; Davis 2012). This is in full accordance with the age deduced from the vertebrate groups discussed above.

STRATIGRAPHICAL RANGES OF ANGEAC-CHARENTE TAXA The recently proposed earliest Cretaceous (Berriasian) age of the Angeac-Charente locality, based on both charophytes and palynomorphs (Benoit *et al.* 2017; Polette *et al.* 2018), is here confirmed by the vertebrate fauna. Consequently, the stratigraphic range of several taxa is now extended, such as *Parvodus celsucuspus*, Archaeopterygidae, and *Dacentrurus*, the last appearance datum of which is slightly extended since the fauna of Angeac-Charente is slightly younger than that of Cherves-de-Cognac. Conversely, as noted above, the ornithomimosaur from Angeac-Charente may be the oldest representative of its group.

COMPARISONS WITH OTHER PURBECKIAN LOCALITIES OF SOUTHWESTERN FRANCE

As previously stated, three "Purbeckian" fossil localities have been exploited over the past two decades in southwestern France, namely Chassiron (Vullo *et al.* 2014), Cherves-de-Cognac (Mazin *et al.* 2008) and Angeac-Charente. A comparison between these different Charente Purbeckian localities is all the easier and more relevant, as the same type of fieldwork was carried out at each of the sites, with particular attention paid to both micro and macroremains (Pouech 2008; Vullo *et al.* 2014). The sampling in all three localities was executed in a relatively homogeneous fashion. If there are differences between the fossil record of these three localities, it is therefore related to a biological signal or, to a lesser extent, to the scope of the excavations, but by no means to the excavation techniques used in the field.

The three localities were dated using several independent methods. The Chassiron bonebed was precisely dated to the early Tithonian using brachiopods, dinoflagellate cysts, calcareous nannofossils and magnetostratigraphy (Schnyder *et al.* 2012; Vullo *et al.* 2014). Cherves-de-Cognac is dated to the early-middle Berriasian based on ostracods, charophytes and dinoflagellates (Colin *et al.* 2004; El Albani *et al.* 2004; Benoit *et al.* 2017). Finally, charophytes, palynomorphs and vertebrate fauna suggest a middle to late Berriasian age for the Angeac-Charente bonebed (see above). Thus, the three Charentese localities have all been fairly solidly dated with different markers and these facilitate the establishment of a chronological succession of continental faunas from the end of the Jurassic to the beginning of the Cretaceous.

The depositional environments of Chassiron, Chervesde-Cognac and Angeac-Charente also show a progressive change in facies, from a paralic coastal environment subject to salinity fluctuations in Chassiron at the end of the Jurassic (Vullo et al. 2014), to a hyper-saline lagoonal environment in Cherves-de-Cognac (Buffetaut et al. 1989; Colin et al. 2004; Mazin et al. 2008; Pouech 2008), to a continental wetland environment in Angeac-Charente (Néraudeau et al. 2012; Rozada et al. 2021). This evolution of the depositional setting is concomitant with the marine regression in Western Europe that characterizes the J/K boundary (Hallam 2001), and is echoed in the paleontological content of each of these localities. The plant assemblages range from small and fragmented, highly transported remains of lycopods (spores), ferns (fronds) and conifers (Agathoxylon and Brachyoxylon wood, twigs, leaves, cones) and seeds at Chassiron, to rare coniferous remains (Agathoxylon wood fragments) at Cherves-de-Cognac, and to abundant native to paranative well-preserved remains of a very diverse flora at Angeac-Charente (Allain et al. 2017). Charophyte assemblages suggest a progressive change from brackish water in Chassiron with abundance of Porocharaceae, to less saline water in Cherves-de-Cognac with the presence of Clavatoraceae, but the dominance of Porocharaceae, and freshwater in Angeac-Charente with the presence of Porocharaceae, but the dominance of Clavatoraceae (Benoit et al. 2017).

The vertebrate faunas show the same range. No coastal or marine taxa are present in Angeac-Charente, which contrasts with what is observed in the other two slightly older Charente localities: Cherves-de-Cognac and Chassiron (Table 1). Chassiron includes remains of bony euryhaline fish and taxa from littoral to paralic environments, such as the hybodont shark Planohybodus, the thalassochelydian turtle Jurassichelon (Pérez-García 2014), and the teleosaurid crocodylomorph Steneosaurus. The rich and diverse, terrestrial, and freshwater vertebrate assemblage mainly consists of isolated meso- and microremains (Vullo et al. 2014). Cherves-de-Cognac also yields a mixture of coastal and continental organisms with abundant chondrichthyan and osteichthyan remains (Pouech 2008). Macroemains are mostly dominated by crocodylomorphs and, to a lesser extent, turtles. Dinosaurs, pterosaurs, lissamphibians, and mammals are far less abundant, but they comprise a diverse assemblage. They are thought to be allochthonous or parautochthonous to the depositional environment (Pouech et al. 2015). This is in stark contrast to the Angeac-Charente ecosystem in which most of the terrestrial taxa, represented by abundant material, have proved to be autochthonous (see above).

Numerous euryhalin, freshwater and/or terrestrial taxa are common to all three localities (Table 1), including Pleurosternidae, *Theriosuchus*, Bernissartiidae, *Goniopholis*, *Pholidosaurus*, Dromaeosauridae, Iguanodontia, Stegosauria, Pinheirodontidae and Cladotheria. This suggests that rather than reflecting faunal turnover between the lower Tithonian and the upper Berriasian, Chassiron, Cherves-de-Cognac and Angeac-Charente vertebrate faunas reflect progressive «terrestrialization» of ecosystems in this area related to the Upper Jurassic regression. Together, they provide a good composite picture of the Purbeckian paleometacommunity of Southwestern France.

PALEOGEOGRAPHIC INSIGHT

During the Jurassic/Cretaceous transition, the European Archipelago is at the crossroad of Asia, North America and Gondwana (Scotese 2014), and this is reflected in the paleogeographic distribution of the taxa that make up the vertebrate assemblages of Charente (Table 1). Only two taxa of this fauna seem to exhibit a certain endemism: the hybodont shark *Parvodus celsucuspus*, which was only known at its type-locality of Cherves-de-Cognac, but is now reported from Angeac-Charente, and the Angeac-Charente ornithomimosaur. Most of the Charente taxa have a European distribution, and what could be called a Purbeckian signature. They are widespread in western Europe and are represented by abundant specimens, such as Belemnobatis, Thrissops, Pleurosternon, Jurassichelon, Hylaeochelys, Goniopholis, Pholidosaurus, Bernissartiidae, Nuthetes, Archaeopterygidae, Echinodon, Dacentrurus, Pinheirodontidae, Triconodon, Sunnyodon and Spalacotherium evansae (Table 1). Nevertheless, at a higher taxonomic rank, most of these taxa have a much more extended distribution,



Fig. 36. - Reconstruction of the Angeac-Charente landscape, 140 Million years ago, © Mazan.

especially in Laurasia, which may suggest a Middle Jurassic Pangaean-inherited distribution, followed by vicariant evolution in western Europe.

The close affinities between the North American (Morrisson Formation), European (Iberian Peninsula) and African (Tendaguru) upper Jurassic macrofauna have already been highlighted (Mateus 2006; Mannion et al. 2019). For some, they are the result of dinosaur faunal exchanges between North America, Europe and Africa in the Late Jurassic (Mateus 2006), but for others they rather reflect a widespread distribution that occurred as early as the Middle Jurassic (Mannion et al. 2019). Faunal interchanges between Europe and North America do not appear to be difficult to envisage during the Jurassic-Cretaceous transition. They are supported by paleogeographic data (Dercourt et al. 2000; Scotese 2014) and corroborated by numerous Late Jurassic taxa, including the theropods Allosaurus, Ceratosaurus and Torvosaurus (Mateus 2006), but also by other Early Cretacous taxa present in Charente, such as Opisthias, Pleurosternidae, Helochelydridae, Hypsilophodontidae and Camptosauridae. Faunal exchanges between Europe and Africa, at the end of the Jurassic and at the beginning of the Cretaceous is less consensual, as there has been no clearly established land connection between the two continents (Mannion et al. 2019). Such a connection between Laurasia and Gondwana cannot be ruled out, however, and seems to have been possible via the Mediterranean Tethyan Sill during low eustatic levels (Vrielynck *et al.* 1995; Dercourt *et al.* 2000; Gheerbrant & Rage 2006). In fact, Lasseron *et al.* (2020) proposed the existence of a terrestrial route across the Tethys, based on several taxa, most of which have been found in Charente.

This is the case for the squamate Paramacellodus, which is known from the Kimmeridgian of United States, and the Tithonian-Berriasian of England, France and Morocco (Hoffstetter 1967; Richter 1994; Broschinski & Sigogneau-Russell 1996; Evans & Chure 1998). It is also worth noting that a paramacellodid dentary from the Late Jurassic of Tanzania has been referred to Becklesius, a second genus otherwise known from the Kimmeridgian-Barremian of Europe (Broschinski 1999). Two mammalian genera, Thereuodon and Peramus, show the same paleogeographic distribution pattern as paramacellodids. The presence of Thereuodon and Peramus at Angeac-Charente first underlines the affinities of this site with those from the Purbeck Group, since the Charente species seem more closely related to English taxa than to any other taxon (see above), but both genera are also present at Anoual, Morocco, on the southern edge of Tethys (Lasseron et al. 2020). It is very likely that the distribution of these taxa, in both Europe and Africa, results from dispersal events from Laurasia to Gondwana via ephemeral land connections during the Late Jurassic and/or the beginning of the Early Cretaceous. Indeed, a sampling bias can hardly be invoked to explain the absence of these taxa in Africa during the Middle Jurassic since the recently discovered Guelb el Ahmar fossiliferous sites in the Bathonian Anoual Formation of Morocco have been intensively sampled and have yielded a riche vertebrate fossil assemblage, including mammals (Haddoumi *et al.* 2016; Lasseron *et al.* 2020). Neither has any of these three taxa been recovered from the Middle Jurassic of Madagascar (Flynn *et al.* 2006).

The Gobiconodontidae from Angeac-Charente represent the first occurrence of this group in France, and only the fourth occurrence in Europe, with a specimen from the Bathonian of Britain (Butler & Sigogneau-Russell 2016) and two others from the Barremian of Britain (Sweetman 2006) and Spain (Cuenca-Bescós & Canudo 2003; Martin et al. 2015). Outside of Europe, Gobiconodontidae are known from Asia, North Africa and North America (Kielan-Jaworowska et al. 2004; Sweetman 2006; Butler & Sigogneau-Russell 2016), in a stratigraphic range comprised between the Lower Jurassic and the Lower Cretaceous (Butler & Sigogneau-Russell 2016). It has been widely accepted that the gobiconodontids originated and diverged in central and eastern Asia (Chow & Rich 1984; Cuenca-Bescós & Canudo 2003). Cuenca-Bescós & Canudo (2003) also concluded that the currently known distribution of this family was the result of at least two dispersal events, both of which originated in central Asia, a first one towards Europe during the Barremian and a second one, later, towards North America. Thus, the gobiconodontids would be the only taxon from the Angeac-Charente fauna to have affinities with Asian taxa. However, the discovery of gobiconodontids in the Upper Jurassic-Lower Cretaceous of Morocco (Sigogneau-Russell 2003) and in the Middle Jurassic and the Lower Cretaceous of Britain (Sweetman 2006; Butler & Sigogneau-Russell 2016) questioned the scenario of their dispersal, and maybe their area of origin. Their presence in the Berriasian of Angeac-Charente confirms that faunal interchanges between Asia and Europe did occur well before the Barremian and suggests that the Gobiconodontidae distribution was at least Laurasian at the beginning of the Cretaceous and maybe since the Middle Jurassic. This is further supported by the presence of the genus Huasteconodon in the Lower Jurassic of Mexico (Montellano et al. 2008), which has been tentatively referred to the Gobiconodontidae.

In summary, the vertebrate fauna from Angeac-Charente has above all a Purberckian character. Many exclusively European genera and species belong to families with an essentially laurasian paleogeographic distribution. Some taxa nevertheless suggest dispersal events between Africa and Europe at the Jurassic/Cretaceous transition. In contrast, none of the fossil taxa in Charente have close affinities with Asian taxa.

IS THERE A TURNOVER OF THE CONTINENTAL FAUNA AT THE J/K BOUNDARY?

The definition of the Jurassic/Cretaceous boundary has been a pervasive problem for many decades and still awaits a widely accepted outcome (Remane 1991; Enay 2020). The main debate focuses on whether the Berriasian represents the upper stage of the Jurassic or the lower stage of the Cretaceous. It has been shown that the J/K boundary does not correspond to any significant faunal change in the marine environment at a small scale, making it difficult to set a clear boundary, unlike the Triassic/Jurassic and Cretaceous/Paleogene boundaries (Remane 1991; Enay 2020). No fewer than eleven possible biological markers for the J/K boundary have thus been proposed (Wimbledon et al. 2011), and with the exception of the disappearance of three species of the calpionellid Crassicolaria, there does not seem to be any real turnover among marine microfossils (Granier 2019a, b). Moreover, the marine fauna shows significant provincialism induced by the Purbeckian regression and a long-range correlation is difficult between the Austral, Tethyan, and Boreal domains (Enay 2020).

The age of the J/K boundary is all the more important since this boundary has long been considered a possible mass-extinction event with a 20% level of extinction (Raup & Sepkoski 1982; 1984). Subsequently, the lack of conclusive evidences of a drastic biotic change and/or a remarkable catastrophic event at the J/K boundary led some authors to downplay the importance of the extinction episode at that time (Hallam 1986; Hallam & Wignall 1997; Bambach *et al.* 2004). Recently, based on a large data set, the hypothesis of an important faunal turnover at the J/K boundary has been resurrected (Tennant *et al.* 2017).

In this context, the Angeac fauna does not in any way allow us to definitively settle the debate on the definition of the Jurassic/Cretaceous boundary, but it does bring, to the continental scale, elements that could be taken into consideration. It also undermines the hypothesis of significant changes in the continental fauna at the J/K boundary as currently defined, (the Tithonian/Berriasian boundary). At the current time, Angeac-Charente is the only known Lagerstätte from the Berriasian and is thus of great importance in the study of extinction/faunal turnover during the J/K transition. The main interest of the Chassiron, Cherves-de-Cognac and Angeac-Charente localities is that they follow one another chronologically from the Lower Tithonian to the Berriasian-Valanginian. Thus, they record continental faunal changes on either side of the J/K boundary in a relatively fine way, especially since the assemblages also include microvertebrates. Most of the taxa recored in Angeac-Charente are known before the Jurassic-Cretaceous boundary. Of the 38 Charente vertebrate taxa that were identified to family or genus rank, none became extinct at the Tithonian/Berriasian boundary, except ramphorhynchid pterosaurs (Table 1). Three taxa at a generic level seem to originate during the Berriasian: the heterodontosaurid dinosaur Echinodon, the multituberculate Sunnyodon and the spalacotheriid Spalacotherium. The Spalacotheriidae is unknown before the Berriasian whereas

the Heterodontosauridae and the Paulchoffatiidae are known respectively from the Early and Late Jurassic. As shown above, rather than reflecting an important faunal turnover between the Tithonian and the Berriasian, the three Charente localities appear to record faunal changes related to local environmental changes resulting from the marine regression that characterized the Jurassic/Cretaceous transition.

The same pattern and close affinities with Middle to Late Jurassic faunas have also been suggested for other less diverse earliest Cretaceous continental faunas around the world, including those from the Berriasian-Hauterivian Kirkwood Formation of South Africa (Rich et al. 1983; McPhee et al. 2016), the Tithonian-Berriasian Ksar Metlili Formation of Morocco (Haddoumi et al. 2016; Lasseron et al. 2020), and the ?Berriasian-Barremian Teete vertebrate locality in Western Yakutia, in Russia (Averianov et al. 2018; Skutschas et al. 2018). Conversely, many groups of large vertebrates are recorded neither in the fauna of the late Jurassic nor in that of Angeac-Charente, although, after the Berriasian, they are known in numerous European localities such as the Wessex Formation in England (Batten 2011) and the Late Barremian Lagerstätte at Las Hoyas in Spain (Poyato-Ariza & Buscalioni 2016). Among these groups are ornithocheirid, tapejarid and istiodactylid pterosaurs; spinosaurid, neovenatorid, and carcharodontosaurid theropods; rebbachisaurid sauropods; and birds.

Paleontological data from Charente thus do not support the hypothesis of a biological turnover at the Tithonian/Berriasian boundary, at least in Europe. This appears to have taken place between the end of the Berriasian and the Barremian, although the fossil record from this interval is very poorly known (Benson *et al.* 2013). Data on the latest Jurassic and earliest Cretaceous continental faunas suggest a reversion of the system boundary to Orbigny's (1841) and Oppel's (1865) historical position who defined the J/K boundary as Berriasian-Valanginian based on faunal turnover at that time (Granier 2019b; Enay 2020).

CONCLUSIONS

Angeac-Charente is an exceptional locality in many ways. In view of the extent and intensity of the fieldwork undertaken there and the number of fossils that have been recovered, it comprises one of the largest and most productive earliest Cretaceous localities in the world. Excavation areas will be further extended over the next decade, since 5000 m² of land has been made available by the Audoin et Fils company.

The vertebrate assemblage of Angeac-Charente consists of at least 40 vertebrate taxa, of which about 15 are represented by microvertebrate remains. The taxa that make up this very diverse fauna are synchronous, and autochtonous or parautochtonous. They comprise freshwater or terrestrial vertebrates that are clearly indicative of a continental wetland ecosystem, with no direct connection to the sea. The functioning and trophic dynamics of this ecosystem, and the associated paleoclimate, are being studied through multi-isotopic geochemical analyses. The taphocoenosis is dominated by an ornithomimosaur herd of at least 70 individuals, but turiasaur sauropods, pleurosternid turtles and goniopholidid crocodyliforms are also abundant. Detailed anatomical and taxonomic descriptions of this material are in preparation, and will be published according to the discoveries in the field, and the degree of completeness of the skeletons of the taxa concerned.

To date Angeac-Charente is the only known Lagerstätte from the Berriasian. It thus fills a gap in the fossil record, providing abundant and varied data on a continental wetland ecosystem at the beginning of the Cretaceous. Combined with data from the Charentese Chassiron and Cherves-de-Cognac localities, these data make it possible to follow the evolution of biodiversity, on a regional scale, at the Jurassic-Cretaceous boundary. The taxonomic composition of the Angeac-Charente locality shows strong similarities to Late Jurassic assemblages. It raises important questions regarding terrestrial faunal turnover across the Jurassic/Cretaceous boundary and the possible redefinition of the Jurassic/Cretaceous boundary placing it at the the base of the Valanginian stage.

Acknowledgements

This work is dedicated to the memory of Jean-Marie Audoin. We are indebted to the Audoin & Fils company for sharing their discovery with us and for providing us with incredible logistical support. We would like to thank Francette Rodet, the Audoin family and the department of La Charente for donating all of the fossil material discovered at Angeac-Charente to the Angoulême museum. We thank the department of La Charente, the community of communes of the Châteauneuf region, the community of agglomeration of Grand Cognac, and the Town Hall of Angoulême for their financial support. We thank Christian Dufront and all the inhabitants of Angeac-Charente for welcoming us every summer since 2011. Finally, we would like to thank the 191 diggers who have taken turns on the Angeac-Charente site every summer since August 2010. Steven Sweetman and Alexander Averianov are deeply thanked for their reviews of the submitted manuscript that prompted significant improvements. We also thank Dominique Abit for sharing his drone pictures with us.

REFERENCES

- ALLAIN R., TYKOSKI R., AQUESBI N., JALIL N. E., MONBARON M., RUSSELL D. & TAQUET P. 2007. — An abelisauroid (Dinosauria: Theropoda) from the Early Jurassic of the High Atlas Mountains, Morocco, and the radiation of ceratosaurs. *Journal of Vertebrate Paleontology* 27 (3): 610-624. https://doi.org/b4f5j9
- ALLAIN R., VULLO R., LEPRINCE A., NERAUDEAU D. & TOURNE-PICHE J.-F. 2011. — An ornithomimosaur-dominated bonebed from the Early Cretaceous of Southwestern France. *Journal of Vertebrate Paleontology* 31 (supp. 6): 61.
- ALLAIN R., ROZADA L. & VULLO R. 2013. A cross-perspective into the Early Cretaceous dinosaur faunas from Europe and Asia: new data on the dinosaurs from the Savannakhet Bassin (Laos) and from the lignitic bonebed of Angeac-Charente (France), *in* VIth International Symposium about Dinosaurs Palaeontology and their Environment, Salas de los Infantes, Spain.

- ALLAIN R., VULLO R., LE LOEUFF J. & TOURNEPICHE J.-F. 2014. European ornithomimosaurs: an undetected record. *Geologica Acta* 12 (2): 127-135. https://doi.org/10.1344/105.000002083
- ALLAIN R., ABIT D., ANQUETIN J., BENOIT R.-A., BOURGEAIS R., DAVIERO-GOMEZ V., DEBUYSSCHERE M., DETHAN I., GOEDERT J., GOMEZ B., GOUSSARD F., HANTZPERGUE P., MARTIN J., MAZAN, MAZIN J.-M., POUECH J., QVARNSTRÖM M., ROZADA L., TOURNEPICHE J.-F. & VULLO R. 2017. — Les géants du vignoble. Eidola, Angoulême, 256 p.
- ALLOUL T., RAGE J.-C., HAMDIDOUCHE R. & JALIL N.-E. 2018. First report on Cretaceous vertebrates from the Algerian Kem Kem Beds. A new procoelous salamander from the Cenomanian, with remarks on African Caudata. *Cretaceous Research* 84: 384-388. https://doi.org/10.1016/j.cretres.2017.11.019
- ANDRADE M. B. & HORNUNG J. J. 2011. A new look into the periorbital morphology of *Goniopholis* (Mesoeucrocodylia: Neosuchia) and related forms. *Journal of Vertebrate Paleontology* 31 (2): 352-368. https://doi.org/10.1080/02724634.2011.550353
- ANDRADE M. B., EDMONDS R., BENTON M. J. & SCHOUTEN R. 2011. — A new Berriasian species of *Goniopholis* (Mesoeucrocodylia, Neosuchia) from England, and a review of the genus: new *Goniopholis* from England. *Zoological Journal of the Linnean Society* 163: S66-S108. https://doi.org/10.1111/j.1096-3642.2011.00709.x
- ANQUETIN J. & ANDRÉ C. 2020. The last surviving Thalassochelydia—A new turtle cranium from the Early Cretaceous of the Purbeck Group (Dorset, UK). *PaleorXiv*: 7pa5c, version 3, peer-reviewed by PCI Paleo. https://doi.org/10.31233/osf. io/7pa5c
- ANQUETIN J., PÜNTENER C. & JOYCE W. G. 2017. A review of the fossil record of turtles of the clade *Thalassochelydia*. Bulletin of the Peabody Museum of Natural History 58: 317-369. https:// doi.org/10.3374/014.058.0205
- ARP G. & MENNERICH C. 2008. Ostracod assemblages, palaeoenvironment and cyclicity of Purbeck-type sediments of the Münder Formation (Lower Cretaceous, Hils Syncline, N-Germany). *Palaeogeography, Palaeoclimatology, Palaeoecology* 264 (3-4): 230-249. https://doi.org/10.1016/j.palaeo.2007.02.051
- AVERIANOV A. O., MARTIN T., EVANS S. E. & BAKIROV A. A. 2006. — First Jurassic Choristodera from Asia. *Naturwissenschaf*ten 93 (1): 46-50. https://doi.org/10.1007/s00114-005-0061-2
- AVERIANOV A. O., MARTIN T., LOPATIN A. V., SKUTSCHAS P. P., SCHELLHORN R., KOLOSOV P. & VITENKO D. 2018. — A high-latitude fauna of mid-Mesozoic mammals from Yakutia, Russia. *PLoS ONE* 13: 1-17. https://doi.org/10.1371/journal. pone.0199983
- BAMBACH R. K., KNOLL A. H., & WANG S. C. 2004. Origination, extinction, and mass depletions of marine diversity. *Paleobiology* 30 (4): 522-542. https://doi.org/cftg4t
- BARRETT P. M., CLARKE J. B., BRINKMAN D. B., CHAPMAN S. D. & ENSOM P. C. 2002. — Morphology, histology and identification of the 'granicones' from the Purbeck Limestone Formation (Lower Cretaceous: Berriasian) of Dorset, southern England. *Cretaceous Research* 23: 279-295. https://doi.org/10.1006/ cres.2002.1002
- BATTEN D. J. (ed.) 2011. *English Wealden Fossils*. The Palaeontological Association, London, 769 p.
- BENOIT R.-A., NERAUDEAU D. & MARTIN-CLOSAS C. 2017. A review of the Late Jurassic-Early Creta-ceous charophytes from the northern Aquitaine Basin in south-west France. *Cretaceous Research* 79: 199-213. https://doi.org/10.1016/j. cretres.2017.07.009
- BENSON R. B., MANNION P. D., BUTLER R. J., UPCHURCH P., GOSWAMI A. & EVANS S. E. 2013. — Cretaceous tetrapod fossil record sampling and faunal turnover: implications for biogeography and the rise of modern clades. *Palaeogeography, Palaeoclimatology, Palaeoecology* 372: 88-107. https://doi. org/10.1016/j.palaeo.2012.10.028

- BENTON M. J. & CLARK J. M. 1988. Archosaur phylogeny and the relationships of the Crocodylia, *in* BENTON M. J. (ed.), *The Phylogeny and Classification of the Tetrapods*. Volume 1. *Amphibians*, *Reptiles, Birds*. Clarendon Press, Oxford: 295-338. (Systematics Association; special volume 35A).
- BILLON-BRUYAT J.-P., MAZIN J.-M. & POUECH J. 2010. A stegosaur tooth (Dinosauria, Ornithischia) from the Early Cretaceous of southwestern France. *Swiss Journal of Geosciences* 103: 143-153. https://doi.org/10.1007/s00015-010-0028-y
- BLOWS W. T. 1987. The armoured dinosaur *Polacanthus foxi* from the Lower Cretaceous of the Isle of Wight. *Palaeontology* 30 (3): 557-580.
- BLOWS W. T. 2001. Dermal armor of the polacanthine dinosaurs, in CARPENTER K. (ed.), *The Armored Dinosaurs*. Indiana University Press, Bloomington and Indianapolis: 363-385.
- BLOWS W. T. & HONEYSETT K. 2014. New teeth of nodosaurid ankylosaurs from the Lower Cretaceous of Southern England. *Acta Palaeontologica Polonica* 59 (4): 835-842. https://doi. org/10.4202/app.2012.0131
- BONAPARTE J. F. 1990. New Late Cretaceous mammals from the Los Alamitos Formation, northern Patagonia. *National Geographic Research* 6 (1): 63-83.
- BONAPARTE J. F. 2002. New Dryolestida (Theria) from the Late Cretaceous of Los Alamitos, Argentina, and paleogeographical comments. *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen* 224: 339-371. https://doi.org/10.1127/njgpa/224/2002/339
- BRITT B. B., SCHEETZ R. D., WHITING M. F. & WILHITE D. R. 2017. — Moabosaurus utahensis, n gen., sp., a new sauropod from the Early Cretaceous (Aptian) of North America. Contributions from the Museum of Paleontology, University of Michigan 32: 189-243. http://hdl.handle.net/2027.42/136227
- BROSCHINSKI A. 1999 Ein Lacertilier (Scincomorpha, Paramacellodidae) aus dem Oberen Jura von Tendaguru (Tansania). *Mitteilungen aus dem Museum für Naturkunde zu Berlin, Geowissenschaftliche Reihe 2*: 155-58. https://doi.org/10.1002/ mmng.1999.4860020111
- BROSCHINSKI A. & SIGOGNEAU-RUSSELL D. 1996. Remarkable lizard remains from the Lower Cretaceous of Anoual (Morocco). Annales de Paléontologie 82: 147-175.
- BRONGNIART A. 1829. Tableau des terrains qui composent l'écorce du globe, ou essai sur la structure de la partie connue de la terre. G. Levrault, Paris, 435 p.
- BUFFETAUT E. 1975. Sur l'anatomie et la position systématique de *Bernissartia fagesii* Dollo, L., 1883, crocodilien du Wealdien de Bernissart, Belgique. *Bulletin de l'Institut royal des Sciences naturelles de Belgique* 51: 1-20.
- BUFFETAUT E. 1986. Une mâchoire de *Goniopholis* (Crocodylia, Mesosuchia) dans le Portlandien supérieur du Boulonnais. *Mémoires de la Société académique du Boulonnais* 1: 64-71.
- BUFFETAUT E. & FORD R. L. E. 1979. The crocodilian *Bernissartia* in the Wealden of the Isle of Wight. *Palaeontology* 22: 905-912.
- BUFFETAUT E. & HUTT S. 1980. Vectisuchus leptognathus, n. g. n. sp., a slender-snouted goniopholid crocodilian from the Wealden of the Isle of Wight. Neues Jahrbuch für Geologie und Paläontologie Monatshefte 7: 385-390. https://doi.org/10.1127/ njgpm/1980/1980/385
- BUFFETAUT E., POUIT D, RIGOLLET L. & ARCHAMBEAU J. P. 1989. Poissons et reptiles continentaux du Purbeckien de la région de Cognac (Charente). Bulletin de la Société géologique de France 5 (5): 1065-1069. https://doi.org/10.2113/gssgfbull.V.5.1065
- BUSCALIONI A. D., ALCALÁ L., ESPILEZ E. & MAMPEL L. 2013. European Goniopholididae from the Early Albian Escucha Formation in Ariño (Teruel, Aragón, Spain). *Spanish Journal of Palaeontology* 28: 103-122. https://doi.org/10.7203/sjp.28.1.17835
- BUSCALIONI Ä. D. & POYATO-ARIZA F. J. 2016. Las Hoyas: a unique Cretaceous ecosystem. Cretaceous Period: Biotic Diversity and Biogeography. *Bulletin of the New Mexico Museum of Natural History and Science* 71: 51-63.

- BUSCALIONI A. D. & SANZ J. L. 1990. The small crocodile Bernissartia fagesii from the Lower Cretaceous of Galve (Teruel, Spain). Bulletin de l'Institut royal des Sciences naturelles de Belgique 60: 129-150.
- BUTLER P. M. & SIGOGNEAU-RUSSELL D. 2016. Diversity of triconodonts in the Middle Jurassic of Great Britain. *Palaeontologia Polonica* 67: 35-65.
- CAMPOS-SOTO S., BENITO M. I., COBOS A., CAUS E., QUIJADA I. E., SUAREZ-GONZALEZ P., MAS R., ROYO-TORRES R. & ALCALÁ L. 2019. — Revisiting the age and palaeoenvironments of the Upper Jurassic-Lower Cretaceous? dinosaur-bearing sedimentary record of eastern Spain: implications for Iberian palaeogeography. *Journal of Iberian Geology* 45: 471-510. https://doi.org/10.1007/ s41513-019-00106-y.
- CARBALLIDO J. L. & SANDER P. M. 2014. Postcranial axial skeleton of *Europasaurus holgeri* (Dinosauria, Sauropoda) from the Upper Jurassic of Germany: implications for sauropod ontogeny and phylogenetic relationships of basal Macronaria. *Journal of Systematic Palaeontology* 12 (3): 335-387. https://doi.org/10.108 0/14772019.2013.764935
- CARPENTER K. & WILSON Y. 2008. A new species of *Camptosaurus* (Ornithopoda: Dinosauria) from the Morrison Formation (Upper Jurassic) of Dinosaur National Monument, Utah, and a biomechanical analysis of its forelimb. *Annals of Carnegie Museum* 76: 227-264. https://doi.org/fkbwqj
- CARRANO M. T., ORESKA M. P. J & LOCKWOOD R. 2016. Vertebrate paleontology of the Cloverly Formation (Lower Cretaceous), II: Paleoecology. *Journal of Vertebrate Paleontology* 36 (2): e1071265. https://doi.org/10.1080/02724634.2015.1071265
- CARRANO M. T., SAMPSON S. D. & FORSTER C. A. 2002. The osteology of *Masiakasaurus knopfleri*, a small abelisauroid (Dinosauria: Theropoda) from the Late Cretaceous of Madagascar. *Journal* of Vertebrate Paleontology 22: 510-534. https://doi.org/cvk6kb
- CARRANO M. T. & WILSON J. A. 2001. Taxon distributions and the tetrapod track record. *Paleobiology* 27: 564-582. https://doi. org/cn2c2m
- CASANOVAS M. L., SANTAFÉ J. V. & SANZ J. L. 2001. Losillasaurus giganteus, un nuevo saurópodo del tránsito Jurásico-Cretácico de la cuenca de "Los Serranos" (Valencia, España). *Paleontologia i Evolució* 32-33: 99-122.
- CERRONI M. A., AGNOLIN F. L., EGLI F. B. & NOVAS F. E. 2019. The phylogenetic position of *Afromimus tenerensis* Sereno, 2017 and its paleobiogeographical implications. *Journal of African Earth Sciences* 159: 103572. https://doi.org/10.1016/j.jafrearsci.2019.103572
- CHOINIERE J. N., FORSTER C. A. & DE KLERK W. J. 2012. New information on *Nqwebasaurus thwazi*, a coelurosaurian theropod from the Early Cretaceous Kirkwood Formation in South Africa. *Journal of African Earth Sciences* 71: 1-17. https://doi.org/10.1016/j. jafrearsci.2012.05.005
- CHOW M. & RICH T. H. V. 1984. A new triconodontan (Mammalia) from the Jurassic of China. *Journal of Vertebrate Paleontology* 3 (4): 226-231. https://www.jstor.org/stable/4522957
- CLARK J. M. 1986. Phylogenetic Relationships of the Crocodylomorph Archosaurs. Unpublished Ph.D, dissertation, University of Chicago, Chicago, 556 p.
- CLEMENS W. A. & MILLS J. R. E. 1971. Review of Peramus tenuirostris Owen (Eupantotheria, Mammalia). Bulletin of the British Museum (Natural History) 20: 89-113. https://www.biodiversitylibrary.org/page/36951234
- COBOS A., ROYO-TORRES R., LUQUE L., ALCALA L. & MAMPEL L. 2010. — An Iberian stegosaurs paradise: The Villar del Arzobispo Formation (Tithonian-Berriasian) in Teruel (Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* 293 (1-2): 223-236. https:// doi.org/10.1016/j.palaeo.2010.05.024
- COCUDE-MICHEL M. 1963. Les rhynchocéphales et les sauriens des calcaires lithographiques (Jurassique supérieur) d'Europe occidentale. *Nouvelles Archives du Muséum d'Histoire naturelle de Lyon* 7: 1-187.

- COLIN J.-P., EL ALBANI A., FÜRSICH F. T., MARTIN-CLOSAS C., MAZIN J.-M. & BILLON-BRUYAT J.-P. 2004. — Le gisement "Purbeckien" de vertébrés de Cherves-de-Cognac, Charente (SW France): nouvelles données biostratigraphiques. *Comptes Rendus Palevol* 3: 9-16. https://doi.org/10.1016/j.crpv.2003.09.013
- COPE E. D. 1875. Check-list of North American Batrachia and Reptilia with a systematic list of the higher groups and an essay on geographical distribution based on the specimens in the U.S. National Museum. *Bulletin of the United States National Museum* 1: 1-104. https://hdl.handle.net/10088/30460
- COPE E. D. 1868. On the origin of genera. Proceedings of the Academy of Natural Sciences of Philadelphia 1868: 242-300.
- COSTA F. & MATEUS O. 2019. Dacentrurine stegosaurs (Dinosauria): A new specimen of *Miragaia longicollum* from the Late Jurassic of Portugal resolves taxonomical validity and shows the occurrence of the clade in North America. *PloS one* 14(11): e0224263. https://doi.org/10.1371/journal.pone.0224263
- CROMPTON A. W. & JENKINS F. A. 1968. Molar occlusion in Late Triassic mammals. *Biological Reviews* 43 (4): 427-458. https://doi.org/10.1111/j.1469-185x.1968.tb00966.x CRONIN T. J. & SHIMADA K. 2019. — New anatomical informa-
- CRONIN T. J. & SHIMADA K. 2019. New anatomical information of the Late Cretaceous bony fish, *Micropycnodon kansasensis* (Actinopterygii: Pycnodontiformes), from the Niobrara Chalk of western Kansas, U.S.A. *Transactions of the Kansas Academy* of Science 122: 19-28. https://doi.org/10.1660/062.122.0102
- CUENCA-BESCÓS G. & CANUDO J. I. 2003. A new gobiconodontid mammal from the Early Cretaceous of Spain and its palaeogeographic implications. *Acta Palaeontologica Polonica* 48 (4): 575-582.
- DAVIS B. M. 2012. Micro-computed tomography reveals a diversity of Peramuran mammals from the Purbeck Group (Berriasian) of England. *Palaeontology* 55 (4): 789-817. https:// doi.org/10.1111/j.1475-4983.2012.01161.x
- D'EMIC M. D. 2012. The early evolution of titanosauriform sauropod dinosaurs. *Zoological Journal of the Linnean Society* 166: 624-671. https://doi.org/10.1111/j.1096-3642.2012.00853.x
- DE CISNEROS C. J. & VERA J. A. 1993. Milankovitch cyclicity in Purbeck peritidal limestones of the Prebetic (Berriasian, southern Spain). *Sedimentology* 40: 513-537. https://doi. org/10.1111/j.1365-3091.1993.tb01348.x
- DERCOURT J., GAETANI M., VRIELYNCK B., BARRIER E., BIJU-DUVAL B., BRUNET M. F., CADET J.-P., CRASQUIN S. & SANDULESCU M. 2000. — Atlas Peri-Tethys Palaeogeographical Maps. Commission de la carte géologique du monde, Paris, 24 p.
- DOLLO L. 1883. Première note sur les crocodiliens de Bernissart. Bulletin du Musée royal d'Histoire naturelle de Belgique 2: 309-338
- DUNKLE D. H. & HIBBARD C. H. 1946. Some comments upon the structure of a Pycnodontid fish from the Upper Cretaceous of Kansas. *The University of Kansas Science Bulletin* 39: 161-181.
- EBERT M. 2018. Cerinichthys koelblae, gen. et sp. nov., from the Upper Jurassic of Cerin, France, and its phylogenetic setting, leading to a reassessment of the phylogenetic relationships of Halecomorphi (Actinopterygii). Journal of Vertebrate Paleontology 38 (1): e1420071. https://doi.org/10.1080/027 24634.2017.1420071
- EBERTH D. A., SHANNON M. & NOLAND B. G. 2007. A bonebeds database: classification, biases, and patterns of occurrence, *in* ROGERS R. R., EBERTH D. A. & FIORILLO A. R. (eds), *Bonebeds: Genesis, Analysis, and Paleobiological Significance.* University of Chicago Press, Chicago, 499 p.
- EL ALBANI A., FÜRSICH F.T., COLIN J.-P., MEUNIER A., HOCHULI P., MARTIN-CLOSAS M., MAZIN J.-M. & BILLON-BRUYAT J.-P. 2004. — Palaeoenvironmental reconstruction of the basal Cretaceous vertebrate bearing beds in the Northern part of the Aquitaine Basin (SW France): sedimentological and geochemical evidence. *Facies* 50: 195-215. https://doi. org/10.1007/s10347-004-0017-6

- ENAY R. 2020. The Jurassic/Cretaceous system boundary is an impasse. Why do not go back to Oppel's 1865 original an historic definition of the Tithonian? *Cretaceous Research*, 106: 104241. https://doi.org/10.1016/j.cretres.2019.104241
- ENSOM P. C. & ŠIGOGNEAU-RUSSELL D. 2000. New symmetrodonts (Mammalia, Theria) from the Purbeck Limestone Group, Lower Cretaceous, southern England. *Cretaceous Research*, 21: 767-779. https://doi.org/10.1006/cres.2000.0227
- ESTES R. 1981. Gymnophiona, Caudata, *in* WELLNHOFER P. (ed.), *Encyclopedia of Paleoherpetology*. Part 2. Gustav Fischer, Stuttgart, 115 p.
- ESTES R. & HOFFSTETTER R. 1976. Les urodèles du Miocène de La Grive-Saint-Alban (Isère, France). *Bulletins du Muséum national d'Histoire naturelle, 3*ème série, *Sciences de la Terre* 398: 297-343. https://www.biodiversitylibrary.org/page/55505209
- EVANS J. & KEMP T. S. 1975. The cranial morphology of a new Lower Cretaceous turtle from Southern England. *Palaeontology* 18 (1): 25-40.
- EVANS S. E. 1991. The postcranial skeleton of the choristodere *Cteniogenys* (Reptilia: Diapsida) from the Middle Jurassic of England. *Geobios* 24 (2): 187-99. https://doi.org/10.1016/ S0016-6995(91)80006-L
- EVANS S. E. & CHURE D. C. 1998. Paramacellodid lizard skulls from the Jurassic Morrison Formation at Dinosaur national monument, Utah. *Journal of Vertebrate Paleontology* 18 (1): 99-114. https://www.jstor.org/stable/4523876
- EVANS S. E. & FRASER N. C. 1992. A sphenodontid jaw (Reptilia: Lepidosauria) from the Upper Jurassic of Dorset. Proceedings of the Dorset Natural History Society 1992: 199-200.
- EVANS S. E. & MILNER Å. R. 1994. Microvertebrate faunas from the Middle Jurassic of Britain, *in* FRASER N. & SUES H.-D. (eds), *In the Shadow of the Dinosaurs: Early Mesozoic Tetrapods*. CambridgeUniversity Press, Cambridge: 303-321.
- EVANS S. E. & SEARLE B. 2002. Lepidosaurian reptiles from the Purbeck Limestone Group of Dorset, southern England. *Special Papers in Palaeontology* 68: 145-159.
- EVANS S. E. & SIGOGNEAU-RUSSELL D. 1997. New sphenodontians (Diapsida: Lepidosauria: Rhynchocephalia) from the Early Cretaceous of North Africa. *Journal of Vertebrate Paleontology* 17 (1): 45-51. https://www.jstor.org/stable/4523785
- EVERS S. W., ROLLOT Y. & JOYCE W. G. 2020. Cranial osteology of the Early Cretaceous turtle *Pleurosternon bullockii* (Paracryptodira: Pleurosternidae). *PeerJ* 8: e9454. https://doi.org/10.7717/peerj.9454
- FLYNN J. J., FOX S. R., PARRISH J. M., RANIVOHARIMANANA L. & WYSS A. R. 2006. — Assessing diversity and paleoecology of a Middle Jurassic microvertebrate assemblage from Madagascar. *New Mexico Museum of Natural History and Science Bulletin* 37: 476-489.
- FOSTER J. R. 2018. A new atoposaurid crocodylomorph from the Morrison Formation (Upper Jurassic) of Wyoming, USA. *Geology* of the intermountain West 5: 287-295. https://doi.org/10.31711/ giw.v5.pp287-295
- FOTH C., RABI M. & JOYCE W. G. 2017. Skull shape variation in extant and extinct Testudinata and its relation to habitat and feeding ecology. Acta Zoologica 98: 310-325. https://doi.org/10.1111/ azo.12181
- FOX R. C. & NAYLOR M. 1982. A reconsideration of the relationships of the fossil amphibian *Albanerpeton. Canadian Journal* of *Earth Sciences* 19: 118-128. https://doi.org/10.1139/e82-009
- GAFFNEY E. S. 1975. A phylogeny and classification of the higher categories of turtles. *Bulletin of the American Museum of Natural History* 155 (5): 387-436. http://hdl.handle.net/2246/614
- History 155 (5): 387-436. http://hdl.handle.net/2246/614 GAFFNEY E. S. 1979. — The Jurassic turtles of North America. Bulletin of the American Museum of Natural History 162 (3): 91-136. http://hdl.handle.net/2246/1275
- GALTON P. M. 1974. The ornithischian dinosaur Hypsilophodon from the Wealden of the Isle of Wight. Bulletin of the British Museum (Natural History) Geology 25 (1): 1-152. https://www. biodiversitylibrary.org/page/2239368

- GALTON P. M. 1980. Armoured dinosaurs (Ornithischia: Ankylosauria) from the Middle and Upper Jurassic of England. *Geobios* 13 (6): 825-837. https://doi.org/10.1016/S0016-6995(80)80038-6
- GALTON P. M. 1982. *Elaphrosaurus*, an ornithomimid dinosaur from the Upper Jurassic of North America and Africa. *Paläontologische Zeitschrift* 56: 265-275. https://doi.org/10.1007/ BF02988803
- GALTON P. M. 1983. Armoured dinosaurs (Ornithischia: Ankylosauria) from the Middle and Upper Jurassic of Europe. *Palaeontographica, Abteilung A* 182: 1-25.
- GALTON P. M. 1985. British plated dinosaurs (Ornithischia, Stegosauridae). *Journal of Vertebrate Paleontology* 5 (3): 211-254. https://doi.org/10.1080/02724634.1985.10011859
- GALTON P. M. 2009. Notes on Neocomian (Lower Cretaceous) ornithopod dinosaurs from England – *Hypsilophodon, Valdosaurus, "Camptosaurus", "Iguanodon"* – and referred specimens from Romania and elsewhere. *Revue de Paléobiologie* 28: 211-273.
- GALTON P. M. & POWELL H. P. 1980. The ornithischian dinosaur *Camptosaurus prestwichii* from the Upper Jurassic of England. *Palaeontology* 23 (2): 411-443.
- GARDNER J. D. 1999a. Redescription of the geologically youngest albanerpetontid (?Lissamphibia): Albanerpeton inexpectatum Estes and Hoffstetter, 1976, from the Miocene of France. Annales de Paléontologie 85: 57-84. https://doi.org/10.1016/ S0753-3969(99)80008-1
- GARDNER J. D. 1999b. The amphibian Albanerpeton arthridion and the Aptian – Albian biogeography of albanerpetontids. Palaeontology 42:529-544. https://doi.org/10.1111/1475-4983.00083
- GARDNER J. D. 2000. Revised taxonomy of albanerpetontid amphibians. *Acta Palaeontologica Polonica* 45 (1): 55-70.
- GARDNER J. D., EVANS S. E. & SIGOGNEAU-RUSSELL D. 2003. New albanerpetontid amphibians from the Early Cretaceous of Morocco and Middle Jurassic of England. *Acta Palaeontologica Polonica* 48 (2): 301-319.
- GASCA J. M., BADIOLA A., CANUDO J. I., MORENO-AZANZA M. & PUÉRTOLAS E. 2012. — La asociación de vertebrados fósiles del yacimiento Pochancalo 1 (Valanginiense-Hauteriviense, Villanueva de Huerva, Zaragoza, España), *in* Actas de V Jornadas Internacionales sobre Paleontología de Dinosaurios y su Entorno. Salas de los Infantes, Burgos: 159-172.
- GHEERBRANT E. & RAGE J.-C. 2006. Paleobiogeography of Africa: how distinct from Gondwana and Laurasia? *Palaeogeography, Palaeoclimatology, Palaeoecology* 241: 224-246. https:// doi.org/10.1016/j.palaeo.2006.03.016
- GILMORE C. W. 1910 A new rhynchocephalian reptile from the Jurassic of Wyoming, with notes on the fauna of "Quarry 9". *Proceedings of the United States National Museum* 37: 35-42. https://doi.org/10.5479/si.00963801.37-1698.35
- GODEFROIT P. & BATTAIL B. 1997. Late Triassic cynodonts from Saint-Nicolas-de-Port (north-eastern France). *Geodiversitas* 19 (3): 567-631.
- GÓNET J., ROZADA L., BOURGEAIS R. & ALLAIN R. 2019. Taphonomic study of a pleurosternid turtle shell from the Early Cretaceous of Angeac-Charente, southwest France. *Lethaia* 52: 232-243. https://doi.org/10.1111/let.12309
- GRANDE L. 2010. An empirical synthetic pattern study of gars (Lepisosteiformes) and closely related species, based mostly on skeletal anatomy. The resurrection of Holostei. *Copeia* 10 (2A): 1-871. https://www.jstor.org/stable/20787269
- GRANDE L. & BEMIS W. E. 1998. A comprehensive phylogenetic study of amiid fishes (Amiidae) based on comparative skeletal anatomy. An empirical search for interconnected patterns of natural history. *Journal of Vertebrate Paleontology* 18 (suppl. 1): 1-696.
- GRANIER B. 2019a. JK2018: International Meeting around the Jurassic/Cretaceous Boundary – Chairperson's Report. *Volumina Jurassica* 17: 1-6.

- GRANIER B. 2019b. Dual biozonation scheme (benthic foraminifera and "calcareous" green algae) over the Jurassic-Cretaceous transition. Another plea to revert the system boundary to its historical Orbigny's and Oppel's definition. *Cretaceous Research* 93: 245-274. https://doi.org/10.1016/j.cretres.2018.08.024
- GUERRERO A. & PÉREZ-GARCÍA A. 2020 On the validity of the British Upper Jurassic turtle "*Pleurosternon portlandicum*" (Paracryptodira, Pleurosternidae). *Journal of Iberian Geology* 46: 419-429. https://doi.org/10.1007/s41513-020-00136-x
- HADDOUMI H., ALLAIN R., MESLOUH S., MÉTAIS G., MONBARON M., PONS D., RAGE J.-C., VULLO R., ZOUHRI S. & GHEERBRANT E. 2016. — Guelb el Ahmar (Bathonian, Anoual Syncline, eastern Morocco): first continental flora and fauna including mammals from the Middle Jurassic of Africa. *Gondwana Research* 29: 290-319. https://doi.org/10.1016/j.gr.2014.12.004
- HAHN G. & HAHN R. 1999. Pinheirodontidae n. fam. (Multituberculata) (Mammalia) aus der tiefen Unter-Kreide Portugals. *Palaeontographica Abteilung A*: 77-222.
- HALLAM A. 1986. The Pliensbachian and Tithonian extinction events. Nature 319: 765-768. https://doi.org/10.1038/319765a0
- HALLAM A. 2001. A review of the broad pattern of Jurassic sealevel changes and their possible causes in the light of current knowledge. *Palaeogeography, Palaeoclimatology, Palaeoecology* 167: 23-37. https://doi.org/10.1016/S0031-0182(00)00229-7
- HALLAM A. & WIGNALL P. B. 1997. Mass Extinctions and their Aftermath. Oxford University Press, Oxford, 328 p.
- HENDRICKX C., MATEUS O. & ARAÚJO R. 2015a. A proposed terminology of theropod teeth (Dinosauria, Saurischia). *Journal* of Vertebrate Paleontology 35: e982797. https://doi.org/10.108 0/02724634.2015.982797.
- HENDRICKX C., MATEUS O. & ARAÚJO R. 2015b. The dentition of megalosaurid theropods. *Acta Palaeontologica Polonica* 60: 627-642. https://doi.org/10.4202/app.00056.2013
- HIRAYAMA R., BRINKMAN D. B. & DANILOV I. G. 2000. Distribution and biogeography of non-marine Cretaceous turtles. *Russian Journal of Herpetology* 7: 181-198.
- HOFFSTETTER R. 1967. Coup d'œil sur les sauriens (= lacertiliens) des couches de Purbeck (Jurassique supérieur d'Angleterre). *Colloques internationaux du Centre national de la Recherche scientifique* 163: 349-371.
- HOLTZ T. R. 1994. The phylogenetic position of the Tyrannosauridae: implications for theropod systematics. *Journal* of *Paleontology* 68: 1100-1117. https://doi.org/10.1017/ S0022336000026706
- HOLTZ T. R. 2004. Tyrannosauroidea, in WEISHAMPEL D. B., DODSON P. & OSMÓLSKA H. (eds), *The Dinosauria*. University of California Press, Berkeley: 111-136.
- HONE D. W. E., TISCHLINGER H., FREY E. & RÖPER M. 2012. A new non-pterodactyloid pterosaur from the Late Jurassic of southern Germany. *PLoS ONE* 7(7): e39312. https://doi. org/10.1371/journal.pone.0039312
- HOPSON J. A. 1994. Synapsid Evolution and the Radiation of Non-Eutherian Mammals, in SPENCER R. S. (ed.), Major Features of Vertebrate Evolution. Short Courses in Paleontology, Cambridge University Press, Cambridge: 109-219.
- HULKE J. W. 1887. VII. Supplemental note on *Polacanthus foxii*, describing the dorsal shield and some parts of the endoskeleton, imperfectly known in 1881. *Philosophical Transactions of the Royal Society of London* B 178: 169-172.
- HUTT S., NAISH D., MARTILL D. M., BARKER M. J. & NEWBERRY P. 2001. — A preliminary account of a new tyrannosauroid theropod from the Wessex Formation (Early Cretaceous) of southern England. *Cretaceous Research* 22: 227-242. https:// doi.org/10.1006/cres.2001.0252
- JÄGER K. R. K., CIFELLI R. L. & MARTIN T. 2021. Tooth eruption in the Early Cretaceous British mammal *Triconodon* and description of a new species. *Papers in Palaeontology* 7 (2): 1065-1080. https://doi.org/10.1002/spp2.1329

- JANENSCH W. 1925. Die Coelurosaurier und Theropoden der Tendaguru-Schichten Deutsch-Ostafrikas. *Palaeontographica* 7 (suppl. I): 1-99.
- JOYCE W. G. 2017. A Review of the Fossil Record of Basal Mesozoic Turtles. *Bulletin of the Peabody Museum of Natural History* 58: 65-113. https://doi.org/10.3374/014.058.0105
- JOYCE W. G. & ANQUETIN J. 2019. A review of the fossil record of non-baenid turtle of the clade *Paracryptodira*. Bulletin of the Peabody Museum of Natural History 60: 129-155. https://doi. org/10.3374/014.060.0204
- JOYCE W. G., CHAPMAN S. D., MOODY R. T. J. & WALKER C. A. 2011. The skull of the solemydid turtle *Helochelydra nopcsai* from the Early Cretaceous of the Isle of Wight (UK) and a review of Solemydidae. *Special Papers in Palaeontology* 86: 75-97.
- JOYCE W. G., RABI M., CLARK J. M. & XU X. 2016. A toothed turtle from the Late Jurassic of China and the global biogeographic history of turtles. *BMC Evolutionary Biology* 16: 236. https://doi.org/10.1186/s12862-016-0762-5
- JOYCE W. G., STERLI J. & CHAPMAN S. D. 2014. The skeletal morphology of the solemydid turtle *Naomichelys speciosa* from the Early Cretaceous of Texas. *Journal of Paleontology* 88 (6): 1257-1287. https://doi.org/10.1666/14-002
- KELLNER A. W. A., CALDWELL M. W., HOLGADO B., DALLA VECCHIA F. M., NOHRA R., SAYÃO J. M. & CURRIE P. J. 2019. — First complete pterosaur from the Afro-Arabian continent: insight into pterodactyloid diversity. *Scientific Reports* 9 (1): 1-9. https://doi. org/10.1038/s41598-019-54042-z
- KIELAN-JAWOROWSKA Z., CIFELLI R. L. & LUO Z.-X. 2004. Mammals from the Age of Dinosaurs: Origins, Evolution, and Structure. Columbia University Press, New York, 640 p.
- KIELAN-JAWOROWSKA Z. & ENSOM P. C. 1992. Multituberculate mammals from the Upper Jurassic Purbeck Limestone Formation of southern England. *Palaeontology* 35 (1): 95-126.
- KLEIN I. T. 1760. Klassification und kurze Geschichte der Vierfüßigen Thiere [translation by F. D. Behn]. Jonas Schmidt, Lübeck, 381 p.
- KOKEN E. 1887. Die Dinosaurier, Crocodiliden und Sauropterygier des norddeutschen Wealden. *Palaeontologische Abhandlungen* 3: 309-419.
- KRAUSE D. W., EVANS S. E. & GAO K.-Q. 2003. First definitive record of Mesozoic lizards from Madagascar. *Journal of Vertebrate Paleontology* 23: 842-856. https://www.jstor.org/stable/4524386
- LAPPARENT DE BROIN F. DE & MURELAGA X. 1996. Une nouvelle faune de chéloniens dans le Crétacé supérieur européen. *Comptes Rendus de l'Académie des Sciences, Série IIa* 323: 729-735.
- LAPPARENT DE BROIN F. DE & MURELAGA X. 1999. Turtles from the Upper Cretaceous of Laño (Iberian Peninsula). Estudios del Museo de Ciencias Naturales de Alava 14: 135-211.
- LASSERON M., ALLAIN R., GHEERBRANT E., HADDOUMI H., JALIL N.-E., MÉTAIS G., RAGE J.-C., VULLO R. & ZOUHRI S. 2020. — New data on the microvertebrate fauna from the Upper Jurassic or lowest Cretaceous of Ksar Metlili (Anoual Syncline, eastern Morocco). *Geological Magazine* 157 (3): 367-392. https://doi. org/10.1017/S0016756819000761
- LAUPRASERT K., LAOJUMPON C., SAENPHALA W., CUNY G., THIRA-KUPT K. & SUTEETHORN V. 2011. — Atoposaurid crocodyliforms from the Khorat group of Thailand: first record of *Theriosuchus* from Southeast Asia. *Paläontologische Zeitschrift* 85: 37-47. https:// doi.org/10.1007/s12542-010-0071-z
- LINDGREN J., REES J., SIVERSON M. & CUNY G. 2004. The first Mesozoic mammal from Scandinavia. *Geologiska Föreningens Förhandlingar* 126: 325-330. https://doi.org/10.1080/11035890401264325
- LONGRICH N. 2008. A new, large ornithomimid from the Cretaceous Dinosaur Park Formation of Alberta, Canada: implications for the study of dissociated dinosaur remains. *Palaeontology* 51 (4): 983-997. https://doi.org/10.1111/j.1475-4983.2008.00791.x
- LÓPEZ-ARBARELLO A. & SFERCO E. 2018. Neopterygian phylogeny: the merger assay. *Royal Society Open Science* 5: 172337. https://doi.org/10.1098/rsos.172337

- LOUCHART A. & POUECH J. 2017. A tooth of Archaeopterygidae (Aves) from the Lower Cretaceous of France extends the spatial and temporal occurrence of the earliest birds. *Cretaceous Research* 73: 40-46. https://doi.org/10.1016/j.cretres.2017.01.004
- LÜ J., XU L. & JI Q. 2008. Restudy of *Liaoxipterus* (Istiodactylidae: Pterosauria), with comments on the Chinese istiodactylid pterosaurs. *Zitteliana B* 28: 229-241. https://doi.org/10.5282/ ubm/epub.12018
- LYDEKKER R. 1889. Catalogue of the Fossil Reptilia and Amphibia in the British Museum (Natural History). Part III, The Order Chelonia. Trustees of the British Museum, London, 239 p.
- LYSON T. R. & JOYCE W. G. 2011. Cranial anatomy and phylogenetic placement of the enigmatic turtle *Compsemys victa* Leidy, 1856. *Journal of Paleontology* 85 (4): 789-801. https:// doi.org/10.1666/10-081.1
- MAIDMENT S. C., NORMAN D. B., BARRETT P. M. & UPCHURCH P. 2008. — Systematics and phylogeny of Stegosauria (Dinosauria: Ornithischia). *Journal of Systematic Palaeontology* 6: 367-407. https://doi.org/10.1017/S1477201908002459
- MANNION P. D., ALLAIN R. & MOINE O. 2017. The earliest known titanosauriform sauropod dinosaur and the evolution of Brachiosauridae. *PeerJ* 5: e3217. https://doi.org/10.7717/ peerj.3217
- MANNION P. D., UPCHURCH P., SCHWARZ D. & WINGS O. 2019. Taxonomic affinities of the putative titanosaurs from the Late Jurassic Tendaguru Formation of Tanzania: phylogenetic and biogeographic implications for eusauropod dinosaur evolution. *Zoological Journal of the Linnean Society*, 185: 784-909. https:// doi.org/10.1093/zoolinnean/zly068
- MANSELL-PLEYDELL J. C. 1888. Fossil reptiles of Dorset. Proceedings of the Dorset Natural History and Antiquarian Field Club 9: 1-40.
- MANTELL G. A. 1844. *The Medals of Creation; or, First Lessons in Geology, and in the Study of Organic Remains.* Vol. 2. Henry G. Bohn, London.
- MARSH O. C. 1895. Art LV. On the Affinities and Classification of the Dinosaurian Reptiles. *American Journal of Science* (1880-1910) 50: 483-498. https://doi.org/10.2475/ajs.s3-50.300.483
- MARTIN J. E., LAUPRASERT K., BUFFETAUT É., LIARD R. & SUTEETHORN V. 2014a. — A large pholidosaurid in the Phu Kradung Formation of north-eastern Thailand. *Palaeontology* 57 (4): 757-769. https://doi.org/10.1111/pala.12086
- MARTIN J. E., RABI M., CSIKI-SAVA Z. & VASILE S. 2014b. Cranial morphology of *Theriosuchus sympiestodon* (Mesoeucrocodylia, Atoposauridae) and the widespread occurrence of *Theriosuchus* in the Late Cretaceous of Europe. *Journal of Paleontology* 88 (3): 444-456. https://doi.org/10.1666/13-106
- MARTIN T., MARUGÁN-LOBÓN J., VULLO R., MARTIN-ABAD H., LUO Z. X. & BUSCALIONI A. D. 2015. — A Cretaceous eutriconodont and integument evolution in early mammals. *Nature* 526: 380-384. https://doi.org/10.1038/nature14905
- MARTIN J. E., DELFINO M. & SMITH T. 2016a. Osteology and affinities of Dollo's goniopholidid (Mesoeucrocodylia) from the Early Cretaceous of Bernissart, Belgium. *Journal of Vertebrate Paleontology* 36 (6): e1222534. https://doi.org/10.1080/0272 4634.2016.1222534
- MARTIN J. E., RASLAN-LOUBATIÉ J. & MAZIN J.-M. 2016b. Cranial anatomy of *Pholidosaurus purbeckensis* from the Lower Cretaceous of France and its bearing on pholidosaurid affinities. *Cretaceous Research* 66: 43-59. https://doi.org/10.1016/j. cretres.2016.05.008
- MARTIN J. E., SMITH T., SALAVIALE C., ADRIEN J. & DELFINO M. 2020. — Virtual reconstruction of the skull of *Bernissartia fagesii* and current understanding of the neosuchian-eusuchian transition. *Journal of Systematic Palaeontology* 18 (13): 1079-1101. https://doi.org/10.1080/14772019.2020.1731722
- MARTIN T. 1999. Dryolestidae (Dryolestoidea, Mammalia) aus dem Oberen Jura-von Portugal. Abhandlungen der senckenbergischen naturforschenden Gesellschaft 550: 1-119.

- MARTIN T. 2002. New stem lineage representatives of Zatheria (Mammalia) from the Late Jurassic of Portugal. *Journal of Vertebrate Paleontology* 22: 322-348. https://doi.org/cb2w55
- MATEUS O. 2006. Late Jurassic dinosaurs from the Morrison Formation (USA), the Lourinha and Alcobaça formations (Portugal), and the Tendaguru Beds (Tanzania): a comparison. *New Mexico Museum of Natural History and Science Bulletin* 36: 223-231.
- MATEUS O., MAIDMENT S. C. & CHRISTIANSEN N. A. 2009. A new long-necked 'sauropod-mimic'stegosaur and the evolution of the plated dinosaurs. *Proceedings of the royal society B: biological sciences* 276: 1815-1821. https://doi.org/10.1098/rspb.2008.1909
- MATSUMOTO R., BUFFETAUT È., ESCUILLIÉ F, HERVET S. & EVANS S. E. 2013. — New material of the choristodere *Lazarussuchus* (Diapsida, Choristodera) from the Paleocene of France. *Journal* of Vertebrate Paleontology 33 (2): 319-39. https://doi.org/10.10 80/02724634.2012.716274
- MATSUMOTO R. & EVANS S. E. 2018. The first record of albanerpetontid amphibians (Amphibia: Albanerpetontidae) from East Asia. *PLoS ONE* 13: e0189767. https://doi.org/10.1371/ journal.pone.0189767
- MATSUMOTO R., MANABE M. & EVANS S. E. 2015. The first record of a long-snouted choristodere (Reptilia, Diapsida) from the Early Cretaceous of Ishikawa Prefecture, Japan. *Historical Biology* 27: 583-594. https://doi.org/10.1080/08912963.2014.898296
- MAZIN J.-M., POUECH J., HANTZPERGUE P. & LENGLET T. 2008. The Purbeckian site of Cherves-de-Cognac (Berriasian, Early Cretaceous, SW France): a first synthesis. *Documents des Laboratoires de Géologie de Lyon, Mid-Mesozoic Life and Environments Symposium* 164: 68-71.
- McGOWAN G. J. 1996. Albanerpetontid amphibians from the Jurassic (Bathonian) of southern England. *Museum of Northern Arizona Bulletin* 60: 227-234.
- MCGOWAN G. J. 1998. The development and function of the atlanto-axial joint in albanerpetontid amphibians. *Journal of Herpetology* 32: 116-122. https://doi.org/10.2307/1565490
- MCGOWAN G. J. 2002. Albanerpetontid amphibians from the Lower Cretaceous of Spain and Italy: a description and reconsideration of their systematics. *Zoological Journal of the Linnean Society* 135: 1-32. https://doi.org/10.1046/j.1096-3642.2002.00013.x
- MCGOWAN G. & ENSOM P. C. 1997. Albanerpetontid amphibians from the Lower Cretaceous of the Isle of Purbeck, Dorset. *Proceedings of the Dorset Natural History and Archaeological Society* 118: 113-117.
- McGowan G. J. & Evans S. E. 1995. Albanerpetontid ampibians from the Cretaceous of Spain. *Nature* 373: 143-145. https:// doi.org/10.1038/373143a0
- MCPHEE B. W., MANNION P. D., DE KLERK W. J. & CHOINIERE J. N. 2016. — High diversity in the sauropod dinosaur fauna of the Lower Cretaceous Kirkwood Formation of South Africa: implications for the Jurassic-Cretaceous transition. *Cretaceous Research* 59: 228-248. https://doi.org/10.1016/j.cretres.2015.11.006
- MEYER H. VON 1841. *Pholidosaurus schaumburgensis*, ein saurus aus dem sandstein der Wald-Formation Nord-Deutschlands. *Neues Jahrbuch für Mineralogie, Geologie, Geognosie und Petrefaktenkunde* 4: 443-445. https://www.biodiversitylibrary.org/page/35788547
- MILNER A. C. 2002. Theropod dinosaurs of the Purbeck limestone group, Southern England. *Special Papers in Palaeontology* 68: 191-202.
- MILNER A. R. 1988. The relationships and origin of living amphibians, in BENTON M. J. (ed.), The Phylogeny and Classification of the Tetrapods. Volume 1. Amphibians, Reptiles, Birds. Clarendon Press, Oxford, 380 p.
- MILNER A. R. 2004. The turtles of the Purbeck Limestone Group of Dorset, southern England. *Palaeontology* 47 (6): 1441-1467. https://doi.org/10.1111/j.0031-0239.2004.00418.x
- MILNER A. R. & BATTEN D. J. 2002. Life and Environments in Purbeck Times. *Special Papers in Palaeontology* 68. The Palaeontological Association, London, 268 p.

- MILNER A. R., EDMONDS R. P. & GODDEN M. 2012. The turtle genus *Hylaeochelys* from the Portland Stone Formation of Dorset. *Symposium of Vertebrate Palaeontology and Comparative Anatomy*: 28-29.
- MOCHO P., ROYO-TORRES R., MALAFAIA E., ESCASO F. & ORTEGA F. 2017. — Sauropod tooth morphotypes from the Upper Jurassic of the Lusitanian basin (Portugal). *Papers in Palaeontology* 3: 259-295. https://doi.org/10.1002/spp2.1075
- MOCHO P., ROYO-TORRES R., ORTEGA F., MALAFAIA E., ESCASO F. & SILVA B. 2016. — Turiasauria-like teeth from the Upper Jurassic of the Lusitanian Basin, Portugal. *Historical Biology* 28 (7): 861-880. https://doi.org/10.1080/08912963.2015.1049948
- MONTELLANO M., HOPSON J. A. & CLARK J. M. 2008. Late Early Jurassic mammaliaforms from Huizachal Canyon, Tamaulipas, México. *Journal of Vertebrate Paleontology* 28 (4): 1130-1143. https://doi.org/10.1671/0272-4634-28.4.1130
- NÉRAUDEAU D., ALLAIN R., BALLÈVRE M., BATTEN D. J., BUFFE-TAUT E., COLIN J. P., DABARD M. P., DAVIERO-GOMEZ V., EL ALBANI A., GOMEZ B., GROSHENY D., LE LŒUFF J., LEPRINCE A., MARTIN-CLOSAS C., MASURE E., MAZIN J.-M., PHILIPPE M., POUECH J., TONG H., TOURNEPICHE J.-F. & VULLO R. 2012. — The Hauterivian-Barremian lignitic bone bed of Angeac (Charente, SW France): stratigraphical, palaeobiological and palaeogeographical implications. *Cretaceous Research* 37: 1-14. https://doi.org/10.1016/j.cretres.2012.01.006
- NOPCSA F. 1928. Palaeontological notes on reptiles. *Geologica* Hungarica, Palaeontology Series 1: 1-84.
- NORELL M. A., CLARK J. M., TURNER A. H., MAKOVICKY P. J., BARSBOLD R. & ROWE T. 2006. — A new dromaeosaurid theropod from Ukhaa Tolgod (Ömnögov, Mongolia). *American Museum Novitates* 3545: 1-51. https://doi.org/d4f9k8
- NORMAN D. B. & BARRETT P. M. 2002. Ornithischian dinosaurs from the lower Cretaceous (Berriasian) of England. *Special Papers in Palaeontology* 68: 161-190.
- NORMAN D. B., SUES H.-D., WITMER L. M. & CORIA R. A. 2004. — Basal Ornithopoda, *in* WEISHAMPEL D. B., DODSON P. & OSMÓLSKA H. (eds), *The Dinosauria*. Edition 2. University of California Press, Berkeley, 880 p.
- NYDAM R. L. & CIFELLI R. L. 2002. Lizards from the Lower Cretaceous (Aptian-Albian) Antlers and Cloverly formations. *Journal of Vertebrate Paleontology* 22 (2): 286-298. https://doi. org/bc5x7j
- NYDAM R. L., ROWE T. B. & CIFELLI R. L. 2013. Lizards and snakes of the Terlingua Local Fauna (late Campanian), Aguja Formation, Texas, with comments on the distribution of paracontemporaneous squamates throughout the Western Interior of North America. *Journal of Vertebrate Paleontology* 33 (5): 1081-1099. https://doi.org/10.1080/02724634.2013.760467
- O'CONNOR P. M. 2007. The postcranial axial skeleton of *Majungasaurus crenatissimus* (Theropoda: Abelisauridae) from the Late Cretaceous of Madagascar. *Journal of Vertebrate Paleontology* 27: 127-163. https://doi.org/bcbvhp
- OPPEL A. 1865. Die Tithonische Etage. Zeitschrift der Deutschen Geologischen Gesellschaft 17: 535-558.
- ORBIGNY A. D. D' 1841. Paléontologie française, terrains crétacés. Tome I. Céphalopodes. Masson, Paris. https://doi.org/10.5962/ bhl.title.50510
- OSMOLSKA H., RONIEWICZ E. & BARSBOLD R. 1972. A new dinosaur, *Gallimimus bullatus* n. gen, n. sp. (Ornithomimidae) from the Upper Cretaceous of Mongolia. *Palaeontologica Polonica* 27: 103-143.
- OWEN R. 1842. Report on British fossil reptiles, part II. Report for the British Association for the Advancement of Science 11: 60-204. https://www.biodiversitylibrary.org/page/33377524
- OWEN R. 1853. Monograph on the Fossil Reptilia of the Wealden and Purbeck Formations. Part 1. Chelonia. Cambridge University Press, Cambridge: 1-12 (Monograph of the Palaeontological Society; 7). https://doi.org/10.1017/CBO9781316146286.003

- OWEN R. 1871. Monograph of the Fossil Mammalia of the Mesozoic Formations. The Palæontographical society, London: 1-115 (Monograph of the Palaeontological Society; 24). https://doi. org/10.5962/bhl.title.14831
- OWEN R. 1875. Monographs on the Fossil Reptilia of the Mesozoic Formations. Part II. (Genera Bothriospondylus, Cetiosaurus, Omosaurus). The Palæontographical society, London: 15-93 (Monograph of the Palaeontological Society; 29).
- OWEN R. 1879. On the association of dwarf crocodiles (Nannosuchus and Theriosuchus pusillus, eg) with the diminutive mammals of the Purbeck Shales. Quarterly Journal of the Geological Society 35: 148-155. https://doi.org/10.1144/GSL.JGS.1879.035.01-04.02
- ÖZDIKMEN H. 2009. Substitute names for some unicellular animal taxa (Protozoa). *Munis Entomology & Zoology* 4: 233e256.
- PEREA D., SOTO M., TORIÑO P., MESA V. & MAISEY J. G. 2018. A Late Jurassic-?earliest Cretaceous ctenochasmatid (Pterosauria, Pterodactyloidea): The first report of pterosaurs from Uruguay. *Journal of South American Earth Sciences* 85: 298-306. https://doi. org/10.1016/j.jsames.2018.05.011
- PEREDA-SUBERBIOLA J. 1994. *Polacanthus* (Ornithischia, Ankylosauria), a transatlantic armoured dinosaur from the Early Cretaceous of Europe and North America. *Palaeontographica Abteilung* A: 133-159.
- PÉREZ-GARCÍA A. 2012. High diversity of pancryptodiran turtles in the Lower Cretaceous of Europe. *Cretaceous Research* 36: 67-82. https://doi.org/10.1016/j.cretres.2012.02.004
- PÉREZ-GARCÍA A. 2014. New data on the poorly-known Late Jurassic European turtles *Thalassemys* and *Enaliochelys* and description of a new basal eucryptodiran taxon. *Journal of Iberian* Geology 41: 21-30. https://doi.org/10.5209/rev_JIGE.2015.v41.n1.48652
- PÉREZ-GARCÍA A. 2015. Revision of the poorly known *Dorsetochelys typocardium*, a relatively abundant pleurosternid turtle (Paracryptodira) in the Early Cretaceous of Europe. *Cretaceous Research* 49: 152-162. https://doi.org/10.1016/j.cretres.2014.02.015
- PÉREZ-GARCÍA Á. & ORTEGA F. 2011. Selenemys lusitanica, gen. et sp. nov., a new pleurosternid turtle (Testudines: Paracryptodira) from the Upper Jurassic of Portugal. Journal of Vertebrate Paleontology 31 (1): 60-69. https://doi.org/10.1080/02724634.2011.540054
- PÉREZ-GARCÍA A. & ORTEGA F. 2014. A new species of the turtle *Hylaeochelys* (Eucryptodira) outside its known geographic and stratigraphic ranges of distribution. *Comptes Rendus Palevol* 13 (3): 183-188. https://doi.org/10.1016/j.crpv.2013.10.009
- PÉREZ-GARCÍA Á., ESPÍLEZ É., MAMPEL L. & ALCALA L. 2015a. A new European Albian turtle that extends the known stratigraphic range of the Pleurosternidae (Paracryptodira). *Cretaceous Research* 55: 74-83. https://doi.org/10.1016/j.cretres.2015.02.007
- PÉREZ-GARCÍA A., ROYO-TORRES R. & COBOS A. 2015b. A new European Late Jurassic pleurosternid (Testudines, Paracryptodira) and a new hypothesis of paracryptodiran phylogeny. *Journal of Systematic Palaeontology* 13 (4): 351-369. https://doi.org/10.108 0/14772019.2014.911212
- PÉREZ-GARCÍA A., ESPÍLEZ E., MAMPEL L. & ALCALÁ L. 2020. A new basal turtle represented by the two most complete skeletons of Helochelydridae in Europe. *Cretaceous Research* 107: 104291. https://doi.org/10.1016/j.cretres.2019.104291
- POLETTE F., BATTEN D. J. & NERAUDEAU D. 2018. Re-examination of the palynological content of the Lower Cretaceous deposits of Angeac, Charente, south-west France: Age, palaeoenvironment and taxonomic determinations. *Cretaceous Research* 90: 204-221. https://doi.org/10.1016/j.cretres.2018.04.017
- POUECH J. 2008. Position des mammifères dans les écosystèmes mésozoïques d'Europe occidentale : Le site de Cherves-de-Cognac (Berriasien, Charente, France). Unpublished PhD Thesis, Université Claude Bernard Lyon 1, 311 p.
- POUECH J., MAZIN J.-M., CAVIN L., POYATO-ARIZA F. J. 2015. A Berriasian actinopterygians fauna from Cherves-de-Cognac, France: biodiversity and palaeoenvironmental implications. *Cretaceous Research* 55: 32-43. https://doi.org/10.1016/j.cretres.2015.01.001

- POYATO-ARIZA F. J. & BUSCALIONI A. D. (eds) 2016. *Las Hoyas: A Cretaceous wetland*. Dr. Friedrich Pfeil Verlag, München, 262 p.
- POYATO-ARIZA F. J. & WENZ S. 2002. A new insight into pycnodontiform fishes. *Geodiversitas* 24: 139-248.
- PRITCHARD P. C. H. 1984. Piscivory in turtles, and evolution of the long-necked Chelidae, *in* FERGUSON M. W. (ed.), *The Structure, Development and Evolution of Reptiles.* Symposia of the Zoological Society of London, 52, Academic Press, London: 87-110.
- PUÉRTOLAS-PASCUAL E., CANUDO J. I. & SENDER L. M. 2015. New material from a huge specimen of Anteophthalmosuchus cf. escuchae (Goniopholididae) from the Albian of Andorra (Teruel, Spain): Phylogenetic implications. Journal of Iberian Geology 41: 41-57. https://doi.org/10.5209/rev_JIGE.2015.v41.n1.48654
- PUNTENER C., ANQUETIN J. & BILLON-BRUYAT J.-P. 2017. The comparative osteology of *Plesiochelys bigleri* n. sp., a new coastal marine turtle from the Late Jurassic of Porrentruy (Switzerland). *PeerJ* 5: e3482. https://doi.org/10.7717/peerj.3482
- PÜNTENER C., BILLON-BRUYAT J.-P., BOCAT L., BERGER J.-P. & JOYCE W.G. 2014. — Taxonomy and phylogeny of the turtle *Tropidemys langii* Rütimeyer, 1873 based on new specimens from the Kimmeridgian of the Swiss Jura Mountains. *Journal* of Vertebrate Paleontology 34 (2): 353-374. https://doi.org/10.1 080/02724634.2013.804412
- RAGE J.-C., MARSHALL L. G. & GAYET M. 1993. Enigmatic Caudata (Amphibia) from the Upper Cretaceous of Gondwana. *Geobios* 26: 515-519. https://doi.org/10.1016/S0016-6995(06)80234-2
- RASLAN-LOUBATIÉ Ĵ. 2007. La place des Goniopholididae et des Pholidosauridae au sein des Neosuchia (Crocodyliformes). Unpublished Master's thesis, Université Claude Bernard Lyon 1 and Muséum national d'Histoire naturelle, 31 p.
- RAUHUT O. W. M. 2003. The interrelationships and evolution of basal theropod dinosaurs. *Special Papers in Palaeontology* 69: 1-213.
- RAUHUT O. W. M. & CARRANO M. T. 2016. The theropod dinosaur *Elaphrosaurus bambergi*, from the Late Jurassic of Tendaguru, Tanzania. *Zoological Journal of the Linnean Society* 178 (3): 546-610. https://doi.org/10.1111/zoj.12425
- RAUHUT O. W. M., MILNER A. C. & MOORE-FAY S.C. 2010. Cranial osteology and phylogenetic position of the theropod dinosaur *Proceratosaurus bradleyi* (Woodward, 1910) from the Middle Jurassic of England. *Zoological Journal of the Linnean Society* 158: 155-195. https://doi.org/10.1111/j.1096-3642.2009.00591.x
- RAUHUT O. W. M., FOTH Č. & TISCHLINGER H. 2018. The oldest Archaeopteryx (Theropoda: Avialiae): a new specimen from the Kimmeridgian/Tithonian boundary of Schamhaupten, Bavaria. *PeerJ* 6: e4191. https://doi.org/10.7717/peerj.4191
- RAUP D. M. & SEPKOSKI J. J. 1982. Mass extinctions in the marine fossil record. *Science* 215: 1501-1503. https://doi.org/10.1126/ science.215.4539.1501
- RAUP D. M. & SEPKOSKI J. J. 1984. Periodicity of extinctions in the geological past. *Proceedings of the National Academy of Sciences of the United States of America* 81: 801-805. https://doi. org/10.1073/pnas.81.3.801
- RAVEN T. J. & MAIDMENT S. C. 2017. A new phylogeny of Stegosauria (Dinosauria, Ornithischia). *Palaeontology* 60: 401-408. https://doi.org/10.1111/pala.12291
- REES J., CUNY G., POUECH J. & MAZIN J. M. 2013. Non-marine selachians from the basal Cretaceous of Charente, SW France. *Cretaceous Research* 44: 122-131. https://doi.org/10.1016/j. cretres.2013.04.002
- REMANE J. 1991. The Jurassic-Cretaceous boundary: problems of definition and procedure. *Cretaceous Research* 12: 447-453. https://doi.org/10.1016/0195-6671(91)90001-S
- RICH T. H. V., MOLNAR R. E. & RICH P. V. 1983. Fossil vertebrates from the Late Jurassic or Early Cretaceous Kirkwood Formation, Algoa Basin, Southern Africa. *Transactions of the Geological Society of South Africa* 86: 281-291. https://hdl.handle. net/10520/AJA10120750_311

- RICHTER A. 1994. Lacertilia aus der Unteren Kreide von Una und Galve (Spanien) und Anoual (Marokko). *Berliner Geowis*senschaftliche Abhandlungen Reihe E (Paläobiologie) 14: 1-147.
- ROČEK Z. 2000. Mesozoic anurans, *in* HEATWOLE H. & CAR-ROLL R. L. (eds), *Amphibian Biology*. Volume 4. *Palaeontology* – *The Evolutionary History of Amphibians*. Surrey Beatty & Sons, Chipping Norton, 1496 p.
- ROYO-TORRES R. & UPCHURCH P. 2012. The cranial anatomy of the sauropod *Turiasaurus riodevensis* and implications for its phylogenetic relationships. *Journal of Systematic Palaeontology* 10: 553-583. https://doi.org/10.1080/14772019.2011.598577
- ROYO-TORRES R., ĈOBOS A. & ALCALÁ L. 2006. A giant european dinosaur and a new sauropod clade. *Science* 314: 1925-1927. https://doi.org/10.1126/science.1132885
- ROYO-TORRES R., UPCHURCH P., KIRKLAND J. I., DEBLIEUX D. D., FOSTER J. R., COBOS A. & ALCALA L. 2017. — Descendants of the Jurassic turiasaurs from Iberia found refuge in the Early Cretaceous of western USA. *Scientific Report* 7: 14311. https://doi. org/10.1038/s41598-017-14677-2
- ROYO-TORRES R., COBOS A., MOCHO P. & ALCALA L. 2021. Origin and evolution of turiasaur dinosaurs set by means of a new "rosetta" specimen from Spain. *Zoological Journal of the Linnean Society* 19: 201-227. https://doi.org/10.1093/zoolinnean/zlaa091
- ROZADA L. 2019. Taphonomie de l'assemblage de macrorestes fossiles de vertébrés du Lagerstätte d'Angeac-Charente (Crétacé inférieur, Charente, France). PhD Thesis, Paris, Muséum national d'histoire naturelle, 283 p.
- ROZADA L., ALLAIN R., VULLO R., LEPRINCE A. & TOURNEPICHE J.-F. 2014. — Taphonomy of the ornithomimosaur dinosaur herd from the Early Cretaceous lignitic bone bed of Angeac-Charente (France), *in* 74th Annual Meeting of the Society of Vertebrate Paleontology, Program and Abstracts. *Journal of Vertebrate Paleontology*, Program and Abstracts 2014: 217.
- ROZADA L., ALLAIN R., VULLO R., AUGIER D., GOEDERT J., JEAN A., ROYO-TORRES R., QVARNSTRÖM M., PEYRE DE FABRÈGUES C. & MARCHAL J. 2021. — A Lower Cretaceous Lagerstätte from Western Europe: taphonomical overview of the vertebrate assemblage of Angeac-Charente site (Western France). *Lethaia* 54: 141-165. https://doi.org/10.1111/let.12394
- SACHS S. & HORNUNG J. J. 2013. Ankylosaur remains from the Early Cretaceous (Valanginian) of northwestern Germany. *PloS* one 8(4): e60571. https://doi.org/10.1371/journal.pone.0060571
- SALISBURY S. W. 2002. Crocodilians from the lower Cretaceous (Berriasian) Purbeck Limestone Group of Dorset, Southern England. *Special Papers in Palaeontology* 68: 121-144.
- SALISBURY S. W. & NAISH D. 2011. Crocodilians, in English Wealden Fossils. The Palaeontological Association, London: 305-369.
- SALISBURY S. W., WILLIS P. M. A., PEITZ S. & SANDER P. M. 1999. The crocodilian *Goniopholis simus* from the lower Cretaceous of north-western Germany. *Special Papers in Palaeontology* 60: 121-148.
- SCHEYER T. M., PÉREZ-GARCÍA A. & MURELAGA X. 2015. Shell bone histology of solemydid turtles (stem Testudines): palaeoecological implications. *Organisms Diversity & Evolution* 15: 199-212. https://doi.org/10.1007/s13127-014-0188-0
- SCHNYDER J., DECONINCK J.-F., BAUDIN F., COLOMBIÉ C., JAN DU CHÊNE R., GARDIN S., GALBRUN B. & DE RAFÉLIS M. 2012. — Purbeck beds (Late Jurassic) in the Phare de Chassiron section (Ile d'Oléron, NW AquitaineBasin, France): refined age-assignment and long-term depositional sequences. *Geobios* 45: 485-499. https://doi.org/10.1016/j.geobios.2012.01.003
- SCHWARZ D. 2002. A New Species of Goniopholis from the Upper Jurassic of Portugal. Palaeontology 45: 185-208. https:// doi.org/10.1111/1475-4983.00233
- SCHWARZ D., RADDATZ M. & WINGS O. 2017. Knoetschkesuchus langenbergensis gen. nov. sp. nov., a new atoposaurid crocodyliform from the Upper Jurassic Langenberg Quarry (Lower Saxony, northwestern Germany), and its relationships to Theriosuchus. PloS ONE 12: e0160617. https://doi.org/10.1371/journal.pone.0160617

- SCHWARZ D. & SALISBURY S. W. 2005. A new species of *Theriosuchus* (Atoposauridae, Crocodylomorpha) from the Late Jurassic (Kimmeridgian) of Guimarota, Portugal. *Geobios* 38: 779-802. https://doi.org/10.1016/j.geobios.2004.04.005
- SCOTESE C. 2014. —Atlas of Early Cretaceous Paleogeographic Maps, PALEOMAP Atlas for ArcGIS, Vol.2. The Cretaceous. Maps 16-22. Mollweide Projection. PALEOMAP Project, Evanston. IL.
- SERENO P. C. & DONG Z. 1992. The skull of the basal stegosaur Huayangosaurus taibaii and a cladistic diagnosis of Stegosauria. Journal of Vertebrate Paleontology 12: 318-343. https://www.jstor. org/stable/4523456
- SERENO P. C. 2012. Taxonomy, morphology, masticatory function and phylogeny of heterodontosaurid dinosaurs. *ZooKeys* 226: 1-225. https://doi.org/10.3897/zookeys.226.2840
- SIGOGNEAU-RUSSELL D. 1989. Découverte du premier symmétrodonte (Mammalia) du continent africain. *Comptes Rendus de l'Académie des Sciences de Paris, Série II* 309: 921-926.
- SIGOGNEAU-RUSSELL D. 1999. Réévaluation des Peramura (Mammalia, Cladotheria) sur la base de nouveaux spécimens du Crétacé inférieur d'Angleterre et du Maroc. *Geodiversitas* 21 (1): 93-127.
- SIGOGNEAU-RUSSELL D. 2003. Diversity of the triconodont mammals from the Early Cretaceous of North Africa – Affinities of the Amphilestids. *Palaeovertebrata* 32 (1): 27-55.
- SIGOGNEAU-RUSSELL D. & ENSOM P. C. 1998. Thereuodon (Theria, Symmetrodonta) from the Lower Cretaceous of North Africa and Europe, and a brief review of symmetrodonts. Cretaceous Research 19: 445-470. https://doi.org/10.1006/cres.1998.0115
- SKUTSCHAS P. P., KOLCHANOV V. V., AVERIANOV A. O., MARTIN T., SCHELLHORN R., KOLOSOV P. N. & VITENKO D. D. 2018. — A new relict stem salamander from the Early Cretaceous of Yakutia, Siberian Russia. *Acta Palaeontologica Polonica* 63: 519-525. https://doi.org/10.4202/app.00498.2018
- SKUTSCHAS P. P. & VITENKO D. D. 2017. Early Cretaceous choristoderes (Diapsida, Choristodera) from Siberia, Russia. *Cretaceous Research* 77: 79-92. https://doi.org/10.1016/j.cretres.2017.05.004
- STERLI J., MÜLLER J., ANQUETIN J. & HILGER A. 2010. The parabasisphenoid complex in Mesozoic turtles and the evolution of the testudinate basicranium. *Canadian Journal of Earth Sciences* 47: 1337-1346. https://doi.org/10.1139/E10-061
- STRASSER A. 1986. Ooids in Purbeck limestones (lowermost Cretaceous) of the Swiss and French Jura. *Sedimentology* 33: 711-727. https://doi.org/10.1111/j.1365-3091.1986.tb01971.x
- STUMPF S., ANSORGE J., PFAFF C. & KRIWET J. 2017. Early Jurassic diversification of pycnodontiform fishes (Actinopterygii, Neopterygii) after the end-Triassic extinction event: evidence from a new genus and species, *Grimmenodon aureum. Journal* of Vertebrate Paleontology 37: e1344679. https://doi.org/10.108 0/02724634.2017.1344679
- SUBALUSKY A. L., DUTTON C. L., ROSI E. J. & POST D. M. 2017. Annual mass drownings of the Serengeti wildebeest migration influence nutrient cycling and storage in the Mara River. *Proceedings of the National Academy of Sciences* 114: 7647-7652. https:// doi.org/10.1073/pnas.1614778114
- SWEETMAN S. C. 2006. A gobiconodontid (Mammalia, Gobiconodontidae) from the Early Cretaceous (Barremian) Wessex Formation of the Isle of Wight, southern Britain. *Palaeontology* 49 (4): 889-897. http://doi.org/10.1111/j.1475-4983.2006.00564.x
- SWEETMAN S. C. 2016. A comparison of Barremian-early Aptian vertebrate assemblages from the Jehol Group, north-east China and the Wealden Group, southern Britain: the value of microvertebrate studies in adverse preservational settings. *Palaeobiodiversity* and Palaeoenvironments 96: 149-167. https://doi.org/10.1007/ s12549-015-0217-9
- SWEETMAN S. C. & EVANS S. E. 2011. Lepidosaurs (lizards), in BATTEN D. J. (ed.), English Wealden fossils. Wiley-Blackwell, London: 264-284 (Palaeontological Association Field Guide to Fossils; 14).

- SWEETMAN S. C. & GARDNER J. D. 2013. A new albanerpetontid amphibian from the Barremian (Early Cretaceous) Wessex Formation of the Isle of Wight, southern England. *Acta Palaeontologica Polonica* 58: 295-324. https://doi.org/10.4202/app.2011.0109
- SWEETMAN S. C. & INSOLE A. N. 2010. The plant debris beds of the Early Cretaceous (Barremian) Wessex Formation of the Isle of Wight, southern England: their genesis and palaeontological significance. *Palaeogeography, Palaeoclimatology, Palaeoecology* 292 (3-4): 409-424. https://doi.org/10.1016/j.palaeo.2010.03.055
- SWEETMAN S. C., GOEDERT J. & MARTILL D. M. 2014. A preliminary account of the fishes of the Lower Cretaceous Wessex Formation (Wealden Group, Barremian) of the Isle of Wight, southern England. *Biological Journal of the Linnean Society* 113: 872-896. https://doi.org/10.4202/app.2011.0109
- SWEETMAN S. C., PEDREIRA-SEGADE U. & VIDOVIC S. U. 2015. A new bernissartiid crocodyliform from the Lower Cretaceous Wessex Formation (Wealden Group, Barremian) of the Isle of Wight, southern England. Acta Palaeontologica Polonica 60: 257-268. https://doi.org/10.4202/app.00038.2013
- TENNANT J. P. & MANNION P. 2014. Revision of the Late Jurassic crocodyliform *Alligatorellus*, and evidence for allopatric speciation driving high diversity in western European atoposaurids. *PeerJ* 2: e599. https://doi.org/10.7717/peerj.599
- TENNANT J. P., MANNION P., UPCHURCH P., SUTTON M. D. & PRICE G. D. 2017. Biotic and environmental dynamics through the Late Jurassic-Early Cretaceous transition: evidence for protracted faunal and ecological turnover. *Biological Reviews* 92: 776-814. https://doi.org/10.1111/brv.12255
- THOMAS S. C. 1996. Differentiation of selected frog (*Rana*) and toad (*Bufo*) using morphological and osteometric traits. *Canadian Zooarchaeology* 10: 2-8.
- THURMOND J. T. 1974. Lower vertebrate faunas of the Trinity Division in north-central Texas. *Geoscience and Man* 8: 103-129.
- UNDERWOOD C. J. & CUMBA S. L. 2010. Chondrichthyans from a Cenomanian (Late Cretaceous) bonebed, Saskatchewan, Canada. *Palaeontology* 53: 903-944. https://doi.org/10.1111/ j.1475-4983.2010.00969.x
- UPCHURCH P., BARRETT P. M. & DODSON P. 2004. Sauropoda, in WEISHAMPEL D. B., DODSON P. & OSMOLSKA H. (eds), *The Dino*sauria. 2nd edition. University of California Press, Berkeley: 259-322.
- VRIELYNCK B., DERCOURT J. & COTTEREAU N. 1995. The Tethys: an ocean broken by seuils lithospheriques, *in* NAIM A. E. M., RICOU L.-E., VRIELYNCK B. & DERCOURT J. (eds), *The Tethys Ocean: The Ocean Basins and Margins*. Volume 8. Plenum Press, New York: 495-511.
- VULLO R., ABIT D., BALLEVRE M., BILLON-BRUYAT J.-P., BOURGEAIS R., BUFFETAUT E., DAVIERO-GOMEZ V., GARCIA G., GOMEZ B., MAZIN J.-M., MOREL S., NERAUDEAU D., POUECH J., RAGE J.-C., SCHNYDER J. & TONG H. 2014. — Palaeontology of the Purbeck-type (Tithonian, Late Jurassic) bonebeds of Chassiron (Oléron Island, western France). *Comptes Rendus Palevol* 13: 421-441. https://doi.org/10.1016/j.crpv.2014.03.003
- VULLO R., LAPPARENT DE BROIN F. DE, NÉRAUDEAU D. & DUR-RIEU N. 2010. — Turtles from the Early Cenomanian paralic deposits (Late Cretaceous) of Charentes, France. *Oryctos* 9: 37-48.
- WANG X., CAMPOS D. A., ZHOU Z. & KELLNER A. W. A. 2008. A primitive istiodactylid pterosaur (Pterodactyloidea) from the Jiufotang Formation (Early Cretaceous), Northeast China. *Zootaxa* 1813: https://doi.org/1-18. 10.5281/zenodo.274343
- WANG S., STIEGLER J., AMIOT R., WANG X., DU G. H., CLARK J. M. & XU X. 2017. — Extreme ontogenetic changes in a ceratosaurian theropod. *Current Biology* 27: 144-148. https://doi. org/10.1016/j.cub.2016.10.043
- WHETSTONE K. N. & WHYBROW P. J. 1983. A "cursorial" crocodilian from the Triassic of Lesotho (Basutoland), southern Africa. Occasional Papers of the Museum of Natural History, University of Kansas 106: 1-37. https://www.biodiversitylibrary. org/page/4488912

- WIBLE J. R., ROUGIER G. W., NOVACEK M. J., MCKENNA M. C. & DASHZEVEG C. 1995. — A mammalian petrosal from the Early Cretaceous of Mongolia: implications for the evolution of the ear region and mammaliamorph interrelationships. *American Museum Novitates* 3149: 1-20. http://hdl.handle. net/2246/3672
- WILSON J. A. 2002. Sauropod dinosaur phylogeny: critique and cladistic analysis. Zoological Journal of the Linnean Society 136: 215-275. https://doi.org/10.1046/j.1096-3642.2002.00029.x
- WILSON J. A. 2005. Overview of Sauropod Phylogeny and Evolution, *in* CURRY ROGERS K. A. & WILSON J. A. (eds), *The Sauropods: Evolution and Paleobiology*. University of California Press, Berkeley: 15-49.
- WIMBLEDON W. A. P., CASELLATO C. E., REHAKOVA D., BULOT L. G., ERBA E., GARDIN S., VERREUSSEL R. M. C. H., MUNSTERMAN D. K. & HUNT C. O. 2011 —. Fixing a basal Berriasian and Jurassic/ Cretaceous (J/K) Boundary – Is there perhaps some light at the end of the tunnel? *Rivista Italiana di Paleontologia e Stratigra*fia 117: 295 e 307. https://doi.org/10.13130/2039-4942/5976
- XU X., CLARK J. M., MO J., ČHOINIERE J., FORSTER C. A., ERICKSON G. M. & ZHAO Q. 2009. — A Jurassic ceratosaur from China helps clarify avian digital homologies. *Nature* 459: 940-944. https://doi.org/10.1038/nature08124
- ZANNO L. E. & MAKOVICKY P. J. 2010. Herbivorous ecomorphology and specialization patterns in theropod dinosaur evolution. *Proceedings of the National Academy of Sciences* 108: 232-237. https://doi.org/10.1073/pnas.1011924108

Submitted on 13 November 2020; accepted on 29 May 2021; published on 21 July 2022.

APPENDIX

APPENDIX 1. — Systematic paleontology of the complete faunal list of the Angeac-Charente locality.

Arthropoda von Siebold, 1848 Crustacea Brünerich, 1772 Ostracoda Latreille, 1802 Cypridea gr. tuberculata Sowerby,1836 Cypridea laevigata Dunker, 1846 Damonella pygmaea Anderson, 1941 Damonella ellipsoidea Wolburg, 1962 Darwinula oblonga Roemer, 1839 Alicenula leguminella Forbes, 1855 Fabanella boloniensis Jones, 1885 Mantelliana sp. Anderson, 1966 Isoptera Brullé, 1832 Family indet. Microcarpolithes hexagonalis Vangerow, 1954 (termite coprolithes) Mollusca Linnaeus, 1758 Gastropoda Cuvier, 1795 Viviparidae Gray, 1847 Viviparidae indet. Bivalvia Linnaeus, 1758 Unionoidea Stoliczka, 1871 cf. Margaritifera sp. Vertebrata Lamarck, 1801 Chondrichthyes Huxley, 1880 Hybodontiformes Owen, 1846 Lonchidiidae Herman, 1977 Parvodus celsucuspus Rees, Cuny, Pouech & Mazin, 2013 Osteichthyes Huxley, 1880 Actinopterygii Klein, 1885 Ginglymodi Cope, 1872 Ginglymodi indet. Ionoscopiformes Grande & Bemis, 1998 Ionoscopiformes indet. Amiiformes Huxley, 1861 Amiiformes indet. Pycnodontiformes Berg, 1937 cf. Micropycnodon sp. Pycnodontidae indet. Amphibia Linnaeus, 1758

Lissamphibia Haeckel, 1866 Lissamphibia indet. Albanerpetontidae Fox and Naylor, 1982 Albanerpetontidae indet. Anura Fischer von Waldheim, 1813 Anura indet. Caudata Scopli, 1777 Caudata indet. Reptilia Laurenti, 1768 Testudinata Klein, 1760 Perichelydia Joyce, 2017 Helochelydridae Nopsca, 1928 (sensu Joyce et al. 2016) Helochelydridae indet. Paracryptodira Gaffney, 1975 Pleurosternidae Cope, 1868 Pleurosternon bullockii (Owen, 1842) Thalassochelydia Anguetin et al., 2017 Hylaeochelys belli? (Mantell, 1844). Lepidosauria Haeckel, 1866 Squamata Oppel, 1811 Scincomorpha Camp, 1923 cf. Paramacellodus sp. Rhynchocephalia Günther, 1867 Sphenodontia Williston, 1925 cf. *Opisthias* sp. Archosauromorpha von Huene, 1946 Choristodera Cope, 1876 cf. Cteniogenys sp. Archosauria Cope, 1870 Crocodyliformes Hay, 1930 Atoposauridae Gervais, 1871 *Theriosuchus* sp. Atoposauridae indet. Bernissartiidae Dollo, 1883 Bernissartiidae indet. Goniopholididae Cope, 1875 Goniopholis sp. Pholidosauridae Eastman, 1902 Pholidosaurus sp. Dinosauria Owen, 1842 Theropoda Marsh, 1881 Megalosauridae Fitzinger, 1843 Megalosauridae indet. Ornithomimosauria Barsbold, 1976 New taxon Tyrannosauroidea Osborn, 1905 Tyrannosauroidea indet. Dromaeosauridae Matthew and Brown, 1922 cf. Nuthetes sp. Owen, 1854 Archaeopterygidae Huxley, 1872 Archaeopterygidae indet. Sauropoda Marsh, 1878 Macronaria indet. Turiasauria Royo-Torres, Cobos & Alcalá, 2006 New taxon Thyreophora Nopcsa, 1915 Stegosauria Marsh, 1877 Dacentrurus. sp. Ankylosauria Osborn, 1923 Ankylosauria indet.
APPENDIX 1. - Continuation.

Heterodontosauridae Kuhn, 1966 *Echinodon* sp. Ornithopoda Marsh, 1881 Hypsilophodontidae Hypsilophodontidae indet. Camptosauridae Camptosauridae indet. Pterosauria Kaup, 1834 Pterosauria indet. Pterodactyloidea Plieninger, 1901 Pterodactyloidea indet. A Pterodactyloidea indet. B

Synapsida Osborn, 1903 Mammalia Linnaeus, 1758 Mammalia indet. Thereuodontidae Sigogneau-Russell & Ensom, 1998 Thereuodon cf. taraktes Eutriconodonta Kermarck, Mussett & Rigney, 1973 Gobiconodontidae Chow & Rich, 1984 Gobiconodon sp. Triconodontidae Marsh, 1887 Triconodon sp. Allotheria Mars, 1880 Multituberculata Cope, 1884 Paulchoffatiidae Hahn, 1969 *Sunnyodon* sp. Pinheirodontidae Hahn et Hahn, 1999 Pinheirodontidae indet. Trechnotheria McKenna, 1975 Spalacotheriidae Marsh, 1887 Spalacotherium sp. Cladotheria McKenna, 1975 Dryolestoidea Butler, 1939 Dryolestidae Marsh, 1879 Dryolestidae indet. Zatheria McKenna, 1975 Peramuridae Kretzoi, 1946 Peramus sp.