



HAL
open science

ACS trace gas detection attempts

Alexander V. Trokhimovskiy, Anna A. Fedorova, Oleg I. Korablev, Denis A. Belyaev, Ashwin Braude, Franck Montmessin, Franck Lefèvre, Juan Alday, Kevin Olsen

► **To cite this version:**

Alexander V. Trokhimovskiy, Anna A. Fedorova, Oleg I. Korablev, Denis A. Belyaev, Ashwin Braude, et al.. ACS trace gas detection attempts. Seventh International Workshop on the Mars Atmosphere: Modelling and Observations, Jun 2022, Paris, France. pp.id.3521. insu-03752489

HAL Id: insu-03752489

<https://insu.hal.science/insu-03752489v1>

Submitted on 16 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ACS TRACE GAS DETECTION ATTEMPTS

A. Trokhimovskiy, A. A. Fedorova, O. I. Korablev, D. A. Belyaev, *Space Research Institute of the Russian Academy of Sciences (IKI RAS), Moscow, Russia (trokh@iki.rssi.ru)*, **A. S. Braude, F. Montmessin, F. Lefèvre**, *LATMOS/IPSL, UVSQ Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France*, **J. Alday**, *Open University, Milton Keynes, U.K.*, **Kevin S. Olsen**, *Department of Physics, University of Oxford, UK*

Introduction: The ACS instrument onboard ExoMars TGO has been developed to meet a variety of science goals, in particular, the sensitive search for trace species in the Martian atmosphere [1]. Regular soundings in solar occultation mode started in March 2018. For methane ACS reported stringent 20 ppt upper limits all over the planet [2,3], which are not correlated with surface measurements by the MSL rover [4]. With the methane lifetime of ~300 years, conventional chemistry cannot explain the disagreement. ACS continues regular measurements to gain statistics and in hope for possible methane spikes reported previously [5]. The presence of methane derivatives like ethane C₂H₆ is very unlikely, yet ACS sets an improved upper limit of 50 ppt.

Recently discovered by ACS [6], hydrogen chloride (HCl) may be involved in many atmospheric processes, so we looked for chloromethane (CH₃Cl) features in the data. The Viking landers detected chloromethane during the search for organic compounds in Martian soils [7]; MSL again noted CH₃Cl during the pyrolysis of soil [8], still these detections are believed to have a contamination source. The ACS upper limit for CH₃Cl in the atmosphere is better than 1 ppb, which is several times better than previous knowledge [9], and extends over seasons.

Sulfur species were searched as an indicator of any volcanic outgassing from the surface of Mars. For the most optimal sensitivity conditions, we determine upper limits of SO₂ at 20 ppbv, H₂S at 15 ppbv, and OCS at 0.4 ppbv [10]; the last value is lower than any previous upper limits imposed on OCS in the literature.

Nitrogen is known to be present in the Martian atmosphere, forming the oxidation products derived from ionospheric dissociation of N₂. The first species to be searched for with ACS is NO₂. It has comparable model predicted abundances with NO in the lower atmosphere. ACS is sensitive to NO₂ with an accuracy of 200 pptv matching model predicted concentrations [11]. Without a confident detection, the median abundance over the entire ACS dataset agrees with this value. In 2021 we started a dedicated search campaign for NH₃ and HCN. Preliminary derived upper limits are 11 ppb and 1 ppb accordingly.

References

[1] Korablev, O. et al. The Atmospheric Chemistry Suite (ACS) of three spectrometers for the ExoMars 2016 Trace Gas Orbiter. *Space Sci. Rev.*

214, 7 (2018).

[2] Korablev, O. et al. No detection of methane on Mars from early ExoMars Trace Gas Orbiter observations. *Nature* 568, 517–520 (2019).

[3] Montmessin, F. et al. A more stringent upper limit of methane on Mars and strong constraints on methane dispersion outside Gale crater. *Astron. Astrophys.* 650, A140 (2021).

[4] Webster, C. et al. Background levels of methane in Mars' atmosphere show strong seasonal variations. *Science* 360, 6393, 1093-1096 (2018).

[5] Giuranna, M. et al. Independent confirmation of a methane spike on Mars and a source region east of Gale Crater. *Nature Geoscience* 12, 326–332 (2019).

[6] Korablev, O. et al. Transient HCl in the atmosphere of Mars. *Sci. Adv.* 7, eabe4386 (2021).

[7] Biemann, K et al. The search for organic substances and inorganic volatile compounds in the surface of Mars. *J. Geophys. Res.*, 82, 4641-4658 (1977).

[8] Glavin, D. et al. Evidence for perchlorates and the origin of chlorinated hydrocarbons detected by SAM at the Rocknest aeolian deposit in Gale Crater. *JGR Planets* 118, 10 (2013).

[9] Villanueva, G. et al. A sensitive search for organics (CH₄, CH₃OH, H₂CO, C₂H₆, C₂H₂, C₂H₄), hydroperoxyl (HO₂), nitrogen compounds (N₂O, NH₃, HCN) and chlorine species (HCl, CH₃Cl) on Mars using ground-based high-resolution infrared spectroscopy. *Icarus* 223(1), 11-27 (2013).

[10] Braude, A. S. et al. No detection of SO₂, H₂S, or OCS in the atmosphere of Mars from the first two Martian years of observations from TGO/ACS. *Astron. Astrophys.* 658, A86 (2022).

[11] Moudden, Y., McConnell, J.C. Three-dimensional on-line chemical modeling in a Mars general circulation model. *Icarus* 188(1), 18-34 (2007).