



HAL
open science

A Four-Year Search of Methane on Mars with ACS onboard ExoMars TGO

Franck Montmessin, Oleg Korablev, Alexander Trokhimovskiy, Kevin Olsen,
Franck Lefèvre, Lucio Baggio, Ashwin Braude, Anna Fedorova, Elise Wright
Knutsen, Abdanour Irbah, et al.

► **To cite this version:**

Franck Montmessin, Oleg Korablev, Alexander Trokhimovskiy, Kevin Olsen, Franck Lefèvre, et al.. A Four-Year Search of Methane on Mars with ACS onboard ExoMars TGO. Europlanet Science Congress 2022, Sep 2022, Granada, Spain. pp.EPSC2022-409. insu-03753094

HAL Id: insu-03753094

<https://insu.hal.science/insu-03753094>

Submitted on 17 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



A Four-Year Search of Methane on Mars with ACS onboard ExoMars TGO

Franck Montmessin¹, Oleg Korablev², Alexander Trokhimovskiy², Kevin Olsen³, Franck Lefèvre¹, Lucio Baggio¹, Ashwin Braude¹, Anna Fedorova², Elise Knutsen¹, Abdanour Irbah¹, Gaétan Lacombe¹, Andrey Patrakeev², and Alexey Shakun²

¹CNRS/IPSL/UVSQ/UPMC, LATMOS, Guyancourt, France (franck.montmessin@latmos.ipsl.fr)

²Research Institute (IKI) RAS, Moscow, Russia

³AOPP, Oxford University, Oxford, United Kingdom

Summary

The mid-infrared channel of the Atmospheric Chemistry Suite (ACS) [1] onboard ESA's Trace Gas Orbiter (TGO) has performed a sensitive search of methane in the Martian atmosphere using the solar occultation technique for more than two Mars years (four terrestrial years). The first two reports [2,3] have concluded on the global absence of methane detection above a mean level of 50 then 20 parts-per-trillion-per-volume (pptv), contrasting with the 0.4 to 45 ppbv concentrations reported from Earth-ground-observations, Mars orbiters, and from Mars' surface by the Curiosity rover [5,6,7,8,9]. Here we present an update of the ACS methane search compiling all the ACS data collected up to summer 2022.

Introduction

For several decades, methane on Mars has been the subject of a continuous and intense search for it may resonate with an ongoing geophysical or biogenic activity on a planet that has long been perceived as inert. After a succession of failed attempts and positive detections between 2004 and 2019 [4,5,6,7,8,9,21,22], TGO started its trace gas detection mission in April 2018 and has continued since then. With its two separate state-of-the-art infrared spectrometers specifically conceived to achieve outstanding detection performances [1,10] for the ultra-sensitive search of a list of trace gases, including methane, TGO was aimed at confirming the presence of methane at Mars, or alternatively to establish the smallest upper limits possible. The Atmospheric Chemistry Suite (ACS) developed by IKI with the support of CNES is one of the spectrometers of TGO. Together with NOMAD, they offer a complementary access to the entire wavelength range from UV to thermal IR. They are also both observing the mid-IR range and therefore provide a redundant and thus reliable exploration of the presence of a variety of trace gases. Among the greatest achievements of these two experiments, we shall mention the joint observation of HDO and H₂O [11,12], the discovery of HCl [13,14,15], the close and joint monitoring of water and ozone [16,17,18], as well as the first CO profiles [19].

Method

The mid-infrared range, in particular the region around the fundamental 3.3 μm band of the C-H bond, provides the most sensitive way to detect methane and higher-order hydrocarbons (Fig. 1).

With ACS, this spectral region has been explored around Mars during the last four years and has revealed an absence of methane above a limit of 20 pptv (Fig. 2), while NOMAD similarly established such limit to 60 pptv [20].

The technique of solar occultation is known to be more sensitive than nadir since the Sun is a source at Mars that is ~600 000 times brighter than Mars itself. While the ubiquitous atmospheric dust prevents observing the near-surface region, the high-resolving power (30 000) and its high Signal-to-Noise Ratio (~10 000 at low level of dust) compensate for this limitation. We attempt to detect CH₄ using the R2 region which features the most intense individual lines of the band system. This region was backed up by parallel search in three other regions to have confirmation in case of a positive detection.

Results

With two Martian years of survey, the dataset consists of more than 1 000 individual altitude profiles. Since [3], improvements in the data processing have been implemented. This 4-year dataset will be presented along with its implications in regard to other observational studies and model attempts to explain reconcile the various observations.

[1] Korablev, O., et al. 2018. Space Sci. Rev.

[2] Korablev, O., et al. 2019. Nature

[3] Montmessin, F., et al. 2021. Astron. & Astrophys.

[4] Lellouch, E., et al. 2000. Astron. & Astrophys.

[5] Formisano, V., et al., 2004. Science

[6] Krasnopolsky, V. et al., 2004. Icarus

[7] Mumma, M. et al., 2009. Science

[8] Webster, C.R., et al., 2015, Science

[9] Webster, C.R., et al., 2018, Science

[10] Vandaele, A.C., et al. 2018. Space Sci. Rev.

[11] Villanueva, G. et al., 2020. Sci. Adv.

[12] Alday, J. et al., 2021. Nat. Ast.

[13] Korablev, O., et al., 2021. Sci. Adv.

[14] Aoki, S., et al., 2021. Geophys. Res. Let.

[15] Olsen, K. et al., 2021. Astron. & Astrophys.

[16] Fedorova et al., A., et al., 2020, Science

[17] Patel, M., et al., 2021, J. of Geophys. Res.

[18] Belyaev, D., et al, 2021., Geophys. Res. Let.

[19] Olsen, K. et al., 2021. Nat. Geo.

[20] Knutsen, E., et al., 2021, Icarus.