



HAL
open science

Methane vertical profile in Titan's atmosphere

Thomas Gautier, Joseph Serigano, Koyena Das, Maélie Coutelier, Sarah Hörst, Sandrine Vinatier, Cyril Szopa, Melissa Trainer

► **To cite this version:**

Thomas Gautier, Joseph Serigano, Koyena Das, Maélie Coutelier, Sarah Hörst, et al.. Methane vertical profile in Titan's atmosphere. Europlanet Science Congress 2022, Sep 2022, Granada, Spain. pp.EPSC2022-431. insu-03753564

HAL Id: insu-03753564

<https://insu.hal.science/insu-03753564>

Submitted on 18 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Methane vertical profile in Titan's atmosphere

Thomas Gautier^{1,2}, Joseph Serigano³, Koyena Das¹, Maélie Coutelier¹, Sarah Hörst³, Sandrine Vinatier², Cyril Szopa¹, and Melissa Trainer⁴

¹LATMOS-IPSL, CNRS, Sorbonne Université, UVSQ, Guyancourt, France (thomas.gautier@latmos.ipsl.fr)

²LESIA-Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université de Paris, Meudon, France

³Department of Earth and Planetary Sciences, Johns Hopkins University, Baltimore, USA

⁴NASA Goddard Space Flight Center, Greenbelt, USA

More than a decade after the arrival of the Cassini-Huygens mission in Saturn's system, data returned by the Huygens probe during its descent remain a unique source of *in-situ* information on the lower atmosphere of Titan. Among the Huygens instrumental suite was the GCMS (Gas Chromatograph Mass Spectrometer) instrument which returned hundreds of mass spectra acquired in the atmosphere below 145 km of altitude.

We will present a reanalysis of GCMS data focusing on the methane vertical profile thanks to recent advances in our knowledge of Titan's atmosphere and in mass spectrometry data treatment ^[1-3].

We retrieved methane mixing ratio slightly lower than the one reported by the original team^[4] and obtained its profile between 145 km and 30 km of altitude with a kilometric vertical resolution, and a sub-kilometric one between 30 km and the surface.

Such a vertical resolution unveiled clear oscillations in the methane mixing ratio in Titan's troposphere and methane vertical concentration diverging from an ideal adiabat. Such features could for example be a sign of small-scale convective zones in the troposphere which could have triggered the gravity waves detected by Huygens in the stratosphere. We hope that the discovery of previously unnoticed features in GCMS data will also enable the reanalysis of data returned by other Huygens instruments such as HASI and DISR on Titan lower atmosphere.

^[1] Gautier et al. *Decomposition of electron ionization mass spectra for space application using a Monte-Carlo approach*. Rapid. Com. Mass Spec. 34(8), e8659 (2020)

^[2] Serigano et al. *Compositional measurements of Saturn's upper atmosphere and rings from Cassini INMS*. JGR:Planets, 125 (8), E006427 (2020)

^[3] Serigano et al. *Compositional Measurements of Saturn's Upper Atmosphere and Rings from Cassini INMS: An extended Analysis of Measurements from Cassini's Grand Finale Orbits*. JGR:Planets, 127, E007238 (2022)

^[4] Niemann et al. *Composition of Titan's lower atmosphere and simple surface volatiles as measured by the Cassini-Huygens probe gas chromatograph mass spectrometer experiment*. JGR 115, E12006, 2010