



HAL
open science

**A fossil flat wasp (Hymenoptera: Bethylidae) from the
early Eocene Green River Formation suggests past
cosmopolitan distribution of the genus *Eupsenella*
Westwood, 1874**

Manuel Brazidec, Vincent Perrichot

► **To cite this version:**

Manuel Brazidec, Vincent Perrichot. A fossil flat wasp (Hymenoptera: Bethylidae) from the early Eocene Green River Formation suggests past cosmopolitan distribution of the genus *Eupsenella* Westwood, 1874. *Palaeoentomology*, 2022, 5 (4), pp.378-384. 10.11646/palaeoentomology.5.4.10 . insu-03764151

HAL Id: insu-03764151

<https://insu.hal.science/insu-03764151>

Submitted on 31 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **A fossil flat wasp (Hymenoptera: Bethyliidae) from the early Eocene Green River**
2 **Formation suggests past cosmopolitan distribution of the genus *Eupsenella* Westwood,**
3 **1874**

4 MANUEL BRAZIDEC^{1,2*}, VINCENT PERRICHOT¹

5 ¹Univ Rennes, CNRS, Géosciences Rennes, UMR 6118, 35000 Rennes, France.

6 manuel.brazidec@gmail.com / <https://orcid.org/0000-0002-0860-8972>;

7 vincent.perrichot@univ-rennes1.fr / <https://orcid.org/0000-0002-7973-0430>

8 ²State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and
9 Palaeontology and Center for Excellence in Life and Palaeoenvironment, Chinese Academy of
10 Sciences, 39 East Beijing Road, Nanjing 210008, China.

11 *corresponding author

12

13 **Abstract**

14 A flat wasp specimen attributed to the genus *Eupsenella* Westwood, 1874 is described
15 and figured from the early Eocene compressions of the Green River Formation. This fossil is
16 the first known outside of the Old World for this genus and documents its wide distribution
17 during the Paleogene, while it is currently confined to Australia and New Zealand. As for
18 several other hymenopteran genera, we assume that this regression is linked to abiotic factors,
19 i.e., the Cenozoic climate cooling, rather than to biotic factors.

20

21 **Keywords:** Hymenoptera; Chrysidoidea; Bethylinae; Ypresian; USA

22 1. Introduction

23 The Green River Formation is an ancient lake system, widespread throughout western
24 United States (Wyoming, Colorado, Texas; Smith *et al.* 2008). The thin lacustrine sediments
25 were deposited during the Early Eocene Climatic Optimum and have exquisitely preserved
26 vertebrates, invertebrates, and plants remains, justifying the use of the term ‘Lagerstätte’
27 (Hellowell & Orr 2012, MacGinitie 1969). Because of the economic importance of the Green
28 River Formation (exploitation of oil-shale layers), its geology and physical setting are well
29 understood (Johnson *et al.*, 2014). In contrast, its biota is far less known and particularly the
30 entomofauna, despite the collection of more than 80,000 specimens (Dlussky & Rasnitsyn
31 2002).

32 The first studies on Green River fossil insects date back to the late 19th and early 20th
33 centuries and were especially conducted by S.H. Scudder and T.D.A. Cockerell (Scudder
34 1890, Cockerell 1907, 1925). Grande (2013) revised the fauna of the deposit, highlighting the
35 greater interest of paleoentomologists for the northern deposits of the Greater Green River
36 Basin and the Fossil Basin. However, in the past few years, some studies focused on the insects
37 from the Piceance Basin, in Colorado (e.g., Vršanský *et al.* 2011, Wappler *et al.* 2015).

38 In this paper, we describe a Bethylidae from the Piceance Basin locality of Denson, the
39 specimen being the first flat wasp known from the Green River Formation. The Bethylidae are
40 a cosmopolitan family, the largest within the Chrysidoidea (Azevedo *et al.* 2018). Of the nine
41 recognized subfamilies (Colombo *et al.* 2020), eight have fossil representatives but mostly
42 from amber deposits (Martynova *et al.* 2019). Bethylids are rarer in compressions, with only a
43 few known specimens (Colombo *et al.* 2021, Peñalver-Mollá 1998, Rasnitsyn 1990), mainly
44 because of the fossorial habits and small size of these wasps.

45

46 **2. Material and methods**

47 The part of a Bethylidae was collected in 2006 by David Kohls at the Denson Site,
48 Garfield County, northwestern Colorado, USA (see map in Wappler *et al.* 2015: fig. 1). The
49 locality corresponds to the lesser-known Piceance Basin, whose insect-bearing stratum is the
50 Parachute Creek Member. This member is the youngest of the basin, dated of the upper
51 Ypresian (Smith *et al.* 2008, Cole & Picard 1978). The specimen is labelled UCM 58339 and
52 housed in the Invertebrate Paleontology collection of the University of Colorado.

53 The images were captured using a Canon EOS 5D camera with 65 mm MPE Canon macro
54 lens attached to a Stackshot macro rail. All images are digitally stacked photomicrographic
55 composite of several focal planes, which were obtained using HeliconFocus. Adobe Illustrator
56 CC2019 and Photoshop CC2019 software were used to compose the figures and ImageJ 1.53
57 for measurements (Schneider *et al.* 2012). The morphological nomenclature follows Lanes *et*
58 *al.* (2020). Main measurements and indices used are as follows: length of forewing (LFW);
59 length of head, measured from top of vertex to apex of clypeus (LH); width of head above
60 eyes (WH); height/width of eye (HE).

61

62 **3. Systematic palaeontology**

63 **Superfamily** Chrysoidea Latreille, 1802

64 **Family** Bethylidae Haliday, 1839

65 **Subfamily** Bethylinae Haliday, 1839

66 **Genus** *Eupsenella* Westwood, 1874

67 *Eupsenella* sp.

68 Figs. 1, 2

69

70

71 Material. Specimen UCM 58339 of unknown sex, part. Right lateral aspect of the body, with
72 legs, left hind wing and parts of the antennae missing.

73 Locality. Denson Site, Green River, Colorado, USA

74 Horizon. Lake Uinta, Piceance Basin, Parachute Creek member; upper Ypresian, Eocene, ca.
75 48 Ma.

76 Description. Body dark brown, not especially flattened, stout (length 3.23 mm). Head ovoid,
77 apparently longer than wide; LH: ca. 0.67 mm; WH: ca. 0.36 mm; HE: 0.21 mm; compound
78 eyes elliptical, longer than high; antennae short, only scape and two apicalmost flagellomeres
79 preserved, apical flagellomere tapering at apex; vertex corners rounded.

80 Mesosoma shorter than metasoma (length 1.04 mm). Fore wing micro-pubescent, without
81 apparent colour patterns (LFW: 1.91 mm); C, Sc+R, M+Cu, 1Rs&1M, Rs+M, Rs and R1
82 preserved and tubular; 1Rs&1M angulate at junction with Rs+M; Rs apically angulate toward
83 wing margin; post-stigmal abscissa of R1 apically arched; [C], [R], [1M] (areolet), [1R1] and
84 [2R1] cells closed; [1M] cell small, subpentagonal, aligned basally with [1R1] cell; [1R1] cell
85 elongate, 1.0 x [2R1] cell; [2R1] cell 0.19 x LFW, evenly wide, as wide as [1R1] cell;

86 pterostigma well-defined, elongate, slightly shorter than [1R1] cell, rectangular. Hind wing
87 hyaline, without apparent venation.

88 Metasoma fusiform (length 1.52 mm); segments not discernible.

89

90

91

92 **4. Discussion**

93 Despite being only observable laterally and missing the legs and part of the antennae,
94 this fossil wasp can confidently be identified as a Bethylinidae of the subfamily Bethylininae,
95 thanks to the unique reduced fore wing venation (five closed cells visible), the Rs+M vein
96 tubular and the 1Rs&1M angled (Azevedo *et al.* 2018). The [1Cu] cell, which is always
97 closed in bethyline wasps, is assumed to be present as well but not preserved, as a stub of the
98 1cu-a vein is visible anteriorly. The comparatively high number of closed cells (six) and their
99 combination (especially the [1M], [1R1] and [2R1] cells) for a Bethylininae strongly indicates
100 either *Lytopsenella* Kieffer, 1911 or *Eupsenella* Westwood, 1874 within the subfamily
101 (*Azevedo et al.* 2018). The distinction between these two genera is based on the relative
102 length of the [1R1] and [2R1] cells and the shape of the [2R1] cell. In our specimen, the [2R1]
103 cell is as long as the [1R1] (while it is longer in *Lytopsenella*) and evenly wide and rounded
104 apically (while it is lanceolate in *Lytopsenella*). The specimen mostly differs from extant
105 species by the size of the [2R1] cell, which is as long as the [1R1] cell and not significantly
106 shorter. This characteristic occurs in all other fossils attributed to *Eupsenella* (Ramos *et al.*
107 2014). Yet, we cannot rule out the possibility that this specimen belongs to a brand new genus

108 of Bethylinae. But it is worth mentioning that no fossil genera of this subfamily are known
109 during the Cenozoic, which is quite uncommon, possibly because of its basal position within
110 the family (Sorg 1988, Ramos *et al.* 2014). Additionally, the fore wing pattern is very
111 characteristic of *Eupsenella* and, except *Lytopsenella*, no other Bethylinidae have such a
112 venation, even in a different subfamily. Therefore, we allocate this specimen in the extant
113 genus *Eupsenella*. However, due to the lack of body features preserved, which prevents from
114 establishing precise diagnosis, we decide to leave it unnamed.

115

116 The genus *Eupsenella* is currently composed of 52 species (Azevedo *et al.* 2018). The
117 45 extant species are restricted to Australia with one species occurring in the South Island of
118 New Zealand (Fig. 3; Ramos & Azevedo 2012). During the late Eocene, four species are
119 known from the contemporaneous Baltic amber and Rovno amber (Ramos *et al.* 2014). Two
120 species originally attributed to *Fushunochrysis* Hong, 2002 and *Sinibethylus* Hong, 2002
121 from the Ypresian Fushun amber (Hong 2002) and one attributed to *Protobethylus* De Ploëg
122 & Nel, 2004 from the Ypresian Oise amber (De Ploëg & Nel 2004) have been synonymized
123 with this genus (Ramos *et al.* 2014). Therefore, *Eupsenella* appears to be widely distributed
124 throughout the Palearctic during the Paleogene but the fossil from Green River is the first
125 known of this genus outside of the Old World. A Paleogene Holarctic distribution for
126 *Eupsenella* implies that it diverged from other bethyline genera when America and Eurasia
127 were still connected (last connection across North Atlantic: ca. 50 Ma; Tiffney, 1985). This
128 reflection also applies to its sister-taxa, *Lytopsenella*, now confined to Chile but with
129 representatives known from the European Baltic and Rovno amber (Azevedo 2009, Ramos *et*
130 *al.* 2014). *Eupsenella* and *Lytopsenella* are the basalmost two genera of the Bethylinae

131 (Ramos & Azevedo 2020), so this scenario is consistent with what is known of the subfamily
132 phylogeny.

133 The regression of *Eupsenella* is likely correlated to the general climate cooling in the
134 last 50 Ma. The Green River Ypresian Mean Annual Temperature (MAT) have been
135 estimated for various sites (e.g., Wilf 2000, Fricke & Wing 2004, Archibald *et al.* 2011) and
136 mid-mesothermal values have been obtained (~16°C; Wolfe 1975). The presence of
137 crocodylians fossil remains is further evidence of a subtropical climate for the Green River
138 Formation (Brochu 2010). The climate of the Paris basin during the early Eocene was
139 certainly paratropical, given its occurrence during the Early Eocene Climatic Optimum and
140 the presence of the Amazon rainforest trees *Hymenaea oblongifolia* (Jossang *et al.* 2008).
141 Similarly, the Eocene Fushun Forest climate was moist subtropical, as suggested by the
142 entomofauna (Wang *et al.* 2014). Finally, the distinct Baltic amber and Rovno amber forests
143 were respectively warm-temperate and subtropical environments ((Sadowski *et al.* 2017,
144 Perkovsky, 2016). This range of climates matches what exists in today Australia, the climate
145 being tropical in the North, subtropical along East and West coasts and warm temperate in the
146 South and Tasmania (Sturman & Tapper 2006). Only *Eupsenella nanda* Ramos & Azevedo,
147 2012 have been collected in a desertic and arid environment, near Alice Springs in the center
148 of Australia.

149 *A contrario*, the *Eupsenella* species for which the biology is known parasitize
150 lepidopterous larvae belonging to the Tortricidae (Gordh & Harris 1996, Berry 1998, Paull &
151 Austin 2006). This family has several fossil representatives in Baltic amber and two in the late
152 Eocene Florissant beds (USA) but still currently shows a worldwide distribution (Heikkilä *et*
153 *al.* 2018) unlike *Eupsenella*. Thus, the host-parasite association of *Eupsenella* and Tortricidae
154 is not sufficient to explain the past and present distribution of the genus.

155 A similar pattern of regression toward tropical or subtropical latitudes has been
156 observed in various hymenopteran families, e.g., Megalyridae (*Megalyra* Westwood, 1832 in
157 Baltic amber; Poinar & Shaw 2007), Formicidae (*Carebara* Westwood, 1840, *Gesomyrmex*
158 Mayr, 1868, *Tetraoponera* Smith, 1852 in Baltic, Rovno and Bitterfeld ambers; Perkovsky
159 2016), Diapriidae (*Doliopria* Kieffer, 1910 in Baltic amber; Brazidec & Vilhelmsen 2022),
160 Mymaridae (*Borneomymar* Huber, 2002 in Baltic amber; Engel *et al.* 2013) or Scolebythidae
161 (*Pristapenesia* Brues, 1933 in Baltic amber; Brues 1933). The regression of *Eupsenella* could
162 be documented by screening younger ambers from southeastern Asia, e.g., Miocene Zhangpu
163 amber, China (Wang *et al.* 2021), Sumatra amber, Indonesia (Ngô-Muller *et al.* 2019), or
164 McGrath Flats, Australia (McCurry *et al.* 2022).

165

166 **Acknowledgments**

167 The images are courtesy of Talia Karim, University of Colorado Museum of Natural
168 History and we warmly thank her. We express our gratitude to the Editor-in-Chief of
169 Palaeoentomology, Prof. Diying Huang for his guidance throughout the submission process,
170 as well as Prof. Alexandr. P. Rasnitsyn and an anonymous reviewer for their useful
171 comments. This work is part of Manuel Brazidec Ph.D. project on “The role of greenhouses
172 on the diversification and evolution of chrysidoid wasps”.

173

174 **References**

175 Archibald, S.B., Johnson, K.R., Mathewes, R.W. & Greenwood, D.R. (2011) Intercontinental
176 dispersal of giant thermophilic ants across the Arctic during early Eocene hyperthermals.
177 *Proceedings of the Royal Society of London (B)*, 278, 3679–3686.
178 <https://doi.org/10.1098/rspb.2011.0729>

179 Azevedo, C.O. (2009) Synopsis of *Lytopsenella* Kieffer (Hymenoptera: Bethylidae). *Zootaxa*,
180 2286, 58–64.
181 <https://doi.org/10.5281/zenodo.191267>

182 Azevedo, C.O., Alencar, I.D.C.C., Ramos, M.S., Barbosa, D.N., Colombo, W.D., Vargas, J.
183 M. & Lim, J. (2018) Global guide of the flat wasps (Hymenoptera, Bethylidae). *Zootaxa*,
184 4489, 1–294.
185 <https://doi.org/10.11646/zootaxa.4489.1.1>

186 Berry, J.A. (1998) The bethyline species (Hymenoptera: Bethylidae: Bethylinae) imported
187 into New Zealand for biological control of pest leafrollers. *New Zealand Journal of*
188 *Zoology*, 25, 329–333.
189 <https://doi.org/10.1080/03014223.1998.9518162>

190 Brazidec, M. & Vilhelmsen, L. (2022) New species of belytines and diapiirine wasps
191 (Hymenoptera: Diapriidae) from Eocene Baltic amber. *European Journal of Taxonomy*,
192 813, 57–86.
193 <https://doi.org/10.5852/ejt.2022.813.1733>

194 Brochu, C.A. (2010) A new alligatoroid from the lower Eocene Green River Formation of
195 Wyoming and the origins of caimans. *Journal of Vertebrate Paleontology*, 30, 1109–1126.
196 <https://doi.org/10.1080/02724634.2010.483569>

- 197 Brues, C.T. (1933) The parasitic Hymenoptera of the Baltic amber, Part 1. *Bernstein-*
198 *Forschungen*, 3, 4–178.
- 199 Cockerell, T.D.A. (1907) A new fly (fam. Mycetophilidae) from the Green River beds.
200 *American Journal of Science*, 136, 285–286.
201 <https://doi.org/10.2475/ajd.s4-23.136.285>
- 202 Cockerell, T.D.A. (1925) Plant and insect fossils from the Green River Eocene of Colorado.
203 *Proceedings of the United States National Museum*, 66 (2556), 1–13.
204 <https://doi.org/10.5479/si.00963801.66-2556.1>
- 205 Cole, R.D. & Picard, M.D. (1978) Comparative mineralogy of nearshore and offshore
206 lacustrine lithofacies, Parachute Creek Member of the Green River Formation, Piceance
207 Creek Basin, Colorado and eastern Uinta Basin, Utah. *Geological Society of America*
208 *Bulletin*, 89 (10), 1441–1454.
209 [https://doi.org/10.1130/0016-7606\(1978\)89<1441:cmonao>2.0.co;2](https://doi.org/10.1130/0016-7606(1978)89<1441:cmonao>2.0.co;2)
- 210 Colombo, W.D., Perkovsky, E.E. & Azevedo, C.O. (2020) Phylogenetic overview of flat
211 wasps (Hymenoptera, Bethyridae) reveals Elektroepyrinae, a new fossil subfamily.
212 *Palaeoentomology*, 3 (3), 296–283.
213 <https://doi.org/10.11646/palaeoentomology.3.3.8>
- 214 Colombo, W.D., Perkovsky, E.E., Waichert, C.C. & Azevedo, C.O. (2021) Synopsis of the
215 fossil flat wasps Epyrinae (Hymenoptera, Bethyridae), with description of three new
216 genera and ten new species. *Journal of Systematic Palaeontology*, 19, 29–89.
217 <https://dx.doi.org/10.1080/14772019.2021.1882593>
- 218 De Ploëg, G.D. & Nel, A. (2004) A new bethylid wasp from the Lowermost Eocene amber of
219 France (Hymenoptera: Bethyridae: Bethylinae). *Geologica acta*, 2, 75–88.

- 220 Dlussky, G.M. & Rasnitsyn, A.P. (2002) Ants (Hymenoptera: Formicidae) of Formation
221 Green River and some other Middle Eocene deposits of North America. *Russian*
222 *Entomological Journal*, 11, 411–436.
- 223 Engel, M.S., McKellar, R.C. & Huber, J.T. (2013) A fossil species of the primitive mymarid
224 genus *Borneomymar* (Hymenoptera: Mymaridae) in Eocene Baltic amber. *Novitates*
225 *Paleoentomologicae*, 5, 1–8.
226 <https://doi.org/10.17161/np.v0i5.4651>
- 227 Fricke, H.C. & Wing, S.L. (2004) Oxygen isotope and paleobotanical estimates of
228 temperature and $\delta^{18}\text{O}$ -latitude gradients over North America during the early Eocene.
229 *American Journal of Science*, 304, 612–635.
230 <https://doi.org/10.2475/ajs.304.7.612>
- 231 Gordh, G. & Harris, A. (1996) New record and a new species of *Eupsenella* (Hymenoptera:
232 Bethyliidae) in New Zealand. *Journal of the Royal Society of New Zealand*, 26, 529–536.
233 <https://doi.org/10.1080/03014223.1996.9517523>
- 234 Grande, L. (2013) *The lost world of fossil lake: snapshots from deep time*. University of
235 Chicago Press, Illinois, 425 pp.
- 236 Heikkilä, M., Brown, J.W., Baixeras, J., Mey, W. & Kozlov, M.V. (2018) Re-examining the
237 rare and the lost: a review of fossil Tortricidae (Lepidoptera). *Zootaxa*, 4394, 41–60.
238 <https://doi.org/10.11646/zootaxa.4394.1.2>
- 239 Hellawell, J. & Orr, P.J. (2012) Deciphering taphonomic processes in the Eocene Green River
240 Formation of Wyoming. *Palaeobiodiversity and Palaeoenvironments*, 92, 353–365.
241 <https://doi.org/10.1007/s12549-012-0092-6>
- 242 Hong, Y. (2002) *Amber insects of China*. Science and Technology Press, Beijing, 653 pp.

243 Johnson, R.C., Mercier, T.J. & Brownfield, M.E. (2014) Spatial and stratigraphic distribution
244 of water in oil shale of the Green River Formation, using Fischer Assay, Piceance Basin,
245 northwestern Colorado. *U.S. Geological Survey Open-File Report*, 1059, 1–59.
246 <https://doi.org/10.3133/ofr20141059>

247 Jossang, J., Bel-Kassaoui, H., Jossang, A., Seuleiman, M. & Nel, A. (2008) Quesnoin, a novel
248 pentacyclic *ent*-Diterpene from 55 million years old Oise amber. *Journal of Organic*
249 *Chemistry*, 73, 412–417.
250 <https://doi.org/10.1021/jo701544k>

251 Lanes, G.O., Kawada, R., Azevedo, C.O. & Brothers, D.J. (2020) Revisited morphology
252 applied for Systematics of flat wasps (Hymenoptera, Bethyridae). *Zootaxa*, 4752, 1–127.
253 <https://doi.org/10.11646/zootaxa.4752.1.1>

254 MacGinitie, H.D. (1969) *The Eocene Green River flora of northwestern Colorado and*
255 *northeastern Utah*. University of California Press, Berkeley, 202 pp.

256 Martynova, K.V., Perkovsky, E.E., Olmi, M. & Vasilenko, D.V. 2019. New records of Upper
257 Eocene chrysidoid wasps from basins of Styr and Stokhod rivers (Rovno amber).
258 *Paleontological Journal*, 53, 998–1023.
259 <https://doi.org/10.1134/S0031030119100125>

260 McCurry, M.R., Cantrill, D.J., Smith, P.M., Beattie, R., Dettmann, M., Baranov, V., Magee,
261 C., Nguyen, J.M.T., Forster, M.A., Hinde, J., Pogson, R., Wang, H., Marjo, C.E.,
262 Vasconcelos, P. & Frese, M. (2022) A Lagerstätte from Australia provides insight into the
263 nature of Miocene mesic ecosystem. *Sciences advances*, 8, eabm1406.
264 <https://doi.org/10.1126/sciadv.abm1406>

- 265 Ngô-Muller, V., Garrouste, R., Pouillon, J., Christophersen, V., Christophersen, A. & Nel, A.
266 (2019) The first representative of the fly genus *Trentepohlia* subgenus *Mongoma* in amber
267 from the Miocene of Sumatra (Diptera: Limoniidae). *Historical Biology*, 33, 1–4.
268 <https://doi.org/10.1080/08912963.2019.1610948>
- 269 Paull, C. & Austin, A.D. (2006) The hymenopteran parasitoids of light brown apple moth,
270 *Epiphyas postvittana* (Walker) (Lepidoptera: Tortricidae) in Australia. *Australian Journal*
271 *of Entomology*, 45, 142–156.
272 <https://doi.org/10.1111/j.1440-6055.2006.00524.x>
- 273 Peñalver-Mollá, E. (1998) *Estudio tafonómico y paleoecológico de los insectos del Mioceno*
274 *de Rubielos de Mora (Teruel)*. Instituto de estudios Turoleses, Teruel, 179 pp.
- 275 Perkovsky, E.E. (2016) Tropical and Holarctic ants in late Eocene ambers. *Vestnik Zoologi*,
276 50, 111–122.
277 <https://doi.org/10.1515/vzoo-2016-0014>
- 278 Poinar, G.O. & Shaw, S.R. (2007) *Megalyra baltica* Poinar and Shaw n. sp. (Hymenoptera:
279 Megalyridae), a long-tailed wasp from Baltic amber. *Zootaxa*, 1478, 65–68.
- 280 Ramos, M.S. & Azevedo, C.O. (2012) Revision of *Eupsenella* Westwood, 1874
281 (Hymenoptera, Bethylidae). *Zootaxa*, 3539, 1–80.
- 282 Ramos, M.S. & Azevedo, C.O. (2020) Revisited phylogeny of Bethylinae (Hymenoptera,
283 Bethylidae) solves basal polytomy. *Insect Systematics & Evolution*, 51, 296–346.
284 <https://doi.org/10.1163/1876312X-00002202>
- 285 Ramos, M.S., Perkovsky, E.E., Rasnitsyn, A.P. & Azevedo, C.O. (2014) Revision of
286 Bethylinae fossils from Baltic, Rovno and Oise amber, with comments on the Tertiary
287 fauna of the subfamily. *Neues Jahrbuch für Geologie und Paläontologie*, 271, 203–228.

288 <https://doi.org/10.1127/0077-7749/2014/0385>

289 Rasnitsyn, A.P. (1990) Hymenoptera. *In*: Ponomarenko, A.G. (Ed.), *Late Mesozoic insects of*
290 *Eastern Transbaikalian*. Nauka Press, Moscow, pp. 177–205.

291 Sadowski, E., Schmidt, E.R., Seyfullah, L.J. & Kunzmann, L. (2017) Conifers of the ‘Baltic
292 amber forest’ and their palaeoecological significance. *Stapfia*, 106, 1–73.

293 Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years of
294 image analysis. *Nature Methods*, 9, 671–675.

295 <https://doi.org/10.1038/nmeth.2089>

296 Scudder, S.H. (1890) The Tertiary insects of North America. *U.S. Geological and*
297 *Geographical Survey of the Territories*, 4, 1–734.

298 <https://doi.org/10.3133/70038967>

299 Smith, M.E., Carroll, A.R. & Singer, B.S. (2008) Synoptic reconstructions of a major ancient
300 lake system: Eocene Green River Formation, western United States. *Geological Society of*
301 *America Bulletin*, 120, 54–84.

302 <https://doi.org/10.1130/B26073.1>

303 Sorg, M. (1988) Zur Phylogenie und Systematik der Bethyilidae (Insecta, Hymenoptera,
304 Chrysidoidea). *Sonderveröffentlichungen des Geologischen Instituts der Universität zu*
305 *Köln*, 63, 1–146.

306 Sturman, A.P. & Tapper, N.J. (2006) *The weather and climate of Australia and New Zealand*.
307 Oxford University Press, 476 pp.

308 Tiffney, B.H. (1985) The Eocene North Atlantic land bridge: its importance in Tertiary and
309 modern phylogeography of the Northern Hemisphere. *Journal of the Arnold Arboretum*,
310 66, 243–273.

- 311 Vršanský, P., Vidlička, L., Čiampor Jr., F. & Marsh, F. (2011) Derived, still living cockroach
312 genus *Carriblatoides* (Blattida: Blattellidae) from the Eocene sediments of Green River in
313 Colorado, USA. *Insect Science*, 19, 143–152.
- 314 Wang, B., Rust, J., Engel, M.S., Szwed, J., Dutta, S., Nel, A., Fan, Y., Meng, F., Shi, G.,
315 Jarzenbowski, E.A., Wappler, T., Stebner, F., Fang, Y., Mao, L., Zheng, D. & Zhang, H.
316 (2014) A diverse paleobiota in Early Eocene Fushun amber from China. *Current Biology*,
317 24, 1606–1610.
318 <https://doi.org/10.1016/j.cub.2014.05.048>
- 319 Wang, B., Shi, G., Xu, C., Spicer, R.A., Perrichot, V., Schmidt, A.R., Feldberg, K., Heinrichs,
320 J., Chény, C., Pang, H., Liu, X., Gao, T., Wang, Z., Ślipiński, A., Solórzano-Kraemer,
321 M.M., Heads, S.W., Thomas, M.J., Sadowski, E., Szwed, J., Azar, D., Nel, A., Liu, Y.,
322 Chen, J., Zhang, Q., Zhang, Q., Luo, C., Yu, T., Zheng, D., Zhang, H. & Engel, M.S.
323 (2021) The mid-Miocene Zhangpu biota reveals an outstandingly rich rainforest biome in
324 East Asia. *Science Advances*, 7, eabg0625.
325 <https://doi.org/10.1126/sciadv.abg0625>
- 326 Wappler, T., Guilber, E., Labandeira, C.C., Hörschemeyer, T. & Wedmann, S. (2015)
327 Morphological and behavioral convergence in extinct and extant bugs: the systematics and
328 biology of an unusual lace bug from the Eocene. *PLoS One*, 10, e0133330.
329 <https://doi.org/10.1371/journal.pone.0133330>
- 330 Wilf, P. (2000) Late Paleocene-early Eocene climate changes in southwestern Wyoming:
331 paleobotanical analysis. *Geological Society of America Bulletin*, 112, 292–307.
332 [https://doi.org/10.1130/0016-7606\(2000\)112<292:LPECCI>2.0.CO;2](https://doi.org/10.1130/0016-7606(2000)112<292:LPECCI>2.0.CO;2)

333 Wolfe, J.A. (1975) Some aspects of plant geography of the Northern Hemisphere during the
334 Late Cretaceous and Tertiary. *Annals of the Missouri Botanical Garden*, 62, 264–279.
335 <https://doi.org/10.2307/2395198>

336

337 **Figure captions:**

338 **Figure 1.** Photograph of *Eupsenella* sp., specimen UCM 58339 (scale bar = 1 mm).

339 **Figure 2.** Line drawing of *Eupsenella* sp., specimen UCM 58339 (dashed lines indicate
340 interpretations; scale bar = 1 mm).

341 **Figure 3.** Distribution map of extant and extinct species of *Eupsenella* showing type localities
342 (green square = extant; orange star = early Eocene; red triangle = late Eocene; question
343 mark = imprecise location).