

The Geotouristic Project "the Geological Adventure" to the Rescue of an Iconic World Heritage Geosite, the Ammonites Slab of Digne-les-Bains (National Geological Nature Reserve of Haute-Provence and Unesco Global Geopark, France)

Didier Bert, Violaine Bousquet, Myette Guiomar, Francesco Bariani, Emmanuel Faure, Jean-Claude Hippolyte, Philippe Bromblet, Jean-Luc Garciaz, Jules Fleury, Patrick Mathieu, et al.

▶ To cite this version:

Didier Bert, Violaine Bousquet, Myette Guiomar, Francesco Bariani, Emmanuel Faure, et al.. The Geotouristic Project "the Geological Adventure" to the Rescue of an Iconic World Heritage Geosite, the Ammonites Slab of Digne-les-Bains (National Geological Nature Reserve of Haute-Provence and Unesco Global Geopark, France). Geoheritage, 2022, 14 (3), pp.101. 10.1007/s12371-022-00739-z . insu-03767996

HAL Id: insu-03767996 https://insu.hal.science/insu-03767996v1

Submitted on 5 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	The geotouristic project 'the Geological Adventure' to the rescue of an
2	iconic world heritage geosite, the Ammonites Slab of Digne-les-Bains
3	(National Geological Nature Reserve of Haute-Provence and Unesco Global
4	Geopark, France)
5	
6	Didier Bert ^{a,b*} , Violaine Bousquet ^c , Myette Guiomar ^a , Francesco Bariani ^a , Jean-Claude Hippolyte ^d ,
7	Philippe Bromblet ^e , Jean-Luc Garciaz ^f , Jules Fleury ^d , Patrick Mathieu ^g , Jean-Simon Pagès ^h , Pierre
8	Tardieu ⁱ , Maud Dupuis-Caillot ^j , Kimberly Leong ^k , Emmanuel Faure ^a
9	
10	* Corresponding author: didier.paleo@gmail.com.
11	^a : Réserve naturelle nationale géologique de Haute-Provence, Conseil départemental des Alpes de
12	Haute-Provence, 13 rue du Docteur Romieu, CS 70216, F-04995 Digne-les-Bains Cedex 9, France.
13	^b : Laboratoire Géosciences, UMR-CNRS 6118, Université de Rennes-1, campus Beaulieu, bâtiment
14	15, F-35042 Rennes cedex, France.
15	^c : Service environnement, Conseil départemental des Alpes de Haute-Provence, 13 rue du Docteur
16	Romieu, CS 70216, F-04995 Digne-les-Bains Cedex 9, France.
17	^d : Aix Marseille Univ, CNRS, IRD, INRAE, Coll France, CEREGE, Aix-en-Provence, France.
18	^e : Centre Interdisciplinaire de Conservation et de Restauration du Patrimoine, 21 rue Guibal, 13003
19	Marseille, France.
20	^f : Laboratoire d'Etudes et de Recherches sur les Matériaux, 23 rue de la Madeleine, CS60136, Arles
21	Cedex 13631, France.
22	^g : Direction du développement, de l'environnement et de l'eau, Conseil départemental des Alpes de
23	Haute-Provence, 13 rue du Docteur Romieu, CS 70216, F-04995 Digne-les-Bains Cedex 9, France.
24	^h : Haute-Provence UNESCO Global Geopark, Musée-promenade, BP 30156, 04990 Digne-les-Bains,
25	France.
26	ⁱ : Bureau ECO ² , 30 avenue St Sylvestre, 06100 Nice, France.

^j: Polymorphe Design, 166 rue du gros bois, 69380 Chazay d'Azergues, France.

- 28 ^k: 42 avenue Demontzey, 04000 Digne-les-Bains, France.
- 29

30 Abstract

Protected under National Nature Reserve, managed by the Departmental Council of the Alpes 31 de Haute-Provence, and labeled UNESCO Geopark, the Ammonites Slab of Digne-les-Bains 32 (SE France) exhibits 1,550 ammonites over a large open-air surface. Despite the virtual 33 absence of special enhancement for the visitors, this spectacular geosite of high scientific and 34 heritage value has played an important role in the years 1990-2000 in the emergence of 35 notions of geological heritage, geological nature reserve, geotourism, and Geopark. Due to its 36 37 exposure to mountain weather conditions, the Slab suffers from significant conservation issues. Several studies have shown that its condition was much worse than expected. Without 38 major restoration work, the Slab was in danger of collapse in the short term. Given the high 39 cost of the needed rescue work, the geotourism project 'The Geological Adventure' was 40 developed under European Interreg ALCOTRA funding. More broadly, this project aims to 41 develop a sustainable experiential geotourism around the geoheritage of the Southwestern 42 Alps cross-border (France-Italy), partly in situ and thanks to digital tools (22 geosites 43 concerned). An important place is given to scientific pedagogy. Concerning the Ammonites 44 45 Slab, the project consisted of: (1) stabilizing and ensuring the sustainability of the fossiliferous layer; (2) planning and developing the overall geosite (parking area, reception 46 area, observation platform); (3) allowing accessibility for people with disabilities; (4) offering 47 progressive and immersive educational mediation to make scientific culture accessible to as 48 many people as possible. 49

50

51 Key-words: Ammonites Slab; Geoconservation; Geotourism; Geological Nature Reserve;
52 UNESCO Geopark; European Funding.

3

53

54 1. Introduction

55

At the beginning of the Jurassic (Sinemurian), 197 million years ago, the sea covered a large 56 part of France. This geological period represents a key moment in connection with the 57 dislocation of Pangea, then the opening of the Tethys Ocean with the flooding of the northern 58 European margin (Fig. 1). The Ammonites Slab of Digne-les-Bains (Southwestern Alps, 59 France) corresponds to an ancient fossilized seabed from this period. This sea floor, formerly 60 horizontal, was then lifted, moved and tilted during the alpine surrection. Today, the result of 61 this long history is an exceptional geological site (Fig. 2), emblematic of the National 62 Geological Nature Reserve of Haute-Provence (RNNGHP by its acronym in French), of the 63 UNESCO Global Geopark of Haute-Provence (UGHP) and of the French Department of the 64 65 Alpes de Haute-Provence (Bert & Pagès 2021). More than 1,550 ammonites are exhibited on an inclined surface of 320 m². The site's location in the open air, along the roadside, allows 66 67 easy access for the public. The Slab is consequently very visited by both the local population and tourists, but also by scientists and schools, including geology students (from 21,000 to 68 30,000 visitors per year according to 2016–2019 data; for comparison, the local population of 69 Digne-les-Bains amounted to 16,068 inhabitants in 2020). The quality and size of this 70 outcrop, the numerous fossils exposed, as well as its scientific and heritage importance, make 71 the Ammonites Slab a unique site in the world. Consequently, it obtained the maximum score 72 of 3 stars (geological site of high heritage value at national/international rank) on the French 73 national inventory of geological heritage managed by the National Museum of Natural 74 History (PAC1989 site: https://inpn.mnhn.fr/site/inpg/PAC1989, accessed in September, 75 2021; De Wever et al. 2014, 2015). 76

Despite its importance and fame, the Ammonites Slab had never been truly developed or 78 landscaped, the bias being to leave it to a certain 'naturalness' (Fig. 3), regardless of a 79 suburban context (Fig. 4). Exposed to weather conditions, the fossiliferous layer can only be 80 deteriorated over time. Since 2014, the Departmental Council of the Alpes de Haute-Provence 81 manages the RNNGHP, and therefore also the Ammonites Slab. Despite the works carried out 82 on several occasions to delay its degradation, a diagnosis carried out in 2016 (Bert et al. 83 2019b and this work) highlighted the imminent risk of partially collapsing of the Slab, 84 involving the need for an emergency response accompanied by major works. The extent of the 85 operations required implied a high cost while taking into account a number of technical and 86 administrative constraints. In this context, the 2015–2016 call for projects of the European 87 Interreg V-A France-Italy Alpes Latines COopération TRAnsfrontalière program 88 (ALCOTRA; Guiomar et al 2018 and this work) was a major opportunity for developing and 89 90 launching a project allowing at the same time: (1) to ensure the protection work of the Ammonites Slab, (2) to enhance this exceptional geosite living up to its international 91 92 reputation, and (3) to go further by networking the geoheritage of the ALCOTRA cross-93 border geographical area with other natural and cultural heritages. All of this with the aim of sustainable local development to meet the objectives of the European program: axis 3 94 'attractiveness of the territory', specific objectives 3.1 ('sustainable tourism by increasing the 95 attractiveness of the territory and preserving the natural and cultural heritage') and 3.2 96 ('improve the management of natural heritage'). It was this project that came into being under 97 the name 'The Geological Adventure'. 98

99

As a possible textbook case about a major conservation threat, through the management and enhancement of an emblematic geosite, which could be of help or interest to be transposed to other geoheritage sites in need, the present work propose to detail (1) the history and context

103	of the Ammonite Slab geosite, (2) the more global geotouristic project that provided the
104	opportunity of a large amount of funding to rescue the site, (3) the rescue operation itself with
105	technical elements for its safeguarding, (4) the new enhancement of the geosite, with
106	structures and scenography elements and (5) some financial benchmarks.
107	
108	2. The Ammonites Slab
109	
110	2.1. Geographic location
111	
112	The Ammonites Slab geosite (44° 07' 10" north; 6° 14' 03" east) is located about 1 km north
113	of the town of Digne-les-Bains (Alpes de Haute-Provence, southeastern France - Fig. 4),
114	along the D900a departmental road on the right edge of the Bléone valley. It is located
115	immediately south of the Isnards farm and was willingly named "la dalle des Isnards" (the
116	Isnards slab) by the local population.
117	
118	2.2. Geological context
119	
120	The Ammonites Slab is representative of the basis of the Digne-les-Bains thrust sheet (Fig. 5 -
121	one of the outermost tectonic units of the southwestern Alps). This thrust sheet is mainly

made of the thick Mesozoic rock sequence of the Subalpine Basin (the pelagic 'dauphinois series'; Haug 1891). The decollement level lies in the Triassic evaporite layers (Goguel 1939), in such a way that the allochthonous thrust sheet has moved over the autochthonous Provençal 'platform' series. Near Digne-les-Bains, the western front of the sheet is a zone of strong tectonic deformation mainly linked to its sliding towards the south. The Ammonites Slab area is located between two dextral strike-slip faults: the Bès fault and the Saint Benoît fault

(Hippolyte et al. 2011; Fig. 5). These two faults have been active in sliding since the upper Miocene at least (10 Ma), with a slip rate estimated to 0.7 mm per year over the last three million years (Hippolyte et al. 2011, 2012). The Ammonites Slab is located in the relay zone between these two faults, in the southern part of the western zone of the thrust sheet, known as the 'La Robine Lobe' (Gidon and Pairis 1988). The eastward dip of the Ammonites Slab results from the sliding of the Digne-les-Bains sheet along these faults.

134

135 **2.3.** Lithology, palaeobiodiversity and biostratigraphy

136

The Ammonites Slab belongs to an alternation of limestone beds and thinner marly interbeds 137 (Fig. 6). The facies suggest subpelagic deposits beneath the storm-wave action limit without 138 any obvious role of currents (Corna et al. 1991; Dommergues and Guiomar 2011). Alongside 139 140 with rare nautiluses of the genus Cenoceras Hyatt, 1884 and a few belemnites, the Slab currently bears more than 1,550 ammonites, often large (up to 70 cm), which make the geosite 141 142 famous. Suspensivorous epibenthic bivalve mollusks and echinoderms (pentacrins in isolated articles) are also very common despite often in debris. This benthic fauna is also clearly more 143 diverse than the nectonic cephalopods, with the presence of at least four bivalve genera 144 (Plagiostoma Sowerby, 1814, Gryphaea Lamarck, 1801, Liostera Douvillé, 1904 and 145 Oxytoma Meek, 1864). In comparison, only three species of ammonites (Arietitidae Hyatt, 146 1874 and Schlotheimiidae Spath, 1923) are present: (1) Coroniceras multicostatum (Sowerby, 147 1824) very largely dominates (99.74% - Fig. 6), (2) Angulaticeras (Sulciferites) cf. 148 charmassei (d'Orbigny, 1844) (0.19%), and (3) Coroniceras (Arietites) sp. (0.07%). This low 149 biodiversity of ammonites, with populations often very largely dominated by a single species, 150 151 was observed throughout the Sinemurian and lower Pliensbachian of the subalpine basin with paleobiogeographic affinities towards the NW of Europe (Burgundy, southern Jura, England; 152

7

see Corna 1985, 1987; Corna and Mouterde 1988; Page 1992, 2003). Despite the more
southerly position of the Ammonites Slab, we can exclude any Tethys influence because of
the total absence of the Phylloceratoidae Zittel, 1884 and Lytoceratoidea Neumayr, 1875.

156

The presence in very large numbers of a single species of ammonite allowed Dommergues 157 and Guiomar (2011) to carry out a detailed paleontological study on the basis of a 3D scan of 158 the Ammonites Slab. This analysis was the first carried out in situ including quantitative 159 160 topics of intraspecific variability in a large Sinemurian ammonite species. Coroniceras multicostatum is a species whose stratigraphic distribution is always very limited, which 161 makes it a reliable index for biostratigraphic purpose. Due to its exceptional exposure 162 conditions, the Ammonites Slab of Digne-les-Bains constitutes to date the best observation 163 point of the Coroniceras multicostatum Interval Horizon, at the extreme top of the 164 165 Coroniceras (Arietites) bucklandi Zone/Subzone (lower Sinemurian, Lower Jurassic, with a numerical age around 197 million years according to Cohen et al. 2013). 166

167

168 2.4. Historical background

169

The precise date of the discovery of the Ammonites Slab was unknown until very recently 170 171 (this work), but it was deemed to be around the middle of the 20th century. No record exists before this period despite the regular visits of many geologists and learned societies in the 172 immediate vicinity (Hebert 1861; Garnier in Vélain and Dieulafait 1872; Haug 1891, 1909; 173 Kilian and Zürcher 1895). Gérard Thomel (palaeontologist specialised in ammonites, 174 formerly curator of the Natural History Museum of Nice, France; pers. comm.) relayed that 175 the local Scientific and Literary Society was at the instigation of the installation of a fence and 176 a sign prohibiting the direct access to the Slab in the 1960s, because of an act of vandalism. 177

Very recently, one of us (JSP) found some new information: actually, the Slab was unearthed in June, 1941 by Mr. Roux, a civil engineer, during work to widen the road of Barles. At that time, less than a quarter of the current surface was exposed. A first excavation work was made in July, 1979 as part of the project to create the National Nature Reserve in the vicinity of Digne-les-Bains (Martini 1979): the surface of the Slab was extended to 160 m² on which there were approximately 600 ammonites. The road has been deviated slightly to leave a little more space for visitors, and a basic layout was proposed (a lectern).

With the birth of the concept of natural heritage and nature reserves in French law (see De 185 Wever et al. 2019), Guy Martini gave rise to the idea of classifying 18 geological sites, 186 including the Ammonites Slab, in the Digne-les-Bains area under this strong legal protection 187 status (Martini 1979; see Guiomar 2009). This project came to realization on October 31, 188 1984 with the creation of the RNNGHP. However, the Ammonites Slab took a long time to 189 190 become known outside the Digne-les-Bains region. The site was officially brought to the attention of the international community only in 1991, following the study by Corna et al. 191 192 (1991) and the holding in Digne-les-Bains of the First International Symposium on the protection of geological heritage (Martini 1994; see also the 1991 International declaration on 193 the rights of the memory of the Earth). The growing notoriety of the Slab, and its diffusion on 194 Japanese television (The Miracle Planet, Chikyû Daikikô, NHK, 1987), led to the realization 195 196 of the molding in 30 pieces of the fossiliferous stratum (the 160 m² then exposed) for the city of Kamaïshi (Iwate Prefecture) in Japan, where it is still displayed at the Iron and Steel 197 History Museum. 198

A second work for securing and clearing the Slab was carried out in July, 1994 with the participation of the University of Bucharest, and led to the doubling of the exposed surface (320 m^2) and the number of visible ammonites (1,550). On this occasion, drilling showed the continuity of the Slab to the north with the presence of the same ammonite density over several thousand square meters; various on-site museum projects did not emerge mainly for financial reasons and the matter ended there. However, under exceptional financing, restoration and security work was carried out in 2009, 15 years after the previous operation. The objective of this stabilization work was to remedy the conservation problems of the Slab in the short term (ideally 2–5 years), still awaiting for a financial opportunity and a political commitment to achieve equipment for lasting protection and enhancement of the site. This latter is presented in this work.

210

211 2.5. Management of the Ammonites Slab by the National Geological Nature Reserve of 212 Haute-Provence (RNNGHP)

213

The RNNGHP is governed by the Environmental Code from French law. Its classification is 214 justified by the scientific interest of the natural heritage, with the objective of preserving the 215 216 integrity of geological sites (including fossils and mineral substances). The RNNGHP now has 18 sites classified as National Nature Reserves (RNN by its acronym in French), including 217 the Ammonites Slab, for a total of 270 ha (currently in the process of being extended - Bert et 218 al. 2019a). These sites are surrounded by a protection perimeter of approximately 2,300 km² 219 (Fig. 7), making it possible to ensure the conservation of the exceptional geological heritage 220 221 in the territory of 59 municipalities of the Provençal Pre-Alps (52 in the Alpes de Haute-Provence and 7 in the Var departments). Throughout the protection perimeter, the removal, 222 destruction or degradation of fossils, minerals and concretions is prohibited. Removals 223 without tools of elements naturally released by erosion are tolerated only if they are carried 224 out in limited quantities. On RNN classified sites, the regulation is more restrictive and 225 prohibit, among other things, any sampling of mineral or fossil substances and any change in 226 227 the character or aspect of the place. Sampling authorisations are issued for scientific purposes

with specific conditions for the two levels of regulation. Concerning the Ammonites Slab, the
strong legal protection of the RNN classification was complemented by land control: first
(1987) by acquisition of the Slab by the township of Digne-les-Bains and then (2018) by its
assignment, extended to a larger part of the geosite bought from its private owner, to the
Departmental Council of the Alpes de Haute-Provence.

233

The Departmental Council of the Alpes de Haute-Provence drew up the third management 234 plan of the RNNGHP (Bert et al. 2019a) through which it is responsible for ensuring (1) the 235 protection of the geoheritage (environmental protection under police missions; physical 236 conservation; management of ex situ collections), (2) the acquisition of knowledge 237 (inventories and scientific research) and (3) the general functioning of the nature reserve. The 238 studies carried out, the rescue work and the opening of the Ammonites Slab to the public, are 239 240 in direct line with these missions. As the RNNGHP manager since 2014, the Departmental Council's first ambition was to regualify the major sites of the nature reserve whose facilities 241 242 were aging. Thus, in order to provide the sites protective and enhancement equipment up to 243 their international reputation, additional funding to the Department Council's investment funds and to the State endowment was sought. 244

245

246 3. The European project 'The Geological Adventure'

247

The Geological Adventure project was carried out within the framework of the European cross-border cooperation program between France and Italy, Interreg V-A ALCOTRA 2014–2020. This project aimed to enhance the great landscapes as well as the remarkable mineral heritage, representative of the history of the Southern Alps, which is illustrated between the Department of the Alpes de Haute-Provence (France) and the Province of Cuneo

(Italy). The project took place over 3 years, from April, 2017 to December, 2020, and 253 mobilized 7 cross-border partners (for France: the Departmental Council of the Alpes de 254 Haute-Provence - project leader and manager of the RNNGHP -, the Provence-Alpes-255 Agglomeration community - manager of the UGHP -, and the municipality of Les Mées; for 256 Italy: the Union of Fossanese municipalities, the municipalities of Cherasco and Frabosa 257 Soprana, and the Federico Sacco Foundation). The Geological Adventure responds de facto to 258 one of the objectives, designated as a priority, of the tourism plan of the Department of the 259 Alpes de Haute-Provence for the period 2016-2020 (ambition renewed for the period 260 2022–2027): to develop experiential and sustainable geotourism. Thus, geoheritage, as a 261 natural quality of territories, is considered as an asset for tourist diversification and a driver of 262 local and economic development (François et al. 2006; Bétard et al. 2017). This is in this 263 context that the financing of the rescue and layout of the Ammonites Slab geosite of Digne-264 les-Bains was made possible. 265

266

267 **3.1. Terms and concepts**

268

The Geological Adventure project is based on several fundamental principles resulting from the reflections on tourism carried out since the end of the 20th century around geotourism, experiential tourism and sustainable tourism. Since there are several definitions around these principles (see Ruban 2015 and bibliography; Gonzalez-Tejada et al. 2017 and bibliography), it is necessary to clarify in which sense the concepts guided our project:

274

Geotourism. Geology and geoheritage (considered here in a very broad sense, including
palaeontology, geochronology, climatic evolution, etc., adapted from Ruban & Ermolaev
2020), are usually considered as a large and complex subject often difficult to access for an

uninitiated audience. Historically, niche tourism in relation to geological heritage rather
concerns geologists (including supervised university internships), enthusiasts, or is based on
the personal appreciation of the mineral landscape from a form of aestheticism (i.e. Gordon
2018; Ruban et al. 2021).

In the 1990s, the term geotourism emerged at a time when notions of sustainability and 282 ecotourism were developing (see Gordon 2018) roughly at the same time as the concept of 283 geoheritage (Gray 2013; de Wever et al 2019). Very early, the RNNGHP clearly stated as an 284 actor in these concepts and initiated the European Geoparks Network with three other 285 European territories and closely collaborated in the development of the Global Geoparks 286 Network (Martini and Zouros 2008). Faced with the need to clarify the concept of geotourism 287 (Gonzalez-Tejada et al. 2017), the 2011 Arouca Declaration (International Congress 288 'Geotourism in Action', 09-13th November, Portugal) defined geotourism as "tourism that 289 290 supports and improves the identity of a territory, taking into account its geology, its environment, its culture, its aesthetic values, its heritage and the well-being of its residents. 291 292 Geological tourism is one of the various components of geotourism". Beyond the scientific component, this growing recognition of the cultural and aesthetic values of geoheritage, in 293 relation to the development of geotourism, is reflected with the emergence and worldwide 294 multiplication of UNESCO Global Geoparks since 2000 (the territory of the RNNGHP was 295 296 one of the first to receive this label; see also Bétard et al. 2017). Their vocation is indeed to explore, develop and celebrate the links between geological heritage and all other aspects of 297 heritage, whether natural, cultural or intangible, and to promote them. This fits with the notion 298 of 'cultural landscapes', a concept adopted by UNESCO in 1992 in which the continuous 299 interaction between natural processes and human activities is emphasized (Gordon 2018). 300

Experiential tourism. The notion of cultural landscapes of geotourism is also part of the 302 principles of experiential tourism described as an emerging trend in the 21st century. 303 Adventure tourism, ecotourism, and heritage or cultural tourism can be considered as 304 experiential tourism (Smith 2006). Experiential tourism develops a participatory aspect that is 305 nowadays highly sought after by tourists (Gordon 2018). In this, the exploration of the 306 geological elements of the landscape through different educational, cultural, even artistic 307 filters, engages in an interactive way the senses, the emotions, the imagination, while 308 309 including the discovery and the learning of new knowledge or the commitment to the history of the place. In all cases, however, didactics must be based on advances in scientific 310 knowledge in the way that the information offered is accurate, reliable and of high quality, 311 even if adapted for an easier understanding. 312

313

314 Sustainable tourism. The objective of sustainable tourism development was defined by the Agenda 21 (1992) in the Sustainable Development plan for the 21th century. The aim is to 315 316 reconcile long-term tourism, whatever its scale, form or type of destination, with the 317 improvement of environmental and social conditions and the maintenance of development capacities (including economic). The principles of sustainable tourism were defined in the 318 Charter for Sustainable Tourism of Lanzarote (UNWTO 1995), then updated in 2004 by the 319 Committee for Sustainable Tourism Development of the Word Tourism Organization 320 (UNWTO 2005, p. 11). While geotourism can offer tourists an enriching experience (and this 321 is clearly also a major issue in sustainable tourism), it can also enable them to address 322 sustainability issues (Bétard et al. 2017). This is also a very present theme in geology through 323 the apprehension of the deep time. In all cases, it is up to the stakeholders (including geosite 324 developers) to guide geotourists towards raising awareness and adopting responsible behavior. 325

14

- 327 **3.2.** The objectives of the Geological Adventure project
- 328

If the southern cross-border Alps (Alpes de Haute-Provence, France and Cuneo region, Italy) 329 have many exceptional geological sites well known to insiders, the Geological Adventure 330 offers to (re)discover some of them through an innovative and modern approach in line with 331 the Arouca Declaration (2011). While other geotourism development projects in the Southern 332 Alps are more oriented towards roaming alone (for example the Via GeoAlpina project, which 333 also provides didactic cards e- see Cayla & Hobléa 2011) or around an ex situ museographic 334 aspect (Cayla 2009), the Geological Adventure proposes to develop sustainable, experiential 335 and joyful tourism through integrated site developments and digital technologies. The 336 geological history of the Southern Alps is the common thread of this geotourism destination, 337 physically or virtually connecting exceptional geosites with other heritages. To do this, the 338 339 Geological Adventure offers various levels of interactivity: emotional, physical or intellectual, if possible in creating the conditions for a lived experience, the aim being to make the 340 341 geological heritage accessible to as many people as possible.

342

343 **3.3.** The achievements of the Geological Adventure

344

The results of Franco-Italian deliveries within the framework of the Geological Adventure are rich in diverse achievements. The present work mainly details those carried out by the Departmental Council of the Alpes de Haute-Provence (France) as responsible and leader of the project. In order to meet the objectives, the actions focused on four axes:

349

350 *The physical development of the geosites.* Five spectacular geosites (3 in France and 2 in 351 Italy) have benefited from specific developments with the dual objective of long-term conservation and heritage enhancement. These are (1) the Ammonites Slab of Digne-lesBains (the main project here developed), (2) the Pénitents des Mées trail (Sensitive Natural
Site of the Department), (3) the belvedere of the Vélodrome geosite at the Vieil Esclangon
(RNNGHP, France), (4) the Cave of Bossea at Frabosa Soprana and (5) the Rio Crosio at
Cherasco (Italy).

357

358 *Scientific mediation.* Twenty-two geosites have been selected to illustrate different steps in 359 the geological history of the Southern Alps (the interactive map of the whole geosites is 360 available at: https://aventuregeologique.com/decouvrir/, accessed in September, 2021). To 361 make this story attractive, the mediation strategy was based on:

- the simplification of the scientific discourse in order to make accessible the concepts
 of geology, often complex, but by limiting the drifts which can be related to the
 simplification of the scientific concepts. This difficult work of science popularization
 required a lot of back and forth between didactic professionals and scientists;
- the design of educational tools offering an original and attractive aesthetic (timelines;
 geosite sheets; lecterns), and promoting knowledge and general understanding of the
 events by all audiences;
- the offer on certain developed sites, and in particular on the Ammonites Slab, of
 interactive didactic instruments (casts, 3D models and reconstructions);

the creation of an educational package around the geological history of the Alps
designed around sequences of playful handling and experimentation for
schoolchildren;

a story-telling in which Federico Sacco and Edouard-Alfred Martel, two illustrious
promoters of geotourism at the start of the 20th century, icons of the project, invite the
public to take a 'journey through time and space';

the production of a four-minute animated fiction retracing the geological history of the
Southern Alps; the elaboration of eight video sequences in augmented / virtual reality
in the way to explain the genesis of current landscapes or to stage fossilised animals in
their original environment (ammonites, ichthyosaurs, sirenians, rudists, etc.);

- the development in Fossano (Italy) of a modern museum space dedicated to the
 enhancement of F. Sacco's paleontological collection;
- the financing of an artist-made fresco representing the living palaeoenvironment of the
 urgonian deposits and displayed at the Municipal Museum of Orgon (Urgonian
 stratotype locality, France).
- 386

Networking. With the aim of inviting the visitors to discover geoheritage, 18 circuits, for one day or over several days, have been created in the territory of the project around 7 themes (sporting, cultural, contemplative, biodiversity, etc.). Each of the circuits includes the discovery of at least one geological site and combines other cultural, natural or intangible heritages.

392

Digital tool. To attract the public attention (including the young public), and to embrace a 393 certain modernity, the digital tool was positioned at the heart of the strategy of mediation, 394 of 395 valuation and promotion the Geological Adventure. А website (https://aventuregeologique.com/, accessed in September, 2021) and a Smartphone application 396 (https://play.google.com/store/apps/details?id=com.easymountain.cd04, accessed in 397 398 September, 2021) were thus deployed. The website allows the public to prepare their tourist trip upstream by providing complete information on the geosites. The Smartphone application 399 400 invites people to live an in situ active experience through thematic circuits and interactive 401 games using geolocation. These games, intended for young audiences, provide scientific

402 concepts and raise awareness of the protection of natural heritage. Tactile tables left for free
403 consultation allow the dissemination of all the digital content of the project; they have been
404 installed in two museums and five tourist offices in the territory of the Alpes de Haute405 Provence.

406

407 **3.4. Discussion**

408

The Geological Adventure has proved very demanding in terms of project management and in 409 particular the mediation aspect. It was necessary to find the right tone: if the discourse 410 remains too technical, we lose part of the lay audience, and conversely, if we simplify the 411 scientific discourse too much, the risk is to deliver erroneous or irrelevant information 412 because of lacking content (Martin et al. 2010). Injecting an emotional and imaginary 413 414 component into didactics via story-telling is also tricky because it can spawn confuse between reality and fiction to a general lay audience. This requires sobriety and unequivocal 415 416 explanations on the media facilities.

417 While the digital tool offers a wide range of possibilities, it also has drawbacks. The experience of the Geological Adventure shows the limits of digital deployment in rural and 418 mountain areas where the telecommunication network does not yet cover the entire territory. 419 The public is faced with uneven signal quality as well as difficulty in downloading the 420 website or Smartphone application in the field. These conditions are obviously not favorable 421 to the systematic downloading of cultural applications by Smartphone users, especially for 422 heavy applications, as the products provided need to be light and quick to download. The 423 Geological Adventure application presents numerous photos and videos, which weigh down 424 425 its content and impose a long download time: some users may therefore become demobilized. In addition, in the digital field where technologies evolve very quickly, the tools developed 426

427 can quickly become obsolete. It is relevant to question the durability of the various
428 informative materials, of which the most classic (lecterns, in situ information panel, etc.)
429 sometimes appear more robust over time and less demanding in terms of management,
430 maintenance and updates.

Finally, as an experiential guarantee, we must not forget to put the human at the heart of the
mediation and promotion processes. An all-digital/technological solution cannot replace the
actors of the territories who are vectors of emotion and authenticity.

It should be noted that almost all the achievements within the framework of the Geological Adventure are free access and free of charge to the public, including the developed geosites. The only realizations with entrance fees are the Bossea Cave site, the F. Sacco Museum and the two museums where the digital tables are displayed. The project as a whole is thus ranked relatively high on the accessibility scale of Mikhailenko et al. (2021), with an overall average score of 15/20 for all the developed geosites (from 8/20 to 20/20 depending on the geosites, including 17/20 for the Ammonites Slab of Digne-les-Bains).

441

442 **4.** The Ammonites Slab rescue operation

443

```
444 4.1. The facts
```

445

The Ammonites Slab geosite is located in a bottom valley subject to mountain climatic conditions. Water infiltrations are present (Fig. 8g) and the limestone rock of the Slab shows many surface desquamation or delamination (Fig. 8a). Some ammonites are fractured due to the opening of joints (Fig. 8b). Ammonite repair, patching or sealing operations have already been carried out using various mortars for years (Fig. 8b). However, more recently, a block of rock (35 X 20 cm – Fig. 8c) illegally collected during May 2016 at the base of the Ammonites

Slab revealed the presence of a large void formed naturally at the interface of the limestone 452 layer (the Slab) and the underlying marlstones. The presence of such other voids could be 453 legitimately suspected. Their formation seemed recent, no cavity of this importance having 454 been observed before (the bottom of the Slab, however, showed a slight swelling - Fig. 8d). 455 Apart from this isolated act of vandalism, the unrestricted accessibility of the fossiliferous 456 layer to the public has resulted in small damage, as well as wear of the fossils due to repeated 457 contact with the hands of visitors (particularly present at children height - Fig. 8e-f); although 458 it was forbidden, some unscrupulous visitors had even taken the risk of climbing the fossil 459 wall! 460 These observations showed the alarming condition of the geosite. The alterations affect the 461 ammonites, their calcareous matrix and, more worryingly, the substratum of the Slab itself. 462 Diagnoses and precise studies were necessary in order to identify the conservation problem as 463 464 a whole and to consider the appropriate treatments to remedy it. 465 4.2. The studies 466 467 4.2.1. The voids 468 469 A geophysical radar study was carried out by the Laboratoire d'Etudes et de Recherches sur 470 les Matériaux (LERM, Arles, France - October, 2016), in order to detect and map 471 heterogeneous zones, or voids, under the fossiliferous stratum. 472 473 *Method.* The principle of geophysical radar uses the reflectometry of electromagnetic waves 474 475 at high rate pulses of short duration. Waves are differentially reflected at the interface of materials with different physical characteristics. The geophysical radar used is a LERM model 476

20

with dual frequency antenna. The measurement was carried out according to the followingparameters:

- double frequency of pulses at 400 MHz and 1 GHz;
- scan duration: 10 ns (1 GHz) and 35 ns (400 MHz), which corresponds to an
 investigation depth of respectively 0.5–1.50 m in the context of the Slab;
- realization of 35 measurement profiles spaced from 50 cm apart along the line of
 greatest slope (length of the profiles 5–24 m);
- vertical resolution: 512 samples per scan (i.e. 1–3 mm per sample); horizontal
 resolution: 50 scans per meter (i.e. 1 measurement every 2 cm).

486 The raw signal (Fig. 9) was processed so as to provide an anomaly mapping at two different
487 depths (10-40 cm and 50-80 cm respectively).

488

Results. The first mapping at 10–40 cm depth (Fig. 10a) shows heterogeneities located mainly in the lower south part of the Slab, around 0.5–3 m from the bottom, for a cumulative surface area of approximately 6 m². The voids are probably not all in contact, but can extend over more than 1 m² continuously. These heterogeneities are distributed over two levels, respectively 10 and 20 cm deep, for an estimated thickness of 2–15 cm. Other heterogeneous areas are located in the upper part of the Slab, in the 2–3 m below the summit, for a varying thickness of 10–50 cm. Only the north lower part of the Slab seems homogeneous.

The second mapping at a 50–80 cm depth (Fig. 10b) shows heterogeneities partly correlated
with the previous ones, but could be linked to inconsistent materials at an interface between
two strata.

499

500 4.2.2. Deformation and fracturing

New 3D photogrammetric acquisitions led to the study of the geometry and fracturing of the
Slab, as well as to produce a digital model, made by the Centre Européen de Recherche et
d'Enseignement des Géosciences de l'Environnement (CEREGE, Aix-Marseille, France 2017–2018).

506

Method. Eight hundred high definition constant distance photos were taken (material used: 507 full frame digital camera of 36 Mpx). These images have a sufficient resolution (0.5 mm) to 508 models of the ammonites (Fig. 6: 509 allow making digital see for example 510 https://sketchfab.com/models/b1299fc681314243a14a577775ca898a, accessed on September, 2021), and they constitute a database for monitoring the condition of the Slab and the 511 ammonites over the long term from the previous data (Dommergues & Guiomar 2011). A 512 513 second drone acquisition of 280 photos was carried out in order to remove acquisition artifacts (cast shadows, safety ropes, etc.). The 3D digital model was produced as a point 514 cloud, which the surface was interpolated and textured (https://skfb.ly/6t9FS, accessed on 515 516 September, 2021). The point cloud was then projected onto the average surface plane of the Slab to constitute a digital terrain model (DTM). The images used for the realization of the 517 DTM were also used to make an ortho-rectified and calibrated photo of the entire Slab 518 allowing precise digital studies (237 million pixels for a ground pixel resolution of 1.67 mm -519 Fig. 11). 520

In addition, 22 geodetic markers were placed on the Slab and its periphery as part of themonitoring of possible deformations of the site over the long term.

523

Results. The DTM and orthophoto were used in 2D and 3D to map the fracturing of the
Ammonites Slab in a 3D reference system. The model now allows a precise measurement of
the Slab:

527	• Surface: 310 m ² ;	
528	• Perimeter: 79.6 m;	
529	• Width: 16.5–11.7 m (the latter at the top of the Slab);	
530	• Height: 24.8 m;	
531	• Tilt: 48 °;	
532	• Orientation: 167°N (due to this slightly north orientation and its inclination, me	
533	is the best for ammonites observation; in the afternoon the site is backlight).	
534 It appears that the Slab has the general shape of a horizontal gutter due to flatness def		
535	5 exceeding 40 cm in amplitude (Fig. 12). This particular shape results essentially fr	
536	overhanging projection of the upper part of the fossiliferous layer (with respect	

norning

ects not om the to the 537 presence of a horizontal fracture) and the bulging of its lower part, where the most voids were observed (see above chapter 4.2.1). This bulge is asymmetrical, present only in the south part 538 of the Slab where it reaches 15 cm in amplitude (Fig. 8d); it is accompanied by a detachment 539 540 of the limestone layer, already observed visually (chapter 4.1 – Fig. 8c), and open star 541 fractures.

542

Analyzes and cartography of fracturing (Fig. 13). Analysis and mapping of the fractures 543 typology allowed reconstructing their history from the deposition of sediments until the 544 present time: 545

Irregular diagenetic joints: without really being fractures, they are sinuous 546 • discontinuities more clayey than the rest of the Slab. These joints are often deep 547 because they are easily eroded by the rainwater flow; they weaken the stratum while 548 contributing to its bumpy appearance. 549

Normal faults and tension joints (Fig. 14a): a dense series of horizontally oriented 550 • joints and faults chop the Slab. The faults almost always show a calcite filling and 551

have small normal slips (therefore perpendicular to the surface plan of the stratum), which can cut and offset ammonites. The tension joints have the same orientation and the presence of calcite suggests that they were formed at the same time than the normal faults in an underwater environment. These structures would therefore be linked to the deformations produced during the rifting phase of the Tethys Ocean, which created the western margin of the Digne-les-Bains thrust sheet basin during the Lower Jurassic.

- Strike-slip faults, calcite joints and associated stylolithic joints (Fig. 14b-d): the largest 559 strike-slip fault cuts the Slab in half through the middle (dextral sliding movement, the 560 upper part having moved to the north $-\frac{\text{Fig. 14c}}{\text{Fig. 14c}}$; calcite joints were formed in the 561 extension relay zones (Fig. 14b). In contrast, stylolithic joints were formed at the end 562 of the strike-slip fault in the compression areas. This strike-slip fault shows a total 563 sliding of 10.2 cm (measured on the early diagenetic joints and the offset ammonites -564 Fig. 14d). The reactivation of these strike-slip cracks is probably linked to the 565 beginning of the activity of the Bès and St Benoît faults, nearly 10 Ma ago, as part of 566 the formation of the Alps. 567
- Joints without calcite: short compression joints without calcite filling are present at the
 end of the strike-slip faults; some appear to be in shear. They probably all resulted
 from compressive deformations during the tilting of the Ammonites Slab.
- <u>Gravity and weathering fractures</u>: they are mainly present in the south and upper parts of the Slab and are associated with the tilting/overhanging of this part. They were probably aggravated by mechanical unloading following the removal of the rock cover, above the fossiliferous layer, during the excavation works of 1979 and 1994.
- 575
- 576 **4.2.3. Discussion and recommendations**

Some of the fractures are filled with calcite (Fig. 14e) and are therefore stable and safe for the 578 sustainability of the Ammonites Slab. On the other hand, other fractures are open and 579 participate in its degradation. After 197 million years of history, the degradation of the Slab 580 appears to be a very recent and rapid phenomenon partly of anthropogenic origin: it results 581 from open air exposure following the withdrawal of the upper layers, which protected it until 582 then. The too rapid mechanical unloading of the rocks associated with exposure to weather 583 conditions favored the opening of fractures and infiltration. This degradation is favored by 584 exposure to ENE and seasonal freezes and thaws alternating. 585

The numerous fillings carried out during the work operation in 2009, to slow down the degradation of the Slab, did not prevent the aggravation of the gravitational bulge and the opening of star fractures that recently appeared in its basal-south part. This deformation is associated, on the one hand, with anomalies detected by geophysical radar (cavities, heterogeneities - Fig. 10), and on the other hand with a dense network of open fractures in the weathered parts (Fig. 13), which allows the infiltration of rainwater.

592 The infiltrated water is blocked by the underlying marly layer and by the shotcrete sprayed in 2009 on the southern edge of the Salb (Fig. 11, left), even though equipped with barbicans for 593 drainage, which was supposed to slow its erosion and lateral collapse. The chemical reactions 594 595 caused by the presence of water in the pyrite-rich marls cause, among other things, the proliferation of small gypsum crystallizations. These crystallizations facilitate the detachment 596 of the marlstone sheets and their swelling in the presence of humidity, which constrains the 597 limestone layer of the fossiliferous stratum. At the same time, the pressurization of the 598 infiltration water causes a reduction in the friction at the base of the limestone layer and 599 600 causes the crawling of the southern half of the Slab (the most altered part), of which the local bulge is the most obvious manifestation. 601

As it stands, these observations suggest a risk of destabilization of the Ammonites Slab in the more or less short term, with the possibility of a sudden rupture of the south part of the limestone layer by a buckling mechanism, which would lead to the irreparable loss of part of the geosite.

606

It therefore appeared that to ensure the protection of the geosite, a rapid intervention was necessary. This operation was intended to limit the percolation of water, but also to ensure the maintaining (by anchoring) of the compartments destabilized by joints and fractures, in order to limit the effects of erosion by raveling and prevent them from falling. The elimination of the shotcrete from the lower/south part was also a necessity to allow a better flow of seepage water and to reduce the hydrostatic pressure.

Note that the problem of anthropogenic erosion, linked to free access to the fossiliferouslayer, should also be resolved.

615

616 **4.3. Sealing device**

617

The installation of reinforcements by sealed metal inclusions (Fig. 15) is particularly complex 618 in a heritage site with such a high density of ammonites to be protected, which considerably 619 limits the methods and possibilities of surface installations. Drilling and coring tests without 620 percussion with a fine implantation were performed to avoid any destabilization and to avoid 621 the ammonites. They made it possible to carry out tensile tests on anchor bars sealed in the 622 fossiliferous limestone using a hollow traction cylinder. The marls were sampled for 623 laboratory analysis in order (1) to test the interactions with the chemical formulation of the 624 envisaged sealing grout, (2) to test the swelling potential and (3) to analyze the corrosivity of 625 the ground. These investigations enabled: 626

to confirm the presence of heterogeneous materials/cavities to define the quantity of
grout to be injected through the boreholes made for the installation of the anchors;

- to show that the soil corrosivity is low (Ph=7.7; chloride ion concentration
 =2.2 mg/kg; sulfate ion concentration =11.8 mg/kg; resistivity at 20°C=5.700 Ω/cm),
 which allows to optimize the composition of the sealing grouts and to define the
 protective treatment for the steel of the anchor bars;
- to define the boreholes diameter and the reinforcement bars diameter to avoid any risk
 of degradation of the fossiliferous layer;
- to define the density and number of anchor bars required (minimum mesh size) while
 avoiding ammonites;
- to test the lateral friction on the limestones during the conformity tensile tests
 according to the actual thickness of the beds;
- to calculate the necessary length for the sealing bars, knowing that the thickness of the
 area to be reinforced is limited to 0.6–0.75 m depth below the surface.

In order to control the movements of the Slab during the drilling and injection of the filling
and sealing grouts (topographic monitoring), sensors of piston extensometer type were
implemented for the duration of the intervention.

644

645 *Choice of the filling/sealing grout.* The main problem is the presence of pyrite in the 646 marlstones underlying the Ammonites Slab, which induces the risk of sulphate reactions with 647 the production of gypsum and sulfuric acid. In such a humid and porous environment, some 648 cement compounds can go into solution and react with pyrite. Sulphates can also lead to the 649 formation of thenardite crystals (anhydrous sodium sulphate - Na_2SO_4) or mirabilite (sodium 650 sulphate decahydrate - Na_2SO_4 , $10H_2O$) further increasing the risk of marls swelling and 651 detachment of the Slab. This means that the sealing grout must have a limited content of sulphates, but also a reduced content of tricalcium aluminate (C_3A) to limit the formation of expansive secondary compounds (secondary ettringite).

For these reasons, the use of portland cements was excluded. In contrast, natural hydraulic lime has low sulphate content and is therefore more stable, but it has a long setting time, low resistance with a risk of going into solution in the presence of water. Air lime/portland cement mixtures would guarantee lower soluble alkali levels than cements alone, but with the danger of a high presence of more soluble portlandite. These solutions were therefore discarded for the treatment of the Ammonites Slab.

According to the method of Zhang et al. (2014; 2018), laboratory tests of several potential grouts were carried out using a 5 kg sample of crushed marls taken directly from the Slab site. Under in situ conditions, the use of resistant sulphate cement should be preferred to avoid the risk of swelling and cracking by delayed hydrate formation, which could jeopardize the integrity of the Slab despite the seals. To fix the lime and avoid efflorescence, compound cement with a low C_3A content based on pozzolan of CEM III type was chosen (Declair 2008). The chemical characteristics of this cement are summarized in Table 1.

667

Table 1. Chemical characteristics of the CEM III cement used for filling voids and sealing anchors on the slab. The main constituents are: clincker (28.5%, of which $C_3A=2.85\%$), blast

furnace slag (71.0%), secondary constituents (siliceous and limestone fly ash, 0.5%).

	Average value (%)
SO ₃	2.8
Cl ⁻	0.14
Loss on ignition (950°C)	0.9
Insoluble	0.3
S ²⁻	0.44
Alkaline equivalents (Na ₂ O + 0.658 K ₂ O)	0.6
Alkaline actives	0.5

Results. The operation was carried out by the company NGE Fondations (Drap, France) 672 between November, 2019 and February, 2020. A total of 262 anchors (Fig. 16-17) were 673 installed perpendicular to the Slab with rods 0.016 m in diameter to a depth of 1 m. Drilling 674 confirmed the average thickness of 0.3 m (0.4–0.13 m) of the Slab with thinner zones located 675 especially at the foot (south part) and in the middle of the Slab. The drilling also confirmed 676 the presence of heterogeneity in the marls, in the altered zones with a clavey tendency, and 677 some of the voids observed during the radar studies. The latter were completely clogged by 678 679 the filling grout. The low corrosivity of the soil, associated with the thinness of the fossiliferous stratum and the need to use small diameter for the tie bars reinforcements, led to 680 the treatment of those rods by galvanization. In order to limit the impact of vibrations during 681 drilling, as well as the visual impact, the drilling diameters have been limited to 0.032 m in 682 diameter. The heads of the anchors were masked and surface protected by a tinted mortar up 683 684 to 0.05 m depth.

Each inclusion provides a minimum resistance capacity of Tr=8.4 kN, thereby improving very significantly the cohesion of the Slab on its support with a resistant effect against possible forces due to swelling (freeze-thaw) and hydrostatic pressures.

During and after the work, the topographic monitoring of the Slab met the expectations anddid not show any significant movement.

690

691 **4.4. Water ingress**

692

693 Whether by infiltration in marls or cracks, by chemical reactions caused by the action of 694 freeze-thaw cycle, or by hydrostatic pressure, water plays a major role in the weathering of 695 rocks and the behavior of the rocky massif. Water is one of the essential elements to be taken 696 into account in studies intended for the conservation of geosites. Consequently, and in addition to the stabilization work on the Ammonites Slab, it was necessary to carry out workaimed at improving the management of seepage water.

699

Water infiltration takes place mainly at the fractures of the fossiliferous stratum, in particular in its upper part (Fig. 13). The work carried out in 2009 to limit this infiltration consisted of the construction of a concrete guttering to drain the upstream water of the Slab, and by filling (resins) the largest open cracks. The guttering, made for a lifespan of around 5 years, was, 7 years later, in very poor condition, leaving some of the water to seep into the marls.

On the south side of the Slab, the shotcrete intended to block the erosion of the marlstones and the fall of overhanging limestone layer elements actually partially blocked the flow of infiltrated water from the surface. The barbicans that were supposed to drain this water never allowed it to be evacuated. In rainy weather, water flowed from the base of the concrete in contact with the lower limestone layer. This installation certainly had a negative impact on water circulation in the southern part of the rock mass, with the consequence of increasing the pressure at the foot of the Slab.

Due to the geomorphology of the site, there is a potential infiltration zone of approximately 100 m² in the rocky massif located immediately above the Slab. It was therefore necessary to remove the 2009 improvements, then to strip and dig a drain behind the Slab over a line of 50 m for a 2 m width. This newly cleared area was thus sealed by a geomembrane covered with C30/37 fiber-reinforced concrete. This new guttering was extended downstream by a rainwater drainage ditch towards the dale.

The edge of the southern part of the Slab was also treated: part of the shotcrete was removed. The remaining concrete was treated to better integrate visually into the geosite. At the same time, the permeable cracks on the surface of the Ammonites Slab were sealed during the anchors and sealing grout installation. 722

30

723 **4.5.** Unsolved problems and perspectives

724

While the Ammonites Slab rescue work ensures the stability of the geosite over the long term, untreated conservation issues remain: in particular, the surface desquamation of the Slab, the shattering of the cracked limestone, the falling rocks, which result from the daily and seasonal temperature contrast and the action of the freeze-thaw cycle in winter.

729 The only effective protection solution would be to shield the site from meteorological agents with strict control of humidity and temperature. Such control would only be possible by 730 complete insertion of the whole Slab inside a regulated building. Beyond its exorbitant cost, a 731 building of the necessary size is currently not feasible for several reasons: (1) the Local 732 Urbanism Plan in force in the municipality of Digne-les-Bains does not allow constructions 733 734 over 10 m in the Ammonites Slab area, while the Slab itself already has an amplitude of 24 m; (2) the proximity of the road would imply its deviation, which is complicated by the 735 736 occupation of land on the other side; (3) the geosite is located into several protection zones 737 forbidding such constructions (national nature reserve and protection zone of the listed site of the neighboring antique plaster industry of Champourcin). The possibility of setting up an 738 intermediate protection structure of courtyard type was also abandoned for the same reasons, 739 740 by very strong technical constraints, by the fact that it would only have reduced direct rain and snow runoff (out of wind) without strict control of environmental conditions, and in fine 741 by the very unfavorable benefit/cost balance. 742

A direct surface protection technique of the fossiliferous stratum was also considered by the installation of a protective synthetic layer, but this technique was very quickly abandoned. Indeed, current techniques allowing sufficient sealing also prevent residual moisture and soluble salts from escaping, which would actually have achieved the opposite effect. Furthermore, the long-term resistance in mountain climatic conditions of this type of coating still needs to be strongly tested before being used on heritage sites. While waiting for new techniques or new materials, the only alternative in addition to the measures already in place (this work) remains the maintenance of the Slab 'by hand' (human interventions using safety ropes) as and when required to limit the ravages of time and bad weather. Beyond the necessity for monitoring the Slab itself to survey the evolution of its condition, for the moment, we are arriving at the limits of the in situ conservation of palaeontological sites.

754

Drilling and field surveys have shown that the extension of the fossiliferous stratum is proven beyond the currently exposed surface of the Slab with the same density of ammonites. On the 7,000 m² where the stratum has been recognized and mapped (Fig. 18), an extrapolation allows to estimate the presence of nearly 34,000 buried ammonites. Physically and legally protected, this exceptional site is left as a legacy for future generations: as long as the latest long-term conservation issues are not resolved, no new large-scale discovery of the ammonites bearing surface is possible.

762

763 **5. Enhancement of the Ammonites Slab geosite**

764

765 5.1. The previous state

766

The Ammonites Slab, exposed to the open air, is located near the road (Fig. 3, 19a). The road
had been rerouted in the late 1970s to free up a safe, but strait, viewing area for the public.
Another derelict part of the road served as a car park linked to the Slab by an old road bridge.
These areas (the parking and the visiting area at the foot of the Slab) were never properly

developed and the surroundings of the site actually looked more like a wasteland than ageosite of international stature.

The mediation was limited to a single lectern with some information and an artistic
reconstruction of the seabed with some organisms from the time of the ammonites present on
the Slab.

776

777 **5.2.** The new visitor structures

778

As part of the Geological Adventure project, and in parallel with the work to ensure the
sustainability of the Ammonites Slab, the entire geosite was enhanced for the public between
November, 2019 and May, 2020. The work was carried out by the company Eiffage routes
Mediterranée-Alpes-Vaucluse.

The old vacant wasteland has been transformed into a parking area (Fig. 19b: 1). It was designed in the shape of a roundabout to allow full accessibility to buses. In order to comply with road safety rules in terms of visibility, it was necessary to shift the road slightly from on the equivalent of a half-roadway in front of the car park entrance (work made by the Departmental Technical House in September, 2019).

The old bridge (Fig. 19b: 2), the deck of which has been raised, is an infrastructure of quality 788 and as such it materializes the new entrance door to the geosite, which opens directly onto a 789 pedestrian reserved reception area (Fig. 19b: 3). Most of the mediation is now developed in 790 this space. From the reception area, a 35 m long (1.60 m wide - 1.45 for the walkway) 791 wooden/metal footbridge now provides access to a new observation platform, raised 2 m 792 above the ground level, facing the Slab and the ammonites (Fig. 19b: 4-5). The slope of the 793 794 footbridge at 4% meets accessibility standards for people with reduced mobility. The interest of the observation platform is threefold: (1) to offer visitors a new panoramic point of view on 795

the Slab partly decontextualized from the road and its nuisances (noise and safety). The foot of the Slab is no longer visible, and the point of view being closer to the wall due to its inclination lead to an increased immersion effect in the geosite (Fig. 20a); (2) the platform allows the neutralization of direct access to the fossil surface by the public, in order to avoid the anthropogenic degradation mentioned in chapter 4.1; (3) keeping the public at a safe distance also helps prevent the risks associated with rock falls, now minimized by the anchoring of the Slab.

803

804 **5.3. Didactic**

805

One of the aims of the geosite enhancement is to allow the public autonomy in their visit. On 806 the one hand in terms of physical accessibility (materialization of paths; controlled slopes and 807 808 drops; seating areas for prolonged station - Fig. 20b); sufficient space to accommodate groups; installation of a public transport stop awaiting to be integrated into the urban mobility 809 810 plan of the city of Digne-les-Bains); on the other hand, by setting up educational furniture 811 (which can also be used as a support for guided tours). The educational path was designed in 8 stages, from the car park to the discovery of the Ammonites Slab itself, following a 812 progressive and immersive approach (Cayla et al. 2010; Martin et al. 2010): 813

Welcome the visitor: the geosite is presented in the context of the 7 other geological or
natural sites to be visited in the Bès Valley, and of which the Ammonites Slab is in
some ways the gateway.

Arouse curiosity: visitors are invited to disconnect from their daily reality in order to
 immerse themselves in the deep time using the geological timeline developed for the
 Geological Adventure project (on the background on Fig. 20c).

- *Amaze:* encounter with the animals that lived on the site 197 million years ago in
 Sinemurian times. Explanations on ammonites with accessible speech are based on the
 latest scientific findings (Fig. 20d).
- *Explore:* what makes the Ammonites Slab a remarkable site? History of the discovery of the Slab with a perspective of its heritage and scientific interest.
- *Learn:* the explanation of the different stages of the ammonite fossilization process on
 the Slab is given using life-size models (Fig. 20e-f).
- *Experiment:* explanation of the formation of strata and then of the folds and erosion
 that shaped the Slab as we see it today.
- *Understand:* the methods in paleontology, investigation of the past, are mentioned.
- *Contemplate, and nothing else:* the last station on the observation platform invites to
 contemplation. A cast of a portion of the Slab allows the public to better take
 ownership of the site and also represents a tactile visit aid for the visually impaired
 (Fig. 20a).
- 834

The new geosite enhancement as a whole takes into account accessibility to people with disabilities according to their specific needs:

People with motor disabilities: the geosite is accessible to visitors in wheelchairs.
 Reserved parking spaces are positioned near the entrance. Seating is arranged
 regularly. The didactic materials are designed to be suitable for both standing and in
 wheelchairs people (or children).

• *Visitors with visual disabilities:* a contrasting tactile guide strip leads from the parking spaces to the entrance to the site. All graphic material is produced in large sans serif characters (>4.5 mm) with contrasting colors. All the themes are backed up by tactile 844 information allowing the understanding of the different concepts: the time scale, the845 animal ammonite, fossilization, geological strata, the molding of a portion of the Slab.

People with mental disabilities: all the facilities provide security by keeping it away
 from the road and the car park. The signage is explicit and all the didactic themes are
 doubled by an insert Easy to Read and Understand (FALC, by its acronym in French)
 identified by the European Easy to Read pictogram.

People with hearing disabilities: moving away from the road allows better hearing
 comfort. The spaces are large enough to allow the animation of groups with a sign
 language interpreter.

853

6. Funding

855

The Geological Adventure project was funded within the framework of the European program ALCOTRA, Interreg V-A 2014-2020 (Alpes Latines COopération TRAnsfrontalière in French). This program has covered the Alpine territory between France and Italy since 1990, where it has co-financed nearly 600 projects for around 550 million euros in European grants. Its general objective is to improve the life quality of populations, the sustainable development of territories and cross-border economic and social systems through cooperation affecting the economy, the environment and services to citizens.

The overall project budget supported by the 7 partners of the Geological Adventure amounts
to 2,331,546 €. Support from Europe, via the European Regional Development Fund (ERDF),
represents 85% of this amount (here the ALCOTRA program). This very high level of public
intervention is an unexpected lever in order to implement ambitious projects.

Regarding the financing of the operation to rescue and enhance the Ammonites Slab, the total
cost of this sole operation amounted to 1,659,673 €, distributed as follows (all taxes included):

- 134,771 € project management and technical and security control missions;
- 594,030 € rescue and reinforcement work;
- 683,247 € enhancing and landscaping work;

• 107,190 € didactic component;

• 140,435 € road shifting.

Fifty-nine percent were self-financed by the Departmental Council of the Alpes de HauteProvence (contracting authority - 979,673 €), 31% by the European Union (ALCOTRA
Interreg V-A 2014–2020 - 510,000€) and 10% by the Sud-Provence-Alpes-Côte d'Azur
Regional Council (170,000 €).

879

880 7. Conclusions

881

Protected since 1984 under the National Geological Nature Reserve of Haute-Provence 882 (RNNGHP), and labeled Geopark since 2000 (UNESCO Global Geopark of Haute-Provence), 883 the Ammonites Slab geosite of Digne-les-Bains (Alpes de Haute-Provence, southeastern 884 France) is the result of 197 million years in Earth history. Exposure to the open air of more 885 than 1,550 large ammonites on this ancient seabed makes it exceptional and spectacular 886 despite the virtual absence of infrastructure for visitors until recently. The Slab has only been 887 exposed to light step by step since the last third of the 20th century, but it has since suffered 888 from the effects of too rapid mechanical unloading of rocks, contrasting mountain climatic 889 conditions, and direct contact with the public. A series of studies (unpublished) have shown 890 that the state of conservation of the fossiliferous layer was much more degraded than 891 892 expected. Without the completion of major works in the relatively short term, the risk of

destabilization by a buckling mechanism would have resulted in the irreparable loss of part ofthe geosite.

895

As a committed manager, the Departmental Council of the Alpes de Haute-Provence has put 896 in place the necessary means to ensure the proper functioning of the RNNGHP, with the 897 desire to recharacterize the geosites to ensure both their physical conservation and public 898 access. Despite the significant financial effort made by the Departmental Council for the 899 900 rescue of the Ammonites Slab, the difficulty lay in mobilizing the significant additional funding required on the sole aspect of conservation. In this context, the 2015–2016 call for 901 projects of the European Interreg V-A France-Italy ALCOTRA program was a major 902 opportunity. It allowed developing the rescue, sustainability and enhancement project that 903 lives up to the international reputation of the Ammonites Slab, within a broader geotourism 904 905 project: the Geological Adventure. In order to respect the ambition and the axes of the European program, this project also concerned the development of other major geosites 906 907 through the territory in a process of experiential and sustainable tourism, their networking by 908 itinerant circuits and the setting up of a quality scientific mediation (including using of digital tools). Mediation was developed from a strategy of allowing making scientific culture 909 accessible to as many people as possible. The overall project budget supported by the 7 910 911 Franco-Italian partners of the Geological Adventure amounted to 2,331,546 €. Support from Europe, via the European Regional Development Fund (ERDF, here the ALCOTRA 912 program), represented 85% of this amount. Such a high level of public intervention is an 913 unexpected lever in order to implement ambitious projects and it is a serious source of 914 funding to be considered by project leaders for the protection and enhancement of European 915 916 geoheritage.

In this context, the conservation works of the Ammonites Slab of Digne-les-Bains (up to 918 1,659,673 €) consisted of (1) anchoring the Slab on its substratum; (2) filling the voids 919 identified during the studies using an inert grout so as to avoid undesirable chemical 920 reactions; (3) dealing with the problem of water infiltration; and (4) neutralizing direct public 921 access to avoid anthropogenic damage. Enhancement work for the entire geosite was carried 922 out in parallel. The new equipment consists of a parking area allowing the circulation of 923 buses; a slight realignment of the road was necessary in order to comply with road safety 924 rules. This new gateway to the geosite opens directly onto a pedestrian reception area, where 925 mediation is developed. A raised platform offers visitors a new panoramic and immersive 926 point of view in front of the ammonites, safe against any risk of rock fall or related to road 927 circulation. A particular effort has been made to provide the public with progressive and 928 immersive educational mediation using different media (lecterns, timeline, models, 3D 929

930 reconstructions, digital tool). The entire geosite, whether through its facilities or adapted 931 teaching methods, is accessible to people with motor, visual, mental or hearing disabilities. 932 The easy access to the Ammonites Slab and its free admission to visitors rank the geosite very 933 high on the accessibility scale of Mikhailenko et al. (2021) with a score of 17/20. This score 934 could be revised and reach 20/20 when the geosite will be integrated into the city's urban 935 traffic plan and served by public transport, a secure pedestrian path and a cycle lane.

936

The RNNGHP is a privileged place to study Geology and Palaeontology for students and scientists from all over the world, and this example highlights the contribution of protected natural areas to the improvement of knowledge and scientific research. Beyond the scientific interest, the project approach of the Geological Adventure is based on the fact that the enhancement of the geological heritage, its protection and conservation, are also factors of sustainable local economic development (Betard et al. 2017). It is also undeniable that the

Ammonites Slab of Digne-les-Bains was a very strong source of inspiration enabling the 943 emergence of concepts of geoheritage, geotourism and Geoparks (Martini 1994). To go further, 944 the Ammonites Slab of Digne-les-Bains was very recently proposed as a candidate to be one of 945 the first 100 IUGS (International Union of Geological Sciences) geological heritage sites. As 946 part of the management of the RNNGHP, the pursuit of actions carried out by the 947 Departmental Council of the Alpes de Haute-Provence for the conservation of geoheritage, the 948 equipment of geosites for the public and the establishment of mediation tools aim to make the 949 Alpes de Haute-Provence, and the UGHP, a leading geotourism destination. 950

951

952 Acknowledgments

953

First of all, the authors are pleased to thank very much the Editor and the both anonymous 954 955 reviewers for the time they spent on the manuscript, their careful read and their very constructive comments. The rescue work for stabilization of the Ammonites Slab and the 956 957 landscaping of the geosite were carried out respectively by the companies NGE Fondations 958 and Eiffage routes Mediterranée-Alpes-Vaucluse. The company Terzatec Ingéniérie et Expertise helped with the interpretation of the drilling data and the chemical laboratory 959 investigations. The company Biotope Communication participated to the didactic aspect. The 960 Office National des Forêts allowed the use of the land under their responsibility. Finally, the 961 road shifting was conducted by the Maison Technique of the Department of the Alpes de 962 Haute-Provence. 963

964

965 **Declarations**

967 Funding: This work was supported by the Departmental Council of the Alpes de Haute-968 Provence, France.

969 **Competing interests:** The authors declare no competing interests.

970 Availability of data and material: Not applicable.

971 **Code availability:** Not applicable.

972

973 **References**

40

974

975 Agenda 21 (1992) Sustainable Development plan for the 21th century, United Nations

976 Conference on Environment & Development, Rio de Janerio, Brazil, 3–14 June 1992.

977 Available online: https://sdgs.un.org/publications/agenda21. Accessed on September 2021

978

979 Arouca Declaration (2011) Available online: http://www.europeangeoparks.org/?p=223.
980 Accessed on September 2021

981

Bert D, Pagès JS (2021) Les Isnards ammonite slab geoheritage site (Digne-les-Bains,
Southeastern France). Geoconservation Research 4(2). https://doi.org/
10.30486/gcr.2021.1917851.1074

985

Bert D, Guiomar M, Bariani F, Faure E (2019a) Réserve Naturelle Nationale géologique de
Haute-Provence - Plan de gestion 2020-2030. Section A, pp 158, Section B, pp 70, Section C,
pp 106, Section D, pp 48. http://www.paca.developpement-durable.gouv.fr/plan-de-gestion2020-2030-de-la-rnn-geologique-de-a12730.html. Accessed on September 2021

991	Bert D, Guiomar M, Hippolyte JC, Bromblet Ph (2019b) Le projet de conservation et
992	d'aménagement de la Dalle aux ammonites de Digne-les-Bains. Congrès APF 2019 - Aix-en-
993	Provence, 16th-20th April 2019:14
994	
995	Bétard F, Hobléa F, Portal C (2017) Les géopatrimoines, de nouvelles ressources territoriales
996	au service du développement local. Annales de Géographie 17:523-543
997	
998	Cayla N (2009), Le patrimoine géologique de l'arc alpin : De la médiation scientifique à la
999	valorization géotouristique. Dissertation, Université de Savoie, pp 307
1000	
1001	Cayla N, Hobléa F (2011) La Via GeoAlpina. Itinérance géologique à travers les Alpes. In:
1002	Reynard E, Laigre L, Kramar N (ed) Les géosciences au service de la société. Actes du
1003	colloque en l'honneur du Professeur Michel Marthaler, 24-26 juin 2010, Lausanne
1004	(Géovisions n° 37). Institut de géographie, Université de Lausanne, pp 119-130
1005	
1006	Cayla N, Hobléa F, Gasquet D (2010) Guide des bonnes pratiques de médiation des
1007	géosciences sur le terrain. Géologie de la France 1:47-55
1008	
1009	Cohen KM., Finney SC, Gibbard PL, Fan JX (2013; updated 2021) The ICS International
1010	Chronostratigraphic Chart. Episodes 36:199-204
1011	
1012	Corna M (1985) Le Lias du Jura méridional, paléontologie biostratigraphique du Sinémurien,
1013	approche paléoécologique. Dissertation, Université Lyon 1, pp 258
1014	

1015	Corna M (1987) Les horizons sinémuriens du Calcaire à gryphées du Jura méridional français
1016	(Zone à Conbeari – Zone à Obtusum). Geobios, 20(4):531-536
1017	
1018	Corna M, Mouterde R (1988) Le Sinémurien de Semur (France): essai de biozonation pour le
1019	Sinémurien inférieur et le Lotharingien inférieur du stratotype et du Jura méridional. 2nd
1020	International Symposium of Jurassic Stratigraphy, Lisboa, pp 101-107
1021	
1022	Corna M, Dommergues JL, Guiffray A, Bulot L (1991) Quelques points remarquables dans le
1023	Sinémurien des Alpes de Haute-Provence (France); précisions biostratigraphiques et
1024	paléontologiques. Géologie Méditerranéenne 17(1): 3-37
1025	
1026	Delair S (2008) Etude de l'atténuation des efflorescences sur les matériaux cimentaires au
1027	moyen de produits pouzzolaniques. Dissertation, Ecole Nationale Supérieure des Mines de
1028	Saint-Etienne
1029	
1030	De Wever P, Egoroff G, Cornée A, Lalanne A. (2014) Géopatrimoine en France. Mémoires
1031	Hors-Série de la Société géologique de France 14: 1-180.
1032	
1033	De Wever P, Alterio I, Egoroff G, Cornée A, Bobrowsky P, Collin G, et al. (2015)
1034	Geoheritage, a national inventory in France. Geoheritage 7(3): 205-247. Doi:
1035	10.1007/s12371-015-0151-2.
1036	
1037	De Wever P, Cornée A, Egoroff G, Collin G, Duranthon F, Lalanne A, De Kermadec C, Lucet
1038	S (2019) Patrimoine géologique : notion, état des lieux, valorisation. Naturae 2019 (1):1-58
1039	

- Dommergues JL, Guiomar M (2011) La "Dalle à ammonites de Digne" (Réserve Naturelle
 Géologique de Haute-Provence, France). Etude d'un site fossilifère d'importance
 patrimoniale. Revue de Paléobiologie 30(1):261-293
- 1043

François H, Hirczak M, Senil N (2006) Territoire et patrimoine: la co-construction d'une
dynamique et de ses ressources. Revue d'Economie Régionale & Urbaine 2006(5):683-700

Gidon M, Pairis JL (1988) La structure des environs de Digne (Chaînes subalpines
méridionales, Alpes-de-Haute-Provence) : un exemple d'interférence entre l'avancée d'une
nappe de charriage épiglyptique et la sédimentation sur son front. C.R.Acad.Sc.Paris
307(II):1283-1288

1051

Goguel J (1939) Tectonique des chaînes subalpines entre la Bléone et la Durance. Bulletin des
Service de la Carte géologique de France et des topographies souterraines 202(41), pp 40

Gonzalez-Tejada C, Du Y, Read M, Girault Y (2017) From nature conservation to geotourism
development: Examining ambivalent attitudes towards UNESCO directives with the global
geopark network. International Journal of Geoheritage 5(2):1-20

1058

Gordon JE (2018) Geoheritage, Geotourim and the Cultural Landscape : Enhancing the visitor
expérience and promoting. Geoheritag, 8(136), pp 25. https://doi.org/
1061 10.3390/geosciences8040136

1062

Gray M (2013) Geodiversity: Valuing and Conserving Abiotic Nature. 2nd edn. John Wiley &
Sons Ltd, Chichester, pp 508

44

1	nr	5
т	υι	5

1066	Guiomar M (2009) Le patrimoine géologique en prise avec son territoire. La lettre de l'OCIM
1067	123 mai-juin 2009, pp 10. https://doi.org10.4000/ocim.238

- 1069 Guiomar M, Bert D, Bousquet V, Mathieu P, Brizio P (2018) The paleontological heritage at
- 1070 the heart of the European cross border Interreg V-A France-Italy Alcotra: the geological
- 1071@dventure. 5th International Paleontological Congress, Paris, 9th-13th July 2018, pp 692

1072

- Haug E (1891) Les Chaînes subalpines entre Gap et Digne. Contribution à l'étude géologique
- 1074 des Alpes françaises. Bulletin du Service de la Carte géologique de France 3(21):1-197
- 1075
- 1076 Haug E (1909) Traité de Géologie. A. Colin ed., Paris, pp 2024
- 1077
- Hebert E (1861) Du terrain jurassique de la Provence: sa division en étages, son indépendance
 des calcaires dolomitiques associés aux gypses. Bulletin de la Société Géologique de France
 19:100-121

1081

Hippolyte JC, Clauzon G, Suc JP (2011) Messinian-Zanclean canyons in the Digne nappe
(south- western Alps): tectonic implications. Bulletin de la Société Géologique de France
182(2):111-132. https://doi.org/10.2113/gssgfbull.182.2.111

1085

Hippolyte JC, Bellier O, Espurt N (2012) Quaternary deformation and stress perturbations
along the Digne thrust front, Southwestern Alpes. Comptes Rendus Geoscience 344:205-213

- 45
- 1089 International declaration on the rights of the memory of the earth (1991) Available online:
- 1090 https://geoheritage-iugs.mnhn.fr/media/1/20120905-
- 1091 declaration_of_the_rights__of_the_earth.pdf. Accessed on September 2021
- 1092
- 1093 Kilian W, Zürcher PH (1895) Réunion extraordinaire de la Société géologique de France dans
- 1094 les Basses-Alpes. Travaux du Laboratoire de Géologie de la Faculté des Sciences de Grenoble
 1095 3:49-106
- 1096
- Martin S, Regolini-Bissig G, Perret A, Kozlik L (2010) Elaboration et évaluation de produits
 géotouristiques. Propositions méthodologiques. Téoros. Revue de recherche en tourisme
- 1099 29(2):55-66. https://oi.org/10.7202/1024871ar
- 1100
- Martini G (1979) Aménagement d'une Réserve géologique dans les Alpes de HauteProvence. Rapport DEA Géologie des formations sédimentaires, Université de Provence,
 1978-1979, pp 51
- 1104
- Martini G (1994) Actes du premier symposium international sur la protection du patrimoine
 géologique, Digne-les-Bains, 11-16 juin 1991, Paris, Société Géologique de France
- 1107
- Martini G, Zouros N (2008) Geopaks, a Vision for the Future. Géosciences 7-8:182-188
- Mikhailenko AV, Ruban DA, Ermolaev VA (2021) Accessibility of geoheritage sites a
 methodological proposals. Heritage 4:1080-1091. https://doi.org/10.3390/heritage4030060
- 1112

- Page K (1992) The sequence of ammonite correlated in the British Sinemurian (Lower
 Jurassic). Newsletters on Stratigraphy 27:129-156
- 1115
- Page K (2003) The Lower Jurassic of Europe: its subdivision and correlation. Geological
 Survey of Denmark and Greenland Bulletin 1:23-59
- 1118
- 1119 Ruban DA (2015) Geotourism A geographical review of the literature. Tourism
 1120 Management Perspectives 15:1-15
- 1121
- Ruban DA, Ermolaev VA (2020) Unique geology and climbing: a literature review.
 Geosciences 10(259), pp 21. https://doi.org10.3390/geosciences10070259
- 1124
- Ruban DA, Mikhailenko AV, Yashalova NN (2021) The power of colour in geoheritage
 studies and marketing: some tentative reflections. Geologos 27(1):57-65. https://doi.org/
 10.2478/logos-2021-0005
- 1128
- Scotese CR (2014) Atlas of Jurassic Paleogeographic Maps, PALEOMAP Atlas for ArcGIS,
 volume 4, The Jurassic and Triassic, Maps 32-42, Mollweide Projection, PALEOMAP
 Project, Evanston, IL.
- 1132
- Smith WL (2006) Experimental Tourism around the World and at Home: Definitions and
 Standards. International journal of Services and Standards 2(1):1-14.
 https://doi.org/10.1504/IJSS.2006.008156
- 1136

- UNWTO, World Tourism Organization (1995) Charter for Sustainable Tourism. UNWTO
- 1138 Declarations 5(4), 16 pp. https://doi.org/10.18111/unwtodeclarations.1995.05.04
- 1139

- UNWTO, World Tourism Organization (2005) Making a tourism more sustainable a guide
 for policy makers. United Nations Environment Programme and World Tourism
 Organization, pp 222
- 1143
- 1144 Vélain CH, Dieulafait L (1872) Compte rendu de la course du 8 septembre à Champourcin.
 1145 Réunion de la Société Géologique de France à Digne (Basses Alpes) du 8 au 18 septembre
 1146 1872. Bulletin de la Société Géologique de France 29:606-617
- 1147
- Zhang Z, Provis JL, Reid A, Wang H (2014) Fly ash-based geopolymers: The relationship
 between composition, pore structure and efflorescence. Cement and Concrete Composite
 journal 64:30-41. http://dx.doi.org/10.1016/j.cemconres.2014.06.004
- 1151
- 1152 Zhang Z, Provis JL, Ma X, Reid A, Wang H (2018) Efflorescence and subflorescence induced
- 1153 microstructural and mechanical evolution in fly ash-based geopolymers. Cement and Concrete
- 1154 Composite journal 92:165-177. https://doi.org/10.1016/j.cemconcomp.2018.06.010
- 1155

1156 Figure captions

- 1157
- Fig. 1: Palaeogeographic map at the Sinemurian age (-200 MY from Scotese, 2014), with
 position of the geosite of the Ammonites Slab (red star).
- 1160
- 1161 Fig. 2: The numerous ammonites present on the Slab.

48

Fig. 3: Stat of the geosite in 2016, before the work. Note the unique lectern and the absence ofparticular enhancing.

1165

1166 Fig. 4: Location map of the Ammonites Slab geosite near the city of Digne-les-Bains (Alpes1167 de Haute-Provence department, SE France).

1168

Fig. 5: Geological and structural context of the Ammonites Slab. The geosite is located alongthe Bès and Saint Benoît faults, in the front of the Digne-les-Bains thrust sheet.

1171

1172 Fig. 6: Section of the Ammonites Slab series and a *Coroniceras* shell reconstruction
1173 (modified from Dommergues & Guiomar, 2011). The digital model of the ammonite is part of
1174 the new 3D photogrammetric acquisitions made by the CEREGE at a 0.5 mm resolution.

1175

Fig. 7: Map of the National Geological Nature Reserve of Haute-Provence (RNNGHP by its French acronym). The 18 green dots are the sites classified under National Nature Reserve (RNN) with strong regulation, and the green area is the protection perimeter were the removal, destruction or degradation of fossils, minerals and concretions is prohibited.

1180

Fig. 8: The damages on the Ammonites Slab; (**a**) surface desquamation and delamination due to climatic conditions; (**b**) open fractures repaired with mortar (visible just beneath the ammonite); (**c**) the void recently identified behind the Slab due to an illegally collected block in May 2016 (35 X 20 cm); (**d**) the slight swelling at the bottom of the Slab, where the most voids were observed. The bulge reaches 15 cm in amplitude (picture reconstructed using 3D photogrammetry); (**e**) unrestricted accessibility of the fossiliferous layer to the public can

result in small damages; (f) a worn ammonite due to repeated contact with the hands ofvisitors; (g) water infiltrations.

1189

Fig. 9: Extract from the raw RADAR signal and its interpretation: voids/heterogeneities are
detected between 10–40 cm below the surface of the Slab (red dot-lines).

1192

1193 Fig. 10: Mapping of the heterogeneities at (a) 10-40 cm, (b) 50-80 cm behind the surface of1194 the Slab.

1195

1196 Fig. 11: Ortho-rectified and calibrated photography projected in the plan of the Slab.

1197

Fig. 12: Digital model of the Ammonites Slab with colorization of the relief and artificialshading. The shades from yellow to red are in front of the average plane of the Slab.

1200

Fig. 13: Fracturing map of the Ammonites Slab showing the different generations of
fractures. The diagenetic joints are in green; normal faults and tension joints are in orange;
joints without calcite are in blue; open fractures (gravity and weathering fractures) are in red.
The yellow areas are concrete parts.

1205

Fig. 14: (a) Small normal fault with vertical throw, which shifts an ammonite; (b) calcite joints in the extension relay zone of the main strike-slip fault; (c) the main strike-slip fault cutting the Slab in half through the middle (dextral sliding movement; total sliding of 10.2 cm); (d) detail of an ammonite cut by the main strike-slip fault (sliding of 2.8 cm at that point); (e) closed joint filled with calcite.

Fig. 15: Principle of the anchoring with sealed metal inclusions. The voids to fill in with groutare represented in orange.

1214

1215 Fig. 16: Location of the 246 anchoring bars (yellow dots) avoiding the ammonites.

1216

1217 Fig. 17: The drilling and anchoring operation showing the anchor metal bars.

1218

Fig. 18: In red is the currently exposed part of the Ammonites Slab; in white is the potential
total unearthed mapped surface of the Slab (7,000 m²); an extrapolation allows to estimate the
presence of nearly 34,000 buried ammonites.

1222

1223 Fig. 19: Surroundings aspect of the Ammonite Slab geosite (a) before the 1224 enhancement/conservative works and (b) just after these works: (1) is the car-park area in the shape of roundabout to allow full accessibility to buses; the bridge that materializes the new 1225 1226 entrance door to the geosite; (3) the pedestrian reserved reception area where most of the 1227 mediation is developed; (4) the wooden footbridge access (the 4% slope meets accessibility standards for people with reduced mobility); and (5) the new observation platform, raised 2 m 1228 above ground level, facing the Slab and the ammonites. 1229

1230

Fig. 20: the new didactic equipments of the geosite: (**a**) the observation platform offering a new point of view to the Slab with the lectern-cast representing a tactile visit aid for the visually impaired; (**b**) detail of the wooden footbridge in front of the strata; (**c**) lecterns that explain stratigraphy concepts and the formation of the Slab; (**d**) ammonite reconstruction with soft-body parts; (**e**) overview on the pedestrian reserved reception area; (**f**) explanations of the fossilization process that occurred on the geosite with 3D life-size models.