

The Far-Ultraviolet Continuum Slope as a Lyman Continuum Escape Estimator at High-redshift

J. Chisholm, A. Saldana-Lopez, S. Flury, D. Schaerer, A. Jaskot, R. Amorín,

H. Atek, S. L. Finkelstein, B. Fleming, H. Ferguson, et al.

▶ To cite this version:

J. Chisholm, A. Saldana-Lopez, S. Flury, D. Schaerer, A. Jaskot, et al.. The Far-Ultraviolet Continuum Slope as a Lyman Continuum Escape Estimator at High-redshift. Monthly Notices of the Royal Astronomical Society, 2022, 10.1093/mnras/stac2874. insu-03839633

HAL Id: insu-03839633 https://insu.hal.science/insu-03839633

Submitted on 6 Jul2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The far-ultraviolet continuum slope as a Lyman Continuum escape estimator at high redshift

J. Chisholm[®],¹* A. Saldana-Lopez[®],² S. Flury,³ D. Schaerer,^{2,4} A. Jaskot,⁵ R. Amorín[®],^{6,7} H. Atek,⁸ S. L. Finkelstein,¹ B. Fleming,⁹ H. Ferguson,¹⁰ V. Fernández,⁶ M. Giavalisco,³ M. Hayes[®],¹¹ T. Heckman,¹² A. Henry,¹⁰ Z. Ji,³ R. Marques-Chaves,² V. Mauerhofer,¹³ S. McCandliss,¹² M. S. Oey,¹⁴ G. Östlin,¹¹ M. Rutkowski,¹⁵ C. Scarlata[®],¹⁶ T. Thuan,¹⁷ M. Trebitsch[®],¹³ B. Wang[®],¹⁸ G. Worseck^{®19} and X. Xu^{®12}

Affiliations are listed at the end of the paper

Accepted 2022 October 4. Received 2022 October 3; in original form 2022 July 20

ABSTRACT

Most of the hydrogen in the intergalactic medium (IGM) was rapidly ionized at high redshifts. While observations have established that reionization occurred, observational constraints on the high-redshift ionizing emissivity remain elusive. Here, we present a new analysis of the Low-redshift Lyman Continuum Survey (LzLCS) and literature observations, a combined sample of 89 star-forming galaxies at redshifts near 0.3 with *Hubble Space Telescope* observations of their ionizing continua (or Lyman Continuum, LyC). We find a strong (6σ significant) inverse correlation between the continuum slope at 1550 Å (defined as $F_{\lambda} \propto \lambda^{\beta_{obs}^{(550)}}$) and both the LyC escape fraction ($f_{esc, LyC}$) and $f_{esc, LyC}$ times the ionizing photon production efficiency (ξ_{ion}). On average, galaxies with redder continuum slopes have smaller $f_{esc, LyC}$ than galaxies with bluer slopes mainly due to higher dust attenuation. More than 5 per cent (20 per cent) of the LyC emission escapes galaxies with $\beta_{obs}^{1550} < -2.1$ (-2.6). We find strong correlations between β_{obs}^{1550} and the [O III]/[O II] flux ratio (at 7.5 σ significance), galaxy stellar mass (at 5.9 σ), the gas-phase metallicity (at 4.6 σ), and the observed far-ultraviolet absolute magnitude (at 3.4 σ). Using previous observations of β_{obs}^{1550} at high redshift, we estimate the evolution of $f_{esc, LyC}$ with both redshift and galaxy magnitude. The LzLCS observations suggest that fainter and lower mass galaxies dominate the ionizing photon budget at higher redshift, possibly due to their rapidly evolving metal and dust content. Finally, we use our correlation between β_{obs}^{1550} and $f_{esc, LyC} \times \xi_{ion}$ to predict the ionizing emissivity of galaxies during the epoch of reionization. Our estimated emissivities match IGM observations, and suggest that star-forming galaxies emit sufficient LyC photons into the IGM to exceed recombinations near redshifts of 7–8.

Key words: galaxies: high-redshift-galaxies: starburst-dark ages, reionization, first stars-ultraviolet: galaxies.

1 INTRODUCTION

Observations of high-redshift quasars indicate that the intergalactic medium (IGM) underwent a large-scale phase change near $z \sim 5-10$ (Becker et al. 2001, 2021; Fan et al. 2006; Bañados et al. 2018). At these redshifts, gas between galaxies rapidly transitions from being largely neutral to being largely ionized. This 'epoch of reionization' seeds the subsequent large-scale structure of the Universe (Gnedin 2000), impacts the interpretation of the cosmic microwave background radiation (Planck Collaboration XLVII 2016), and sets the IGM temperature (Miralda-Escudé & Rees 1994).

Determining the structure and evolution of reionization requires uncovering what emitted the first hydrogen ionizing, or Lyman Continuum (LyC), photons. Whether the sources of ionizing photons are broadly distributed spatially, or more concentrated impacts the morphology of reionization (Kulkarni et al. 2019a). The timing and duration of reionization strongly depend upon whether massive stars, active galactic nuclei (AGNs), or evolved stellar remnants are key contributors to reionization (Robertson et al. 2013, 2015; Madau & Dickinson 2014; Rosdahl et al. 2018).

The initial debate on the sources of reionization centered around whether AGNs or star-forming galaxies reionized the early IGM (Hopkins et al. 2008; Faucher-Giguère et al. 2009; Ouchi et al. 2009; Finkelstein et al. 2012; Robertson et al. 2013; Madau & Haardt 2015). However, current observations and simulations find too few AGNs during the epoch of reionization to dominate the ionizing photon budget (Fontanot et al. 2007; Ricci et al. 2017; Trebitsch et al. 2017, 2021; Matsuoka et al. 2018; Kulkarni, Worseck & Hennawi 2019b; Daval et al. 2020; Faucher-Giguère 2020; Shen et al. 2020; Yung et al. 2021; Jiang et al. 2022). The observational landscape of highredshift AGNs may dramatically change as the sensitivity of the James Webb Space Telescope (JWST) opens a new window on to the epoch of reionization, but the current observational picture suggests that star-forming galaxies reionized the IGM. Coupled with the recent success of LyC observations from local galaxies (see below), the current debate has shifted to whether bright massive galaxies that are clustered in over densities or widely distributed low-mass galaxies emitted the requisite ionizing photons (Finkelstein et al. 2019; Mason et al. 2019; Naidu et al. 2020, 2022; Matthee et al. 2021).

^{*} E-mail: chisholm@austin.utexas.edu

Determining the ionizing emissivity (\dot{n}_{ion}) of a given population (e.g. AGNs or star-forming galaxies) at high redshift requires estimating the number density of sources and the number of ionizing photons that each source emits into the IGM. This can be cast in terms of the UV luminosity density ($\rho_{\rm UV}$), the production efficiency of ionizing photons (ξ_{ion} , or the number of ionizing photons per FUV luminosity), and the fraction of LyC photons that escape each source ($f_{\rm esc, LyC}$). Numerically, \dot{n}_{ion} is the product of these three quantities as

$$\dot{n}_{\rm ion} = \rho_{\rm UV} \times \xi_{\rm ion} \times f_{\rm esc, LyC}.$$
(1)

All of these quantities likely vary with redshift and with galaxy property [e.g. stellar mass or far-ultraviolet (FUV) absolute magnitude], but each quantity can in principle be estimated from galaxies in the epoch of reionization.

Deep *Hubble Space Telescope* observations have estimated ρ_{UV} out to redshifts as high as 10 (Bouwens et al. 2015, 2021, 2022; Finkelstein et al. 2015; McLeod, McLure & Dunlop 2016; Livermore, Finkelstein & Lotz 2017; Mehta et al. 2017; Atek et al. 2018; Oesch et al. 2018; Yue et al. 2018). Similarly, deep rest-frame FUV spectroscopy (Stark et al. 2015; Mainali et al. 2018; Hutchison et al. 2019; Tang et al. 2019) and *Spitzer* rest-frame optical (Bouwens et al. 2016; De Barros et al. 2019; Endsley et al. 2021) observations of epoch of reionization-era systems find hard radiation fields indicative of high ξ_{ion} [~10^{25,3-26} (s⁻¹/erg s⁻¹ Hz⁻¹)]. Combined, these observations hint that star-forming galaxies can reionize the IGM if $f_{esc, LyC} > 5$ -20 per cent (Ouchi et al. 2009; Robertson et al. 2013, 2015; Finkelstein et al. 2019; Mason et al. 2019; Naidu et al. 2020, 2022; Matthee et al. 2021).

The neutral IGM precludes direct LyC detections at z > 4 in a statistical sense (Inoue et al. 2014; Becker et al. 2021), outside of rare ionized regions (Endsley et al. 2021; Endsley & Stark 2022). However, $f_{\rm esc, LyC}$ can be directly measured at z < 4 (Bergvall et al. 2006; Leitet et al. 2011; Borthakur et al. 2014; Izotov et al. 2016a, b, 2018a, b, 2021; Leitherer et al. 2016; Shapley et al. 2016; Vanzella et al. 2016; Steidel et al. 2018; Fletcher et al. 2019 ; Rivera-Thorsen et al. 2019; Wang et al. 2019; Ji et al. 2020; Davis et al. 2021; Marques-Chaves et al. 2021; Xu et al. 2022). To overcome the neutral high-redshift IGM, recent work has correlated indirect observables with $f_{esc, LVC}$ at these low redshifts. Tentative correlations, with appreciable scatter and significant non-detections (Rutkowski et al. 2017), have been found between the Ly α emission properties (Verhamme et al. 2015; Rivera-Thorsen et al. 2017; Izotov et al. 2018b, 2021; Steidel et al. 2018; Gazagnes et al. 2020), ISM absorption properties (Reddy et al. 2016; Chisholm et al. 2017, 2018; Gazagnes et al. 2018; Steidel et al. 2018), Mg II resonant emission lines (Henry et al. 2018; Chisholm et al. 2020; Witstok et al. 2021; Xu et al. 2022), and optical emission-line ratios (Jaskot & Oey 2014; Nakajima & Ouchi 2014; Wang et al. 2019). Recent simulations have tested these indirect indicators and have found modest efficacy with strong impacts from geometry and dust (Mauerhofer et al. 2021; Katz et al. 2022). These first glimpses of small samples of LyC emitters at lower redshifts provide a blueprint to estimate $f_{esc, LvC}$ during the epoch of reionization.

While there are promising LyC tracers, LyC-emitting galaxies have been elusive and the sample sizes of confirmed LyC detections was still modest. The small LyC samples may obscure trends or introduce false trends in LyC escape detections. The Low-Redshift Lyman Continuum Survey (LzLCS; Flury et al. 2022a) is a 134 orbit *Hubble Space Telescope* project that used the Cosmic Origins Spectrograph (COS) to target 66 new $z \sim 0.3$ LyC-emitting candidates. The LzLCS was developed to explore the galaxy parameter range relevant to cosmic reionization to assess the validity of a variety of indirect LyC tracers (Flury et al. 2022a). The LzLCS sample has a dynamic range in three purported LyC diagnostics: H β equivalent width, the [O III]/[O II] flux ratio, and the slope of the FUV stellar continuum (β). This range of parameters stringently tests indirect trends of LyC escape, establishing whether they scale with $f_{esc, LyC}$ at low redshifts (Flury et al. 2022b).

The combination of the LzLCS and additional galaxies with literature HST/COS observations of the LyC, a sample we call the LzLCS+ sample, is a resounding success. The LzLCS+ consists of 89 LyC-emitting candidates with HST observations. More than half, 55 per cent, of these galaxies have LyC emission detected above the background at the >97.7 per cent confidence level and the other 45 per cent have robust upper limits (Flury et al. 2022a). This provides a definitive local sample to explore indirect diagnostics in f_{esc} IxC. Flury et al. (2022b) found that many of the classical LyC diagnostics have appreciable scatter, but in general more highly ionized, vigorously star-forming, and compact galaxies are more likely to emit ionizing photons. The Ly α equivalent width, escape fraction, and velocity separation strongly correlate with $f_{esc, LvC}$ (Flury et al. 2022b). LyC emitters within the LzLCS+ are deficient in low-ionization optical metal emission lines, specifically [S II], indicating that LyC emitters are relatively deficient in low-ionization gas (Wang et al. 2021). The COS FUV spectral range also provides invaluable probes of the H I in the galaxies with the Lyman Series absorption lines. These HI lines suggest that LyC emission escapes through regions in the galaxy with low neutral gas covering fraction and low dust attenuation (Saldana-Lopez et al. 2022). As such, $f_{esc, LyC}$ can be robustly predicted using the combination of HI absorption and dust attenuation.

Here, we extend the previous LzLCS+ work to a detailed analysis of the FUV spectral slope, β_{obs}^{1550} , derived from stellar population fits and their relation to the $f_{\rm esc, LyC}$. $\beta_{\rm obs}^{1550}$ traces both the stellar population parameters and the dust attenuation in the LzLCS+ galaxies, and has been theoretically predicted to correlate with $f_{esc, LvC}$ (e.g. Zackrisson, Inoue & Jensen 2013). In Section 2, we describe the LzLCS+ sample and observations. Section 3 describes the modelling of the FUV stellar continua, and the derivation of stellar continuum properties. We explore correlations with β_{obs}^{1550} (Section 4) before introducing the strong relation between β_{obs}^{1550} and the $f_{esc, LyC}$ (Section 5). The physical picture presented by the β_{obs}^{1550} - $f_{esc, LyC}$ relation is explored in Section 6. We conclude in Section 7 using the β_{obs}^{1550} - $f_{esc, LyC}$ relation to make predictions for $f_{\rm esc, LyC}$ and $\dot{n}_{\rm ion}$ during the epoch of reionization to predict which galaxies heavily contribute to reionization. In Appendix A, we also detail the impact of different assumed extinction laws. Throughout this paper we use AB magnitudes; a standard cosmology with $H_0 = 70$ km s⁻¹, $\Omega_m = 0.3$, and $\Omega_{\Lambda} = 0.7$; and stellar metallicities that are relative to a solar value of 0.02.

2 OBSERVATIONS

Here, we describe the observations of the local LyC-emitting galaxies. We use three subsets of data: the LzLCS (Section 2.1), literature observations of 23 LyC emitters (Section 2.2; Izotov et al. 2016a,b, 2018a,b; Wang et al. 2019; Izotov et al. 2021), and finally a coaddition of all the LzLCS plus literature spectra. We combine the LzLCS and literature samples into one cohesive sample that we refer to as the LzLCS+ sample.

2.1 The LzLCS

We use data from the LzLCS (PI: Jaskot, *HST* Project ID: 15626; Flury et al. 2022a). The LzLCS is a large *Hubble Space Telescope*

program consisting of 134 orbits targeting 66 star-forming galaxies at redshifts in the range 0.219–0.431. We chose these redshifts to shift the LyC on to the sensitive spectral portion of the COS detector. These 66 galaxies were selected using optical observations from the 16th Data Release of the Sloan Digital Sky Survey (SDSS; Ahumada et al. 2020), and non-ionizing FUV observations from the *GALaxy Evolution eXplorer* (GALEX; Martin et al. 2005). The LzLCS was designed to test the efficacy of three potential indicators of LyC escape by creating three separate, but overlapping, samples with a large dynamic range of star formation rate surface density (as probed by the Σ_{SFR} measured from H β emission), nebular ionization state (as probed by the [O III]/[O II] optical emission-line ratio), and UV continuum slopes (as measured by the *GALEX* FUV-NUV slope). The resultant sample probes the full property range expected at high redshift to test the impact of various properties on $f_{esc, LyC}$.

The details of the data reduction are given in Flury et al. (2022a), but here we summarize the procedure. Each galaxy was observed for between one and five orbits with the low-resolution G140L grating of the COS (Green et al. 2012). We used the G140L/800 mode (Redwine et al. 2016), which has observed wavelength coverage in the range 800–1950 Å at a spectral resolution of $R \sim 1000$ at 1100 Å. This extremely blue-optimized COS configuration places the LyC on to the sensitive portion of the COS detector and affords rest-frame coverage of 600–1480 Å for the median redshift of the LzLCS. In addition to the crucial LyC, the LzLCS observations sample the very blue of the non-ionizing FUV continuum.

The COS observations were flat-fielded, wavelength-calibrated, and flux-calibrated using CALCOS v3.3.9. The individual exposures were co-added in the raw counts using FAINTCOS (Makan et al. 2021). FAINTCOS directly uses many time-baselines of archival COS observations to re-estimate the dark current, uses the scattered light model from Worseck et al. (2016) to improve the flux calibration at faint flux levels, and accounts for the Poisson uncertainties of the photon-counting COS FUV detector. Flury et al. (2022a) also details the accounting of the geocoronal emission to address possible LyC contamination.

The individual COS spectra were corrected for the foreground Milky Way reddening using the Green et al. (2018) dust maps and the Milky Way extinction curve (Fitzpatrick 1999). We then measured the observed counts in 20 Å regions in the rest-frame LyC. The inferred LyC counts and background levels, along with a Poisson distribution, are used in a survival function to determine the probability that the signal arises through chance realizations from the background distribution. If the survival analysis confirms that the flux is above the background counts at the >97.7 per cent confidence level, we classify these as LyC detections, and we refer to them as galaxies with 'Detected LyC' for the rest of the paper. We also checked the two-dimensional spectra by eye to ensure that these detections are not spikes in background counts. The galaxies with LyC flux <97.7 per cent above the background are called 'non-detected LyC' galaxies throughout this paper. These detection thresholds do not mean that the non-detected galaxies do not emit LyC photons, rather the LyC is not significantly detected given the observed backgrounds. Of the 66 galaxies in the LzLCS, 35 have LyC photons detected at >97.7 per cent confidence level.

All LzLCS galaxies have SDSS optical spectroscopy (Ahumada et al. 2020). These optical emission lines describe the nebular emission line properties and the gaseous conditions. In Section 4, we use these emission lines to explore the relation between nebular conditions and the FUV slope. The nebular reddening values, $E (B - V)_{\text{gas}}$, are determined in an iterative way where the temperature,

density, and extinction (assuming the extinction law from Cardelli, Clayton & Mathis 1989) are all iterated until the Balmer emission line flux ratios converge (Flury et al. 2022a). In this way, the temperature and density dependence of the Balmer line ratios are accounted for when determining the internal reddening of the galaxies. The nebular electron temperatures (T_e) are directly measured using the [O III] 4363 Å auroral line for 54 galaxies. The extinction-corrected emission line fluxes are used to determine line ratios of certain ions (e.g. [O III] 5007/[O II] 3726 + 3729 Å) which serve as a proxy of the gas ionization. Finally, the [S II] 6717, 6731 Å Å doublet is used to determine the electron density of the gas and the metallicities are measured from the inferred electron temperatures, densities, and attenuation-corrected fluxes. Metallicities of the 12 galaxies without auroral line detections were determined using the strong-line calibration from Pilyugin, Vílchez & Thuan (2006).

2.2 Literature data

In combination with the LzLCS, we also use a compilation of 23 galaxies drawn from the literature (Izotov et al. 2016a,b, 2018a,b, 2021; Wang et al. 2019). These galaxies are also observed with COS on *HST* and their reduction is handled in the same exact way as the rest of the LzLCS. This produces a homogeneous sample of 89 galaxies with LyC observations called the LzLCS+ sample. Of the literature sample, we detect the LyC above the background at a >97.7 per cent confidence level in 15 galaxies. This leads to a total of 50 detections in the LzLCS+ sample.

2.3 Spectral co-additions

In order to illustrate the changes of the stellar features at high significance, we also coadd the spectra to increase the resultant signal to noise. The details of this process are found in Flury et al. (in preparation). We coadded all of the LzLCS+ sample and made two separate composite spectra: (1) of all galaxies with detected LyC emission and (2) of all galaxies without LyC detections. We take care to mask intervening ISM, Milky Way absorption features, as well as the Ly α and OI 1302 and 1306 Å geocoronal emission lines, in each individual galaxy before co-adding the spectra. The individual spectra are then normalized to the median flux density at 1100 Å and the spectra are coadded in the rest frame. The uncertainties on the flux density are determined by bootstrapping the individual fluxes that comprise the stack. This high signal-to-noise stack is used in the next section to compare to stellar population synthesis models.

3 STELLAR CONTINUUM FITTING

3.1 Fitting procedure

In Saldana-Lopez et al. (2022), we fit the observed stellar continua of the LzLCS+, following Chisholm et al. (2019), as a linear combination of multiple single-age, single-metallicity stellar continuum models (see Section 3.2 for a discussion of the models used). Here, we use those continuum fits and briefly describe the fitting procedure, but refer the reader to Chisholm et al. (2019) and Saldana-Lopez et al. (2022) for in-depth discussions on the procedure and the associated assumptions.

We assume that the observed stellar continuum $(F(\lambda))$ is the product of the intrinsic stellar continuum $(F_i(\lambda))$ attenuated by a uniform dust screen. The treatment of the dust screen geometry impacts the interpretation of the stellar continuum fitting (Calzetti, Kinney & Storchi-Bergmann 1994; Vasei et al. 2016; Gazagnes et al.

2018), but it is challenging to distinguish the appropriate model with many degenerate effects (Calzetti 2013). Therefore, we assume a uniform dust screen. The intrinsic flux is modelled as a linear combination of a set of *j* stellar continuum models with different ages and metallicities $(M_j(\lambda))$ times a weight (X_j) that describes the contribution of each M_j to the total $F_i(\lambda)$. These X_j can have values greater than or equal to 0. Mathematically, this is defined as

$$F(\lambda) = 10^{-0.4 \text{E}(\text{B}-\text{V})k(\lambda)} \times F_i(\lambda)$$

= 10^{-0.4 \text{E}(\text{B}-\text{V})k(\lambda)} \times \Sigma_j \left(X_j \times M_j(\lambda)\right). (2)}

We use the reformulated Reddy et al. (2016) attenuation curves that is prescribed with a dust law, $k(\lambda)$, and the continuum colour excess (E (B - V)). The Reddy et al. (2016) law is similar in shape to the Calzetti et al. (2000) law, but with a lower normalization [the E (B - V) is inferred to be 0.008 mag redder than the Calzetti attenuation law; see fig. 5 of Reddy et al. 2016]. In Appendix A, we introduce analytic relations to convert between different attenuation laws. The Reddy et al. (2016) law is observationally defined in the observed wavelength regime down to 950 Å, while the Calzetti et al. (2000) is only defined to 1250 Å. We use the python package LMFIT (Newville et al. 2014) to determine the best-fitting values for each of the X_j linear coefficients and the E (B - V)values.

To fit the spectra, we place the galaxies into the rest frame using the SDSS redshifts (Ahumada et al. 2020) and normalize the flux density to the median rest-frame flux density in the range 1070-1100 Å. The rest-frame FUV contains many non-stellar features, such as foreground Milky Way absorption lines, geocoronal emission features, and ISM absorption lines from the targeted galaxies. These features do not arise from the stellar populations and should not be included in the fitting process. We mask these regions of the spectra by hand. We also mask out regions of low signal-to-noise ratios (less than one) to avoid fitting continuum noise. We tested whether including low signal-to-noise data improves the fitting and found a poorer match between models and observations when including lower quality data. We fit the rest-frame wavelengths in the range 950–1345 Å to avoid heavy contamination of the Lyman Series lines at bluer wavelengths. The median reduced χ^2 is 1.08 (Saldana-Lopez et al. 2022). Detailed comparisons of models and individual spectra are given in figs 1 and 2 of Saldana-Lopez et al. (2022).

Errors on the stellar continuum fits were calculated by modulating the observed flux density with the error spectra at each pixel, refitting the spectrum, and then tabulating the associated stellar continuum properties (Saldana-Lopez et al. 2022). We repeated this process 500 times for each galaxy to build a sample distribution of stellar population fits. We measured the standard deviations of these Monte Carlo distributions to determine the standard deviations on all fitted parameters (X_i and E(B - V)) and the other inferred parameters.

3.2 Stellar continuum models

To fit the observed LzLCS+ FUV spectra, we use fully theoretical single-age bursts of star formation to model the very young stellar continua. As in Saldana-Lopez et al. (2022), we include stellar models with a range of single ages of 1, 2, 3, 4, 5, 8, and 10 Myr with metallicities in the range of 0.05, 0.2, 0.4, and 1.0 Z_{\odot} . These ages were carefully chosen to sample the spectral variations of the FUV stellar continuum (Chisholm et al. 2019). Each stellar model (defined by an age and metallicity) includes a modelling of the nebular continuum by placing each single-age model into CLOUDY

v17.01 (Ferland et al. 2017), assuming that all of the ionizing photons are absorbed by the nebular gas, and tabulating the resultant nebular continuum. The nebular continuum is then added back to original stellar model (Chisholm et al. 2019). The nebular continuum does not heavily affect the wavelengths covered by the LzLCS+ observations, but contributes to the continuum slope at 1550 Å (Raiter, Schaerer & Fosbury 2010). We tested including older ages (up to 40 Myr) and found that they do not significantly change the inferred stellar population properties of the very young LzLCS+ spectra. However, the fully theoretical, high-resolution STARBURST99 models do not densely sample the H-R diagram at effective temperatures below 15000 K (Leitherer et al. 2010). The cooler, high-resolution models have artificially bluer continuum slopes that makes older stellar populations appear much bluer than expected for their temperatures. A similar resolution issue is found in the BPASS models (Chisholm et al. 2019). Thus, we choose the maximum model age range of 10 Myr that includes high-resolution models that accurately probe the stellar spectral features and the continuum shape. We fully describe the observed stellar continuum using 28 free parameters (each X_i) for the stellar models and one for the dust (for a total of 29 free parameters). We used the supersolar (2 Z_{\odot}) models for the three galaxies from Wang et al. (2019) which require higher metallicities.

We use the STARBURST99 models, convolved to the G140L spectral resolution, as our base stellar population models (Leitherer et al. 1999, 2010). We also fit the observed stellar continua with BPASS v2.2.1 stellar continuum models that include binary evolution (Eldridge et al. 2017; Stanway & Eldridge 2018) to test the impact of the assumed stellar continuum model on the stellar population results. To do this, we create a nebular continuum for the BPASS models using the same method as the STARBURST99 models.

The 1010-1060 Å wavelength range contains a host of stellar spectral features that are sensitive to the stellar population properties. In particular, the top panel of Fig. 1 compares the age-sensitive O VI P-Cygni stellar wind feature that is strong in both the LyC-detected (blue line) and non-detected (gold line) coadded spectra. Overlaid on these coadded spectra are single-age and single-metallicity STAR-BURST99 (red solid line) and BPASS (blue dotted lines) models. We use a 5 Myr and 0.2 Z_{\odot} model because this is the median fitted values of the sample. The STARBURST99 model roughly matches the observed broad O VI absorption, while the BPASS model does not match the observations. The bottom panel of Fig. 1 expands the age range of the models and shows that the absorption and emission component of O VI strongly depends on the stellar population age. By the time the STARBURST99 stellar population models reach 10 Myr there is very little O VI emission present in either model, indicating that the average observed LzLCS+ FUV stellar population is dominated by <10 Myr stellar populations. Other studies (e.g. Izotov et al. 2018b, 2021) have also found that LyC-emitting galaxies have strong O VI profiles which likely indicates the presence of a very young stellar population. This suggests that our choice of young stellar models is adequate to match the observed stellar continuum.

The STARBURST99 models, in general, more closely reproduce the observed O VI feature which is likely due to difference in the treatment of X-rays in the stellar winds of the two models. Since the O VI profile is the most obvious stellar feature in the very blue wavelength regime of the LzLCS+ spectra, we hereafter use the STARBURST99 models as our models. However, the inferred FUV stellar continuum slopes, the key observable used here, do not significantly change when using BPASS models.



Figure 1. Upper panel: the O VI 1032,1038 Å doublet stellar wind feature for the co-added LzLCS+ galaxies with LyC detections (blue line) and nondetections of the LyC (gold line). The O VI profile has two main components: broad absorption that extends to 1025 Å and broad emission that extends out to 1050 Å. Strong ISM absorption lines are marked by the grey vertical dashed lines. Overlaid are two fully theoretical stellar population models from STARBURST99 (red line) and BPASS (blue dotted line), with similar parameters as the median of our fits (age of 5 Myr and metallicity of $0.2 Z_{\odot}$). The BPASS model does not match either the observed P-Cygni absorption or emission, while the STARBURST99 model matches the absorption profile but not the full extent of the emission. Lower panel: stellar population synthesis models at a range of ages: 3 (blue), 4 (gold), and 5 Myr (red). We include both STARBURST99 (solid lines) and BPASS (dotted lines) models. The O VI profile weakens with increasing age for the STARBURST99 models, indicating that O VI is sensitive to the stellar population age. The O VI region of the BPASS models hardly varies with stellar population age. The strong O VI P-Cygni profile suggests that the LzLCS+ galaxies have very young stellar populations.

3.3 Inferred stellar population properties

Here, we describe the various properties derived from these stellar fits. These properties come in two different types: (1) directly fitted to the observations and (2) inferred from the stellar population fits. The directly fitted parameters are the 29 values in equation (2), including the X_j and the E (B - V) values. All of these parameters describe the stellar continuum in the observed FUV (near 1100 Å) and should be considered light-weighted properties at these wavelengths. All the parameters are found in tables A1–A4 of the appendix of Saldana-Lopez et al. (2022).

3.3.1 Properties derived from fitted parameters

From the individual fitted parameters (X_j and E(B - V)) we provide light-weighted estimates of the stellar population properties. We scale the individual model parameters by the fit weights to infer the light-

weighted age as

$$Age = \frac{\sum_{i} X_{j} Age_{j}}{\sum_{i} X_{j}}$$
(3)

and stellar metallicity (Z_*) as

$$Z_* = \frac{\sum_i X_j Z_j}{\sum_j X_j}.$$
(4)

The median Age and Z_* of the LzLCS+ sample is 4.6 Myr and 0.22 Z_{\odot} , respectively.

3.3.2 Properties derived from stellar continuum fit

We derive stellar continuum properties from the best-fitting stellar continuum. In particular, we derive the stellar continuum slope (β), the absolute magnitude ($M_{\rm UV}^{\rm obs}$), $f_{\rm esc, LyC}$, and the ionizing photon production efficiency ($\xi_{\rm ion}$).

The most important property that we measure in this paper is the slope of the FUV stellar continuum at a given wavelength, β^{λ} . We derive the slope between rest-frame 1300 and 1800 Å (an average wavelength of 1550 Å) assuming that

$$F_{\lambda} \propto \lambda^{\beta^{1550}}.$$
 (5)

We fit for β^{1550} by extrapolating the best-fitting stellar continuum model of the LzLCS+ COS observations described in Section 3 to redder wavelengths. We then fit for β_{obs}^{1550} assuming a power-law model in the 1300–1800 Å wavelength range of the stellar continuum fit. We tested two other methods to determine β^{1550} using (1) several small wavelength regions in the stellar continuum models without strong stellar or ISM features and (2) fitting for the slope between two points (1300 and 1800 Å). We found that all three models produced consistent β^{1550} estimates. Since we fit both the stellar population properties (giving the intrinsic continuum slope) and dust attenuation (giving the observed spectral slope), we estimate both the observed continuum slope, β_{obs}^{1550} , and the intrinsic slope, β_{int}^{1550} , which is the continuum slope without dust attenuation.

We use β_{obs}^{1550} from the stellar continuum modelling instead of the β observed in the *HST*/COS data (β^{COS} , or $\approx \beta^{1150}$; Flury et al. 2022a) because it probes the redder wavelengths (\sim 1550 Å) that are observed at higher redshift (Finkelstein et al. 2012; Dunlop et al. 2013; Bouwens et al. 2014; Wilkins et al. 2016; Bhatawdekar & Conselice 2021; Tacchella et al. 2022). The $\beta_{\rm obs}^{1550}$ values scale strongly (Kendall's rank coefficient of $\tau = 0.46$ and p-value of 1×10^{-10}) with β_{COS} from Flury et al. (2022a). However, β_{COS} values are 14 per cent redder than β_{obs}^{1550} measured here, largely because the reddening curve is steeper at bluer wavelengths (see the discussion in Section 4) and the intrinsic stellar continuum is slightly flatter between 900 and 1200 Å than at 1500 Å (Leitherer et al. 1999). Fig. 2 shows the same stellar continuum model attenuated by three different E(B - V) values. These three E(B - V) values produce three different β_{obs}^{1550} values that bracket the observed LzLCS+ range. Fig. 2 shows unique and distinguishable β^{1150} values for the three different E (B – V), but β^{1150} is much flatter, especially for high attenuation cases, than β ¹⁵⁵⁰. Thus, spectral slopes estimated at 1150 Å are not the same as at 1550 Å. To compare low-redshift observations to high-redshift observations, we must measure β at similar wavelengths.

How well do the extrapolated fits recover the actual β_{obs}^{1550} ? Recent *HST*/STIS observations by Schaerer et al. (2022) provide redder observations to test the extrapolated and actual β_{obs}^{1550} values. These redder observations are unavailable for the full LzLCS+ sample, but



Figure 2. Illustration of the impact of dust attenuation on the observed stellar continuum slope (β_{obs}^{1550}) keeping a constant underlying stellar population. All three spectra have the same underlying stellar population and are normalized near 1200 Å, but have a varying uniform dust attenuation using the Reddy et al. (2016) law (*E* (*B* - *V*) = 0.05, 0.15, 0.25 for the light blue, blue, and dark blue lines, respectively). This leads to three markedly different stellar continuum slopes (β_{obs}^{1550} =-2.4, -1.8, -1.2, respectively).



Figure 3. Typical LzLCS+ data (grey; from Izotov et al. 2018b) only probe the blue portion of the spectrum, but recent observations from Schaerer et al. (2022) of J1243+4646 (blue data) compare the extrapolated slope from the LzLCS+ data (gold line) to the observed continuum slope at 1550 Å. The extrapolated LzLCS+ continuum slope accurately matches the slope of the redder observations.

provide a test of our extrapolation for eight LyC emitters. Fig. 3 shows the extrapolated continuum slope from the fit to the LzLCS+ data in gold. The gold line was estimated by fitting the stellar population model to the G140L data between rest-frame wavelengths of 950–1200 Å (grey data). The redder STIS data, shown in dark blue, have a similar slope as the LzLCS+ fit even though these observations were not included in the fit ($\beta_{obs}^{1550} = -2.56$ from the models and -2.50 from the STIS observations). This demonstrates that the extrapolated β_{obs}^{1550} matches observed continuum slopes of LzLCS+ galaxies.

We estimated the observed, or uncorrected for internal extinction, FUV luminosity, L_{UV} , of the galaxies as the mean flux density of the best-fitting stellar model in a 100 Å region centred on 1500 Å. We

then converted this into an AB absolute magnitude $(M_{\rm UV}^{\rm obs})$ using the redshift from the SDSS spectra.

Flury et al. (2022a) took the ratio of the observed LyC emission to the fitted stellar continuum to estimate $f_{esc, LyC}$ in a few ways. We predominantly use the $f_{esc, LyC}$ derived from fitting the stellar continuum, but $f_{esc, LyC}$ can also be derived using the observed nebular Balmer emission lines to define the intrinsic LyC emission. The different $f_{esc, LyC}$ estimates scale with each other along a one-toone relation, although there does exist significant scatter within the relation (see fig. 19 in Flury et al. 2022a). Flury et al. (2022a) also derived the relative escape fraction, $f_{esc, LyC}^{ell}$, by comparing the dustattenuated stellar continuum models to the observed LyC flux. This quantifies the LyC absorption only due to H I within the galaxy.

Finally, we inferred the production efficiency of ionizing photons as

$$\xi_{\rm ion} = \frac{Q}{L_{\rm UV}},\tag{6}$$

where Q is the intrinsic number of ionizing photons produced by the stellar population and L_{UV} is the intrinsic mono-chromatic UV luminosity density of the stellar population at 1500 Å. We dereddened the best-fitting stellar continuum models, divided each ionizing flux density by the respective photon energy, and integrated over the entire ionizing continuum (21–912 Å) to determine Q (although the *number* of hydrogen-ionizing photons is dominated by photons with wavelengths near 912 Å). We then divided Q by the dereddened stellar population flux density at 1500 Å from the models. The LzLCS+ log(ξ_{ion} [s⁻¹/(erg s⁻¹ Hz⁻¹)]) ranges from 24.94 to 25.86, with a median value of 25.47. These ξ_{ion} values bracket the canonical high-redshift value of 25.3 (see Section 7), agree with those used in semi-analytical models (Yung et al. 2020), and are similar to the large range of high-redshift ξ_{ion} values (Bouwens et al. 2016; Harikane et al. 2018; Maseda et al. 2020; Stefanon et al. 2022).

4 THE FUV CONTINUUM SLOPE

FUV bright stellar populations are typically characterized by very blue colours because their spectral energy distributions peak in the extreme-ultraviolet (Leitherer et al. 1999; Raiter et al. 2010; Eldridge et al. 2017). The blackbody nature of massive stars leads to a very steep negative power-law index at 1500 Å with β_{int}^{1550} between -2.8and -2.5, after accounting for the nebular continuum, which depends on the age, metallicity, and star formation history of the stellar population (Leitherer et al. 1999). β_{int}^{1550} evolves very little over the first 10 Myr after a starburst. This is because 1550 Å is still on the Rayleigh–Jeans portion of the blackbody curve for late O-stars. The non-ionizing FUV light from massive stars propagates out of the galaxy, where it is absorbed and attenuated by the same gas and dust that absorbs LyC photons. This dust attenuation flattens the observed spectrum and makes β_{obs}^{1550} more positive.

The stellar population fits allow us to explore the range of β_{int}^{1550} within the LzLCS+. The β_{int}^{1550} has a fairly narrow range of values between -2.8 and -2.5, with a median of -2.71, which does not significantly change with stellar population age (left-hand panel of Fig. 4). These values are fully consistent with the narrow range of β_{int}^{1550} expected from young stellar populations that include the nebular continuum (Leitherer et al. 1999; Raiter et al. 2010). Fig. 4 shows that the β_{int}^{1550} spread does not statistically scale with age or E $(B - V)^{R16}$ (Kendall's τ p-values of 0.062 and 0.37, respectively), and there is also not a statistically consistent correlation with Z_* (p-value of 0.065). When we refer to E (B - V), we denote the attenuation law that we use as a subscript after E (B - V). This emphasizes that



Figure 4. The impact of fitted parameters on the intrinsic slope of the stellar continuum at 1550 Å (β_{int}^{1550}) for LyC-detected (blue circles) and non-detected LyC galaxies (gold squares). Left-hand panel: β_{int}^{1550} does not scale strongly with the inferred light-weighted ages. Some scatter is introduced due to various stellar evolution that happens between 3 and 5 Myr. Right-hand panel: The β_{int}^{1550} is uncorrelated with the continuum colour excess ($E (B - V)^{R16}$).

the E(B - V) value sensitively depends on the assumed attenuation law [i.e. $E(B - V)^{R16}$ is the E(B - V) derived using the Reddy et al. (2016) attenuation law]. In Appendix A, we describe how the following results can be recast to any attenuation law.

The non-dependence of β_{int}^{1550} on age may seem counter-intuitive. However, this is expected from the stellar plus nebular continuum models. As a stellar population ages the stellar continuum becomes redder, but the nebular continuum decreases to make β_{int}^{1550} bluer as there are fewer ionizing photons. This combined effect means that β_{int}^{1550} for very young stellar populations remains within a narrow range (between -2.8 and -2.5) for stellar populations with ages in the range 2–10 Myr. This relatively narrow scatter about a single value arises from β_{int}^{1550} depending on the population age, evolutionary sequence, star formation history, and stellar metallicity (in that rough order of importance). The fits to the LzLCS+ continua suggest that these very young star-forming galaxies occupy a narrow parameter range of age and star formation history.

The β_{obs}^{1550} does not scale significantly with the very young lightweighted stellar population age of the LzLCS+ (left-hand panel of Fig. 5), but scales strongly with the fitted $E (B - V)^{R16}$ using a Reddy et al. (2016) attenuation law (right-hand panel of Fig. 5). We use LINMIX, a hierarchical Bayesian linear regression routine that accounts for errors on both variables (Kelly 2007), to find the relation between $E (B - V)^{R16}$ and β_{obs}^{1550} to be

$$E(B-V)^{R16} = (0.47 \pm 0.01) + (0.171 \pm 0.007) \times \beta_{obs}^{1550}.$$
 (7)

This strong correlation and small scatter suggests that there is a simple analytical relation between E (B - V) and β_{obs}^{1550} . If we use the assumption that the observed continuum is a power law with wavelength (equation 5) and that the observed flux is equal to the intrinsic flux times the dust attenuation factor (equation 2), we find an analytic relation between E (B - V) and the continuum slope at the average of two wavelengths (λ^1 and λ^2 , respectively) as

$$E(B-V) = \frac{\log_{10} \frac{\lambda^{1}}{\lambda^{2}}}{0.4\Delta k} \left(\beta_{\text{int}}^{<1,2>} - \beta_{\text{obs}}^{<1,2>} \right), \tag{8}$$

where Δk is the difference in the reddening law at λ^1 and λ^2 . This equation emphasizes why it is crucial to compare β_{obs}^{λ} values at similar wavelengths: if E(B - V) is constant, the estimated β_{obs}^{λ} depends on the logarithm of the ratio of the two wavelengths used to measure the spectral slope. If we use $\lambda^1 = 1300$ Å and $\lambda^2 = 1800$ Å (the wavelength range used to determine β_{obs}^{1550}), the Δk from the

Reddy et al. (2016) law between 1300 and 1800 Å (2.08), and the median $\beta_{\text{int}}^{1550}$ of -2.7, we find that E(B - V) is related to the $\beta_{\text{obs}}^{1550}$ as

$$E(B-V)^{\text{R16}} = 0.46 + 0.17 \,\beta_{\text{obs}}^{1550}.$$
(9)

This agrees with the fitted relationship between $E (B - V)^{R16}$ and β_{obs}^{1550} (equation 7). This relation will change based upon the wavelength range that β_{obs}^{1550} is determined and the attenuation law that is used (see Appendix A).

Thus, the tight correlation between β_{obs}^{1550} and E(B - V) arises due to the shape of the attenuation law and β_{int}^{1550} . This also explains why there is relatively little scatter within the relationship between E(B - V) and β_{obs}^{1550} : the extremes of the fitted β_{int}^{1550} only change the inferred E(B - V) by 0.1 mag, equivalent to the scatter of E(B - V) at fixed β_{obs}^{1550} (Fig. 5). Next, we correlate β_{obs}^{1550} with parameters like $f_{esc, LyC}$, but equation (8) stresses that the inferred stellar attenuation, E(B - V), does depend on how well β_{int}^{1550} can be constrained. A 0.1 mag scatter is introduced to the E(B - V) observations due to β_{int}^{1550} . The relatively narrow range of β_{int}^{1550} suggests that β_{obs}^{1550} is largely set by the dust reddening with a moderate impact from the β_{int}^{1550} .

With this analytic understanding of β_{obs}^{1550} , it is useful to explore how β_{obs}^{1550} depends on other observables. The left-hand panel of Fig. 6 shows that the internal attenuation-corrected $O_{32} = [O \text{ III}]5007 \text{ Å}/[O \text{ II}] 3727 \text{ Å}$ flux ratio has an inverse, logarithmic, and highly significant (Kendall's τ rank coefficient of -0.55, p-value of 2×10^{-14} , or 7.5σ significant) trend with β_{obs}^{1550} . The grey line in Fig. 6 shows this best-fitting relation of

$$\beta_{\rm obs}^{1500} = (-1.3 \pm 0.1) \times \log_{10} O_{32} - (1.0 \pm 0.1) \,. \tag{10}$$

O₃₂ is often quoted as an ionization parameter indicator (Vilchez & Pagel 1988; Skillman 1989; McGaugh 1991; Izotov et al. 2016a), and does not have any direct observational connection to β_{obs}^{1550} (although there is a connection to the optical extinction through the Balmer dust correction). Similarly, there is a 6.7 σ (p-value of 8 × 10⁻¹²) significant correlation between β_{obs}^{1550} and the H β equivalent width, another measure of the nebular ionization state.

 $\beta_{\rm obs}^{1550}$ also scales with the gas-phase metallicity, 12 + log(O/H), with a 4.6 σ significance (Kendall's $\tau = 0.34$ and p-value of 2×10^{-6}). This trend suggests that $\beta_{\rm obs}^{1550}$ becomes redder for more metal-enriched systems, and that the bluest $\beta_{\rm obs}^{1550}$ values are found in low-metallicity systems. This relation is more scattered than the O₃₂ and EW(H β) relations, however, all eight of the galaxies



Figure 5. The correlation of the observed continuum slope at 1550Å (β_{obs}^{1550}) with fitted stellar population parameters for LyC-detected (blue circles) and non-detected LyC galaxies (gold squares). Left-hand panel: β_{obs}^{1550} does not strongly correlate with the light-weighted age of the stellar population. Right-hand panel: The observed FUV continuum slope strongly correlates with the stellar colour excess using Reddy et al. (2016) extinction law, $E (B - V)^{R16}$. These two plots strongly suggest that β_{obs}^{1550} is largely set by the dust attenuation.



Figure 6. The observed continuum slope at 1550Å (β_{obs}^{1550}) correlates strongly with galaxy properties derived from the optical spectra for both LyC-detected (blue circles) and non-detected galaxies (gold squares). Left-hand panel: β_{obs}^{1550} correlates with the flux ratio of the internal attenuation-corrected [O III] 5007 Å and [O II] 3727 Å flux ratio (O₃₂) at the 5.9 σ significance. O₃₂ is often used as a proxy of the ionization state of the nebular gas. Galaxies with bluer FUV slopes are more highly ionized. The grey line shows the best-fitting relation between β_{obs}^{1550} and O₃₂ (equation 10). Right-hand panel: The β_{obs}^{1550} scales with the oxygen abundance (12 + log(O/H)) at the 4.6 σ significance. The reddest galaxies ($\beta_{obs}^{1550} > -0.75$) do not have direct metallicities and their 12 + log(O/H) values are more uncertain. This suggests that more metal-rich galaxies have redder continuu and that dust attenuation increases with metallicity.

with $\beta_{\rm obs}^{1550}$ greater than -0.75 do not have observations of the temperature-sensitive [O III] 4363 Å line (Flury et al. 2022a). As such, the metallicities of the reddest galaxies may be uncertain by as much as 0.4 dex (Kewley et al. 2006), introducing significant scatter into this relation. Even with this significant calibration uncertainty, there is a nearly 5σ trend between $\beta_{\rm obs}^{1550}$ and the gas-phase metallicity.

Finally, we find a 5.9 σ and 3.4 σ significant relationship between β_{obs}^{1550} and the galaxy stellar mass (M_{*}) and observed (uncorrected for dust attenuation) FUV absolute magnitude (M_{UV}^{obs}), respectively. These relations suggest that lower mass and fainter galaxies have bluer stellar continuum slopes. We fully introduce and discuss the importance of these relations in Section 7.

In this section, we have explored relationships between both β_{int}^{1550} and β_{obs}^{1550} . We found that β_{int}^{1550} has a narrow range of values, but does not scale significantly with other properties. We found significant correlations between β_{obs}^{1550} and the fitted $E (B - V)^{R16}$, the observed optical [O II]/[O II] flux ratio, the H β equivalent width, M_{*}, M_{UV}^{obs} , and the gas-phase 12 + log(O/H). With these correlations in mind, the next section explores the correlation between β_{obs}^{1550} and the LyC escape fraction.

5 PREDICTING THE LYMAN CONTINUUM ESCAPE FRACTION WITH β

The previous section found that β_{obs}^{1550} scales strongly with the continuum colour excess, E(B - V). Similarly, Saldana-Lopez et al. (2022) found a strong correlation between E(B - V) and $f_{esc, LyC}$. Thus, Fig. 5 suggests that β_{obs}^{1550} likely correlates with $f_{esc, LyC}$ (Flury et al. 2022b). Fig. 7 shows the relationship between $f_{esc, LyC}$ and β_{obs}^{1550}



Figure 7. The LyC escape fraction $(f_{esc, LyC})$ strongly scales with the observed stellar continuum slope at 1550 Å (β_{obs}^{1550}) . The LyC detections are shown as circles and the non-detections as downward pointing arrows. Representative error bars for the LyC detections are shown in the bottom left corner. Both detections and non-detections are colour coded by their relative escape fractions $(f_{esc, LyC}^{rel})$, or the fraction of ionizing photons absorbed only by neutral hydrogen. At fixed β_{obs}^{1550} there is a strong vertical gradient in the colour of the plotted points, suggesting that neutral hydrogen introduces significant scatter at fixed dust properties. The grey dashed line shows the best-fitting relation for $f_{esc, LyC}$ in terms of β_{obs}^{1550} (equation 11) which accounts for both the $f_{esc, LyC}$ non-detections and the errors on both β_{obs}^{1550} and $f_{esc, LyC}$. The grey dotted line shows the theoretical relation assuming that only dust attenuation removes ionizing photons (equation 16). This acts as an upper envelope for the observations.

with the LyC-detected sample as circles and the non-detected LyC galaxies as downward pointing arrows at their corresponding 1σ $f_{esc, LyC}$ upper limits.

 $f_{\rm esc, LyC}$ upper limits. $\beta_{\rm obs}^{1550}$ scales strongly with $f_{\rm esc, LyC}$. The median $\beta_{\rm obs}^{1550}$ of the LyC-detected sample is -2.00 and -1.63 for the non-detected sample, indicating that LyC-emitting galaxies have substantially bluer continuum slopes. Using a Kendall's τ test for censored data (the *R* package CENKEN), we find that the $\beta_{\rm obs}^{1550}$ scales with $f_{\rm esc, LyC}$ at the 5.7 σ significance (Kendall's τ = -0.42 and a p-value of 5×10^{-9}). We use LINMIX (Kelly 2007), a hierarchical Bayesian linear regression routine which accounts for the errors on both $f_{\rm esc, LyC}$ and $\beta_{\rm obs}^{1550}$ as well as for the upper limits on $f_{\rm esc, LyC}$, to find the analytic relation between $f_{\rm esc, LyC}$ and $\beta_{\rm obs}^{1550}$ (the grey dashed line on Fig. 7) of

$$f_{\rm esc,LyC} = (1.3 \pm 0.6) \times 10^{-4} \times 10^{(-1.22 \pm 0.1)\beta_{\rm obs}^{1550}}.$$
 (11)

Many theoretical models for reionization suggest that star-forming galaxies must emit more than 5, 10, or 20 per cent of their ionizing photons to reionize the IGM (Ouchi et al. 2009; Robertson et al. 2013, 2015; Finkelstein et al. 2015, 2019; Rosdahl et al. 2018; Naidu et al. 2020). Equation (11) suggests that these $f_{\rm esc, LyC}$ occur on average in the LzLCS+ for $\beta_{\rm obs}^{1550}$ values less than -2.11, -2.35, and -2.60, respectively. We also find a 2.9 σ and 3.4 σ significant correlation (related to a 0.0003, respectively) between $\beta_{\rm obs}^{1550}$ and $f_{\rm esc, LyC}^{\rm rel}$ and the escape fraction calculated using the H β equivalent width (Flury et al. 2022a). Thus, $\beta_{\rm obs}^{1550}$ correlates with complementary $f_{\rm esc, LyC}$ measures.



Figure 8. The LyC escape fraction $(f_{esc, LyC})$ times the ionizing photon production efficiency $(\xi_{ion} = Q/L_{UV})$ strongly correlates with the observed continuum slope at 1550 Å (β_{obs}^{1550}) . The grey dashed line shows the best-fitting relation for $f_{esc, LyC} \times \xi_{ion}$ in terms of β_{obs}^{1550} (equation 12) that accounts for both the $f_{esc, LyC}$ upper limits and errors.

Equation (11) describes the population average relation between $\beta_{\rm obs}^{1550}$ and $f_{\rm esc, LvC}$. This relation contains real and significant scatter that must be addressed when applying it to higher redshift observations. With the appreciable scatter in this relation, equation (11) more robustly estimates population-averaged escape fractions rather than individual galaxy-by-galaxy escape fractions. How much of the scatter is physical? The LzLCS+ $f_{\rm esc, LyC}$ detections have a median signal-to-noise of 3 (Flury et al. 2022a). Thus, there are significant observational uncertainties in the $f_{\rm esc, LyC}$ measurements (see the representative error bars in the lower left of Fig. 7). LINMIX estimates that about 0.15 dex of the scatter in Fig. 7 is intrinsic to the relationship. A possible source of intrinsic scatter is the neutral gas column density, the neutral gas covering fraction, or the geometry. These have been observed to scale strongly with $f_{esc, LVC}$ in the LzLCS+ (Saldana-Lopez et al. 2022). The points in Fig. 7 are colour-coded by $f_{\rm esc,LvC}^{\rm rel}$ or the escape fraction in the absence of dust. While $f_{\text{esc,LyC}}^{\text{rel}}$ only moderately scales with $\beta_{\text{obs}}^{1550}$ (3.4 σ), at fixed $\beta_{\text{obs}}^{1550}$ there is a strong vertical $f_{\rm esc, LyC}^{\rm rel}$ gradient, which may suggest that galaxy-to-galaxy variations in neutral gas sets the scatter in Fig. 7. This strong secondary scaling with $f_{\rm esc,LyC}^{\rm rel}$ echoes the findings in Section 4 where β_{obs}^{1550} strongly scales with O₃₂ and H β EW, both of which trace the ionization state. In a future paper (Jaskot et al. in preparation), we will explore multivariate correlations with $f_{\rm esc, LyC}$ and β_{obs}^{1550} , O₃₂, H β equivalent width, metallicity, or star formation rate surface density.

Finally, Fig. 8 shows the scaling relation between $\beta_{\rm obs}^{1550}$ and the emitted ionizing efficiency ($\xi_{\rm ion} \times f_{\rm esc}$). Using LINMIX we determine this relation to be

$$\xi_{\rm ion} \times f_{\rm esc} = (4 \pm 2) \times 10^{21} [\rm s^{-1} / erg \ s^{-1} \ Hz^{-1}] \times 10^{(-1.2 \pm 0.1)\beta_{\rm obs}^{1550}}.$$
 (12)

This has a similar $\beta_{\rm obs}^{1550}$ dependence as the $f_{\rm esc, LyC}$ relation given in equation (11) with a power-law exponent of -1.2. The scatter on the emitted ionizing efficiency relation is similar to the scatter in equation (11), consistent with the insignificant relationship found

between $\beta_{\rm obs}^{1550}$ and $\xi_{\rm ion}$ (p-value of 0.155). Thus, the ionizing emissivity scales the relation between $\beta_{\rm obs}^{1550}$ and $f_{\rm esc, LyC}$ to include the production efficiency of ionizing photons. In Section 7, we use equation (12) to estimate the ionizing emissivity of galaxies during the epoch of reionization.

6 IMPLICATIONS FOR THE ESCAPE OF IONIZING PHOTONS

The FUV spectral slope strongly correlates with $f_{esc, LvC}$. Figs 4 and 5 demonstrate that the major determinant of the FUV spectral slope is the dust attenuation derived from the observed continuum shape. Dust connects the gas-phase metallicity, through the dust-to-gas ratio, and to the ionization state of the gas because metals are the predominate nebular coolant. The metallicity of the gas strongly depends on the stellar mass of the galaxies through the mass-metallicity relation (Tremonti et al. 2004; Berg et al. 2012) and the mass-luminosity relation. Therefore, β_{obs}^{1550} , M_* , O_{32} , H β equivalent width, M_{UV}^{obs} , and E(B-V) are all linked together through the gas-phase metallicity. Gas-phase metallicity physically explains why Section 4 found all of these parameters to correlate. This connection between LyC and dust echoes relations between Ly α and dust (Hayes et al. 2011). Here, we explore the physical connection between $\beta_{\rm obs}^{1550}$ and $f_{\rm esc, LyC}$ and how the observed β_{obs}^{1550} - $f_{esc, LyC}$ relation informs on how ionizing photons escape galaxies.

To explore the connection between $\beta_{\rm obs}^{1550}$ and $f_{\rm esc, LyC}$, we revisit the absolute escape fraction definition (see Saldana-Lopez et al. 2022). There are two main sinks of ionizing photons: dust (with an optical depth of $\tau_{\rm dust}$) and H I (Chisholm et al. 2018). Thus, $f_{\rm esc, LyC}$ is determined as the product of the attenuation of dust and $f_{\rm esc, LyC}^{\rm rel}$ as

$$f_{\rm esc,LyC} = 10^{-0.4\tau_{\rm dust}} \times f_{\rm esc,LyC}^{\rm rel},\tag{13}$$

where

$$\tau_{\rm dust}(912) = k(912) \,\mathrm{E(B-V)}.\tag{14}$$

A possible explanation for the significant correlation between β_{obs}^{1550} and $f_{esc, LyC}$ is that dust is the only sink of ionizing photons. This can be tested assuming that the H I transmits all of the ionizing photons ($f_{esc, LyC}^{rel} = 1$). Equation (13) then only depends on the dust as

$$\log(f_{\rm esc,LvC}^{\rm dust}) \approx -0.4\tau_{\rm dust}(912) = -0.4E(B-V)k(912).$$
(15)

Using equation (8), equation (9), and the fact that k(912) = 12.87 for the Reddy et al. (2016) extinction law, this equation can be redefined in terms of β_{obs}^{1550} as

$$\log(f_{\rm esc, LyC}^{\rm dust}) \approx -k(912) \frac{\log_{10} \frac{\lambda^{1}}{\lambda^{2}}}{\Delta k} (\beta_{\rm int} - \beta_{\rm obs})$$

= -2.4 - 0.9 $\beta_{\rm obs}^{1550}$. (16)

We overplot this relation as the dotted line in Fig. 7 and see that this relation provides a strong upper envelope for $f_{\rm esc, LyC}$. There are no observed points above this 'dust only' line. All the points lie below the 'dust only' line because neutral gas also decreases $f_{\rm esc, LyC}$. As emphasized in equation (13), $f_{\rm esc, LyC}$ is the product of the dust and $f_{\rm esc, LyC}^{\rm rel}$. This naturally explains why there is a strong gradient of $f_{\rm esc, LyC}^{\rm rel}$ in Fig. 7 at fixed $\beta_{\rm obs}^{\rm 1550}$: the absolute escape fraction is a product of both dust and H I opacity. Saldana-Lopez et al. (2022) used the Lyman Series absorption lines to find that the points close to the 'dust only' line have very low neutral gas covering fractions and H I absorbs little LyC, explaining why $f_{\rm esc, LyC}^{\rm rel} \sim 100$ per cent for these points.



Figure 9. The observed absolute FUV magnitude $(M_{\rm UV}^{\rm obs})$ of the LyC detections (coloured circles) and non-detections (grey squares) for the LzLCS+ sample scales with the observed stellar continuum slope at 1550 Å ($\beta_{\rm obs}^{1550}$) at the 3.4 σ significance. Brighter LzLCS+ galaxies are redder and possibly emit a smaller fraction of their ionizing photons. The $\beta_{\rm obs}^{1550}$ - $M_{\rm Obs}^{\rm obs}$ observed at $z \sim 7$ from Bouwens et al. (2014) is included as a dashed line. This $z \sim 7$ relation matches the general trend of the LyC detections in the LzLCS+. The black triangles show individual z > 8 galaxies from Wilkins et al. (2016), Bhatawdekar & Conselice (2021), and Tacchella et al. (2022). The LyC detections are coloured by the observed LyC escape fraction ($\log(f_{\rm esc, LyC})$) to emphasize galaxies with $f_{\rm esc, LyC} > 5$ per cent as blue points. We use equation (11) and $\beta_{\rm obs}^{1550}$ to estimate population averages of $f_{\rm esc, LyC}$.

7 IMPLICATIONS FOR COSMIC REIONIZATION

The goal of the LzLCS is to provide local examples of LyC escape that are directly transferable to high-redshift observations to determine their contribution to reionization. To reionize the early IGM, starforming galaxies must have an ionizing emissivity, \dot{n}_{ion} , much larger than the cosmic baryon density at high-redshift. Equation (1) infers \dot{n}_{ion} if the integrated FUV luminosity function (ρ_{UV}), the ionizing efficiency (ξ_{ion}), and $f_{esc, LyC}$ are observed (Ouchi et al. 2009; Robertson et al. 2013, 2015; Finkelstein et al. 2015, 2019; Mason et al. 2019; Naidu et al. 2020). Previously, we have provided an empirical route to estimate $f_{esc, LyC}$ and $\xi_{ion} \times f_{esc, LyC}$ using the FUV continuum slope, a common observable of high-redshift galaxies (Finkelstein et al. 2012; Dunlop et al. 2013; Bouwens et al. 2014; Wilkins et al. 2016; Bhatawdekar & Conselice 2021; Roberts-Borsani et al. 2022; Tacchella et al. 2022). If we assume that these LzLCS+ relations apply to high-redshift galaxies, do early star-forming galaxies emit sufficient ionizing photons to reionize the early IGM?

Fig. 9 compares β_{obs}^{1550} to the LzLCS+ observed (not corrected for dust) M_{UV}^{obs} . We colour-code the LzLCS+ points as grey squares for non-detections and detected galaxies with the observed $f_{esc, LyC}$. The colourbar is shifted such that blue points represent the so-called 'cosmologically relevant' escape fractions (>5 per cent). Using the relationship between β_{obs}^{1550} and $f_{esc, LyC}$ (equation 11) we shade the regions of this figure that correspond to LzLCS+ population averages of 5, 10, and 20 per cent escape fractions. This shading emphasizes that the LzLCS+ galaxies with $f_{esc, LyC}$ greater than 5 per cent are predominately found at fainter M_{UV}^{obs} (Flury et al. 2022b).



Figure 10. The observed absolute FUV magnitude $(M_{\rm UV}^{\rm obs})$ versus the observed LyC escape fraction $(f_{\rm esc, LyC})$ for galaxies with detected (blue circles) and non-detected (downward pointing arrows) LyC emission. The inferred relationship at $z \sim 7$ is shown as a light blue line using equation (18) and the observed relation between $\beta_{\rm obs}^{1550}$ and $M_{\rm UV}^{\rm obs}$ from Bouwens et al. (2014). The best-fitting relation from the LzLCS+, accounting for the upper limits on $f_{\rm esc, LyC}$, is shown as the dashed grey line. The LzLCS+ and $z \sim 7$ relations have statistically similar slopes (0.34 \pm 0.10 and 0.25 \pm 0.09 for the LzLCS+ and $z \sim 7$ relation, respectively) and normalizations (5.27 \pm 2.1 and 5.14 \pm 0.3, respectively).

Many previous studies have explored the median relationship between β_{obs}^{1550} and M_{UV}^{obs} at high redshift (Finkelstein et al. 2012; Dunlop et al. 2013; Bouwens et al. 2014). Bouwens et al. (2014) fit the $\beta_{obs}^{1550}-M_{UV}^{obs}$ relation at different redshifts as

$$\beta_{\rm obs}^{1550} = b + a(M_{\rm UV}^{\rm obs} + 19.5). \tag{17}$$

Table 3 in Bouwens et al. (2014) gives the *a* and *b* values for six different redshifts from 2.8 to 8. We note that the $z = 8 \beta_{obs}^{1550} - M_{UV}^{UV}$ relation from Bouwens et al. (2014) has a fixed slope (*a* value) due to the uncertain *Y*-band data. The dashed line in Fig. 9 shows the z = 7 relation over the LzLCS+ observations. The high-redshift relation matches the LzLCS+ across most M_{UV}^{obs} . This suggests that the LzLCS+ and $z \sim 7$ galaxies have a similar relation between their FUV brightnesses and continuum slopes. Combining the observed $z \sim 7 \beta_{obs}^{1550} - M_{UV}^{obs}$ relation (equation 17) and the LzLCS+ $\beta_{obs}^{1550} - f_{esc, LyC}$ at various redshifts as

$$f_{\rm esc,LyC} = 1.3 \times 10^{-4} \times 10^{-1.22(b+a[M_{\rm UV}^{\rm obs}+19.5])}.$$
 (18)

In Fig. 10, we overplot this predicted $z \sim 7$ relationship using the *a* and *b* values from Bouwens et al. (2014) as a light blue line. The dashed grey line illustrates the best-fitting LzLCS+ relation using LINMIX (Kelly 2007) to account for the $f_{\rm esc, LyC}$ upper limits and the errors on both $f_{\rm esc, LyC}$ and $M_{\rm UV}^{\rm obs}$. The LzLCS+ and $z \sim 7$ relations have relatively large error bars, but their slopes and normalizations are statistically similar (a slope in log-space of 0.34 ± 0.10 and 0.25 ± 0.09 for the LzLCS+ and $z \sim 7$ relations, respectively). Thus, the $z \sim 7$ relation roughly reproduces the observed LzLCS+ trend. The observed log-space trend is very shallow (Flury et al. 2022b) and suggests only a factor of $10 f_{\rm esc, LyC}$ change in over four orders of magnitude of $M_{\rm UV}^{\rm obs}$. The large LzLCS+ $M_{\rm UV}^{\rm obs}$ dynamic range likely explains why smaller previous samples did not find a statistically significant trend between $M_{\rm UV}^{\rm obs}$ and $f_{\rm esc, LyC}$ (Izotov et al. 2021).



Figure 11. LzLCS+ galaxies that are blue and low stellar mass (M_{*}) tend to have larger LyC escape fractions. The $\beta_{\rm obs}^{150}$ of the LzLCS+ detections (coloured circles) and non-detections (grey squares) scale with the M_{*} at the 5.9 σ significance. We include the $z \sim 7$ relation from Finkelstein et al. (2012) as the dashed line and individual z > 8 galaxies from Wilkins et al. (2016), Bhatawdekar & Conselice (2021), and Tacchella et al. (2022) as black triangles. The LzLCS+ LyC detections are coloured by the observed LyC escape fraction (log($f_{\rm esc, LyC}$)), such that galaxies with $f_{\rm esc, LyC} > 5$ per cent are blue. We use equation (11) to shade regions of the plot that correspond to 5, 10, and 20 per cent $f_{\rm esc, LyC}$. The overall trend suggested by the LzLCS+ for the high-redshift galaxies is that lower M_{*} galaxies have higher $f_{\rm esc, LyC}$.

Using the Bouwens et al. (2014) observations, equation (18) predicts that $z \sim 7$ galaxies emit > 5per cent (> 20per cent) of their ionizing photons if they are fainter than $M_{\rm UV}^{\rm obs} > -19.2$ mag (-16.7 mag; see the shading in Fig. 9). Other works have made similar β_{obs}^{1550} observations at high redshift. Dunlop et al. (2013) find $\beta_{\rm obs}^{1550} = -2.08$ at $z \sim 7$ for $M_{\rm UV}^{\rm obs} = -18.5$ mag galaxies and -1.81for $M_{\rm UV}^{\rm obs} = -19.5$ mag galaxies. This leads to a population average $f_{\rm esc, LyC} \simeq 5$ per cent for faint galaxies, while brighter galaxies have an $f_{\rm esc, LyC}$ of 2 per cent. The triangles in Fig. 9 show individual z > 8 galaxies (Wilkins et al. 2016; Bhatawdekar & Conselice 2021; Tacchella et al. 2022). While sparsely sampled at the moment, z> 8 galaxies fainter than -20.5 mag have a median $\beta_{obs}^{1550} = -2.1$ (suggesting an $f_{\rm esc, LyC}$ near 5 per cent) and galaxies brighter than -20.5 have $\beta_{obs}^{1550} = -1.9$. Since β_{obs}^{1550} largely tracks the dust contribution to $f_{esc, LyC}$ (Section 6), we can use equation (16) to put an upper limit of $f_{\rm esc, LyC} < 30$ per cent for the individual $M_{\rm UV}^{\rm obs} > -20.5$ $z \sim 8$ galaxies. This is the 'dust free' scenario and modest $f_{\rm esc,LvC}^{\rm rel}$ can reduce $f_{esc, LyC}$ to population averages of 5 per cent. Finally, we note that the LzLCS+ β_{obs}^{1550} - M_{UV}^{obs} relations are entirely consistent with semi-analytical models of β_{obs}^{1550} at high redshift, including the large scatter to higher β_{obs}^{1550} values (Yung et al. 2019).

The stellar mass, M_* , represents the integrated star formation of a given galaxy. Fig. 11 shows the 5.9σ significant relationship between M_* and β_{obs}^{1550} . More massive galaxies have redder continuum slopes than their lower mass counterparts, likely as their gas-phase metallicity (Tremonti et al. 2004; Mannucci et al. 2010; Berg et al. 2012) and dust content increases (Rémy-Ruyer et al. 2013; Popping, Somerville & Galametz 2017; Popping & Péroux 2022). The relationship between β_{obs}^{1550} and M_* for a sample of $z \sim 7$ galaxies from Finkelstein et al. (2012) is overplotted as a black dashed line



Figure 12. The inferred evolution of $f_{\rm esc, LyC}$ at high redshift, assuming the relation between $\beta_{\rm obs}^{1550}$ and $f_{\rm esc, LyC}$ (Fig. 7) applies for the $M_{\rm UV}^{\rm obs} - \beta_{\rm obs}^{1550}$ relations from Bouwens et al. (2014) (equation 17). We estimate $f_{\rm esc, LyC}$ at six different redshifts and provide an estimate of the evolution of $f_{\rm esc, LyC}$ at with $M_{\rm UV}^{\rm obs}$. The $f_{\rm esc, LyC}$ varies shallowly with $M_{\rm UV}^{\rm obs}$ for galaxies brighter than -20 mag. In general, these observationally motivated relations suggest that fainter galaxies at higher redshifts emit a larger fraction of their ionizing photons.

in Fig. 11. We choose the $z \sim 7$ range from Finkelstein et al. (2012) because $z \sim 7$ is squarely within the epoch of reionization and it is the same redshift used for $M_{\rm UV}^{\rm obs}$ from Bouwens et al. (2014). Similar to Fig. 9, the correspondence between the high-redshift samples and the LzLCS+ is striking: the Finkelstein et al. (2012) relation tracks through the centre of the LzLCS+ points (both detections and nondetections). This suggests that the LzLCS+ β_{obs}^{1550} varies with host galaxy properties in similar ways as high-redshift galaxies. We use equation (11) to shade regions in Fig. 11 of expected $f_{esc, LVC}$ values. On average, $z \sim 7$ galaxies with $\log(M_*/M_{\odot}) < 8.7$ (7.6) emit more than 5 per cent (20 per cent) of their ionizing photons. Fig. 11 also includes individual z > 8 galaxies from the literature (Wilkins et al. 2016: Bhatawdekar & Conselice 2021: Tacchella et al. 2022). Of all the individual z > 8 galaxies, only GOODSN-35589 and EGS-68560, both at $\log(M_*/M_{\odot}) = 9.1$, are blue enough to emit more than 5 per cent of their ionizing photons. These galaxies are weak LyC emitters with a predicted $f_{\rm esc, LyC} \sim 6$ per cent. The other z > 8sources are too red to emit appreciable LyC. This is likely because most of the individual literature sources are biased towards bright and massive galaxies. If the LzLCS+ relationship between β_{obs}^{1550} and $f_{\rm esc, LvC}$ holds for galaxies within the epoch of reionization, faint and low stellar mass galaxies likely dominate the ionizing photon budget.

The LzLCS+ relations and the high-redshift β_{obs}^{1550} observations allow for some of the first indirect estimates of $f_{esc, LyC}$ during the epoch of reionization. We convert the Bouwens et al. (2014) β_{obs}^{1550} relations in equation (18) into population averages of $f_{esc, LyC}$ using the LzLCS+ relation. The predicted $f_{esc, LyC}$ curves for M_{UV}^{obs} down to -16 mag, often the current detection limits, are plotted in Fig. 12 for six different redshifts. These observationally motivated relations suggest that only galaxies fainter than -16 mag have $f_{esc, LyC} > 10$ per cent at z < 5. This is broadly consistent with current samples of LyC emitters at $z \sim 3$: galaxies brighter than $L_{z=3}^*$ ($M_{UV}^{obs} = -21$ mag) have non-detected LyC, while $f_{esc, LyC}$ increases with decreasing luminosity and is $f_{esc, LyC} \sim 12$ per cent for galaxies fainter than $L_{z=3}^*$ (Pahl et al. 2021). During the epoch of reionization (blue curves), all galaxies brighter than $L_{z=3}^*$ ($M_{UV}^{obs} \lesssim -21$ mag) have populationaveraged escape fractions less than 5 per cent. $f_{esc, LyC}$ increases for fainter galaxies as their β_{obs}^{1550} becomes bluer. During the epoch of reionization, the population-averaged $f_{esc, LyC}$ exceeds 10 per cent for M_{UV}^{obs} values fainter than -18 mag. These $f_{esc, LyC}$ trends are in slight tension with recent $f_{esc, LyC}$ estimates using the Ly α luminosity function (Matthee et al. 2022). This work found the $z \sim 7 f_{esc, LyC}$ distribution to peak near $f_{esc, LyC} = 11$ per cent for galaxies with $M_{UV}^{obs} = -19.5$ (a value two times higher than predicted in Fig. 12) and decreases for both brighter and fainter galaxies. The difference could be due to a β_{obs}^{150} tracing the evolving metal content of galaxies, while Matthee et al. (2022) determine $f_{esc, LyC}$ from the Ly α properties.

Can metallicity evolution explain the dramatic increase of $f_{esc, LyC}$ from z = 5 to z = 6 observed in Fig. 12 as reionization concludes? The increase in $f_{\text{esc, LyC}}$ at z > 6 for faint galaxies in Fig. 12 is due to the *b* value in equation (17) from Bouwens et al. (2014)becoming more negative. The b measures the β_{obs}^{1550} zero-point at fixed $M_{\rm UV}^{\rm obs}$. As b becomes more negative, from -1.9 ± 0.02 at z = 5 to -2.13 ± 0.44 at z = 8, galaxies become bluer at fixed $M_{\rm UV}^{\rm obs}$. What causes this blue shift? Figs 6 and 11 show that changes to gas-phase ionization, metallicity, and M_{*} heavily impacts β_{obs}^{1550} . Galaxies that vigorously form stars (and simultaneously produce ionizing photons) also rapidly synthesize metals. As galaxies grow in M_{*} they retain more of the these metals (Tremonti et al. 2004), allowing for them to create more dust (Rémy-Ruyer et al. 2013; Popping et al. 2017; Popping & Péroux 2022). Thus, the rapid buildup of stellar mass and metallicity during the epoch of reionization could lead to redder β_{obs}^{1550} and push $f_{esc, LyC}$ to lower values at later times.

The LzLCS+ β_{obs}^{1550} - $f_{esc, LyC}$ relationship and the empirical scaling between β_{obs}^{1550} and M_{UV}^{obs} can estimate \dot{n}_{ion} at high redshift. We use the luminosity function from Bouwens et al. (2015) and integrate from -24 to -16 mag (or approximately 0.01L*) for z = 4, 5, 6, 7, and 8 respectively. We use the luminosity functions from Bouwens et al. (2015) largely for consistency with the observed $M_{UV}^{obs} - \beta_{obs}^{1550}$ relations (equation 17), but other luminosity functions provide similar results at these M_{UV}^{obs} values (Livermore et al. 2017; Atek et al. 2018).

First, we follow previous work and assume a constant $\log(\xi_{ion}) =$ 25.27 (Bouwens et al. 2016) measured for a population of galaxies at $z \sim 4-5$. This ξ_{ion} is broadly consistent with values used in theoretical studies (Robertson et al. 2013, 2015; Naidu et al. 2020). In this base case, we assume a constant $f_{esc, LyC}$ for all galaxies of 10 per cent (dark gold line in Fig. 13) and 20 per cent (light gold line in Fig. 13), again similar to previous work (Robertson et al. 2013, 2015). The grey line in Fig. 13 estimates the \dot{n}_{ion} needed to balance hydrogen ionization and recombinations in the IGM at a given redshift (Madau et al. 1999; Ouchi et al. 2009), assuming a clumping factor that evolves with redshift (Shull et al. 2012). If \dot{n}_{ion} is greater than the grey line, ionizations occur more frequently than recombinations and the ionization state of the IGM increases. When \dot{n}_{ion} is less than this grey line, recombinations occur more frequently than photoionizations, and the ionization state of the IGM becomes more neutral (or does not change if the IGM is already neutral). Future work will refine these prescriptions and use semi-analytic models to solve the differential equations required to quantify the impact of the LzLCS+ prescriptions on the IGM reionization history (Trebitsch in preparation).

The gold lines roughly intersect with the grey line in the range $z \sim$ 7–8. This suggests that star-forming galaxies with constant properties produce enough ionizing photons to increase the IGM ionization fraction near redshifts of 8 (Ouchi et al. 2009; Robertson et al. 2013, 2015). The ionizing emissivity of star-forming galaxies continues to rise at lower redshifts as the number density of galaxies increases.



Figure 13. An observationally motivated estimate of the ionizing emissivity (\dot{n}_{ion}) during the epoch of reionization. The light blue circles are the values predicted using the LzLCS+ relation between β_{obs}^{1550} and $\xi_{ion} \times f_{esc, LyC}$ (equation 12) and the observed $M_{\rm UV}^{\rm obs} - \beta_{\rm obs}^{\rm 1550}$ relation (equation 17) at redshifts 4, 5, 6, 7, and 8 from Bouwens et al. (2014). The dark and light gold curves are estimates assuming a constant fesc. LvC of 10 and 20 per cent, respectively. The grey squares are observations of the ionizing emissivity (Becker & Bolton 2013; Becker et al. 2021). The grey line is the ionizing emissivity required for ionization to balance recombination at a given redshift (Madau, Haardt & Rees 1999; Ouchi et al. 2009), assuming a redshift-dependent clumping factor (Shull et al. 2012). The values above the grey line will increase the hydrogen ionization fraction in the IGM. The red line is a theoretical estimate of the AGN emissivity from Finkelstein et al. (2019). The dark blue circles are the sum of the AGN and the empirical star-forming ionizing emissivities. Star-forming galaxies dominate the ionizing emissivity at z > 5. The observationally motivated \dot{n}_{ion} is statistically consistent with the observed redshift 4 and 5 points, in marginal statistical tension with the $z \sim$ 6 data point, and suggests that star-forming galaxies began to emit sufficient ionizing photons to shift the IGM ionization state near redshifts of 8.

The LzLCS+ and high-redshift data suggest that $f_{\rm esc, LyC}$ is not constant, but rather varies with $\beta_{\rm obs}^{1550}$, M_{*}, and $M_{\rm UV}^{\rm obs}$. Does the LzLCS+ $\beta_{\rm obs}^{1550}$ - $f_{\rm esc, LyC}$ relation and high-redshift $\beta_{\rm obs}^{1550}$ observations suggest a similar production of ionizing photons as these constant assumptions? Using the Bouwens et al. (2014) $\beta_{\rm obs}^{1550}$ - $M_{\rm UV}^{\rm obs}$ relations (equation 17), we test two LzLCS+ relations: (1) the LzLCS+ $\beta_{\rm obs}^{1550}$ - $f_{\rm esc, LyC}$ relation (equation 18) with a constant $\log(\xi_{\rm ion}) = 25.27$ and (2) the LzLCS+ $\beta_{\rm obs}^{1550}$ - $\xi_{\rm esc, LyC}$ relation (equation 12). We integrate the product of these relations and the luminosity functions from -23 to -16 mag. These two predicted $\dot{n}_{\rm ion}$ values track each other within 0.2 dex (the constant $\xi_{\rm ion}$ has lower $\dot{n}_{\rm ion}$), and we only include the relation using $\xi_{\rm ion} \times f_{\rm esc, LyC}$ in Fig. 13 for clarity (the light blue points in Fig. 13).

The empirically motivated $\dot{n}_{\rm ion}$ values exceed the value to balance recombinations at $z \sim 7-8$. Near these redshifts, star-forming galaxies produce sufficient ionizing photons to increase the IGM ionization fraction. While we currently do not have statistical populations of $\beta_{\rm obs}^{1550}$ at z > 8, recent work suggests that the normalization of the luminosity function continues to decline at higher redshifts (Oesch et al. 2018; Bouwens et al. 2021, 2022; Finkelstein et al. 2022). Unless z > 8 galaxies are significantly bluer than $z \sim 7$ galaxies, the z > 8 galaxy populations will emit an insufficient number of ionizing photons to exceed recombinations in the IGM.

Between redshifts of 3 and 8, $\dot{n}_{\rm ion}$ from star-forming galaxies varies by less than a factor of 2. This constant $\dot{n}_{\rm ion}$ occurs even though $\rho_{\rm UV}$ increases by nearly a factor of 10 from $z \sim 8$ to z

 \sim 3. The dwindling $f_{\rm esc, LyC}$ at lower redshift (Fig. 12) balances this increasing number of galaxies to keep \dot{n}_{ion} relatively constant. Meanwhile, \dot{n}_{ion} from AGNs increases at lower redshifts (red line), such that star-forming galaxies and AGNs have similar \dot{n}_{ion} at z < z5. At $z \sim$ 3–4 star-forming galaxies and AGNs contribute nearly equally to the HI ionizing photon budget (Steidel et al. 2018; Dayal et al. 2020; Trebitsch et al. 2021). The total (AGNs plus starforming galaxies) \dot{n}_{ion} (dark blue circles) match both the low-redshift observations of the ionizing emissivity (grey squares) and the \dot{n}_{ion} value required for ionizations to exceed recombinations in the IGM (grey line) without overproducing ionizing photons at lower redshift. The observationally motivated $\beta_{\rm obs}^{1550}$ prescription of $f_{\rm esc, LyC}$ provides an empirically motivated estimation of the ionizing emissivity of high-redshift star-forming galaxies and suggests that star-forming galaxies produced sufficient ionizing photons to increase the IGM ionization fraction $z \sim 7-8$.

Inferring $\dot{n}_{\rm ion}$ at high redshifts using the LzLCS+ $\beta_{\rm obs}^{1550}$ - $f_{\rm esc, LyC}$ relations relies on a few critical assumptions extending from $z \sim$ 0.3 to $z \sim 6-9$. First, the dust extinction law sensitively impacts the shape of the FUV continuum, which connects β_{obs}^{1550} to the absorption of ionizing photons at 912 Å (see Appendix A). Dust properties may evolve strongly over time as different elements - specifically Fe, C, Si, and Mg - have different formation mechanisms and are produced on different time-scales. The strong correspondence between $\beta_{\rm obs}^{1550}$ and both M_{IIV}^{obs} and M_* suggests that there are similarities between galaxy properties and dust properties at high and low redshifts, but there could be significant differences. For instance, Saldana-Lopez et al. (2022) found $f_{esc, LVC}$ to be a median factor of 1.08 higher using the Reddy et al. (2016) extinction law than the SMC law. This factor would slightly impact the ionizing emissivity of star-forming galaxies, but would be insufficient to qualitatively change their ability to reionize the early IGM. If the high-redshift dust extinction law is observed to significantly deviate from the Reddy et al. (2016) law, Appendix A and Section 4 illustrate how the $f_{esc, LvC}$ relations can be updated in the future to include these reformulated high-redshift dust laws. Second, observing the range of β_{int}^{1550} of high-redshift samples will determine whether β_{int}^{1550} strongly deviates from the LzLCS+ distribution. Equation (8) demonstrates that $\beta_{\text{int}}^{1550}$ is required to connect E (B - V) and $\beta_{\text{obs}}^{1550}$; even though $\beta_{\text{int}}^{1550}$ has a narrow range in the LzLCS+, it must also be constrained at high redshift. Third, the current high-redshift $\beta_{\rm obs}^{1550} - M_{\rm UV}$ relations have appreciable uncertainties. For instance, the slope of the $z \sim 7 \beta_{\rm obs}^{1550} - M_{\rm UV}$ from Bouwens et al. (2014) is 0.2 ± 0.07 , implying that the relative scaling of the $f_{\rm esc, LyC}$ - $\beta_{\rm obs}^{1550}$ at high redshift is still modestly uncertain. Large samples from JWST observations will dramatically improve these relations and tighten our understanding of $f_{\rm esc, LyC}$ during the epoch of reionization. Finally, using the LzLCS + β_{obs}^{150} to determine $f_{esc, LyC}$ relies on high- and low-redshift galaxies having similar neutral gas properties. The ionization states of high-redshift galaxies are currently underconstrained and cannot be compared to the LzLCS+. Future LzLCS studies will attempt to disentangle additional correlations between $f_{\rm esc, LyC}$, $\beta_{\rm obs}^{1550}$, and the ionization properties of the LzLCS (Jaskot et al. in preparation). As the JWST makes similar observations at high redshift, the correlations presented here can be tailored to the discovered conditions at high redshift.

8 SUMMARY AND CONCLUSIONS

Here, we presented a new analysis of the FUV stellar continuum slope (β) of a combined sample that we refer to as the LzLCS+ sample, which consists of 66 galaxies from the LzLCS and 23 galaxies from literature low-redshift studies. We fit 89 galaxies that have *Hubble*

place, the land. The he moved an Indian n or have s piece of hents that ugh grant Institute. ASA/ESA e Science niversities t NAS 5s) 13744,

Space Telescope LyC observations with stellar population synthesis models to estimate the intrinsic (β_{int}^{1550}) and observed (β_{obs}^{1550}) stellar continuum slope at 1550 Å. These stellar population fits are critical to determine β_{obs}^{1550} at the non-ionizing wavelengths that are typically observed at high redshifts. Comparing the observed O VI stellar wind profiles to stellar population models demonstrates that the LzLCS+ is characterized by young stellar populations (Fig. 1). We calculated and the stellar population stellar population stellar wind stellar by the stellar population stellar population (Fig. 1). We calculated

many properties from the stellar population fits, but focused on $\beta_{\rm obs}^{1550}$ because it is currently commonly observed at high redshift. We then explored trends between $\beta_{\rm obs}^{1550}$ and various properties of

we then explored trends between β_{obs}^{**} and various properties of the LzLCS+. Our main findings are:

(i) The β_{obs}^{1550} scales with the [O III]/[O II] flux ratio (7.5 σ significance; Fig. 6, left-hand panel), 12 + log(O/H) (3.4 σ ; Fig. 6, right-hand panel), H β equivalent width (6.7 σ), stellar mass (5.9 σ ; Fig. 11), and the observed FUV absolute magnitude (M_{UV}^{obs} ; 3.4 σ ; Fig. 9). In general, this implies that lower metallicity, higher ionization, lower mass, and fainter galaxies have bluer FUV continuum slopes.

(ii) β_{obs}^{1550} is largely set by the colour excess, E(B - V) (Fig. 5), and does not scale with intrinsic stellar properties (Fig. 4). Thus, we suggest that the LzLCS+ β_{obs}^{1550} is largely driven by dust attenuation.

(iii) β_{obs}^{1550} scales strongly with the LyC escape fraction ($f_{esc, LyC}$) at the 5.7 σ significance (Fig. 7). Galaxies with $\beta_{obs}^{1550} = -2.11$, -2.35, -2.60 have population-averaged $f_{esc, LyC} = 5$, 10, 20 per cent, respectively. We provide a scaling relation between β_{obs}^{1550} and $f_{esc, LyC}$ in equation (11).

(iv) $\beta_{\text{obs}}^{1550}$ also scales with the product of $f_{\text{esc, LyC}} \times \xi_{\text{ion}}$ (Fig. 8 and equation 12).

(v) There is appreciable scatter in the $f_{\rm esc, LyC}$ - $\beta_{\rm obs}^{1550}$ relations that scales strongly with $f_{\rm esc, LyC}^{\rm rel}$ (see colour-coding of the points in Fig. 7). These $\beta_{\rm obs}^{1550}$ - $f_{\rm esc, LyC}$ relations are well suited to estimate population averages, rather than $f_{\rm esc, LyC}$ from individual galaxies.

Assuming these above findings transfer to high-redshift galaxies, and combining them with previous observations of β_{obs}^{1550} at high redshift, we infer the emission of ionizing photons into the IGM during the epoch of reionization (Section 7). Specifically we find that:

(i) The β_{obs}^{1550} of high-redshift and low-redshift galaxies have similar relations between both M_{UV}^{obs} (Fig. 9) and stellar mass (Fig. 11). Low-mass galaxies are bluer at both high and low redshift (Fig. 9). Faint galaxies likely dominated the emission of ionizing photons during the epoch of reionization.

(ii) $f_{\rm esc, LyC}$ increases for fainter $M_{\rm UV}^{\rm obs}$ at higher redshifts (Fig. 12). Only galaxies with $M_{\rm UV}^{\rm obs} > -16$ mag at z < 5 have populationaveraged $f_{\rm esc, LyC}$ greater than 10 per cent. Galaxies during the epoch of reionization are estimated to have higher $f_{\rm esc, LyC}$ than their lower redshift counterparts (compare the blue and golds curves in Fig. 12). This implies that relatively bright galaxies ($M_{\rm UV}^{\rm obs} > -18$ mag) have $f_{\rm esc, LyC}$ population averages greater than 10 per cent.

(iii) β_{obs}^{1550} (and $f_{esc, LyC}$) likely evolves with time and galaxy mass as galaxies synthesize and retain metals and dust.

We combine the LzLCS+ β_{obs}^{1550} - $f_{esc, LyC}$ relation, the high-redshift β_{obs}^{1550} observations, and the observed luminosity functions to make empirical estimates of the ionizing photon emissivity at high redshift (Fig. 13). These empirically motivated prescriptions suggest that galaxies near $z \sim 7-8$ first emitted a sufficient number of ionizing photons to increase the IGM ionization fraction. \dot{n}_{ion} of star-forming galaxies flattens at lower redshifts and varies by less than a factor of 2 from z = 8 to z = 4. As such, $z \sim 4$ star-forming galaxies and AGNs have similar \dot{n}_{ion} . The star-forming plus AGN emissivities are consistent with IGM observations at $z \sim 4-5$.

Soon *JWST* will compile larger and more robust samples of faint galaxies at z > 5. The empirical relations provided here will reveal whether these galaxy populations produced sufficient ionizing photons to reionize the IGM. Jaskot et al. (in preparation) will explore whether there exists strong secondary correlations between $f_{\text{esc, LyC}}$ and parameters such as $\beta_{\text{obs}}^{1550}$, O₃₂, H β equivalent width, and stellar mass. These new relations will take advantage of the upcoming spectroscopic capabilities of *JWST* to provide even more robust constraints on the ionizing emissivity of high-redshift galaxies and their impact on cosmic reionization.

ACKNOWLEDGEMENTS

We acknowledge that the location where this work took place, the University of Texas at Austin, sits on a stolen indigenous land. The Tonkawa live in central Texas and the Comanche and Apache moved through this area. We pay our respects to all the American Indian and Indigenous Peoples and communities who have been or have become a part of these lands and territories in Texas, on this piece of Turtle Island.

We thank the referee for insightful and useful comments that significantly improved the clarity and scope of this paper.

Support for this work was provided by NASA through grant number HST-GO-15626 from the Space Telescope Science Institute. This research is based on observations made with the NASA/ESA *Hubble Space Telescope* obtained from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These observations are associated with program(s) 13744, 14635, 15341, 15626, 15639, and 15941. STScI is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract NAS 5-26555. ASL acknowledge support from Swiss National Science Foundation. RA acknowledges support from ANID Fondecyt Regular 1202007. MH is fellow of the Knut and Alice Wallenberg Foundation.

DATA AVAILABILITY

All data in this paper are made possible through previous publications.

REFERENCES

- Ahumada R. et al., 2020, ApJS, 249, 3
- Atek H., Richard J., Kneib J.-P., Schaerer D., 2018, MNRAS, 479, 5184
- Bañados E. et al., 2018, Nature, 553, 473
- Becker G. D., Bolton J. S., 2013, MNRAS, 436, 1023
- Becker R. H. et al., 2001, AJ, 122, 2850
- Becker G. D., D'Aloisio A., Christenson H. M., Zhu Y., Worseck G., Bolton J. S., 2021, MNRAS, 508, 1853
- Berg D. A. et al., 2012, ApJ, 754, 98
- Bergvall N., Zackrisson E., Andersson B.-G., Arnberg D., Masegosa J., Östlin G., 2006, A&A, 448, 513
- Bhatawdekar R., Conselice C. J., 2021, ApJ, 909, 144
- Borthakur S., Heckman T. M., Leitherer C., Overzier R. A., 2014, Science, 346, 216
- Bouwens R. J. et al., 2014, ApJ, 793, 115
- Bouwens R. J. et al., 2015, ApJ, 803, 34
- Bouwens R. J., Smit R., Labbé I., Franx M., Caruana J., Oesch P., Stefanon M., Rasappu N., 2016, ApJ, 831, 176
- Bouwens R. J. et al., 2021, AJ, 162, 47
- Bouwens R. J., Illingworth G. D., Ellis R. S., Oesch P. A., Stefanon M., 2022, preprint (arXiv:2205.11526)

- Calzetti D., 2013, in Falcón-Barroso J., Knapen J. H., eds, Secular Evolution of Galaxies. Cambridge University Press, Cambridge, UK, p. 419
- Calzetti D., Kinney A. L., Storchi-Bergmann T., 1994, ApJ, 429, 582
- Calzetti D., Armus L., Bohlin R. C., Kinney A. L., Koornneef J., Storchi-Bergmann T., 2000, ApJ, 533, 682
- Cardelli J. A., Clayton G. C., Mathis J. S., 1989, ApJ, 345, 245
- Chisholm J., Tremonti C. A., Leitherer C., Chen Y., 2017, MNRAS, 469, 4831
- Chisholm J. et al., 2018, A&A, 616, A30
- Chisholm J., Rigby J. R., Bayliss M., Berg D. A., Dahle H., Gladders M., Sharon K., 2019, ApJ, 882, 182
- Chisholm J., Prochaska J. X., Schaerer D., Gazagnes S., Henry A., 2020, MNRAS, 498, 2554
- Davis D. et al., 2021, ApJ, 920, 122
- Dayal P. et al., 2020, MNRAS, 495, 3065
- De Barros S., Oesch P. A., Labbé I., Stefanon M., González V., Smit R., Bouwens R. J., Illingworth G. D., 2019, MNRAS, 489, 2355
- Dunlop J. S. et al., 2013, MNRAS, 432, 3520
- Eldridge J. J., Stanway E. R., Xiao L., McClelland L. A. S., Taylor G., Ng M., Greis S. M. L., Bray J. C., 2017, Publ. Astron. Soc. Aust., 34, e058
- Endsley R., Stark D. P., 2022, MNRAS, 511, 6042
- Endsley R., Stark D. P., Chevallard J., Charlot S., 2021, MNRAS, 500, 5229
- Fan X. et al., 2006, AJ, 132, 117
- Faucher-Giguère C.-A., 2020, MNRAS, 493, 1614
- Faucher-Giguère C.-A., Lidz A., Zaldarriaga M., Hernquist L., 2009, ApJ, 703, 1416
- Ferland G. J. et al., 2017, Rev. Mex. Astron. Astrofis., 53, 385
- Finkelstein S. L. et al., 2012, ApJ, 756, 164
- Finkelstein S. L. et al., 2015, ApJ, 810, 71
- Finkelstein S. L. et al., 2019, ApJ, 879, 36
- Finkelstein S. L. et al., 2022, ApJ, 928, 52
- Fitzpatrick E. L., 1999, PASP, 111, 63
- Fletcher T. J., Tang M., Robertson B. E., Nakajima K., Ellis R. S., Stark D. P., Inoue A., 2019, ApJ, 878, 87
- Flury S. R. et al., 2022a, ApJS, 260, 1
- Flury S. R. et al., 2022b, ApJ, 930, 126
- Fontanot F., Cristiani S., Monaco P., Nonino M., Vanzella E., Brandt W. N., Grazian A., Mao J., 2007, A&A, 461, 39
- Gazagnes S., Chisholm J., Schaerer D., Verhamme A., Rigby J. R., Bayliss M., 2018, A&A, 616, A29
- Gazagnes S., Chisholm J., Schaerer D., Verhamme A., Izotov Y., 2020, A&A, 639, A85
- Gnedin N. Y., 2000, ApJ, 542, 535
- Green J. C. et al., 2012, ApJ, 744, 60
- Green G. M. et al., 2018, MNRAS, 478, 651
- Harikane Y. et al., 2018, ApJ, 859, 84
- Hayes M., Schaerer D., Östlin G., Mas-Hesse J. M., Atek H., Kunth D., 2011, ApJ, 730, 8
- Henry A., Berg D. A., Scarlata C., Verhamme A., Erb D., 2018, ApJ, 855, 96
- Hopkins P. F., Hernquist L., Cox T. J., Kereš D., 2008, ApJS, 175, 356
- Hutchison T. A. et al., 2019, ApJ, 879, 70
- Inoue A. K., Shimizu I., Iwata I., Tanaka M., 2014, MNRAS, 442, 1805
- Izotov Y. I., Schaerer D., Thuan T. X., Worseck G., Guseva N. G., Orlitová I., Verhamme A., 2016a, MNRAS, 461, 3683
- Izotov Y. I., Orlitová I., Schaerer D., Thuan T. X., Verhamme A., Guseva N. G., Worseck G., 2016b, Nature, 529, 178
- Izotov Y. I., Schaerer D., Worseck G., Guseva N. G., Thuan T. X., Verhamme A., Orlitová I., Fricke K. J., 2018a, MNRAS, 474, 4514
- Izotov Y. I., Worseck G., Schaerer D., Guseva N. G., Thuan T. X., Fricke A. V., Orlitová I., 2018b, MNRAS, 478, 4851
- Izotov Y. I., Worseck G., Schaerer D., Guseva N. G., Chisholm J., Thuan T. X., Fricke K. J., Verhamme A., 2021, MNRAS, 503, 1734
- Jaskot A. E., Oey M. S., 2014, ApJ, 791, L19
- Ji Z. et al., 2020, ApJ, 888, 109
- Jiang L. et al., 2022, Nat. Astron., 6, 850
- Katz H. et al., 2022, MNRAS, 515, 4265
- Kelly B. C., 2007, ApJ, 665, 1489

MNRAS 517, 5104-5120 (2022)

- Kewley L. J., Groves B., Kauffmann G., Heckman T., 2006, MNRAS, 372, 961
- Kulkarni G., Keating L. C., Haehnelt M. G., Bosman S. E. I., Puchwein E., Chardin J., Aubert D., 2019a, MNRAS, 485, L24
- Kulkarni G., Worseck G., Hennawi J. F., 2019b, MNRAS, 488, 1035
- Leitet E., Bergvall N., Piskunov N., Andersson B.-G., 2011, A&A, 532, A107 Leitherer C. et al., 1999, ApJS, 123, 3
- Leitherer C., Ortiz Otálvaro P. A., Bresolin F., Kudritzki R.-P., Lo Faro B., Pauldrach A. W. A., Pettini M., Rix S. A., 2010, ApJS, 189, 309
- Leitherer C., Hernandez S., Lee J. C., Oey M. S., 2016, ApJ, 823, 64
- Livermore R. C., Finkelstein S. L., Lotz J. M., 2017, ApJ, 835, 113
- Madau P., Dickinson M., 2014, ARA&A, 52, 415
- Madau P., Haardt F., 2015, ApJ, 813, L8
- Madau P., Haardt F., Rees M. J., 1999, ApJ, 514, 648
- Mainali R. et al., 2018, MNRAS, 479, 1180
- Makan K., Worseck G., Davies F. B., Hennawi J. F., Prochaska J. X., Richter P., 2021, ApJ, 912, 38
- Mannucci F., Cresci G., Maiolino R., Marconi A., Gnerucci A., 2010, MNRAS, 408, 2115
- Marques-Chaves R., Schaerer D., Álvarez-Márquez J., Colina L., Dessauges-Zavadsky M., Pérez-Fournon I., Saldana-Lopez A., Verhamme A., 2021, MNRAS, 507, 524
- Martin D. C. et al., 2005, ApJ, 619, L1
- Maseda M. V. et al., 2020, MNRAS, 493, 5120
- Mason C. A., Naidu R. P., Tacchella S., Leja J., 2019, MNRAS, 489, 2669
- Matsuoka Y. et al., 2018, ApJ, 869, 150
- Matthee J. et al., 2021, MNRAS, 505, 1382
- Matthee J. et al., 2022, MNRAS, 512, 5960
- Mauerhofer V., Verhamme A., Blaizot J., Garel T., Kimm T., Michel-Dansac L., Rosdahl J., 2021, A&A, 646, A80
- McGaugh S. S., 1991, ApJ, 380, 140
- McLeod D. J., McLure R. J., Dunlop J. S., 2016, MNRAS, 459, 3812
- Mehta V. et al., 2017, ApJ, 838, 29
- Miralda-Escudé J., Rees M. J., 1994, MNRAS, 266, 343
- Naidu R. P., Tacchella S., Mason C. A., Bose S., Oesch P. A., Conroy C., 2020, ApJ, 892, 109

Downloaded from https://academic.oup.com/mnras/article/517/4/5104/6753210 by CNRS user on 06 July 2022

- Naidu R. P. et al., 2022, MNRAS, 510, 4582
- Nakajima K., Ouchi M., 2014, MNRAS, 442, 900
- Newville M., Stensitzki T., Allen D. B., Ingargiola A., 2014, LMFIT: Non-Linear Least-Square Minimization and Curve-Fitting for Python
- Oesch P. A., Bouwens R. J., Illingworth G. D., Labbé I., Stefanon M., 2018, ApJ, 855, 105
- Ouchi M. et al., 2009, ApJ, 696, 1164
- Pahl A. J., Shapley A., Steidel C. C., Chen Y., Reddy N. A., 2021, MNRAS, 505, 2447
- Pilyugin L. S., Vílchez J. M., Thuan T. X., 2006, MNRAS, 370, 1928
- Planck Collaboration XLVII, 2016, A&A, 596, A108
- Popping G., Péroux C., 2022, MNRAS, 513, 1531
- Popping G., Somerville R. S., Galametz M., 2017, MNRAS, 471, 3152 Prevot M. L., Lequeux J., Maurice E., Prevot L., Rocca-Volmerange B., 1984,
 - A&A, 132, 389
- Raiter A., Schaerer D., Fosbury R. A. E., 2010, A&A, 523, A64
- Reddy N. A., Steidel C. C., Pettini M., Bogosavljević M., 2016, ApJ, 828, 107
- Redwine K. et al., 2016, PASP, 128, 105006
- Rémy-Ruyer A. et al., 2013, A&A, 557, A95
- Ricci F., Marchesi S., Shankar F., La Franca F., Civano F., 2017, MNRAS, 465, 1915
- Rivera-Thorsen T. E., Östlin G., Hayes M., Puschnig J., 2017, ApJ, 837, 29
- Rivera-Thorsen T. E. et al., 2019, Science, 366, 738
- Roberts-Borsani G., Morishita T., Treu T., Leethochawalit N., Trenti M., 2022, ApJ, 927, 236
- Robertson B. E. et al., 2013, ApJ, 768, 71
- Robertson B. E., Ellis R. S., Furlanetto S. R., Dunlop J. S., 2015, ApJ, 802, L19
- Rosdahl J. et al., 2018, MNRAS, 479, 994
- Rutkowski M. J. et al., 2017, ApJ, 841, L27
- Saldana-Lopez A. et al., 2022, A&A, 663, A59

- Schaerer D. et al., 2022, A&A, 658, L11
- Shapley A. E., Steidel C. C., Strom A. L., Bogosavljević M., Reddy N. A., Siana B., Mostardi R. E., Rudie G. C., 2016, ApJ, 826, L24
- Shen X., Hopkins P. F., Faucher-Giguère C.-A., Alexander D. M., Richards G. T., Ross N. P., Hickox R. C., 2020, MNRAS, 495, 3252
- Shull J. M., Harness A., Trenti M., Smith B. D., 2012, ApJ, 747, 100
- Skillman E. D., 1989, ApJ, 347, 883
- Stanway E. R., Eldridge J. J., 2018, MNRAS, 479, 75
- Stark D. P. et al., 2015, MNRAS, 454, 1393
- Stefanon M., Bouwens R. J., Illingworth G. D., Labbé I., Oesch P. A., Gonzalez V., 2022, ApJ, 935, 94
- Steidel C. C., Bogosavljević M., Shapley A. E., Reddy N. A., Rudie G. C., Pettini M., Trainor R. F., Strom A. L., 2018, ApJ, 869, 123
- Tacchella S. et al., 2022, ApJ, 927, 170
- Tang M., Stark D. P., Chevallard J., Charlot S., 2019, MNRAS, 489, 2572
- Trebitsch M., Blaizot J., Rosdahl J., Devriendt J., Slyz A., 2017, MNRAS, 470, 224
- Trebitsch M. et al., 2021, A&A, 653, A154
- Tremonti C. A. et al., 2004, ApJ, 613, 898
- Vanzella E. et al., 2016, ApJ, 825, 41
- Vasei K. et al., 2016, ApJ, 831, 38
- Verhamme A., Orlitová I., Schaerer D., Hayes M., 2015, A&A, 578, A7
- Vilchez J. M., Pagel B. E. J., 1988, MNRAS, 231, 257
- Wang B., Heckman T. M., Leitherer C., Alexandroff R., Borthakur S., Overzier R. A., 2019, ApJ, 885, 57
- Wang B. et al., 2021, ApJ, 916, 3
- Wilkins S. M., Bouwens R. J., Oesch P. A., Labbé I., Sargent M., Caruana J., Wardlow J., Clay S., 2016, MNRAS, 455, 659
- Witstok J., Smit R., Maiolino R., Curti M., Laporte N., Massey R., Richard J., Swinbank M., 2021, MNRAS, 508, 1686
- Worseck G., Prochaska J. X., Hennawi J. F., McQuinn M., 2016, ApJ, 825, 144
- Xu X. et al., 2022, ApJ, 933, 202
- Yue B. et al., 2018, ApJ, 868, 115
- Yung L. Y. A., Somerville R. S., Finkelstein S. L., Popping G., Davé R., 2019, MNRAS, 483, 2983
- Yung L. Y. A., Somerville R. S., Popping G., Finkelstein S. L., 2020, MNRAS, 494, 1002

Yung L. Y. A., Somerville R. S., Finkelstein S. L., Hirschmann M., Davé R., Popping G., Gardner J. P., Venkatesan A., 2021, MNRAS, 508, 2706

Zackrisson E., Inoue A. K., Jensen H., 2013, ApJ, 777, 39

APPENDIX A: IMPACT OF THE EXTINCTION LAW

The fitted E(B - V) has one of the most statistically significant correlations with $f_{\text{esc, LyC}}$ (Saldana-Lopez et al. 2022). E(B - V) also strongly determines the $\beta_{\text{obs}}^{1550}$ of the LzLCS (see Section 4), but the E(B - V) parameter depends on the assumed extinction law. As such an important diagnostic quantity, here we discuss the assumptions underpinning the chosen extinction law and its impact on the inferred spectral shape of the stellar continuum models and the E(B - V)parameter.

In Saldana-Lopez et al. (2022), we tested the impact of the assumed extinction law. We found that varying the extinction law from the Reddy et al. (2016) to an extrapolation of the SMC law (Prevot et al. 1984) did not impact the model parameters that set the intrinsic shape: age and metallicity. Therefore, the intrinsic stellar population properties do not appreciably change with different extinction laws. However, the absolute values of the individual fitted X_j values do change because the SMC extinction law is both steeper and has larger $k(\lambda)$ values than the Reddy et al. (2016) law. This is likely because the FUV stellar spectral features (e.g. Fig. 1) constrain the intrinsic properties of the stellar population. This sets the intrinsic shape of the stellar continuum, β_{int}^{1550} , and E (B - V) is varied, using the assumed reddening law, to reconcile β_{int}^{1550} to β_{obs}^{1550} (see Section 4).

Comparing the different extinction laws provides a prescription to convert β_{obs}^{1550} into E(B - V) for different extinction laws. The left-hand panel of Fig. A1 shows the ratio of the E(B - V) assuming a Reddy et al. (2016) $(E(B - V)^{R16})$ to the E(B - V) assuming an extrapolation of the SMC law $(E(B - V)^{SMC})$. The ratio of the two E(B - V) values peaks strongly with a median value of 3.09 and a standard deviation of 0.09, much smaller than the median 0.7 mag uncertainty on the measured ratio. The right-hand panel of Fig. A1 shows the strong correlation between the two E(B - V)values.

The E(B - V) differences derived with different reddening laws can be estimated analytically. First, we approximate the ratio of the observed flux at two wavelengths (F(λ^1) and F(λ^1)) as the ratio of



Figure A1. The impact of the assumed extinction law on the fitted attenuation parameter. Left-hand panel: Histogram of the ratio of the attenuation parameters derived using the Reddy et al. (2016) (R16) and the SMC laws. This ratio peaks strongly at a value of 3.1. Right-hand panel: The correlation of the attenuation parameter for the R16 and SMC extinction laws for the LyC-detected (blue circles) and non-detected (gold squares) samples. Overplotted as a grey dashed line is an analytical fit to this scaling (equation A4).

5120 J. Chisholm et al.

the intrinsic flux times an unknown attenuation factor as

$$\frac{F(\lambda^1)}{F(\lambda^2)} = \frac{F_i(\lambda^1)}{F_i(\lambda^2)} e^{-E(B-V)\left[k(\lambda^1) - k(\lambda^2)\right]} = R e^{-E(B-V)\Delta k},$$
(A1)

where we have defined *R* as the intrinsic flux ratio at wavelengths 1 and 2, and $\Delta k = k(\lambda^1) - k(\lambda^2)$ is the difference of the extinction law at two wavelengths (λ^1 and λ^2). Regardless of the assumed law, the observed flux densities must be equal at the two wavelengths. Numerically, this means that

$$\frac{F^{\text{R16}}(\lambda^1)}{F^{\text{R16}}(\lambda^2)} = \frac{F^{\text{SMC}}(\lambda^1)}{F^{\text{SMC}}(\lambda^2)}.$$
(A2)

Substituting in equation (A1) then suggests that $E (B - V)^{R16}$ is related to $E (B - V)^{SMC}$ as

$$E(B-V)^{R16} = \frac{\Delta k^{SMC}}{\Delta k^{R16}} E(B-V)^{SMC} + \frac{1}{\Delta k^{R16}} \ln \frac{R^{R16}}{R^{SMC}}$$
(A3)

If we use 950 Å as wavelength 1 and 1200 Å as wavelength 2 (typical wavelengths used to fit the models for the LzLCS+), we find that $\Delta k^{R16} = 2.1$ and $\Delta k^{SMC} = 6.7$ such that the E (B - V) of the two laws is related as

$$E(B - V)^{R16} = 3.14 \times E(B - V)^{\text{SMC}} + 0.47 \ln \frac{R^{R16}}{R^{\text{SMC}}}$$
 (A4)

The slope of this line matches the median 3.09 found from the ratio of the two attenuation parameters (left-hand panel of Fig. A1). If we fix the slope of the line in Fig. A1 to the theoretical ratio of $\Delta k^{\text{SMC}}/\Delta k^{\text{R16}} = 3.14$, we find that $0.47 \ln (R^{\text{R16}}/R^{\text{SMC}}) = 0.004$, implying that assuming an SMC law leads to an intrinsic FUV flux that is 1.01 times that of the Reddy et al. (2016) law. Thus, a good approximation for the conversion of the E (B - V) determined in the 950–1200 Å region with the Reddy et al. (2016) law to the SMC law is a constant factor of 3.1. This exercise illustrates how the individual reddening laws can be converted between each other, depending on the Δk values of the individual extinction laws at their specific wavelengths. While we use the Reddy et al. (2016) law for the paper, equation (A3) can be used to convert E (B - V) to other dust extinction laws and derive the respective $\beta_{\text{obs}}^{1550}$ values.

¹Department of Astronomy, The University of Texas at Austin, 2515 Speedway, Stop C1400, Austin, TX 78712-1205, USA

²Department of Astronomy, University of Geneva, 51 Chemin Pegasi, CH-1290 Versoix, Switzerland

³Department of Astronomy, University of Massachusetts Amherst, Amherst, MA 01002, USA

⁴CNRS, IRAP, 14 Avenue E. Belin, F-31400 Toulouse, France

 ⁵Department of Astronomy, Williams College, Williamstown, MA 01267, USA
 ⁶Departamento de Astronomía, Universidad de La Serena, Av. Juan Cisternas 1200 Norte, La Serena, Chile

⁷ Instituto de Investigación Multidisciplinar en Ciencia y Tecnología, Universidad de La Serena, Raul Bitran 1305 La Serena, Chile

⁸Institut d'astrophysique de Paris, CNRS UMR7095, Sorbonne Université, 98bis Boulevard Arago, F-75014 Paris, France

⁹Laboratory for Atmospheric and Space Physics, Boulder, Colorado 80303, USA

¹⁰Space Telescope Science Institute, 3700 San Martin Drive Baltimore, MD 21218, USA

¹¹The Oskar Klein Centre, Department of Astronomy, Stockholm University, AlbaNova, SE-10691 Stockholm, Sweden

¹²Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

¹³Kapteyn Astronomical Institute, University of Groningen, PO Box 800, NL-9700 AV Groningen, the Netherlands

¹⁴Department of Astronomy, University of Michigan, Ann Arbor, MI 48109, USA

¹⁵Department of Physics and Astronomy, Minnesota State University, Mankato, MN, 56001, USA

¹⁶Minnesota Institute for Astrophysics, School of Physics and Astronomy, University of Minnesota, 316 Church Street SE, Minneapolis, MN 55455, USA

¹⁷Astronomy Department, University of Virginia, Charlottesville, VA 22904, USA

¹⁸Department of Astronomy & Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA

¹⁹Institut für Physik und Astronomie, Universität Potsdam, Karl-Liebknecht-Str 24/25, D-14476 Potsdam, Germany

This paper has been typeset from a TEX/LATEX file prepared by the author.