

Classical novae with CUBES

Luca Izzo, Paolo Molaro, Piercarlo Bonifacio, Gabriele Cescutti, Massimo Della Valle, Pierluigi Selvelli

► To cite this version:

Luca Izzo, Paolo Molaro, Piercarlo Bonifacio, Gabriele Cescutti, Massimo Della Valle, et al.. Classical novae with CUBES. Experimental Astronomy, 2022, 10.1007/s10686-022-09876-3. insu-03849370

HAL Id: insu-03849370 https://insu.hal.science/insu-03849370

Submitted on 19 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Classical novae with CUBES

```
Luca Izzo · Paolo Molaro · Piercarlo
Bonifacio · Gabriele Cescutti · Massimo
Della Valle · Pierluigi Selvelli
```

Received: date / Accepted: date

Abstract Among the main science cases that have motivated the proposal of CUBES, a new high-resolution spectrograph for the Very Large Telescope at the European Southern Observatory, there is the study and the characterisation of the nucleosynthesis of beryllium. Classical novae have been proposed since the '70s as one of the main factories of lithium in the Galaxy, but this hypothesis has been demonstrated on empirical basis only recently thanks to the direct identification of lithium in V1369 Cen and through the observations of the resonance transition of ⁷Be II, the ⁷Li parent, at 313.0 nm in the near-UV range. CUBES is then the ideal instrument to quantify the amount of ⁷Be and therefore of ⁷Li produced by the different novae types hosted in the different components of the Milky Way and also in its nearby satellite galaxies. As important by-product of high resolution spectroscopic observa-

L. Izzo

E-mail: luca.izzo@nbi.ku.dk

P. Molaro

INAF-Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, I-34143 Trieste, Italy Institute of Fundamental Physics of the Universe, Via Beirut 2, Miramare, I-34143 Trieste, Italy

P. Bonifacio

G. Cescutti

Dipartimento di Fisica, Sezione di Astronomia, Università di Trieste, Via G. B. Tiepolo 11, 34143 Trieste, Italy

INAF-Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, I-34143 Trieste, Italy M. Della Valle

INAF-Osservatorio Astronomico di Napoli, Salita Moiariello 16, I-80131 Napoli, Italy P. Selvelli

DARK, Niels Bohr Institute, University of Copenhagen, Jagtvej 128, 2200 Copenhagen, Denmark

GEPI, Observatoire de Paris, Université PSL, CNRS, Place Jules Janssen, 92195 Meudon, France

INAF-Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, I-34143 Trieste, Italy

tions obtained with CUBES, there are the study of the properties of nova ejecta abundances, the shocks evolution in novae and their connection with the high-energy emission observed in these transients, from satellites as Fermi and Swift.

Keywords nuclear reactions \cdot nucleosynthesis \cdot abundances – stars: novae

1 Introduction

Classical novae (CNe) originate in interacting binary systems composed by a primary white dwarf (WD) that accretes matter from a late-type main sequence, or a red giant companion [1]. The accreted gas piles up onto the WD surface until thermo-nuclear reactions ignite [2]. The following production of unstable e^+ -isotopes provides the fuel for the expansion of the external layers into the interstellar medium and then contributing to its enrichment [3]. In the '70s, it was proposed that, during the thermo-nuclear runaway (TNR), CNe could form lithium through the formation of the ⁷Be isotope from the ³He (α, γ) ⁷Be reaction [4,5]. This isotope decays indeed into ⁷Li via electron capture with a half-life time decay of ~ 53 days [6]. The study of the processes that lead to the production of lithium in the Universe, and its evolution, is one of the hot topics of modern astrophysics, both for cosmology [7] and for the chemical evolution of the Galaxy [8–11]. The present lithium abundance, $A(Li) \sim 3.3$ [12], as measured from meteorites or from young T Tauri stars, is a factor of four larger than the BBN primordial value, $A(Li) \sim 2.7$ [7] and one order of magnitude larger than the abundance observed in old metal-poor stars [13,14]. Then, a Galactic source is required to explain the observed overabundance, and it is then crucial to look for the $^{7}\text{Be}/^{7}\text{Li}$ features in the early phase spectra of CNe to confirm them as lithium factories in our Galaxy.

2 The identification of ⁷Be in nova spectra

The time-scale of a CN outburst mainly depends on the mass density of the ejecta and its expanding velocity [15], which in turn are a proxy for the underlying WD mass [16]. Based on the above information and on the time-scale of the beryllium decay, our search for ⁷Be-⁷Li features must focus on the epochs immediately after the maximum brightness of the nova outburst, when the ejecta is still cool and optically thick. A relatively high cadence monitoring will allow to identify the nova composition and its kinematics from the analysis of the evolution of P-Cygni absorption line profiles, originating from the nova ejecta emission. This timeline generally corresponds to 1-2 weeks after the outburst for very fast CNe, to two-three months for slow outbursts. However, despite CN spectra have been observed for almost a hundred years, it was only in recent times that we have been able to search for ⁷Be/Li evidences in CN spectra, in particular in the near-UV range where the resonance

line of the ⁷Be II 313.0 nm is observed, thanks to the use of advanced highresolution spectrographs, such as the proposed Cassegrain U-Band Efficient Spectrograph (CUBES) [17,18] for the Very Large Telescope (VLT) in Chile. The first identifications of ⁷Be II in CN spectra have indeed been reported using specific spectrographs mounted on 8-meter class telescopes, and that are capable to observe the near-UV range, such as the High Dispersion Spectrograph (HDS) at the SUBARU telescope at Hawaii, US, and the Ultraviolet and Visual Echelle Spectrograph (UVES) currently operating at the VLT [19, 20].

A monitoring program is then necessary to quantify the Li yield from CNe. Since 2015 we have started a target-of-opportunity (ToO) program at the VLT to observe the outburst of bright novae with the UVES instrument [20]. A ToO program is necessary, given the transient nature of this class of astrophysical sources: we have been able to identify the P-Cygni absorptions for both the doublet components of the ⁷Be II 313.0 nm feature in the early spectra of fast and slow novae [20–22], see as an example in Fig. 1 the recent detection in V6595 Sgr [23]. We have monitored the spectral evolution until the ⁷Be features resulted unaffected by saturation effects. The $^7\mathrm{Be}$ abundance is estimated by comparison with the resonance Ca II H,K lines. Calcium is not synthesised in the TNR and provides an estimate for the ⁷Be, or ⁷Li, synthesized in the TNR of log $N(BeII)/N(HI) + 12 = 7.38 \pm 0.49$ [23]. Using these yields, chemical abundance simulations of the evolution of lithium in our Galaxy suggest that CNe can account for a large fraction of the lithium observed in the Milky Way [24,25,10,9]. However, the over-abundance of lithium estimated in CN outbursts are also in disagreement (by a factor of ~ 10 in N(Li)) with the expectations from theoretical simulations of CN explosions: while these still confirm the main role played by CNe as main factories of lithium in the Galaxy, at the same time they do not reproduce the observed lithium yield value using different nova models and accreted matter composition onto the primary WD [25]. On the other hand, recent and more accurate Galactic chemical evolution simulations that take into account the formation and evolution of binary systems giving rise to CNe explosions, are in agreement with the yield inferred from observations [26]. Consequently, more data, and of high quality, are still needed to: 1) increase the current statistics of events in the Galaxy and nearby systems (being currently composed of 12 Galactic and two SMC novae [23]), to infer a more accurate lithium yield by building a statistically-significant nova sample, and then solve the "tension" between the observational and the theoretical results; 2) to characterize the different CN progenitor populations in terms of their lithium yield (see also below); and 3) to verify if ionization effects and line saturation [27,28] can have an important role in the lithium yield estimate, thanks to a better signal-to-noise value (>> 50) in the 300-400 nm range; detailed analysis made with the 1-dim CLOUDY code have excluded this possibility in some events [23], but more detailed spectral synthesis models that takes into account 3-dim effects, non-local thermodynamic equilibrium and the effects of ejecta inhomogeneities must be used to observed spectra in order to provide a more accurate modelling and lithium abundance inference.



Fig. 1 The recent detection of $^7\mathrm{Be}$ II in V6595 Sgr [23], compared with other absorption lines in the nova ejecta.

An important, and yet unexplored field, is also represented by the characterization of CN populations in the Milky Way. There is nowadays a clear evidence that CNe originates from two, or maybe more, progenitor populations: the main indication comes from the evidence that fast novae are more concentrated on the Galactic plane while slow novae are observed also at higher Galactic latitudes [15]. The speed class is mainly driven by the ejected mass [16], with fast novae characterised by smaller expelled masses, and then more massive progenitor WDs, than slow novae. This result could have important implications for the chemical enrichment of different regions of the MW: recent simulations suggest that CNe could be the main factories of lithium for the thin disk component of the Milky Way [9]. The identification of a massive ONe progenitor WD is generally associated with the presence of bright forbidden lines of neon, which are visible in the wavelength range 330-390 nm, which is available to CUBES. We still lack a more complete sample of Galactic novae at high spectral resolution to understand the properties of nova progenitors, if these can be ascribed to a different ejecta composition/metallicity, or to other properties [15], and their relationship with lithium production. All this can be accomplished with future near-UV spectrographs, such as CUBES.

The recent ⁷Be detection in CNe observed in nearby local galaxies, such as the two Magellanic clouds [29] has also confirmed the important role of CNe as lithium factories in galaxies that are characterised by a metal poor stellar population and with an unresolved lithium enrichment problem [30]. CUBES at the VLT will definitely allow to investigate the near-UV spectral region for CNe in the nearby galaxies such as the Magellanic clouds (where CNe peak at $V \sim 11 - 12$ mag), and pushing our search to fainter events with $V \sim 16$ mag, given its more than > ten-fold flux sensitivity compared to UVES. We have indeed simulated the Day 1.6 spectrum of the recurrent nova RS Oph (Molaro et al. submitted), see also Fig. 2, using the CUBES exposure time calculator software [31], and the high-resolution mode, obtaining a signal-tonoise of ~ 120 using an exposure time which is $\sim 1/10$ the one used for the UVES spectrum. The high spectral resolution ($R \sim 22,000$, corresponding to a velocity resolution of $\sim 12 \text{ km/s}$ provided by CUBES will clearly allow us to identify and measure the properties of the ⁷Be features in CNe, and disentangle the doublet components, which are separated by ~ 62 km/s. Consequently, the lower spectral resolution compared with UVES does not limit our science goals, being high enough to distinguish narrow lines in the ejecta of CNe. This is clearly evident in our simulation of the Day 1.6 spectrum of RS Oph, which has been corrected for the telluric O3 Huggins bands [32]. To this purpose, we used the E2E CUBES simulator described in [31]. The results of our simulation are shown in Fig. 2, where it is clearly shown the potential of CUBES in detecting and reconstructing the line profile of the ⁷Be II absorption feature, as well as to identify narrow metal lines during the early stages of a CN outburst. The wavelength coverage provided by CUBES is ideal for this purpose: so far, ⁷Be abundance has been measured using Ca II H&K lines as reference, and both lines are visible in the wavelength range provided by CUBES. Last, while a flux calibration precision of less than 5% is required for late nebular epochs, when from the analysis of emission lines we will infer some physical properties of the ejecta like the abundance and ejected mass, for the early phases it is not required, given that to estimate relative abundances we only need equivalent width measurements, which are not strictly dependent on the level of the flux calibration.

3 Further insights on nova phenomenology with CUBES.

An interesting by-product from the monitoring of CNe with the spectral resolution provided by CUBES is the characterisation of the abundance pattern derived from narrow absorption lines of heavy elements observed during the bright optically thick phases of CNe, also named THEA (transient heavy element absorptions - [33]). The study and the abundance characterisation of these features in the near-UV range can definitely shed light on their origin (if associated with the progenitor WD or from the secondary companion), their evolution within the first days/weeks of a nova outburst, how they relate with the new proposed paradigm for nova explosions [34], and finally to check if the components where these transitions take origin include lithium/beryllium.

Only a high-cadence follow-up program will permit to study the dynamics of the ejecta and its correlation with the high-energy radiation recently reported by gamma-ray detectors [35]. The energetic emission was found to be correlated with rebrightenings observed in the optical light curves as well as with the simultaneous appearance of additional higher velocity components in the spectra [36]. This evidence supported the proposal of a new scenario for the CN explosions, characterised by the presence of multiple ejecta components that interact with previously expelled gas, thus producing the observed shocked emission [34,37].

4 Conclusions

High-resolution spectrographs with near-UV capabilities opened a new era for the study of the TNR in CN. Multi-wavelength observations have provided evidence for shocks, aspherical ejecta and particle acceleration. The possibility to combine these observations with the high-resolution at optical and near-UV wavelengths with CUBES will be relevant in understanding the origin of the high-energy radiation and of the shocked emission. At the same time, CUBES will be one of the few spectrographs that will allow to study the abundance of the ⁷Be II and therefore shed a definitive light on the role of CNe as main lithium factories in the MW and in the Local Group galaxies.

Acknowledgements Data presented in this work will be made available on reasonable request. LI was supported by grants from VILLUM FONDEN (project number 16599 and 25501).

Conflict of interest

The authors declare that they have no conflict of interest.

References

- 1. M.F. Bode, A. Evans, Classical Novae, vol. 43 (2008)
- 2. J.S. Gallagher, S. Starrfield, ARA&A16, 171 (1978). DOI 10.1146/annurev.aa.16.090178.001131
- R.D. Gehrz, J.W. Truran, R.E. Williams, S. Starrfield, PASP110, 3 (1998). DOI 10.1086/316107
- 4. M. Arnould, H. Norgaard, A&A42, 55 (1975)
- S. Starrfield, J.W. Truran, W.M. Sparks, M. Arnould, ApJ222, 600 (1978). DOI 10.1086/156175
- 6. A.G.W. Cameron, W.A. Fowler, ApJ164, 111 (1971). DOI 10.1086/150821
- 7. B.D. Fields, P. Molaro, S. Sarkar, ArXiv e-prints (2014)



Fig. 2 (Upper panel) The spectrum of the Be star HD149757 as observed by CUBES (blue curve) using the dedicated E2E code [31], and used for the estimate of the O_3 correction (green curve). The orange curve corresponds to the stellar continuum used for the estimate of the O_3 bands. Narrow stellar/interstellar features have been opportunely excluded using a smoothing median filter on the final O_3 correction spectrum. (Lower panel) The simulated spectrum of the recurrent nova RS Oph on Day 1.6 as if it were observed by CUBES (blue curve). The original spectrum was observed with UVES, and it was given as input of the E2E code. The ⁷Be II feature in the nova ejecta would have been clearly detected also with CUBES (gray mark), including several other narrower absorptions originating in the interaction of the nova ejecta with the secondary red giant wind (Molaro et al. submitted). The upper green curve corresponds to the telluric correction for the O_3 Huggins bands, derived from HD149757. The orange curve is the result of the correction for the Huggins bands from the simulated observed spectrum: note how the absorption feature associated with ⁷Be II still persists in the corrected spectrum, while other "fake" absorptions disappear, being originated in O_3 telluric features. To test the quality of the CUBES simulations, we have measured a pseudo equivalent width for the $^7\mathrm{Be}$ II feature of $pEW_{Be,C}=1.043\pm0.133$ Å (SNR ~ 120 for a 3s exposure) while from UVES we got $pEW_{Be,U} = 1.065 \pm 0.086$ Å.

- 8. F. D'Antona, F. Matteucci, A&A248, 62 (1991)
- 9. G. Cescutti, P. Molaro, MNRAS482, 4372 (2019). DOI 10.1093/mnras/sty2967
- D. Romano, L. Magrini, S. Randich, G. Casali, P. Bonifacio, R.D. Jeffries, F. Matteucci, E. Franciosini, L. Spina, G. Guiglion, C. Chiappini, A. Mucciarelli, P. Ventura, V. Grisoni, M. Bellazzini, T. Bensby, A. Bragaglia, P. de Laverny, A.J. Korn, S.L. Martell, G. Tautvaisiene, G. Carraro, A. Gonneau, P. Jofré, E. Pancino, R. Smiljanic, A. Vallenari, X. Fu, M.L. Gutiérrez Albarrán, F.M. Jiménez-Esteban, D. Montes, F. Damiani, M. Bergemann, C. Worley, arXiv e-prints arXiv:2106.11614 (2021)
- 11. F. Matteucci, A&A Rev.29(1), 5 (2021). DOI 10.1007/s00159-021-00133-8
- K. Lodders, H. Palme, H.P. Gail, Landolt Börnstein 4B, 712 (2009). DOI 10.1007/978-3-540-88055-4_34

- 13. F. Spite, M. Spite, A&A115, 357 (1982)
- L. Sbordone, P. Bonifacio, E. Caffau, H.G. Ludwig, N.T. Behara, J.I. González Hernández, M. Steffen, R. Cayrel, B. Freytag, C. van't Veer, P. Molaro, B. Plez, T. Sivarani, M. Spite, F. Spite, T.C. Beers, N. Christlieb, P. François, V. Hill, A&A522, A26 (2010). DOI 10.1051/0004-6361/200913282
- 15. M. Della Valle, L. Izzo, A&A Rev.28(1), 3 (2020). DOI 10.1007/s00159-020-0124-6
- 16. M. Livio, ApJ**393**, 516 (1992). DOI 10.1086/171524
- B. Barbuy, V. Bawden Macanhan, P. Bristow, B. Castilho, H. Dekker, B. Delabre, M. Diaz, C. Gneiding, F. Kerber, H. Kuntschner, G. La Mura, W. Maciel, J. Meléndez, L. Pasquini, C.B. Pereira, P. Petitjean, R. Reiss, C. Siqueira-Mello, R. Smiljanic, J. Vernet, Ap&SS354(1), 191 (2014). DOI 10.1007/s10509-014-2039-z
- A. Zanutta, S. Cristiani, D. Atkinson, V. Baldini, A. Balestra, B. Barbuy, V.B.P. Macanhan, A. Calcines, G. Calderone, S. Case, B.V. Castilho, G. Cescutti, R. Cirami, I. Coretti, S. Covino, G. Cupani, V. De Caprio, H. Dekker, P. Di Marcantonio, V. D'Odorico, H. Ernandes, C. Evans, T. Feger, C. Feiz, M. Franchini, M. Genoni, C.D. Gneiding, M. Kałuszyński, M. Landoni, J. Lawrence, D. Lunney, C. Miller, K. Molaverdikhani, C. Opitom, G. Pariani, S. Piranomonte, A. Quirrenbach, E.M.A. Redaelli, M. Riva, D. Robertson, S. Rossi, F. Rothmaier, W. Seifert, R. Smiljanic, J. Stürmer, I. Stilz, A. Trost, O. Verducci, C. Waring, S. Watson, M. Wells, W. Xu, T. Zafar, S. Zorba, Experimental Astronomy (2022). DOI 10.1007/s10686-022-09837-w
- A. Tajitsu, K. Sadakane, H. Naito, A. Arai, W. Aoki, Nature**518**, 381 (2015). DOI 10.1038/nature14161
- P. Molaro, L. Izzo, E. Mason, P. Bonifacio, M. Della Valle, MNRAS463, L117 (2016). DOI 10.1093/mnrasl/slw169
- L. Izzo, P. Molaro, P. Bonifacio, M. Della Valle, Z. Cano, A. de Ugarte Postigo, J.L. Prieto, C. Thöne, L. Vanzi, A. Zapata, D. Fernandez, MNRAS478, 1601 (2018). DOI 10.1093/mnras/sty435
- P. Molaro, L. Izzo, P. Bonifacio, M. Hernanz, P. Selvelli, M. della Valle, MNRAS492(4), 4975 (2020). DOI 10.1093/mnras/stz3587
- P. Molaro, L. Izzo, V. D'Odorico, E. Aydi, P. Bonifacio, G. Cescutti, E.J. Harvey, M. Hernanz, P. Selvelli, M. della Valle, MNRAS509(3), 3258 (2022). DOI 10.1093/mnras/stab3106
- L. Izzo, M. Della Valle, E. Mason, F. Matteucci, D. Romano, L. Pasquini, L. Vanzi, A. Jordan, J.M. Fernandez, P. Bluhm, R. Brahm, N. Espinoza, R. Williams, ApJ808, L14 (2015). DOI 10.1088/2041-8205/808/1/L14
- S. Starrfield, M. Bose, C. Iliadis, W.R. Hix, C.E. Woodward, R.M. Wagner, ApJ895(1), 70 (2020). DOI 10.3847/1538-4357/ab8d23
- A.J. Kemp, A.I. Karakas, A.R. Casey, B. Côté, R.G. Izzard, Z. Osborn, ApJ933(2), L30 (2022). DOI 10.3847/2041-8213/ac7c72
- N.N. Chugai, A.D. Kudryashov, Astronomy Letters 46(2), 92 (2020). DOI 10.1134/S1063773720020036
- S.N. Shore, I. De Gennaro Aquino, A&A639, L12 (2020). DOI 10.1051/0004-6361/202038599
- L. Izzo, P. Molaro, G. Cescutti, E. Aydi, P. Selvelli, E. Harvey, A. Agnello, P. Bonifacio, M. Della Valle, E. Guido, M. Hernanz, MNRAS510(4), 5302 (2022). DOI 10.1093/mnras/stab3761
- J.C. Howk, N. Lehner, B.D. Fields, G.J. Mathews, Nature489, 121 (2012). DOI 10.1038/nature11407
- M. Genoni, M. Landoni, G. Cupani, M. Franchini, R. Cirami, A. Zanutta, C. Evans, P. Di Marcantonio, S. Cristiani, A. Trost, S. Zorba, Experimental Astronomy (2022). DOI 10.1007/s10686-022-09844-x
- 32. J. Schachter, PASP103, 457 (1991). DOI 10.1086/132842
- 33. R. Williams, E. Mason, M. Della Valle, A. Ederoclite, ApJ685(1), 451 (2008). DOI 10.1086/590056
- 34. K. Mukai, J.L. Sokoloski, Physics Today 72(11), 38 (2019). DOI 10.1063/PT.3.4341
- A. Franckowiak, P. Jean, M. Wood, C.C. Cheung, S. Buson, A&A609, A120 (2018). DOI 10.1051/0004-6361/201731516

- E. Aydi, K.V. Sokolovsky, L. Chomiuk, E. Steinberg, K.L. Li, I. Vurm, B.D. Metzger, J. Strader, K. Mukai, O. Pejcha, K.J. Shen, G.A. Wade, R. Kuschnig, A.F.J. Moffat, H. Pablo, A. Pigulski, A. Popowicz, W. Weiss, K. Zwintz, L. Izzo, K.R. Pollard, G. Handler, S.D. Ryder, M.D. Filipović, R.Z.E. Alsaberi, P. Manojlović, R. Lopes de Oliveira, F.M. Walter, P.J. Vallely, D.A.H. Buckley, M.J.I. Brown, E.J. Harvey, A. Kawash, A. Kniazev, C.S. Kochanek, J. Linford, J. Mikolajewska, P. Molaro, M. Orio, K.L. Page, B.J. Shappee, J.L. Sokoloski, Nature Astronomy 4, 776 (2020). DOI 10.1038/s41550-020-1070-y
- E. Aydi, L. Chomiuk, L. Izzo, E.J. Harvey, J. Leahy-McGregor, J. Strader, D.A.H. Buckley, K.V. Sokolovsky, A. Kawash, C.S. Kochanek, J.D. Linford, B.D. Metzger, K. Mukai, M. Orio, B.J. Shappee, L. Shishkovsky, E. Steinberg, S.J. Swihart, J.L. Sokoloski, F.M. Walter, P.A. Woudt, ApJ905(1), 62 (2020). DOI 10.3847/1538-4357/abc3bb