

# Consolidating the radiative transfer code to analyze VenSpec-H measurements

Séverine Robert, Paul Simon, Justin Erwin, Filip Vanhellemont, Emmanuel Marcq, Bruno Bézard, Jörn Helbert, Ann Carine Vandaele

### ▶ To cite this version:

Séverine Robert, Paul Simon, Justin Erwin, Filip Vanhellemont, Emmanuel Marcq, et al.. Consolidating the radiative transfer code to analyze VenSpec-H measurements. Europlanet Science Congress 2022, Sep 2022, Granada, Spain. 10.5194/epsc2022-899. insu-03850394

### HAL Id: insu-03850394 https://insu.hal.science/insu-03850394

Submitted on 13 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



EPSC Abstracts

Vol. 16, EPSC2022-899, 2022

https://doi.org/10.5194/epsc2022-899

Europlanet Science Congress 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



## Consolidating the radiative transfer code to analyze VenSpec-H measurements

**Séverine Robert**<sup>1</sup>, Paul Simon<sup>1,2</sup>, Justin Erwin<sup>1</sup>, Filip Vanhellemont<sup>1</sup>, Emmanuel Marcq<sup>3</sup>, Bruno Bézard<sup>4</sup>, Jörn Helbert<sup>5</sup>, and Ann Carine Vandaele<sup>1</sup>

<sup>1</sup>Royal Belgian Institute for Space Aeronomy, Brussels, Belgium (severine.robert@aeronomie.be)

The VenSpec-H instrument is part of the EnVision payload. EnVision is a medium class mission to determine the nature and current state of geological activity on Venus, and its relationship with the atmosphere, to understand how Venus and Earth could have evolved so differently. The launch of the spacecraft is planned in 2031 for a 4 years mission duration.

VenSpec-H is part of the VenSpec suite [1], including also an IR mapper and a UV spectrometer [2] suite. The science objectives of this suite are to search for temporal variations in surface temperatures and tropospheric concentrations of volcanically emitted gases, indicative of volcanic eruptions; and study surface-atmosphere interactions and weathering by mapping surface emissivity and tropospheric gas abundances. Recent and perhaps ongoing volcanic activity has been inferred in data from both Venus Express and Magellan. Maintenance of the clouds requires a constant input of  $H_2O$  and  $SO_2$ . A large eruption would locally alter the composition by increasing abundances of  $H_2O$ ,  $SO_2$  and CO and perhaps decreasing D/H ratio. Observations of changes in lower atmospheric  $SO_2$ , CO and CO and perhaps decreasing D/H ratio. Observation, and mesospheric CO, are therefore required to link specific volcanic events with past and ongoing observations of the variable and dynamic mesosphere, to understand both the importance of volatiles in volcanic activity on Venus and their effect on cloud maintenance and dynamics.

To contribute to this investigation, VenSpec-H is designed to measure  $H_2O$ , HDO, CO, OCS and  $SO_2$  both on the nightside and on the dayside. To ensure the reliability of the analysis, the BIRA-IASB radiative transfer code, ASIMUT-ALVL [3], has been scrutinized making sure all contributions were properly modelled. The radiances of the nightside atmosphere of Venus originate from the thermal emission of the surface and atmosphere while on the dayside they originate from the sunlight penetrating the atmosphere and bouncing back on the cloud cover.

While VenSpec-H will be able to measure 4 spectral ranges between 1 to 2.5 microns, we focused on the upper wavelengths' range, from 2.35 to 2.5 microns, identified as Band#2 in VenSpec-H's design and enabling simultaneous measurements of the different molecular species. A sensitivity study was performed on nightside spectra. The impacts of the molecular species (line by line and collision induced absorption) and of the aerosols were quantified. We will discuss the choices we

<sup>&</sup>lt;sup>2</sup>Université Paris-Saclay, Paris, France

<sup>&</sup>lt;sup>3</sup>LATMOS/Univ. de Versailles Saint-Quentin-en-Yvelines, Paris, France

<sup>&</sup>lt;sup>4</sup>LESIA, Observatoire de Paris, Université PSL, Sorbonne Université, Université Paris Cité, CNRS, France

<sup>&</sup>lt;sup>5</sup>Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR), German Aerospace Center, Institute for Planetary Research, Germany

have made in terms of absorption line parameters and clouds' physical properties

### Acknowledgements

This work has been performed with the support of the Belgian Science Policy Office (BELSPO), with the financial and contractual coordination by the ESA Prodex Office.

#### References

- [1] J. Helbert et al., "The VenSpec suite on the ESA EnVision mission to Venus", *Proc. SPIE 11128, Infrared Remote Sensing and Instrumentation XXVII*, 1112804 (2019).
- [2] E. Marcq et al., "Instrumental requirements for the study of Venus' cloud top using the UV imaging spectrometer VeSUV", *Advances in Space Research*, 68 (2021) 275-291.
- [3] A.C. Vandaele, M. Kruglanski and M. De Mazière, "Modeling and retrieval of atmospheric spectra using ASIMUT", *Proc. of the First 'Atmospheric Science Conference'*, ESRIN, Frascati, Italy, 2006.