



HAL
open science

Two new chrysidoid wasps (Hymenoptera: Bethyliidae, Chrysididae) from mid-Miocene Zhangpu amber

Manuel Brazidec, Vincent Perrichot

► To cite this version:

Manuel Brazidec, Vincent Perrichot. Two new chrysidoid wasps (Hymenoptera: Bethyliidae, Chrysididae) from mid-Miocene Zhangpu amber. *Palaeoworld*, 2023, 32 (4), pp.699-708. 10.1016/j.palwor.2022.11.003 . insu-03862285

HAL Id: insu-03862285

<https://insu.hal.science/insu-03862285>

Submitted on 21 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proofs

Two new chrysidoid wasps (Hymenoptera: Bethyliidae, Chrysididae) from mid-Miocene Zhangpu amber

Manuel Brazidec, Vincent Perrichot

PII: S1871-174X(22)00092-0
DOI: <https://doi.org/10.1016/j.palwor.2022.11.003>
Reference: PALWOR 725


To appear in: *Palaeoworld*

Received Date: 3 October 2022
Revised Date: 2 November 2022
Accepted Date: 11 November 2022

Please cite this article as: M. Brazidec, V. Perrichot, Two new chrysidoid wasps (Hymenoptera: Bethyliidae, Chrysididae) from mid-Miocene Zhangpu amber, *Palaeoworld* (2022), doi: <https://doi.org/10.1016/j.palwor.2022.11.003>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Elsevier B.V. and Nanjing Institute of Geology and Palaeontology, CAS.



Two new chrysidoid wasps (Hymenoptera: Bethylidae, Chrysididae) from mid-Miocene Zhangpu amber

Manuel Brazidec^{1,2*}, Vincent Perrichot¹

¹Univ Rennes, CNRS, Géosciences Rennes, UMR 6118, 35000 Rennes, France.

² State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology and Center for Excellence in Life and Palaeoenvironment, Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008, China.

*corresponding author

E-mail addresses: manuel.brazidec@gmail.com (M. Brazidec),
vincent.perrichot@univ-rennes1.fr (V. Perrichot)

Abstract

The Miocene fossil record of Chryridoidea is particularly scarce despite numbers of insect-bearing deposits available. This record is here summarized, highlighting the overwhelming proportion of fossils from Dominican Republic amber while other deposits, e.g., Zhangpu, Mexican or Ethiopian ambers, seem promising for the study of the group but suffer from the lack of interest for this period. We describe and figure two new species from Zhangpu, China, a rich amber biota that formed in the Asian rainforest during the Mid-Miocene Climatic Optimum (MMCO): *Sierola colomboi* n. sp. (Bethylidae, the second from Zhangpu amber) and *Hedychridium rosai* n. sp. (Chrysididae). *Sierola* currently has a circum-pacific distribution with two fossils from European Eocene deposits. Four species have been recorded from China but *Sierola colomboi* n. sp. is the first fossil species from the country. *Hedychridium* is a cosmopolitan genus, the second largest within its family and *Hedychridium rosai* n. sp. represents its first known fossil species.

Two new chrysidoid wasps (Hymenoptera: Bethylidae, Chrysididae) from mid-Miocene Zhangpu amber

Manuel Brazidec^{a,b*}, Vincent Perrichot^a

^a Univ. Rennes, CNRS, Géosciences Rennes, UMR 6118, 35000 Rennes, France

^b State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology and Center for Excellence in Life and Paleoenvironment, Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008, China

* Corresponding author.

E-mail addresses: manuel.brazidec@gmail.com (M. Brazidec), vincent.perrichot@univ-rennes1.fr (V. Perrichot)

Abstract

The Miocene fossil record of Chrysidoidea is particularly scarce despite numbers of insect-bearing deposits available. Their record is here summarized, highlighting the overwhelming proportion of fossils from Dominican Republic amber while other deposits, e.g., Zhangpu, Mexican or Ethiopian ambers, seem promising for the study of the group but suffer from the insufficient interest in this period. We describe and figure two new species from Zhangpu, China, a rich amber biota that formed in the Asian rainforest during the Mid-Miocene Climatic Optimum (MMCO): *Sierola colomboi* n. sp. (Bethylinidae, the second from Zhangpu amber) and *Hedychridium rosai* n. sp. (Chrysididae). *Sierola* currently has a circum-pacific distribution with two fossils from European Eocene deposits. Four species have been recorded from China but *Sierola colomboi* n. sp. is the first fossil species from the country. *Hedychridium* is a cosmopolitan genus, the second largest within its family and *Hedychridium rosai* n. sp. represents its first known fossil species.

Keywords: Chrysidoidea; Bethylinidae; Chrysidinae; Cenozoic; Miocene; China

1. Introduction

The Chrysidoidea are the basalmost superfamily within stinging Hymenoptera (Aculeata). They are composed of six extant and two extinct families, their oldest representative dating back to the Lower Cretaceous (early Barremian) of the Isle of Wight, UK (Perkovsky et al., 2020a). Even with disparities between each family, Chrysidoidea are not infrequent in the fossil record, with about 270 formally named species (Martynova et al., 2019, and later summaries of Brazidec and Perrichot, in press for Bethylinidae; Lucena and Almeida, 2021 for Chrysididae; Martynova et al., 2020 for Dryinidae; Perkovsky et al., 2020a for Embolemidae; Perkovsky et al., 2020b for Sclerogibbidae; Lepeco and Melo, 2022 for Scolebythidae; and Rasnitsyn and Brothers, 2020 for Plumalexiidae). The extant Bethylinidae, Chrysididae and

Dryinidae are much more speciose than the rest of chrysidoid families and are more commonly encountered in the fossil record.

The Bethylidae, dubbed ‘flat wasps’, are the most diverse of all chrysidoid families, with 3000 species described (Azevedo et al., 2018). The bethylids for which biology is known are external parasitoids of lepidopteran and coleopteran larvae in cryptic habitats (Evans, 1964; Rubink and Evans, 1979). Nine subfamilies are recognized within the Bethylidae (Colombo et al., 2020), four of them being exclusively fossil.

The 2500 valid species of Chrysididae display a greater diversity in temperate desert but are distributed worldwide (Finnamore and Brothers, 1993). Chrysidids are sometimes referred to as ‘gold wasps’ because of the predominantly metallic coloration of the body, which makes them highly appreciated by collectors, or ‘cuckoo wasps’ because of the cleptoparasitic habits of some species (Kimsey and Bohart, 1991). The current consensus on the chrysidid taxonomy acknowledges four main subfamilies: the Cleptinae, parasitic on prepupal larvae of sawfly wasps; the Amiseginae and Loboscellidinae, which oviposit into previously nipped stick-insects (Phasmatodea) eggs; and the Chrysidinae, the true cuckoo wasps, specialized cleptoparasites attacking aculeate wasp nests (Kimsey and Bohart, 1991; Lucena and Almeida, 2021).

In this contribution, one new species of Bethylidae and another of Chrysididae are described from the recently-discovered amber deposit of Zhangpu, in Fujian Province, China, where paleobiota is for now largely unknown.

2. Material and methods

Zhangpu amber is found with plant impressions in two layers of sandy mudstone interbedded with coal seams that belong to the Fotan Group, a geological unit that occurs widely in Zhangpu County, Fujian Province, southeastern China (Wang, B. et al., 2021, fig. 1). Under- and overlying basalt layers allow constrained dating of the amber between 14.8 ± 0.6 Ma and 14.7 ± 0.4 Ma, which corresponds to the middle Miocene (Langhian; Zheng et al., 2019).

This study is based on two individuals in separate pieces kindly lent by Prof. Bo Wang (Nanjing, China) to the authors for study, and housed in the Nanjing Institute of Geology and Palaeontology (NIGP), Chinese Academy of Sciences, Nanjing, China. The fragile amber pieces were embedded in a block of epoxy resin (Araldite® 2020) for consolidation and posteriorly polished to facilitate the observation of the specimens, using thin silicon carbide papers on a grinder polisher (Buehler MetaServ 3000). The examination and photographs

were conducted with a Leica DMC4500 camera attached to a Leica M205C stereomicroscope or a Canon EOS 250D camera attached to a Leica M80 stereomicroscope. All images are digitally stacked photomicrographic composites of several focal planes, which were obtained using Helicon Focus 6.7 software. Adobe Illustrator CC2019 and Photoshop CC2019 software were used to compose the figures and ImageJ 1.53 for measurements. The description of the surface sculpturing follows the nomenclature of Harris (1979); bethylid and chrysidid morphological characters respectively follow the nomenclatures of Lanes et al. (2020) and Kimsey and Bohart (1991). Main measurements and indices used are as follows: the maximum length of fore wing from the apex of the axillary sclerite to the wing apex (LFW); the length of the head capsule excluding the mandibles, measured from posteriormost point of vertex to anteriormost point of clypeus (LH); the width of the head measured immediately behind the eyes in dorsal view (WH); the width of frons measured in dorsal view (WF); the maximum width of eye in lateral view (HE); ocello-ocular line, the minimum length from the posterior ocellus to the ocular margin (OOL); the maximum width of the ocellar triangle, measured in dorsal view (WOT); the maximum diameter of anterior ocellus (DAO); vertex-ocular line, the minimum length from the vertexal margin to posteriormost eye margin, measured in lateral view (VOL).

This published work and its new nomenclatural acts are registered in ZooBank with the following LSID (reference): [urn:lsid:zoobank.org:pub:BB079D89-C9FE-4F27-AF99-E94506E891E5](https://zoobank.org/pub:BB079D89-C9FE-4F27-AF99-E94506E891E5).

3. Systematic palaeontology

Superfamily Chrysoidea Latreille, 1802

Family Bethyloidea Haliday, 1839

Subfamily Bethyloinae Haliday, 1839

Genus *Sierola* Cameron, 1881

Sierola colomboi n. sp.

(Figs. 1, 2)

LSID: [urn:lsid:zoobank.org:act:2786F41D-6B8E-440D-A7EE-8CDBF3FECAE0](https://zoobank.org/act:2786F41D-6B8E-440D-A7EE-8CDBF3FECAE0).

Etymology: We dedicate this species to Dr. Wesley D. Colombo, specialist on living and fossil Bethyloidea. The specific epithet is to be treated as a noun in a genitive case.

Material: Holotype, NIGP200740, a complete male with some features hidden by small bubbles.

Diagnosis: Body stout (Fig. 2A); head globose; frons coriaceous with sparse long setae; median clypeal lobe obtuse triangular; mandibles large, ventral margin outcurved; posterior ocelli very close to occiput; dorsal pronotal area 2.3 times wider than long, transverse pronotal carina raised, humeral angles sharp; anteromesoscutum 0.78 length of dorsal pronotal area (Fig. 2B); fore wing uniformly micropubescent (Fig. 1); [C], [R], [1Cu], [1M] and [2R1] cells closed by tubular veins (Fig. 2D); meso- and metafemora thick (Fig. 2A, C).

Type locality: Zhangpu County, Zhangzhou Prefecture, Fujian Province, China.

Horizon: Sedimentary layer II, Fotan Group, middle Miocene: Langhian, 14.7 Ma.

Description: Body not especially flattened (length 1.90 mm). Head prognathous, globular, slightly longer than wide; LH: 0.42 mm, WH: 0.43 mm; WF: 0.26 mm; HE: 0.23 mm; OOL: 0.16 mm; WOT: 0.06 mm; DAO: 0.04 mm; VOL: ca. 0.12 mm; frons slightly convex, bearing sparse long setae, longer than flagellomere width; compound eyes almond-shaped, longer than high, glabrous; frons and vertex finely coriaceous; clypeus short with median lobe obtuse triangular, clypeal carina extending on frons for short distance; mandibles large, ventral margin outcurved, four short teeth along masticatory margin; antenna short, not extending beyond pronotum; scape 3.6 times longer than pedicel (length 0.18 mm); pedicel longer than flagellomere 1 (length 0.05 mm versus 0.03 mm); 11 flagellomeres hardly longer than wide, cylindrical; vertex crest rounded; ocellar triangle compact, posteriad on head, posterior ocelli distant from occiput by less than their diameter; occipital carina absent.

Mesosoma shorter than metasoma (length 0.66 mm), dorsum sparsely pubescent; dorsal pronotal area wider than long (length 0.14 mm; width 0.32 mm), transverse pronotal carina protruding, forming sharp humeral angles, posterior margin straight medially; anteromesoscutum 0.78 length of dorsal pronotal area; notauli absent; parapsidal signum present; mesopleuron not prominent, barely visible dorsally; mesoscutellum with posterior margin blunted, not contacting propodeum; metanotum reduced; dorsal area of metapectal-propodeal complex coriaceous, bordered by marginal carinae; median metapostnotal and metapostnotal-propodeal suture absent; propodeal declivity without carina. Fore wing hyaline and without particular patterns of micropubescent (LFW: 1.53 mm); C, Sc+R, M+Cu, A, Rs&M and cu-a tubular; Rs&M curved; Rs+M tubular; prestigma triangular; pterostigma rounded; 2r-rs&Rs contiguous, widely arched toward anterior wing margin distally and reaching it; [C], [R], [1Cu], [1M] (areolet) and [2R1] cells closed by tubular veins; [2R1] cell

wider distally than basally. Legs slightly pubescent; meso- and metafemora thick; tarsal claws angled.

Metasoma fusiform (length 0.82 mm), smooth.

Family Chrysididae Latreille, 1802

Subfamily Chrysidinae Latreille, 1802

Tribe Elampini Dahlbom, 1854

Genus *Hedychridium* Abeille de Perrin, 1878

Hedychridium rosai n. sp.

(Figs. 3–5)

LSID: urn:lsid:zoobank.org:act:9589AF1B-BE56-4DC3-B69E-FB7195148F65.

Etymology: We dedicate this species to Dr. Paolo Rosa, specialist on Chrysididae. The specific epithet is to be treated as a noun in a genitive case.

Material: Holotype, NIGP200741, a complete female.

Diagnosis: Body metallic gold except metallic black mesoscutum, mesoscutellum, metanotum, propodeum and flagellomeres 4–10 (Fig. 3); vertex and frons punctate; scapal basin flat, not depressed, with minute, continuous and parallel striations (Fig. 5B); scape 5.5 times longer than pedicel; first flagellomere 4 times longer than second flagellomere (Fig. 4B); compound eyes with sparse setae (Fig. 5A); pronotal collar distinct and two-thirds as long as rest of pronotum, swollen, foveolate; pronotum with median sulcus; propodeum areolate (Figs. 3, 4C, 5C); metafemur enlarged dorsoventrally (Fig. 4D); fore wing cu-a postfurcad M (Fig. 5D).

Type locality: Zhangpu County, Zhangzhou Prefecture, Fujian Province, China.

Horizon: Sedimentary layer II, Fotan Group, middle Miocene: Langhian, 14.7 Ma.

Description: Body length 3.20 mm; integument shining; body mostly metallic gold except mesoscutum, mesoscutellum, metanotum, propodeum and flagellomeres 4–10, black with metallic reflections.

Head as long as wide (length and width 0.64 mm), hypognathous; vertex and frons punctate; vertex with a few long setae, without median ridge; ocellar triangle compact, on top of head; scapal basin flat, not depressed, with minute, continuous and parallel striations; clypeus short and wide; scape elongate, 5.5 times longer than pedicel (length 0.33 mm); pedicel shorter than flagellomeres (length 0.06 mm); 11 flagellomeres; flagellomere 1

elongate, 4 times longer than following flagellomeres (length 0.28 mm versus 0.07 mm); antennae bearing numerous short sensilla, shorter than one third the length of the apical flagellomere; compound eye large, covering most of head length, bulging, with some sparse setae, inner ocular margin rather diverging downward; malar space shorter than one third of eye length, convergent.

Mesosoma shorter than metasoma (length 1.25 mm); pronotal collar swollen, easily distinguishable from pronotum, two-thirds as long as rest of pronotum (dorsal length 0.21 mm), minutely foveolate, posteriorly delimited by transverse groove; dorsal area of pronotum box-like, punctate with long setae along lateral margins, (length 0.32 mm), divided by thin median sulcus, lateral lobe nearly contacting tegula but not reaching it; mesoscutum punctate (length 0.33 mm), notauli present and deep; mesopleuron punctate, rounded, scrobal sulcus and omaulus indistinct; propodeum short, dorsal surface areolate. Legs bearing erect long setae; protibial spur long and curved; probasitarsomere basally curved; metacoxa elongate; metafemur enlarged dorsoventrally; tarsal claws curved, with one small subsidiary tooth; arolium present. Fore wing hyaline, uniformly micropubescent (length 1.31 mm); C, Sc+R, M+Cu, A, M, cu-a and Rs tubular; cu-a slightly postfurcad M; Cu nebulous; Rs+M spectral; Rs curved toward wing margin, tubular on half length of marginal cell; pterostigma poorly defined, shorter than M; R1 as long as pterostigma. Hind wing without visible venation; 4 hamuli.

Metasoma smooth (length 1.31 mm); three external segments visible; sternum concave; tergite 2 longest; tergite 3 without apical row or pits or posterior denticulate margin; ovipositor exerted, curved and well-sclerotized (length 0.52 mm).

4. Discussion

The Bethyridae specimen can confidently be identified as a Bethylinae, given the presence of numerous features reminiscent of the subfamily: the clypeal carina extending on frons, the tarsal claws angled, the fore wing with Rs+M tubular and Rs&M angled. Following the key to the genera of Bethylinae by Azevedo et al. (2018), the specimen keys in *Sierola* Cameron, 1881 for the following characters: antennae with 11 flagellomeres, dorsal pronotal area regularly long, posterior margin outcurved medially, fore wing with five closed cells, metapectal-propodeal complex without metapostnotal-propodeal suture and mesopleuron not prominent, fore wing with [2R1] cell closed. The absence of notauli and median propodeal carina, the [1M] cell closed and the [2R1] cell ‘triangular’ (distalmost part of the cell wider than basalmost part) are additional diagnostic characters of *Sierola*, which strengthen the

attribution of the fossil to this genus. However, *Sierola colomboi* n. sp. also displays some unusual features for a Bethylinae. The anterior part of the dorsal pronotal area is bordered by a raised carina, known as the transverse pronotal carina, producing the anterior pronotal corners into sharp humeral angles. This configuration is similar to what is known in the epyrine genus *Anisepyris* Kieffer, 1905 in Kieffer and Marshall, 1904–1906 (Barbosa and Azevedo, 2018, fig. 1J, K) while *Sierola* usually has more rounded corners (Wang, C. et al., 2021, figs. 1D, 2F, 3F). The presence of the transverse pronotal carina is a novel pattern for *Sierola* although some species possesses more or less produced anterior pronotal corners. Nevertheless, it does not question the generic position of the fossil.

Sierola is the largest genus within its subfamily, with 249 species, but no less than 232 have been described from Hawaiian Islands (Fullaway, 1920; Azevedo et al., 2018; Magnacca, 2020). It has a circum-pacific distribution, with species recorded from Marques Islands, Fiji (Fullaway, 1920), California (Evans, 1978), Australia (Gordh, 1998), Thailand, India (Terayama, 2004) and Japan (Terayama, 2006). Gordh (1998) assumed that *Sierola* originates from continental Asia, before invading Hawaiian Islands, and rapidly radiated in a few million years while Magnacca (2020) proposed an Australian origin of the genus. Four species are known from China (Zhejiang Province and Macao, Wang, C. et al., 2021). Despite being apparently endemic of the circum-Pacific belt, two species have been described from the Upper Eocene Baltic and Rovno ambers: *Sierola hastata* Sorg, 1988 and *Sierola rovniana* Ramos and Azevedo in Ramos et al., 2014. This pattern of regression toward tropical latitudes of Australasia (maybe correlated to Cenozoic global climate cooling) is also found in the extant Australian *Eupsenella* Westwood, 1874 (Ramos and Azevedo, 2012; Ramos et al., 2014). *Sierola colomboi* n. sp. differs from the two other fossil species as follows: clypeus obtuse rather than angulate, notauli absent, [R], [1Cu] and [1M] cells formed by tubular veins rather than nebulous (*S. rovniana*; Ramos et al., 2014); [1M] cell pentagonal rather than hexagonal; distal segment of 2r-rs&Rs strongly angulate rather than gently bent toward anterior wing margin (*S. hastata*; Sorg, 1988).

Sierola species for which the biology is known mainly parasitize microlepidopteran caterpillars (e.g., Gelechiidae, Batrachedridae, Tortricidae, Gracillariidae, Magnacca, 2020 and references therein), which is considered to be the ground plan for the genus. Some species would also present unusual habits such as attacking insects in non-lepidopteran-induced galls (Ashmead, 1901; Santosh, 2017) or beetle larvae (Swezey, 1954; Magnacca, 2020) but these records are yet to be confirmed. The extant *Sierola mawarajo* Terayama, 2004 was reared in the tropical rainforest of the Khao Yai National Park (Thailand), where vegetation is

dominated by members of the Dipterocarpaceae (Smitinand, 1968). Today's southeast Asian dipterocarp forests are the closest extant relatives of the mid-Miocene Zhangpu amber forest in terms of floral assemblage and climate (Wang, B. et al., 2021, fig. 2). Furthermore, the lepidopteran families mentioned above occur widely in these environments (e.g., Ujiye et al., 1996; Park and Ponomarenko, 1999, 2004; Brown et al., 2019). Therefore, both for biogeographical and biological reasons, it is not surprising to discover *Sierola* in the Zhangpu biome, simultaneously with the megathermal climate of the Mid-Miocene Climatic Optimum that allowed the tropical forest to expand toward higher latitudes (Kasbohm and Schoene, 2018).

Following the key to the subfamilies of Chrysididae by Kimsey and Bohart (1991), the chrysidid specimen keys in the Chrysidinae for the gaster with three external segments, the sternum rather concave and the propodeum short dorsally. Chrysidinae are further divided in four tribes: the Allocoeliini, Parnopini, Chrysidini and Elampini. The specimen keys in the latter tribe due to the gaster with three terga, the body partly metallic, the tegula not covering both fore and hind wing bases and the tergite 3 without pit row. Following the key to the genera by Elampini of Kimsey and Bohart (1991), it keys in *Hedychridium* Abeille de Perrin, 1878 because of the following characters: hind tarsal claws with one tooth, scutellum anteriorly without tubercles, fore wing Rs more than half as long as M, face flat with at least a narrow zone of fine cross-ridging in scapal basin. *Hedychridium rosai* n. sp. is easily recognizable within the genus for the hypertrophied pronotal collar, which differs from the rest of the pronotum by its sculpture.

Hedychridium is the second largest genus of the Chrysididae and the largest within its tribe with more than 300 species described (Rosa et al., 2022). This genus occurs in all zoogeographic regions except the Australia, its diversity being highest in arid parts of the Holarctic and southern Africa (Kimsey and Bohart, 1991). Seven species are known in China, exclusively from the Palearctic part of the country: Xinjiang, Inner Mongolia, Gansu, Shanxi, Hebei and Heilongjiang regions (Rosa et al., 2014). Chrysididae are scarcer than Bethyridae or Dryinidae in the fossil record (Martynova et al., 2019), for current equivalent diversity. Thus, even if *Hedychridium* contains numerous species, *Hedychridium rosai* n. sp. is its first fossil species reported to date.

Little is known about the *Hedychridium* biology, the few records suggesting a parasitic behavior on ground-nesting bees and Sphecidae (e.g., Mocsáry, 1889). *Hedychridium monochroum* Buysson, 1888 is the only representative of the genus in southeast Asia, from Lamphun, North Thailand (Tsuneki, 1961), where the vegetation is partly composed of

dipterocarp forests (Maxwell et al., 1995). Additionally, this species has been reported to lay eggs into nests of Crabronidae (Martynova, 2017). It is not possible to evaluate the possible hosts for *Hedychridium rosai* n. sp., as no aculeate wasps other than Bethylidae, ants (Perrichot, in press) and stingless bees (Engel et al., 2021) have been described from this deposit. However, similar to *Sierola colomboi* n. sp., the presence of *Hedychridium* in the Zhangpu amber does not denote with what is known of the tropical palaeoenvironment.

The Miocene Chrysoidea are poorly known, possibly because of the increasing interest for Cretaceous material that has become much more accessible in the past few years. *Hedychridium rosai* n. sp. is only the second formally described Chrysididae from this period, after *Ceratochrysis dominica* Engel, 2006 and the second unambiguous fossil species belonging to an extant genus. A few non-amber fossils have been attributed to *Chrysis* Linnaeus, 1761 but without enough characters preserved to ensure their generic attribution (e.g., Förster, 1891; Rohwer, 1909). The Bethylidae are more common, with eight species previously named, *Sierola colomboi* n. sp. being the ninth one. Thanks to the work of Olmi (1989, 1995, 2005a), the Dryinidae received more consideration, with 17 species reported in amber. Finally, the less diverse Sclerogibbidae and Scolebythidae are both known by two species while only two indetermined Embolemidae are mentioned in the literature (Table 1). However, the vast majority of the chrysidoid wasps from this period have been described from Dominican Republic. This is not surprising, considering that Dominican amber has drawn attention for at least six decades (Sanderson and Farr, 1960) but it is much less than in Burmese amber, where renewed interest is more recent (e.g., Olmi et al., 2014; Melo and Lucena, 2019; Martynova et al., 2020). Several fossiliferous sites worldwide, notably Zhangpu, Mexican, Ethiopian or New Zealand ambers, could enrich the knowledge on Miocene Chrysoidea (Solórzano Kraemer et al., 2015; Schmidt et al., 2018; Bouju and Perrichot, 2020; Wang, B. et al., 2021). With more than 25000 inclusions identified in Zhangpu amber, this deposit is very likely to reveal some additional chrysidoid wasps in the upcoming years.

Acknowledgments

We warmly thank the guest editor Prof. Bo Wang for inviting us to contribute this special issue of Palaeoworld. We are grateful to reviewers, Profs. Celso O. Azevedo and Alexandr P. Rasnitsyn, for their valuable comments on the manuscript. Research on the Zhangpu amber was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB26000000) and National Natural Science Foundation of China

(42125201, 41688103). This work is part of Manuel Brazidec's Ph.D. project CHRYSIS: "The role of greenhouses on the diversification and evolution of chrysidoid wasps".

References

- Abeille de Perrin, E., 1878. Diagnoses de chrysidides nouvelles. Published by the author, Marseille, 6 pp.
- Ashmead, W.H., 1901. Notes on some New Zealand and Australian parasitic Hymenoptera, with descriptions of new genera and new species. *Proceedings of the Linnean Society of New South Wales* 25, 327–360.
- Azevedo, C.O., Alencar, I.D.C.C, Ramos, M.S., Barbosa, D.N., Colombo, W.D., Vargas, J.M., Lim, J., 2018. Global guide of the flat wasps (Hymenoptera, Bethylidae). *Zootaxa* 4489, 1–294.
- Barbosa, D.N., Azevedo, C.O., 2018. Revision of *Anisepyris* Kieffer (Hymenoptera, Bethylidae), with description of 135 new species. *Zootaxa* 4416, 1–258.
- Bouju, V., Perrichot, V., 2020. A review of amber and copal occurrences in Africa and their paleontological significance. *BSGF – Earth Science Bulletin* 191, Article Number 17, doi: 10.1051/bsgf/2020018.
- Brazidec, M., Perrichot, V., 2022. The first flat wasps (Hymenoptera: Bethylidae) from Miocene Mexican amber. *Journal of South American Earth Sciences* 116, 103838. 10.1016/j.jsames.2022.103838
- Brazidec, M., Perrichot, V., in press. The first fossil *Parascleroderma* (Hymenoptera: Bethylidae): a new species in mid-Miocene Zhangpu amber. *Palaeoworld*, doi: 10.1016/j.palwor.2022.01.009.
- Brown, J.W., Basset, Y., Panmeng, M., Putnau, S., Miller, S.E., 2019. Host records for Tortricidae (Lepidoptera) reared from seeds and fruits in a Thailand rainforest. *Proceedings of the Entomological Society of Washington* 121, 544–556.
- Buysson, R., 1888. Descriptions de Chrysidides nouvelles. *Revue d'Entomology* 6, 167–201.
- Cameron, P., 1881. Notes on Hymenoptera, with descriptions of new species. *Transactions of the Royal Entomological Society of London* 1881, 555–577.
- Colombo, W.D., Perkovsky, E.E., Azevedo, C.O., 2020. Phylogenetic overview of flat wasps (Hymenoptera, Bethylidae) reveals Elektroepyrinae, a new fossil subfamily. *Palaeoentomology* 3, 296–283.

- Colombo, W.D., Perkovsky, E.E., Azevedo, C.O., 2022. Synopsis of the rare genus *Alongatepyris* (Hymenoptera, Bethylidae), with description of the first extinct species from Dominican amber. *Palaeoentomology* 5, 57–65.
- Currado, I., Olmi, M., 1983. Primo reperto di Driinide fossile in ambra della Repubblica Dominicana (Hymenoptera, Dryinidae). *Bollettino del Museo Regionale di Scienze Naturali Torino* 1, 329–334
- Dahlbom, A.G., 1854. *Hymenoptera Europea praecipue borealia etc.* Vol. 2, xxiii. Lundbergiana, Lund, 411 pp.
- Engel, M.S., 2006. A new cuckoo wasp of the genus *Ceratochrysis* in amber from the Dominican Republic (Hymenoptera: Chrysididae). *Polskie Pismo Entomologiczne* 75, 499–504.
- Engel, M.S., 2015. The wasp genus *Clytopsenella* in Early Miocene amber from the Dominican Republic (Hymenoptera: Scolebythidae). *Novitates Paleoentomologicae* 12, 1–14.
- Engel, M.S., Herhold, H., Davis, S., Wang, B., Thomas, J., 2021. Stingless bees in Miocene amber of southeastern China (Hymenoptera: Apidae). *Journal of Melittology* 105, 1–83.
- Evans, H.E., 1964. A synopsis of the American Bethylidae (Hymenoptera, Aculeata). *Bulletin of the Museum of Comparative Zoology* 132, 1–222.
- Evans, H.E., 1978. The Bethylidae of America North of Mexico. *Memoirs of the American Entomologist Institute* 27, 1–332.
- Finnamore, A.T., Brothers, D.J., 1993. Superfamily Chryridoidea. In: Goulet, H., Huber, J.T. (Eds.), *Hymenoptera of the World: An Identification Guide to Families*. Research Branch Agriculture Canada Publication, Ottawa, pp. 130–160.
- Förster, B., 1891. Die Insekten der “Plattigen Steinmergels” von Brunstatt. *Abhandlungen zur Geologischen Specialkarte von Elsass-Lothringen*, Band 3, 1, 335–593.
- Fullaway, D.T., 1920. New species of *Sierola* with explanatory notes. *Occasional Papers of the Bernice Pauahi Museum of Polynesian Ethnology and Natural History* 7, 57–159.
- Gordh, G., 1998. A new species of *Sierola* parasitic on moth larvae in Western Australia (Hymenoptera: Bethylidae). *Proceedings of the Hawaiian Entomological Society* 33, 83–88.
- Haliday, A.H., 1839. *Hymenoptera Britannica*, Part 1. Oxyura. Addendum to Vol. 1. *Hymenopterorum synopsis ad methodum fallenii ut plurimum accomodata*. Hippolytas Bailliere, London, 28 pp.

- Harris, R.A., 1979. A glossary of surface sculpturing. *Occasional Papers in Entomology* 28, 1–31.
- Kasbohm, J., Schoene, B., 2018. Rapid eruption of the Columbia River flood basalt and correlation with the mid-Miocene climate optimum. *Sciences Advances* 4, eaat8223, doi: 10.1126/sciadv.aat8223.
- Kieffer, J.J., Marshall, T.A., 1904–1906. Proctotrypides. In: André, E. (Ed.), *Species des Hyménoptères d'Europe & d'Algérie*. Tome IX. Librairie Scientifique A. Hermann, Paris, pp. 145–288 [publication dates of the different parts: pages 1–64, 1st January 1904; 65–144, 1st April 1905; 145–288, 1st November 1905; 289–368, 31st January 1906; 369–551, 1st October 1906].
- Kimsey, L.S., Bohart, R.M., 1991. *The Chrysidid Wasps of the World*. Oxford University Press, New York, 652 pp.
- Lanes, G.O., Kawada, R., Azevedo, C.O., Brothers, D.J., 2020. Revisited morphology applied for systematics of flat wasps (Hymenoptera, Bethyridae). *Zootaxa* 4752, 1–127.
- Latreille, P.A., 1802. *Histoire naturelle, générale et particulière des Crustacés et des insectes*. Vol. 3. Dufort, Paris, 348 pp.
- Lepeco, A., Melo, G.A.R., 2022. Mid-Cretaceous amber from Myanmar reveals a rich diversity of extinct scolebythid wasps (Hymenoptera: Chrysoidea). *Cretaceous Research* 137, 105232.
- Linnaeus, C., 1761. *Fauna Svecica: Sistens Animalia Sveciae Regni: Mammalia, Aves, Amphibia, Pisces, Insecta, Vermes, Distributa per Classes & Ordines, Genera & Species, cum Differentiis Specierum, Synonymis Auctorum, Nominibus Incolarum, Locis Natalium, Descriptionibus Insectorum*. Sumtu & Literis Direct. Laurentii Salvii, Stockholmiae, 578 pp.
- Lucena, D.A.A., Almeida, E.A.B., 2021. Morphology and Bayesian tip-dating recover deep Cretaceous-age divergences among major chrysidid lineages (Hymenoptera: Chrysididae). *Zoological Journal of the Linnean Society* 194, 36–79.
- Magnacca, K.N., 2020. Review of *Sierola* Cameron (Hymenoptera: Bethyridae) of the Hawaiian Islands. Part I: O'ahu. Bishop Museum Press, Honolulu, 311 pp.
- Martins, A.L., Melo, G.A.R., 2020. Revision of the fossil species of *Thaumatodryinus* Perkins from Dominican amber, with a new combination and description of a new species (Hymenoptera, Dryinidae). *Journal of Hymenoptera Research* 79, 77–88.
- Martynova, K.V., 2017. *Hedychridium monochroum* and *Chrysis lanceolata* (Hymenoptera: Chrysididae) — two species of cuckoo wasps reared from nests of *Solierella compedita*

- (Hymenoptera: Crabronidae) in southeastern Ukraine. *Turkish Journal of Zoology* 41, 397–407.
- Martynova, K.V., Perkovsky, E.E., Olmi, M., Vasilenko, D.V., 2019. New records of Upper Eocene chrysidoid wasps from basins of Styr and Stokhod rivers (Rovno amber). *Paleontological Journal* 53, 998–1023.
- Martynova, K.V., Zhang, Q., Olmi, M., Müller, P., Perkovsky, E.E., 2020. Revision of the genus *Dryinus* Latreille (Hymenoptera: Dryinidae) from mid-Cretaceous Kachin (Myanmar) amber. *Cretaceous Research* 106, 104217.
- Maxwell, J.F., Elliott, S., Palee, P., Anusarnsunthorn, V., 1995. The vegetation of Doi Kuhn Tan National Park, Lamphun-Lampang provinces, Thailand. *Natural History Bulletin of the Siam Society* 43, 185–205.
- Melo, G.A.R., Lucena, D.A.A., 2019. †Chrysobythidae, a new family of chrysidoid wasps from Cretaceous Burmese amber (Hymenoptera, Aculeata). *Historical Biology* 32, 1143–1155.
- Mocsáry, A., 1889. *Monographia Chrysididarum orbis terrarium universi*. Hungarian Academy of Science, Typis Societatis Franklinian, Budapest, 643 pp.
- Olmi, M., 1987. New species of Dryinidae, with description of a new subfamily from Florida and a new species from Dominican amber (Hymenoptera: Chryridoidea). *Bollettino del Museo Regionale di Scienze Naturali Torino* 5, 211–238.
- Olmi, M., 1989. Supplement to the revision of the world Dryinidae (Hymenoptera Chryridoidea). *Frustula Entomologica* 12, 109–395.
- Olmi, M., 1995. Dryinids and Embolemids in amber (Hymenoptera Dryinidae et Embolemidae). *Redia* 78, 253–271.
- Olmi, M., 1998. New oriental and neotropic Dryinidae (Hymenoptera Chryridoidea). *Frustula Entomologica* 20, 152–167.
- Olmi, M., 2005a. A new fossil dryinid in amber from the Dominican Republic: *Harpactosphēcion scheveni* n. sp. (Hymenoptera Dryinidae). *Frustula Entomologica* 26-27, 15–19.
- Olmi, M., 2005b. A revision of the world Sclerogibbidae (Hymenoptera Chryridoidea). *Frustula Entomologica* 26-27, 46–193.
- Olmi, M., Guglielmino, A., 2011. Revision of fossil species of *Dryinus* belonging to *lamellatus* group, with description of a new species (Hymenoptera, Dryinidae). *ZooKeys* 130, 505–514.

- Oلمي, M., Rasnitsyn, A.P., Brothers, D.J., Guglielmino, A., 2014. The first fossil Embolemidae (Hymenoptera: Chrysoidea) from Burmese amber (Myanmar) and Orapa kimberlitic deposits (Botswana) and their phylogenetic significance. *Journal of Systematic Palaeontology* 12, 623–635.
- Park, K.T., Ponomarenko, M.G., 1999. Genus *Hypatima* Hübner and allied genera from Thailand (Lepidoptera, Gelechiidae). *Species Diversity* 4, 321–337.
- Park, K.T., Ponomarenko, M.G., 2004. New species of the genus *Dichomeris* Hübner (Lepidoptera, Gelechiidae) from Thailand. *Transactions of the Lepidopterological Society of Japan* 55, 147–159.
- Peñalver-Mollá, E., 1998. Estudio tafonómico y paleoecológico de los insectos del Mioceno de Rubielos de Mora (Teruel). Instituto de estudios Turolenses, Teruel, 179 pp.
- Perkovsky, E.E., Oلمي, M., Müller, P., Guglielmino, A., Jarzenowski, E.A., Capradossi, L., Rasnitsyn, A.P., 2020a. A review of the fossil Embolemidae (Hymenoptera: Chrysoidea), with description of seven new species and history of the family. *Cretaceous Research* 121, 104708.
- Perkovsky, E.E., Martynova, K.V., Mita, T., Oلمي, M., Zheng, Y., Müller, P., Zhang, Q., Gantier, F., Perrichot, V., 2020b. A golden age for ectoparasitoids of Embioidea: Cretaceous Sclerogibbidae (Hymenoptera, Chrysoidea) from Kachin (Myanmar), Charentes (France) and Choshi (Japan) ambers. *Gondwana Research* 87, 1–22.
- Perrichot, V., in press. The first fossil occurrence of *Odontoponera* (Formicidae: Ponerinae), from Middle Miocene Zhangpu amber. *Palaeoworld*, doi: 10.1016/j.palwor.2022.09.007.
- Prentice, M.A., Poinar Jr., G.O., Milki, R., 1996. Fossil scolybythids (Hymenoptera: Scolybythidae) from Lebanese and Dominican amber. *Proceedings of the Entomological Society of Washington* 98, 802–811.
- Ramos, M.S., Azevedo, C.O., 2012. Revision of *Eupsenella* Westwood, 1874 (Hymenoptera: Bethyloidea). *Zootaxa* 3539, 1–80.
- Ramos, M.S., Perkovsky, E.E., Rasnitsyn, A.P., Azevedo, C.O., 2014. Revision of Bethyloinae fossils from Baltic, Rovno and Oise amber, with comments on the Tertiary fauna of the subfamily. *Neues Jahrbuch für Geologie und Paläontologie* 271, 203–228.
- Rasnitsyn, A.P., 1996. New Early Cretaceous Embolemidae (Vespida = Hymenoptera: Chrysoidea). *Memoirs of the Entomological Society of Washington* 17, 183–187.
- Rasnitsyn, A.P., Brothers, D.J., 2020. The first plumalexiid wasp (Hymenoptera: Chrysoidea, Plumalexiidae) from the mid-Cretaceous Burmese amber. *Cretaceous Research* 115, 104568.

- Rohwer, S.A., 1909. Three new fossil insects from Florissant, Colorado. *Bulletin of the American Museum of Natural History* 23, 605–616.
- Rosa, P., Wei, N., Xu, Z., 2014. An annotated checklist of the chrysidid wasps (Hymenoptera, Chrysididae) from China. *ZooKeys* 455, 1–128.
- Rosa, P., Heim, R., Bernasconi, M.V., 2022. The Palaearctic types of Chrysididae (Insecta, Hymenoptera) deposited in the Linsemaier collection. Part 2. Elampini: genus *Hedychridium* Abeille de Perrin, 1878. *Zootaxa* 5090, 1–94.
- Rubink, W.L., Evans, H.E., 1979. Notes on the nesting behavior of the bethylid wasps, *Epyris eriogoni* Kieffer, in southern Texas. *Psyche* 86, 313–320.
- Sanderson, M.W., Farr, T.H., 1960. Amber with insect and plant inclusions from the Dominican Republic. *Science* 131, 1313–1314.
- Santosh, S., 2017. A taxonomic revision of *Sierola* Cameron (Hymenoptera: Bethyridae) from Oriental region. In: Santosh, S., Nasser, M., Sudheer, K. (Eds.), *Insect Diversity and Taxonomy*. Prof. TC Narendran Trust for Animal Taxonomy, Calicut, pp. 307–317.
- Schmidt, A.R., Kaulfuss, U., Bannister, J.M., Baranov, V., Beimforde, C., Bleile, N., Borkent, A., Busch, A., Conran, J.G., Engel, M.S., Harvey, M., Kennedy, E.M., Kerr, P.H., Kettunen, E., Kiecksee, A.P., Lengeling, F., Lindqvist, J.K., Maraun, M., Mildenhall, D.C., Perrichot, V., Rikkinen, J., Sadowski, E.M., Seyfullah, L.J., Stebner, F., Szwed, J., Ulbrich, P., Lee, D.E., 2018. Amber inclusions from New Zealand. *Gondwana Research* 56, 135–146.
- Smitinand, T., 1968. Vegetation of Khao Yai National Park. *Natural History Bulletin of the Siam Society* 22 (3–4), 289–305.
- Solórzano Kraemer, M.M., Kraemer, A.S., Stebner, F., Bickel, D.J., Rust, J., 2015. Entrapment bias of arthropods in Miocene amber revealed by trapping experiments in a tropical forest in Chiapas, Mexico. *PLoS ONE* 10, e0118820, doi: 10.1371/journal.pone.0118820.
- Sorg, M., 1988. Zur Phylogenie und Systematik der Bethyridae (Insecta, Hymenoptera, Chryridoidea). *Sonderveröffentlichungen des Geologischen Instituts der Universität zu Köln* 63, 1–146.
- Swezey, O.H., 1954. Forest entomology in Hawaii. *Bernice P. Bishop Museum Special Publication* 44, 1–266.
- Sziráki, G., Dulai, A., 2002. Sarmatian (Late Miocene) arthropods from Tállya and neighbouring localities (Tokaj Mts, Hungary): A preliminary report. *Annales Historico-Naturales Musei Nationalis Hungarici* 94, 31–44.

- Terayama, M., 2004. Descriptions of new taxa and distribution records of the family Bethylidae (Insecta, Hymenoptera) II. Subfamily Bethylinae and fossil taxa. Academic Reports of the Faculty of Engineering of Tokyo Polytechnic University 27, 39–52.
- Terayama, M., 2006. Insects of Japan (Vol. 1): Bethylidae (Hymenoptera). Touka Shobo Co., Fukuoka, 317 pp.
- Tsuneki, K., 1961. Chrysididae collected by the Osaka City University Biological Expedition to southeast Asia, 1957–1958. Nature and Life in Southeast Asia 1, 367–382.
- Ujiye, T., Kamiyo, K., Morakote, R., 1996. Species composition of parasitoids and rate of parasitism of the citrus leafminer (CLM), *Phyllocnistis citrella* Stainton (Lepidoptera: Gracillariidae) in central and northern Thailand, with key to parasitoids of CLM collected from Japan, Taiwan and Thailand. Bulletin of the Fruit Tree Research Station 29, 79–106.
- Wang, B., Shi, G., Xu, C., Spicer, R.A., Perrichot, V., Schmidt, A.R., Feldberg, K., Heinrichs, J., Chény, C., Pang, H., Liu, X., Gao, T., Wang, Z., Ślipiński, A., Solórzano-Kraemer, M.M., Heads, S.W., Thomas, M.J., Sadowski, E., Szwed, J., Azar, D., Nel, A., Liu, Y., Chen, J., Zhang, Q., Zhang, Q., Luo, C., Yu, T., Zheng, D., Zhang, H., Engel, M.S., 2021. The mid-Miocene Zhangpu biota reveals an outstandingly rich rainforest biome in East Asia. Science Advances 7, eabg0625, doi: 10.1126/sciadv.abg0625.
- Wang, C., He, J., Chen, X., 2021. Taxonomy of *Sierola* Cameron (Hymenoptera, Bethylidae) from China with three new species. Journal of Hymenoptera Research 84, 405–415.
- Westwood, J.O., 1874. Thesaurus Entomologicus Oxoniensis. Clarendon Press, Oxford, 205 pp.
- Zheng, D., Shi, G., Hemming, S.R., Zhang, H., Wang, W., Wang, B., Chang, S.C., 2019. Age constraint on a Neogene tropical rainforest in China and its relation to the Middle Miocene Climatic Optimum. Palaeogeography, Palaeoclimatology, Palaeoecology 518, 82–88.

Figure captions


Fig. 1. *Sierola colomboi* n. sp., holotype, male, NIGP200740, habitus in dorsal view. Scale bar = 0.5 mm.





Fig. 2. *Sierola colomboi* n. sp., holotype, male, NIGP200740. (A) Habitus in lateral view. (B) Head and mesosoma in dorsal view. (C) Propodeum in laterodorsal view. (D) Line drawing of fore wing. Scale bar: 0.5 mm for (A, D); 0.25 mm for (B, C).


Fig. 3. *Hedychridium rosai* n. sp., holotype, female, NIGP200741, habitus in right lateral view. Scale bar = 1 mm.

Fig. 4. *Hedychridium rosai* n. sp., holotype, female, NIGP200741. (A) Habitus in left lateral view. (B) Habitus in ventral view. (C) Habitus (centred on mesosoma) in dorsal view. (D) Habitus (centered on anterior metasoma) in dorsal view. Scale bar = 1 mm.



Fig. 5. *Hedychridium rosai* n. sp., holotype, female, NIGP200741. (A) Head and mesosoma in left lateral view. (B) Head in frontal view. (C) Mesosoma in dorsal view (white arrows: hamuli). (D) Line drawing of fore wing. Scale bar = 0.5 mm.







Journal Pre-proofs






Table 1. Summary of the Miocene record of Chrysidoidea.

Taxon	Deposit	Reference
Bethylidae		
<i>Apenesia miki</i> (Terayama, 2004)	Dominican Republic amber	Terayama, 2004
<i>Alongatepyris pedrocai</i> Colombo and Azevedo, 2022	Dominican Republic amber	Colombo et al., 2022
<i>Anisepyris gradatus</i> Sorg, 1988	Dominican Republic amber	Sorg, 1988
<i>Goniozus respectus</i> Sorg, 1988	Dominican Republic amber	Sorg, 1988
<i>Bakeriella nanyhelae</i> Brazidec and Perrichot, 2022	Mexican amber	Brazidec and Perrichot, 2022
<i>Goniozus cotyi</i> Brazidec and Perrichot, 2022	Mexican amber	Brazidec and Perrichot, 2022

<i>Solepyris electromexicanus</i> Brazidec and Perrichot, 2022	Mexican amber	Brazidec and Perrichot, 2022
<i>Goniozus</i> sp.	Rubielos de Mora, Spain	Peñalver-Mollá, 1998
<i>Parascleroderma palaeosinica</i> Brazidec and Perrichot	Zhangpu amber, China	Brazidec and Perrichot, in press
<i>Sierola colomboi</i> n. sp.	Zhangpu amber, China	This paper
Chrysididae		
<i>Ceratochrysis dominica</i> Engel, 2006	Dominican Republic amber	Engel, 2006
Chrysididae indet.	Tállya, Hungary	Sziráki and Dulai, 2002
<i>Hedychridium rosai</i> n. sp.	Zhangpu amber, China	This paper
Dryinidae		
<i>Aphelopus poinari</i> Olmi, 1998	Dominican Republic amber	Olmi, 1998
<i>Bocchus vetustus</i> Olmi, 1989	Dominican Republic amber	Olmi, 1989
<i>Dryinus alamellatus</i> Olmi and Guglielmino, 2011	Dominican Republic amber	Olmi and Guglielmino, 2011
<i>Dryinus grimaldii</i> Olmi, 1995	Dominican Republic amber	Olmi, 1995
<i>Dryinus hymenaeophilus</i> Olmi, 1995	Dominican Republic amber	Olmi, 1995
<i>Dryinus palaeodominicanus</i> Currado and Olmi, 1983	Dominican Republic amber	Currado and Olmi, 1983
<i>Dryinus poinari</i> Olmi, 1998	Dominican Republic amber	Olmi, 1998
<i>Dryinus priscus</i> Olmi, 1998	Dominican Republic amber	Olmi, 1998
<i>Dryinus pristinus</i> Olmi, 1998	Dominican Republic amber	Olmi, 1998
<i>Dryinus rasnitsyni</i> Olmi and Guglielmino, 2011	Dominican Republic amber	Olmi and Guglielmino, 2011
<i>Dryinus vetustus</i> Olmi, 1995	Dominican Republic amber	Olmi, 1995
<i>Harpactospecion scheveni</i> Olmi, 2005	Dominican Republic amber	Olmi, 2005a
<i>Harpactospecion sucinum</i> (Olmi, 1987)	Dominican Republic amber	Olmi, 1987
<i>Thaumatodryinus fuscescens</i> Martins and Melo, 2020	Dominican Republic amber	Martins and Melo, 2020
<i>Thaumatodryinus miocenicus</i> Olmi, 1995	Dominican Republic amber	Olmi, 1995
<i>Thaumatodryinus priscus</i> (Olmi, 1998)	Dominican Republic amber	Olmi, 1998
<i>Dryinus palaeomexicanus</i> Olmi, 1995	Mexican amber	Olmi, 1995
Embolemidae		
Embolemidae indet.	Dominican Republic amber	Rasnitsyn, 1996
Sclerogibbidae		
<i>Probethylus poinari</i> Olmi, 2005	Dominican Republic amber	Olmi, 2005b

<i>Pterosclerogibba antiqua</i> Olmi, 2005	Dominican Republic amber	Olmi, 2005b
Scolebythidae		
<i>Clytopsenella mirabilis</i> Engel, 2015	Dominican Republic amber	Engel, 2015
<i>Pristapenesia inopinata</i> (Prentice and Poinar, 1996)	Dominican Republic amber	Prentice et al., 1996