



**HAL**  
open science

## Early evolution of the Hunga – Tonga Volcanic Plume from Lidar. Observations at Reunion Island (Indian Ocean, 21°S, 55°E)

Alexandre Baron, Guillaume Payen, Valentin Duflot, Patrick Chazette, Sergey Khaykin, Yann Hello, Nicolas Marquestaut, Marion Ranaivombola, Nelson Bègue, Thierry Portafaix, et al.

### ► To cite this version:

Alexandre Baron, Guillaume Payen, Valentin Duflot, Patrick Chazette, Sergey Khaykin, et al.. Early evolution of the Hunga – Tonga Volcanic Plume from Lidar. Observations at Reunion Island (Indian Ocean, 21°S, 55°E). EGU General Assembly 2022, EGU, May 2022, Vienna, Austria. 10.5194/egusphere-egu22-13599 . insu-03873420

**HAL Id: insu-03873420**

**<https://insu.hal.science/insu-03873420>**

Submitted on 26 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

EGU22-13599

<https://doi.org/10.5194/egusphere-egu22-13599>

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



## Early evolution of the Hunga – Tonga Volcanic Plume from Lidar Observations at Reunion Island (Indian Ocean, 21°S, 55°E)

**Alexandre Baron**<sup>1</sup>, Guillaume Payen<sup>2</sup>, Valentin Duflot<sup>1,2</sup>, Patrick Chazette<sup>3</sup>, Sergey Khaykin<sup>4</sup>, Yann Hello<sup>2</sup>, Nicolas Marquestaut<sup>2</sup>, Marion Ranaivombola<sup>1</sup>, Nelson Bègue<sup>1</sup>, Thierry Portafaix<sup>1</sup>, and Jean-Pierre Cammas<sup>2</sup>

<sup>1</sup>Laboratoire de l'Atmosphère et des Cyclones (LACy), UMR 8105, Université de La Réunion - CNRS - Météo-France, Saint-Denis de La Réunion, France, E-mail: alexandre.baron@univ-reunion.fr

<sup>2</sup>Observatoire des Sciences de l'Univers de La Réunion (OSU-R), UMS 3365, Saint-Denis de La Réunion, France

<sup>3</sup>Laboratoire des Sciences du Climat et de l'Environnement, LSCE-IPSL, UMR 8212, Université Paris-Saclay - CEA - CNRS - UVSQ, Gif-sur-Yvette, France

<sup>4</sup>Laboratoire Atmosphère Milieux Observations Spatiales, LATMOS-IPSL, UMR 8190, Sorbonne Université - UVSQ - CNRS, Paris, France

Explosive volcanism periodically induces disturbances of the upper troposphere and low stratosphere. These injections of massive amount of aerosols, ash and gases perturb locally the physico-chemical balance of the impacted atmospheric layers, in particular the ozone concentration via heterogeneous chemistry on particles. On a larger scale some exceptional eruption can have a significant influence on the Earth radiative budget as it was the case following eruptions of El Chichon in 1982 and Mount Pinatubo in 1991.

On January 15, 2022, the Hunga-Tonga volcano erupted in the Tonga archipelago (20.5°S, 175.4°W). The Plinian eruption was of a rare intensity, especially because of the depth of the underwater caldera. The first estimates indicate a power between 10 and 15 Mt TNT, probably the most powerful since the eruption of Krakatoa in 1883. This short (~ 8min) but intense explosion whose pressure wave was observed all around the globe injected about 400 kt of material into the atmosphere (to be compared to the 20 Mt injected during the Mount Pinatubo eruption). The Volcano Stratospheric Plume (VSP) quickly moved westwards and then overflowed the island of La Réunion (21°S, 55°E), located at ~12000 km away from Tonga.

In order to monitor the evolution of the VSP, lidar observations were performed at the *Observatoire de Physique de l'Atmosphère de La Réunion* (OPAR). This observatory is equipped with three lidars capable of stratospheric aerosols measurements at two wavelengths (355 nm and 532 nm). First observations were performed every night from 19 to 27 January 2022 when the first passage of the VSP occurred. The plume structures appeared to be highly variable along time, with altitudes ranging from 19 km to 36 km above the mean sea level while plume thicknesses were ranging from ~1 km to more than 3 km. Remarkable aerosol optical depth were associated with these stratospheric aerosol layers, up to 0.8 at 532 nm on January 21.

The temporal evolution of the VSP structure and optical properties will be presented and discussed.