



**HAL**  
open science

## Vertical distribution of atmospheric temperature and density from the solar occultation instruments NOMAD and ACS on board the Trace Gas Orbiter

Miguel Angel Lopez-Valverde, Bernd Funke, Adrian Brines, Aurélien Stolzenbach, Ashimananda Modak, Francisco Gonzalez-Galindo, Shohei Aoki, Loic Trompet, Ian Thomas, Gerónimo Villanueva, et al.

### ► To cite this version:

Miguel Angel Lopez-Valverde, Bernd Funke, Adrian Brines, Aurélien Stolzenbach, Ashimananda Modak, et al.. Vertical distribution of atmospheric temperature and density from the solar occultation instruments NOMAD and ACS on board the Trace Gas Orbiter. Europlanet Science Congress 2022, Sep 2022, Granada, Spain. pp.EPSC2022-779, 10.5194/epsc2022-779 . insu-03881618

**HAL Id: insu-03881618**

**<https://insu.hal.science/insu-03881618v1>**

Submitted on 1 Dec 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Vertical distribution of atmospheric temperature and density from the solar occultation instruments NOMAD and ACS on board the Trace Gas Orbiter

**Miguel Angel Lopez-Valverde**<sup>1</sup>, Bernd Funke<sup>1</sup>, Adrian Brines<sup>1</sup>, Aurélien Stolzenbach<sup>1</sup>, Ashimananda Modak<sup>1</sup>, Francisco Gonzalez-Galindo<sup>1</sup>, Shohei Aoki<sup>3</sup>, Loic Trompet<sup>2</sup>, Ian Thomas<sup>2</sup>, Gerónimo Villanueva<sup>4</sup>, Giuliano Liuzzi<sup>5</sup>, Denis Belyaev<sup>6</sup>, Kevin Olsen<sup>7</sup>, Alexander Trokhimovsky<sup>6</sup>, Jose Juan Lopez-Moreno<sup>1</sup>, Ann Carine Vandaele<sup>2</sup>, Manish Patel<sup>10</sup>, Giancarlo Bellucci<sup>9</sup>, Oleg Korablev<sup>6</sup>, and Franck Montmessin<sup>8</sup>

<sup>1</sup>Instituto de Astrofísica de Andalucía / CSIC, Departamento Sistema Solar, Granada, Spain (valverde@iaa.es)

<sup>2</sup>Royal Belgian Institute for Space Aeronomy, Brussels, Belgium

<sup>3</sup>The University of Tokyo, Kashiwa, Japan

<sup>4</sup>NASA Goddard Space Flight Center, USA

<sup>5</sup>American University, Washington DC, USA

<sup>6</sup>Space Research Institute (IKI RAS), Moscow, Russia

<sup>7</sup>Department of Physics, Oxford University, UK

<sup>8</sup>LATMOS / CNRS, France

<sup>9</sup>Institute for Space Astrophysics and Planetology, Italy

<sup>10</sup>Open University, Milton Keynes, UK

One of the goals of the ESA and Roscosmos Exomars Trace Gas Orbiter (TGO) is the exploitation of the two solar occultation instruments NOMAD [1] and ACS [2] to characterize the thermal structure of the Martian atmosphere with unprecedented vertical resolution, and from the ground up to the thermosphere. Specifically for the upper atmosphere this is a unique opportunity [3] and we present here an on-going effort to retrieve CO<sub>2</sub> abundance and temperature profiles simultaneously from each of these instruments and to combine them for a mutual validation and for obtaining a most complete mapping from the TGO orbit.

We developed a retrieval suite common for both instruments, comprising: (a) a cleaning/pre-processing module to build vertical profiles of calibrated transmittances which computes and correct for residual calibration and instrumental effects like spectral shifts, bending of the continuum and variations in the instrument line shape; (b) a state-of-the-art retrieval scheme designed originally for Earth atmospheric remote sensing [4,5,6] and applied to Mars [7], in order to derive simultaneous density and temperature profiles allowing for hydrostatic adjustments during the internal iteration.

Recently the first application of this retrieval scheme focused on measurements by the NOMAD SO channel at altitudes below 100 km, and for the first year of TGO operations, from April 2018 to March 2019 (second half or “perihelion” season of MY34), and revealed very interesting results [8]. The thermal structure is strongly affected by the MY34 global dust storm at all altitudes, a cold mesosphere (in comparison to global climate models) was found during the post-GDS period, and

wavy structures at mesospheric altitudes in the morning terminator seem to reveal very strong thermal tides at low-mid latitudes.

In this presentation we also focus on the Martian troposphere and mesosphere and build upon the above mentioned work during the 1<sup>st</sup> year of TGO by extending the study to a full Mars year and adding retrievals from ACS MIR channel. Both NOMAD/SO and ACS/MIR channels observe the strong CO<sub>2</sub> ro-vibrational band at 2.7 micron in the same spectral region with some differences in spectral resolution and noise level, in addition to very different instrument characteristics, which are included in our retrieval approach. In both cases we use calibrated atmospheric transmittances to tackle three targets, CO<sub>2</sub> density, temperature, and dust loading, in a simultaneous global-fit inversion, including contaminant species like H<sub>2</sub>O. The contamination by aerosol can severely limit the ability to sound low tangent altitudes with both instruments, when the gas absorption lines become hidden within the aerosol continuum. But also these measurements permit a good characterization of aerosol properties [9]. Propagation of measurement noise, tuning of regularization, and computation of averaging kernels are performed with the same code and the comparison of the retrieval performance is a first step towards a common validation between these two instruments, considering that a complete correlation is not possible since the two instruments' individual solar occultation scans are non-coincident in time and space. Our results will be discussed and compared to parallel retrieval efforts by other teams within the NOMAD and ACS consortia using the same datasets [10, 11, 12]. We will also compare them with specific runs of the LMD-Mars GCM (see also a companion presentation in this conference on this specific topic [13]). One of the important applications of our inversion is to supply true-field and a.priori inputs for the inversion of trace species from collocated NOMAD spectra, as we are doing for water vapour [14] and carbon monoxide [15].

### **Acknowledgments**

The IAA/CSIC team acknowledges financial support from the State Agency for Research of the Spanish MCIU through the 'Center of Excellence Severo Ochoa' award for the Instituto de Astrofísica de Andalucía (SEV-2017-0709) and funding by grant PGC2018-101836-B-100 (MCIU/AEI/FEDER, EU). ExoMars is a space mission of the European Space Agency (ESA) and Roscosmos. The NOMAD experiment is led by the Royal Belgian Institute for Space Aeronomy (IASB-BIRA), assisted by Co-PI teams from Spain (IAA-CSIC), Italy (INAF-IAPS), and the United Kingdom (Open University). US investigators were supported by the National Aeronautics and Space Administration.

### **References**

- [1] Vandaele et al., *Space Science Reviews* 214, 5, 2018
- [2] Koroblev et al., *Space. Sci. Rev.* 214, 7 (2018).
- [3] Lopez-Valverde et al., *Space Sci Rev*, 214, 29 (2018)
- [4] Funke, B., et al. , *Atmos. Chem. Phys.*, 9(7), 2387–2411 (2009).
- [5] Stiller et al., *JQSRT*, 72, 249–280 (2002)
- [6] von Clarmann et al., *J. Geophys. Res.* 108, 4746 (2003)
- [7] Jimenez-Monferrer et al., *Icarus*, 353, 113830 (2020), doi.org/10.1016/j.icarus.2020.113830.
- [8] Lopez-Valverde et al., *JGR-Planets* (submitted, 2022)
- [9] Stolzenbach et al., *JGR-Planets* (submitted, 2022) and Stolzenbach et al., *EPSC 2022* (this conference)
- [10] Trompet et al., *JGR-Planets* (submitted, 2022)
- [11] Belyaev et al., *GRL*

[12] Belyaev et al., JGR-Planets (submitted, 2022)

[13] Gonzalez-Galindo et al., EPSC 2022 (this conference)

[14] Brines et al., JGR-Planets (submitted, 2022) and Brines et al., EPSC 2022 (this conference)

[15] Modak et al., JGR-Planets (submitted, 2022) and Modak et al., EPSC 2022 (this conference)