



HAL
open science

Upper Jurassic (TITHONIAN) belemnites from the Neuquén basin, Argentina

Verónica V Vennari, Beatriz Aguirre-Urreta, Luciana S Marin, Pierre Pellenard, Mathieu Martínez, Maisa Tunik

► **To cite this version:**

Verónica V Vennari, Beatriz Aguirre-Urreta, Luciana S Marin, Pierre Pellenard, Mathieu Martínez, et al.. Upper Jurassic (TITHONIAN) belemnites from the Neuquén basin, Argentina. *Journal of South American Earth Sciences*, 2023, 124, pp.104200. 10.1016/j.jsames.2023.104200 . insu-03999990

HAL Id: insu-03999990

<https://insu.hal.science/insu-03999990v1>

Submitted on 22 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proof

Upper Jurassic (TITHONIAN) belemnites from the Neuquén basin, Argentina

Verónica V. Vennari, Beatriz Aguirre-Urreta, Luciana S. Marin, Pierre Pellenard, Mathieu Martínez, Maisa Tunik



PII: S0895-9811(23)00011-1

DOI: <https://doi.org/10.1016/j.jsames.2023.104200>

Reference: SAMES 104200

To appear in: *Journal of South American Earth Sciences*

Received Date: 4 July 2022

Revised Date: 27 December 2022

Accepted Date: 15 January 2023

Please cite this article as: Vennari, Veró.V., Aguirre-Urreta, B., Marin, L.S., Pellenard, P., Martínez, M., Tunik, M., Upper Jurassic (TITHONIAN) belemnites from the Neuquén basin, Argentina, *Journal of South American Earth Sciences* (2023), doi: <https://doi.org/10.1016/j.jsames.2023.104200>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Ltd.

1 **UPPER JURASSIC (TITHONIAN) BELEMNITES FROM THE NEUQUÉN**
2 **BASIN, ARGENTINA**

3

4 VERÓNICA V. VENNARI*^{1,3}, BEATRIZ AGUIRRE-URRETA^{2,3}, LUCIANA S. MARIN^{2,3}, PIERRE
5 PELLENARD⁴, MATHIEU MARTÍNEZ⁵ AND MAISA TUNIK⁶

6 ¹ Instituto de Evolución, Ecología Histórica y Ambiente (IDEVEA), Universidad
7 Tecnológica Nacional, Facultad Regional San Rafael (UTN-FRSR). Gral. J.J. Urquiza
8 314, M5602GCH, San Rafael, Mendoza, Argentina. vvennari@mendoza-conicet.gob.ar

9 ² Instituto de Estudios Andinos "Don Pablo Groeber"(IDEAN), Universidad de Buenos
10 Aires, Facultad de Ciencias Exactas y Naturales. Intendente Güiraldes 2160 Ciudad
11 Universitaria, C1428EGA, CABA, Argentina. aguirre@gl.fcen.uba.ar;
12 lmartin@gl.fcen.uba.ar

13 ³ Consejo Nacional de Investigaciones científicas y Tecnológicas (CONICET)

14 ⁴ Biogéosciences, UMR 6282 CNRS/uB/EPHE 6 Bd Gabriel, Université Bourgogne
15 Franche-Comté, Dijon, Francia. pierre.pellenard@u-bourgogne.fr

16 ⁵ Univ Rennes, CNRS, Géosciences Rennes - UMR 6118, F-35000 Rennes, Francia.
17 mathieu.martinez@univ-rennes1.fr

18 ⁶ Instituto de Investigación en Paleobiología y Geología, CONICET & Universidad de
19 Río Negro, Sede Alto Valle, 8332 General Roca, Río Negro, Argentina.
20 mtunik@unrn.edu.ar

21

22 * Corresponding author. e-mail address: vvennari@conicet-mendoza.gob.ar

23

24 **Abstract.** Belemnites were a group of Mesozoic coleoids with a global distribution
25 from the Early Jurassic to the Late Cretaceous. During the Late Jurassic–Early
26 Cretaceous, Gondwana's marine shelves were dominated by *Belemnopsis* and *Hibolithes*
27 populations. In contrast to the Austral Basin, where both genera are quite common, their
28 record in the Neuquén Basin (Argentina) is rather scarce and poor, probably due to a
29 combination of locally unfavorable environmental conditions and predation pressure.
30 Here we describe a new belemnite association from the Tithonian of the Vaca Muerta
31 Formation from southern Mendoza and northern Neuquén provinces. Belemnite's
32 findings are assignable to the Tithonian *Windhausenicerias internispinosum* and
33 *Corongoceras alternans* Andean ammonoid Biozones. The studied rostra are identified
34 as *Belemnopsis* cf. *B. patagoniensis*, *Belemnopsis* cf. *B. launceloti*, and *Hibolithes*
35 *argentinus*? Affinities of the Neuquén Basin belemnite fauna are referred to the
36 Madagascan-Eastern Africa associations (Ethiopian Subprovince, Tethyan Province),
37 adding evidence to an established Trans-Gondwana seaway enabling the connection
38 between those areas from the Tithonian onwards.

39

40 **Keywords.** Fossil coleoidea. *Belemnopsis*. *Hibolithes*. Vaca Muerta Formation. Trans-
41 Gondwana seaway.

42

43 1.1 Introduction

44 Belemnites are an extinct group of Mesozoic nektonic cephalopods considered in the
45 most recent contributions as stem-decachabrachian coleoids (Fuchs et al., 2013; Hoffmann
46 and Stevens, 2020). Usually reconstructed as externally morphologically close to
47 loginid squids (Naef, 1922; Stevens, 1965), belemnites had a mantle muscle structure
48 similar to that of sepiids (Monks et al., 1996). Their mineralized inner shell was
49 generally divided into three parts: proostracum, phragmocone, and rostrum. Of these,
50 the low-Mg calcitic rostrum is the most commonly preserved part due to its resistant
51 structure of radially arranged calcite fibers that, combined with variable amounts of
52 organic matter, form concentric rings in cross-section (Stolley, 1911; Jeletzki, 1966;
53 Riegraf, 1980; Hoffmann et al., 2016; Hoffmann and Stevens, 2020). Belemnite
54 taxonomy mostly relies on morphological characters of the rostrum: general
55 morphology, both in outline and profile view, shape of the transverse section, form and
56 orientation of the apex, and presence and extension of grooves (apical, ventral, dorsal,
57 and/or lateral). The orientation and form of the apical line and alveolus-related
58 characters (*i.e.*, where the phragmocone lies inside the rostrum) are usually accessorially
59 described (Stevens, 1965; Jeletzky, 1966; Doyle and Kelly, 1988; Challinor, 1999).

60 Belemnites are recorded from the Late Triassic (Iba et al., 2014), but it is from
61 the Toarcian onwards that they attain a worldwide distribution (Doyle, 1987). By the
62 Late Jurassic–Early Cretaceous Gondwana shelf-areas were dominated by *Belemnopsis*
63 and *Hibolithes* (Challinor, 1991; Doyle and Pirrie, 1999), both included in the
64 "Belemnopseina," a group characterized by alveolar furrows and whose taxonomic
65 validity has been recently challenged (Stevens, in press). *Hibolithes* was a cosmopolitan
66 and ecologically wide-ranged genus whose species are mostly differentiated based on
67 the ventral groove characters (Stevens, 1965; Mutterlose, 1988; Doyle and Kelly, 1988;

68 Challinor, 1991). *Belemnopsis* is a more variable genus (Stevens, 1965; Combémoré,
69 1988; Howlett, 1989; Challinor, 1991, 2001, 2003; Challinor and Hikuroa, 2007) mostly
70 represented in Gondwana by robust, cylindroconical to slightly depressed rostra with a
71 variably strong ventral groove (Challinor, 1991; Doyle et al., 1996, 1997; Doyle and
72 Pirrie, 1999). In Argentina, both *Hibolithes* and *Belemnopsis* had been long recognized
73 from the Upper Jurassic and, mostly, from the Lower Cretaceous of the Austral Basin in
74 south Patagonia (e.g., Dana, 1848; Favre, 1908; Feruglio, 1936, 1949) and have been
75 revised or cited in the literature from there on (e.g., Aguirre-Urreta and Suárez, 1985;
76 Riccardi, 1976, 1977, 1988). In contrast, Neuquén Basin belemnite records of the same
77 age are significantly less frequent, and taxonomic-focused contributions are lacking
78 (Stevens, 1965; Doyle et al., 1996; 1997).

79 Therefore, the objective of this work is to fully document new belemnite rostra
80 from the Upper Jurassic (Tithonian) of the Neuquén Basin to start filling the gap in the
81 knowledge of this group in west-central Argentina. We also aim to refer the findings to
82 the Andean ammonoid-based biostratigraphic framework, and to discuss the
83 paleobiogeographic, biostratigraphic, and broad paleoecological implications of their
84 occurrences.

85 **1.2. Previous Neocomian belemnite records in Argentina and nearby regions**

86 Upper Jurassic–Lower Cretaceous belemnite records from Argentina are currently
87 concentrated in the Austral Basin (Fig. 1). *B. patagoniensis* (Favre) was originally
88 described from the Hauterivian–Barremian of Lago Belgrano area and later restricted to
89 the Hauterivian (Favre, 1908; Riccardi, 1988). Findings around Lago San Martín, at the
90 top of the Springhill Formation, were alternatively ascribed to the Tithonian (Bonarelli
91 and Nágera, 1921; Feruglio, 1949), the Hauterivian–Barremian (Riccardi, 1971), or the
92 Berriasian (Riccardi, 1976, 1977). Finally, they were assigned to a Berriasian–

93 Valanginian age (Medina and Riccardi, 2005; Riccardi, 1988). From the same levels,
94 Riccardi (1977) reported *B. cf. B. madagascariensis* (Besairie) and *Hibolites* aff. *H.*
95 *jaculum* (Phillips). Also from the Lago San Martín area, at the base of the Río Mayer
96 Formation, Medina and Riccardi (2005) reported *B. patagoniensis*, *B. cf. gladiatoris*
97 Willey, and *Belemnopsis* sp. together with the ammonoids *Favrella americana* (Favre)
98 and *F. wilckensi* (Favre), thus indicating a Hauterivian age for the association. To the
99 south, in the Lago Argentino area, Estancia La Cristina, Feruglio (1936) introduced a
100 supposedly Valanginian belemnite association composed of *B. patagoniensis*,
101 *Hibolites argentinus* Feruglio, *H. aff. H. jaculum*, and a specimen openly assigned as
102 *Belemnites* sp. I, which was reinterpreted as a *Belemnopsis* sp. by Stevens (1965, p.
103 159) and alternatively as *Parahibolites* by Challinor (1991, p. 316). The specimen
104 assigned to *B. patagoniensis* was reassigned to *B. gladiatoris* Willey due to
105 morphological differences with the type material (Stevens, 1965; Willey, 1973) and
106 subsequently to *B. launzeloti* Howlett (Howlett, 1989). Feruglio (1949, p. 169) also
107 mentioned, but not illustrated, *B. patagoniensis* from another locality (Cerro de Los
108 Fósiles) within the Lago Argentino area, there associated with Tithonian to Berriasian
109 ammonoid taxa (cf. Leanza, A., 1967). A reassessment of the Indo-Pacific affinity of the
110 whole assemblage (Stevens, 1965) as well as a revision of the ammonoid-related fauna,
111 pointed to an upper Tithonian age for Feruglio's original belemnite collection (Leanza,
112 A., 1967). The same seems valid for related specimens from the Springhill Formation
113 near Lago Argentino, where *Hibolites* ranges into the Berriasian and younger beds of
114 the Río Mayer Formation *sensu* Doyle et al. (1996).

115 Other illustrated records of *B. patagoniensis* or *B. cf. patagoniensis* come from
116 the southernmost end of South America, from Berriasian–Hauterivian beds of the
117 Yaghán Formation, Bahía Nassau, Tierra del Fuego (Dana, 1848; Stevens, 1965;

118 Stolley, 1928) and from the Valanginian?–Hauterivian Beauvoir Formation in the
119 western region of the Isla de los Estados (Harrington, 1943; Caminos and Nullo, 1979).
120 Also, Aguirre-Urreta and Suárez (1985) presented new Berriasian–Hauterivian records
121 of *B. madagascariensis* from several sections within the Magallanes Archipelago.
122 Recently, Ippolitov et al. (2015) described a single specimen from the Berriasian of
123 Crimea as *Parabelemnopsis* cf. *patagoniensis*, indicating that its presence was probably
124 the result of a rare immigration episode to the Northern Hemisphere. Aside from the
125 later record, *Belemnopsis patagoniensis* appears to be a well-distributed species of the
126 Austral Basin during the Lower Cretaceous.

127 Other records of *Belemnopsis* in nearby areas are known from the Valanginian–
128 Hauterivian of the Rocas Verdes Basin, Chile, where robust forms were ascribed to the
129 Antarctic species *B.* cf. *B. launceloti* Howlett (Stinnesbeck et al., 2014). Additional
130 material retrieved from a drill hole at the Falkland/Malvinas Plateau was identified as
131 *Belemnopsis* spp. and assigned a New Zealand to Himalayan affinity (Jeletzky, 1983).

132 Within the Neuquén Basin (Fig. 1), Upper Jurassic–Lower Cretaceous belemnite
133 records are sparser and poorly known. Oxfordian *Belemnopsis* with European affinities
134 have been mentioned from La Manga Formation in southern Neuquén (Doyle et al.,
135 1996; Riccardi et al., 2011) and from the Callovian–Oxfordian of Vega de la Veranada,
136 in northern Neuquén (Alberti et al., 2020). Younger not-illustrated records from
137 Mendoza Province include an indeterminate rostrum fragment from the Tithonian of
138 Arroyo Serrucho (Krantz, 1928) and some thin incomplete rostra from the upper
139 Tithonian of the Vaca Muerta Formation in Bardas Blancas (Gerth, 1925; here referred
140 as Cañada de Leiva). Additional fragmentary belemnites were reported from a
141 Berriasian interval in the Arroyo La Manga area (Gerth, 1925). Other Tithonian records
142 from Neuquén Province were either identified as *Hibolithes* sp. (e.g., Alberti et al.,

143 2020, from Pampa Tril) or not specified (*e.g.*, Kietzmann and Vennari, 2013, from
144 Cerro Domuyo, here revised). Besides, Stevens (1965, p. 158) indicated a possible
145 Kimmeridgian to Tithonian age for ?*Hibolithes* sp., *B. cf. B. patagoniensis* and
146 ?*Dicoelites* sp. from Cerro Lotena, and for ?*Hibolithes* sp. from Arroyo Chacay
147 Melehue, but unfortunately, none of those records were illustrated.

148 FIGURE 1 AROUND HERE

149 **2. Geological setting**

150 The Neuquén Basin (Fig. 1) is a retro-arc basin situated over the western margin of
151 Gondwana, between 32° to 40° South Latitude, extending over central-western
152 Argentina and central Chile (Legarreta and Uliana, 1991; Howell et al., 2005). Its nearly
153 continuous Mesozoic to lower Cenozoic sedimentary infill, intercalated with volcanic
154 episodes, and the high quality and richness of its fossil record arouse the interest of the
155 international scientific community over the basin.

156 Tithonian successions bearing the belemnite fauna here presented are included in
157 the Vaca Muerta Formation (Weaver, 1931), a marine geographically extended unit
158 mainly composed by an alternation of shale/marls with a variety of limestone beds
159 deposited over a carbonate ramp to basin setting from the lower Tithonian to the lower
160 Valanginian (Kietzmann et al., 2014; Legarreta and Uliana, 1991). Ammonoids are the
161 most common megascopic fossils of the Vaca Muerta Formation, including both
162 endemic and more extended distributed taxa. Their correlation with Tethyan faunas and
163 with calcareous nanofossils and microfossils bioevents provide the current Upper
164 Jurassic–Lower Cretaceous biostratigraphic scheme for the Neuquén Basin (Leanza, H.
165 et al., 2020 and references therein; Lescano et al., 2022).

166 **3. Material and methods**

167 **3.1. Localities**

168 The twenty-one belemnites herein studied come from Tithonian beds of the Vaca
169 Muerta Formation in four localities situated over the northern sector of the Neuquén
170 Basin (Fig. 2). From the northeast to the southwest, they are Cañada de Leiva (fourteen
171 specimens), Las Loicas (one specimen), Las Alcantarillas (one specimen), and Cerro
172 Domuyo (five specimens).

173 *3.1.1. Cañada de Leiva*

174 It is a classic locality encompassing the Upper Jurassic–Lower Cretaceous transition in
175 the Neuquén Basin (35° 48' S, 69° 48' W), 6 km northeast of the Bardas Blancas
176 settlement (Dessanti, 1973; Gerth, 1925; Gulisano and Gutiérrez-Pleimling, 1994). Here
177 the Vaca Muerta Formation spans nearly 240 m and is dominated by tempestite-facies
178 deposited over a middle to proximal outer carbonate ramp setting (Kietzmann and
179 Palma, 2011; Kietzmann et al., 2018). Belemnites from this section come from up to 40
180 cm thick, very consolidated hummocky cross-bedding bioclastic packstone limestones
181 (CLBe) intercalated with laminated marls/siltstone intervals, and a decimetric massive
182 to laterally concretionary wackestone limestone with ammonoids, and located about 24.5
183 m from the base of the logged section in Figure 2. The CLBe sample is a densely
184 packed, poorly sorted, and oriented (Fig. 3) packstone with micrite and organic matter.
185 The grains are 70% echinoderm fragments (Figs. 3.1–3.4), 10% of bone fragments
186 (Figs. 3.1, 3.4), 10% of assorted fragmented bioclasts, probably bivalve shells (Fig. 3.2),
187 bryozoan fragments and small gastropods (Fig. 3.4), and 10% of siliciclastic material
188 such as volcanic lithic fragments, detrital mono and polycrystalline quartz, feldspar and
189 abundant pyrite. Among the echinoderm fragments were differentiated: star-shaped
190 crinoid columnals (Fig. 3.1), cross and longitudinal sections of stems, crinoid arm
191 elements, some with lunate patterns (Figs. 3.1, 3.3), and scarce echinoid spines. Most
192 echinoid fragments are highly reworked, replaced and recrystallized, and some of them

193 have microborings (Fig. 3.4). Most of the bone fragments have evidence of microbial
194 boring activity as well. The matrix is micritic and some pore spaces are filled with
195 authigenic quartz. The belemnite-bearing interval is the same identified by Gerth in
196 1911 and 1913 (Gerth, 1925, fig. 10).

197 FIGURE 2 and FIGURE 3 AROUND HERE

198 **3.1.2. Las Loicas**

199 It is located close to the 145 National Road heading towards the Pehuenche
200 International Pass to Chile, approximately 1 km to the west of the homonymous
201 settlement (35° 47' S, 70° 09' W). The marine sediments of the Vaca Muerta Formation
202 are punctuated by several volcanic layers, some of which have provided precise U/Pb
203 absolute ages (Lena et al., 2019; Vennari et al., 2014). Belemnites from this section
204 were retrieved from decimetric to metric laminated to massive dark wackestones
205 bearing ammonoid and brachiopod impressions, intercalated with dark laminated marls.
206 Based on the facies associations interpreted by Kietzmann et al. (2021) in the nearby
207 Las Tapaderas section, this interval is assignable to the transition between basinal to
208 distal outer ramp settings.

209 **3.1.3. Las Alcantarillas**

210 Located about 20 km west of the Las Loicas section, it is well exposed over the right
211 margin of the Arroyo Pehuenche that crosses 145 National Road (35° 57' S, 70° 14' W).
212 The single belemnite rostrum from this locality comes from a 12 cm thick, dark
213 laminated wackestone limestone showing a thin pyritized band, which is intercalated
214 within an interval dominated by calcareous-nodule bearing dark laminated marls.
215 Likewise, the previous locality, the facies association indicates a distal outer ramp
216 setting (Kietzmann et al., 2021).

217 **3.1.4. Cerro Domuyo**

218 This locality is at the foothills of the Cerro Domuyo, at the southern tip of the Cordillera
219 del Viento (36° 41' S, 70° 26' W). Here the Vaca Muerta Formation is exposed over the
220 left margin of the Arroyo Covunco, and all belemnites were recorded from centimetric
221 to decimetric grey to dark laminated wackestones, intercalated between large intervals
222 of laminated shales and marls. Facies association analyses indicate a basinal to distal
223 outer ramp setting for this interval, occurring just below slumped calcareous sandstone
224 deposits assigned to the Huncal Member of the Vaca Muerta Formation by Kietzmann
225 and Vennari (2013) and reinterpreted by Naipauer et al. (2020).

226 **3.2. Measurements and illustrations**

227 Belemnite rostra's study and description follow the outlines of Stevens (1965) and
228 Challinor (1999, 2003). Challinor (1999) redefined the flattening index (A) of Stevens
229 (1965) as a percentage ratio:

$$230 \quad A = \text{rostrum transverse diameter} / \text{rostrum sagittal diameter} \times 100$$

231 Rostra linear measurements were acquired using a vernier digital caliper (± 1 mm
232 accuracy). Angular measurements were obtained with a protractor ($\pm 1^\circ$) or by digital
233 means. Pictures of the belemnite rostra covered with sublimated ammonium chloride
234 were taken with a Canon SX60 semi-reflex camera. Linear illustrations and plates were
235 prepared with CorelDraw 2020 Graphics Suite and Adobe Photoshop 2020.

236 **3.3. Abbreviations**

237 ***Institutional abbreviations.*** **BAS KG**, British Antarctic Survey Collection, Cambridge,
238 United Kingdom; **CPBA**, Paleontological Collection of the University of Buenos Aires,
239 Buenos Aires, Argentina; **MCNAM-PI**, Colección de Paleontología de Invertebrados
240 del Museo de Ciencias Naturales y Antropológicas "Juan Cornelio Moyano," Mendoza,
241 Argentina.

242 **Anatomical abbreviations.** **A**, index of flattening; **dtM**, maximum transverse diameter;
 243 **dtm**, minimum transverse diameter.

244 **Stratigraphical abbreviations.** **BB**, Cañada de Leiva section; **DO**, Cerro Domuyo
 245 section; **LA**, Las Alcantarillas section; **LL**, Las Loicas section.

246 **4. Systematic paleontology**

247 Order BELEMNOPSEINA Jeletzky, 1965

248 Family BELEMNOPSEIDAE Naef, 1922, amend Jeletzky, 1946

249 Genus *Belemnopsis* Bayle, 1878

250 **Type species.** *Belemnopsis apiciconus* Blainville, 1827 (pl. 2, figs. 2, 2a). ?Upper
 251 Aalenian–Bajocian, England, France, and Southern Germany (*cf.* Riegraf, 1999).
 252 Lectotype subsequently designated by Riegraf (1999, p. 66). A reassessment of the
 253 validity of the genus was recently undertaken by Mitchell (2015) and followed here.

254 ***Belemnopsis* sp. cf. *Belemnopsis patagoniensis* (Favre, 1908)**

255 Figures 4.1–4.6; 5.1–5.2

256 *cf.* 1908 *Belemnites patagoniensis* n. sp., Favre, p. 640, pl. 37, figs. 6–7

257 **Material.** Six specimens. One from the Las Alcantarillas section (MCNAM-PI 24881);
 258 five fragments which probably correspond to five individuals from the Cerro Domuyo
 259 section (CPBA 22924.1–22924.6).

260 **Geographic and stratigraphic occurrence.** Las Alcantarillas section, bed LA7, base of
 261 *Corongoceras alternans* Biozone (upper Tithonian) and Cerro Domuyo section, bed DO
 262 30 and located within the upper portion of *Windhauseniceras internispinosum* Biozone
 263 (upper lower Tithonian), Mendoza Province, Argentina.

264 **Description.** Based on the best-preserved specimen MCNAM-PI 24881 (Figs. 4.1–4.3;
265 5.1–5.2). Rostrum cylindroconical, elongated and slender, 98 mm long, only preserves
266 the apical portion and part of the stem region. The maximum transverse diameter (dtM=
267 14 mm) is attained at about 60 mm from the apex. Outline symmetrical, non-hastate
268 with margins tapering into an acute dorsally oriented apex. Profile somewhat
269 asymmetrical, dorsally flattened but slightly ventrally inflated around the apical region.
270 Cross-section elliptical and depressed throughout (A c. 115 over the stem region),
271 although this character might be accentuated by sedimentary flattening. A narrow,
272 shallow ventral groove with angular margins runs over the mid-region of the stem and
273 extends posteriorly at least up to 18 mm before the apex. The groove is 1 mm deep over
274 the apical region, where it shows a "v" profile in cross-section and appears to deepen
275 anteriorly. The apical line is slightly ventral situated, and no lateral lines were observed.

276 **Remarks.** The Tethyan genus *Belemnopsis* (Bayle, 1878) groups more than one
277 hundred European and Indo-Pacific belemnite species ranging from the Middle Jurassic
278 up to the Lower Cretaceous (Riegraf, 1999). Although we agree with Riegraf (1980),
279 Howlett (1989), and Ippolitov et al. (2015) that such a broad taxonomic concept is
280 problematic, there is still no agreement among belemnite workers on the validity of the
281 proposed subgeneric categories (*e.g.*, Challinor, 1991; Challinor and Hikuroa, 2007) and
282 the poor preservation state of our material does not enable further insight on the matter,
283 so we follow the classic taxonomic assignation to the genus *Belemnopsis*.

284 The outline and profile general forms, as well as the length and distinctiveness of
285 the ventral groove, are important taxonomic characters (Stevens, 1965) that justify the
286 assignation of the studied material to *Belemnopsis*. The rostrum cross-section appears to
287 be a more variable trait since different authors have described *Belemnopsis* species with
288 either circular to elliptical cross-sections (*e.g.*, Doyle and Kelly, 1988; Howlett, 1989;

289 Challinor and Hikuroa, 2007). More specifically, the elongated rostrum, with an acute
290 apex and a narrow ventral groove with angular margins, enables comparing the
291 Neuquén Basin material with *Belemnopsis patagoniensis* (Favre, 1908), a well-known
292 Lower Cretaceous species frequently recorded in the Austral Basin from the Berriasian
293 to the Hauterivian (see section 1.2). Upper Tithonian (cf. Leanza, A., 1967) *B.*
294 *patagoniensis* described by Feruglio (1936) from Santa Cruz differ from the type
295 material of the species (Stevens, 1965) and has been repeatedly reassigned (see section
296 1.2). Independently of their systematic position, Feruglio's *Belemnopsis* and *Hibolites*
297 *argentinus* show morphological affinities with Indonesian and New Zealand
298 assemblages cf. Stevens (1965) and to elements from Madagascar, cf. Challinor (1991).
299 Nonetheless, Patagonian specimens are typically more robust, almost twice wider and
300 longer than the material described here (e.g., Favre, 1908, pl. 37, figs. 6–7; Riccardi,
301 1977, figs. 6a–c, g–i). A similar slender Berriasian specimen was described from
302 Crimea by Ippolitov et al. (2015), who assigned it to *Parabelemnopsis* cf. *patagoniensis*
303 and interpreted its record in the Northern Hemisphere as a singular short invasion
304 episode. The Crimean specimen differs from the Neuquén Basin material in having a
305 much shorter ventral groove. Antarctic *Belemnopsis*, on the other hand, seem to be
306 closely related to those from New Zealand-Indonesian of the "*uhligi* Complex"
307 (Stevens, 1965; Challinor, 1991), though more recently Challinor and Hikuroa (2007)
308 indicated some affinities with Madagascan forms.

309 *Belemnopsis patagoniensis*, *B. madagascariensis* (Besairie, 1930), and *B.*
310 *casterasi* Besairie, 1936 are recognized as a cluster of closely related Madagascan-
311 South American Late Jurassic–Early Cretaceous large-sized, narrow-grooved
312 belemnites (Stevens, 1965; Howlett, 1989). Howlett (1989), Challinor (1991), and
313 Ippolitov et al. (2015) have pointed out the strong similitude of some specimens

314 assigned to *B. patagoniensis*, *B. aff. madagascariensis* (Riccardi, 1977, 1988) and *B.*
315 *madagascariensis* (Aguirre-Urreta and Suarez, 1985). Even more, Combémoré (1988)
316 suggested that *B. madagascariensis* and *B. casterasi* might conform to a dimorphic pair.
317 Phylogenetic relationships between them await a detailed revision to assess the extent of
318 ontogenetic and population variability (Challinor, 2003; Challinor and Hikuroa, 2007).

319 FIGURE 4 AND FIGURE 5 AROUND HERE

320 ***Belemnopsis* sp. cf. *Belemnopsis launceloti* Howlett, 1989**

321 Figures 4.7–4.13; 5.3–5.4

322 **Material.** 14 specimens from the Cañada de Leiva section (MCNAM-PI 24647.1–
323 24647.14).

324 **Geographic and stratigraphic occurrence.** Cañada de Leiva section, bed BB19, upper
325 *Corongoceras alternans* Biozone (upper Tithonian), Mendoza Province, Argentina.

326 **Description.** Based on the most complete specimen, MCNAM-PI 24647.1, unless
327 otherwise stated. MCNAM-PI 24647.1 is most probably a juvenile individual due to the
328 presence of larger and stouter specimens in the assemblage (Figs. 4.10–4.13; 5.3–5.4).
329 Moderately preserved, 119 mm long, slender (nearly 22 times longer than thicker),
330 symmetrical, and markedly hastate rostrum both in outline and profile. The maximum
331 transverse diameter (dtM= 9.84 mm) is attained over the apical region, at about 25 mm
332 from the apex, while the minimum transverse diameter (dtm= 5.4 mm) is placed over
333 the broken anterior portion where no traces of the alveolus is observed. Apex
334 moderately obtuse. Cross-section circular to slightly depressed over the apical and
335 middle-stem region, where the venter is almost flattened, and more compressed to the
336 anterior-stem region where the venter is more excavated. A medium-width, shallow
337 ventral groove with rounded and weakly pronounced edges extends up to two-thirds

338 over the stem. In a probable adult specimen (MCNAM-PI 24647.2; Figs. 4.7–4.9), the
339 groove still has rounded margins, but it is broader (3 mm wide) and runs into the apical
340 region, at least up to 30 mm from the apex. Specimen MCNAM-PI 24647.2 also shows
341 a slightly asymmetrical profile due to inflation of the venter, and the outline appears to
342 be cyliandroconical (dtM= 12.2 mm) with a circular cross-section (A= 100). No dorsal or
343 lateral grooves are observed in any of the specimens. The rest of the rostra recorded in
344 the same bed, 15.4 mm long on average (dtM= 1.4), are too poorly preserved to provide
345 any further insights into ontogenetic traits.

346 **Remarks.** The studied specimens were retrieved from the same interval where Gerth
347 (1925) collected some material which was then revised by Stolley and identified as
348 *Hibolithes* sp. (cf. Gerth, 1925, p. 42, footnote). However, some rostra traits, like the
349 slender and long stem region and the shallow, medium to broad ventral groove, make
350 them akin to *Belemnopsis launceloti* Howlett (1989, p. 40, pl. 6, figs. 1–2, 6–7; pl. 8,
351 figs. 5–6) from the Alexander Island. Still, the Antarctic material is less hastate, has a
352 maximum transverse diameter over the middle stem region, and the ventral groove
353 narrows anteriorly (pictures of the type material of *B. launceloti* are available at the
354 BAS website:
355 [https://legacy.bas.ac.uk/bas_research/data/access/fossildatabase/filter.php?taxonKeywor](https://legacy.bas.ac.uk/bas_research/data/access/fossildatabase/filter.php?taxonKeyword=BELEMNITIDA)
356 [d=BELEMNITIDA](https://legacy.bas.ac.uk/bas_research/data/access/fossildatabase/filter.php?taxonKeyword=BELEMNITIDA)). Some of the mentioned differences might have an ontogenetic
357 explanation. Stevens (1965, p. 40) indicated that *Belemnopsis* suffered changes in its
358 degree of hastation in the early growth stages of the rostrum. Challinor (2001, p. 234)
359 also pointed out that *Belemnopsis aucklandica* presents the point of maximum
360 transverse diameter nearer the apex in juvenile specimens but higher up in adults, a trait
361 that affects the degree of hastation of the rostrum. He also noted that in more hastated
362 individuals, the ventral groove tends to weaken at about the mid-rostrum and extend

363 apically as a wide shallow depression. Unfortunately, the poor preservation state of the
 364 material hinders further insight into its ontogeny, so its systematic assignation is left
 365 open.

366 Howlett (1989) included in the *B. launceloti* synonymy list two specimens
 367 described by Feruglio (1936, pl. 10, figs. 3–4) from the Austral Basin, formerly
 368 reassigned to *B. gladiatoris* by Willey (1973), both of which are more robust than the
 369 Neuquén Basin material. Howlett (1989), also renamed as *B. launceloti* a specimen
 370 from Madagascar figured as *B. africanus* Tate, 1867 by Besairie (1936, p. 23, figs. 6–7).

371 Genus *Hibolithes* Montfort, 1808

372 **Type species.** *Hibolithes hastatus* Montfort, 1808 (text-fig. 47), by monotypy.
 373 Bajocian–Aptian of some Boreal and most Tethyan regions (Doyle and Kelly, 1988;
 374 Howlett, 1989).

375 *Hibolithes argentinus?* (Feruglio, 1936)

376 Figure 4.14–4.16

377 **Material.** One fragmentary specimen from the Las Loicas section (CPBA 23224).

378 **Geographic and stratigraphic occurrence.** Las Loicas section, bed LL062–063, upper
 379 *Windhausenicerias internispinosum* Biozone (upper lower Tithonian), Mendoza
 380 Province, Argentina.

381 **Description.** Rostrum fragment of 27.68 mm of maximum length, probably hastate both
 382 in outline and profile. The maximum transverse diameter (dtM= 12.2 mm) is recorded
 383 over the broken anterior, and the cross-section is almost circular (A= 103). No ventral
 384 or lateral grooves are observed. The apical line seems ortholineate.

385 **Remarks.** The circular form of the rostrum cross-section and the absence of a ventral
386 groove in the preserved portion leads us to consider this fragment as part of the lower
387 stem region of an *Hibolithes* representative. The most frequent *Hibolithes* species in
388 Argentina is *H. argentinus* (Feruglio, 1936), which is characterized by a slender hastate
389 rostrum with circular to slightly depressed transverse sections and a shallow ventral
390 groove that extends no further than the upper stem region. Howlett (1989, p. 52)
391 designated one of the specimens figured by Feruglio (1936, pl. 10, fig. 17) as lectotype
392 and added new material from Alexander Island. The fragment here described agrees
393 well both with the lower stem regions of the lectotype and the with the specimen BAS
394 KG 2909.18 figured by Howlett (1989, pl. 6, figs. 4, 10), but as no other features are
395 preserved, the assignation to *H. argentinus* is stated with doubts.

396 **5. Discussion**

397 **5.1. Paleobiogeographic considerations**

398 Several authors differentiated an Indo-Pacific Province within the Tethyan Realm from
399 the Kimmeridgian onwards over the basis of belemnite geographic distribution,
400 particularly of *Belemnopsis* and *Hibolithes* regional assemblages (*e.g.*, Combémoré,
401 1988; Doyle and Howlett, 1989; Mutterlose, 1986, 1992, Stevens, 1965, 1973).
402 Alternatively, Challinor (1991; Challinor in Grant-Mackie et al., 2000) proposed a
403 different subdivision of the Tethyan Realm into two provinces, 1) Tethyan Province,
404 including a Mediterranean, Ethiopian and Tethyan Subprovinces (the last partly
405 equivalent to the Indo-Pacific Province), and 2) South Pacific Province, extending along
406 the Pacific coasts of Gondwana, from southern South America (including the Neuquén
407 Basin) through West Antarctica to New Zealand, and probably including New
408 Caledonia. The South Pacific Province was not completely isolated from the Tethyan
409 Province and at least from the latest Jurassic, some belemnite migrations took place

410 from the Ethiopian Subprovince into Antarctica and South America (Challinor, 1991;
411 Mutterlose, 1992). From the Oxfordian to the Valanginian, a long-ranging lineage of
412 narrowly-grooved *Belemnopsis* developed in Madagascar and Eastern Africa, forming
413 the *B. madagascariensis*-*B. casterasi* group (Besairie, 1930; Stevens, 1965;
414 Combémoré, 1988). Some components of that cluster, as well as *Hibolithes*, might
415 have reached the South Pacific coast from the Tithonian onwards through the Trans-
416 Gondwana or Indo-Madagascan seaway (*e.g.*, Cecca, 1999; Mutterlose, 1986; Fig. 6).
417 Even more, during the Berriasian, that marine connection is supposed to have held a
418 "relict" population of *Belemnopsis* of the *madagascariensis-casterasi* group, since
419 *Belemnopsis* is unknown or poorly represented from European and most Indo-Pacific
420 regions at that time (Doyle and Howlett, 1989; Stevens, 1965). That remnant population
421 is thought to be related to the *Belemnopsis patagoniensis* assemblages of the Austral
422 Basin (Riccardi, 1977, 1988) and now from the upper Tithonian in the Neuquén Basin.
423 The same is valid for *B. launceloti*, recorded in Madagascar, Alexander Island, and in
424 the Austral Basin. *Belemnopsis* records here described add new elements connecting the
425 Neuquén Basin assemblages with those of Madagascar and Eastern Africa, passing
426 through Antarctica during the Upper Jurassic. These basins are known to share several
427 ammonoid genera among other marine invertebrate and vertebrate taxa (Mutterlose,
428 1986; Riccardi, 1991; Vennari and Aguirre-Urreta, 2019 and references therein;
429 Campos et al., 2021).

430 FIGURE 6 AROUND HERE

431 **5.2. Biostratigraphic considerations**

432 Belemnites' biostratigraphic potential has long been acknowledged for Jurassic and
433 Cretaceous successions, though belemnite-based correlations are more widely used for
434 the Cretaceous of the Northern Hemisphere, as summarized by Christensen (1990) and

435 Mutterlose (1990). Doyle and Bennet (1995) complemented those contributions by
436 reviewing Jurassic belemnite occurrences of the Southern Hemisphere, and then Doyle
437 et al. (1996, 1997) focused on the Antarctic and the Argentinean Jurassic records. The
438 former includes the Kimmeridgian–Tithonian *Belemnopsis* fauna described from the
439 Antarctic Peninsula and Alexander Island (Crame and Howlett, 1988; Howlett, 1989;
440 Mutterlose, 1986; Riley et al., 1997). Upwards, the Tithonian/Berriasian transition is
441 encompassed by the *Hibolithes* fauna (or Biozone according to Howlett, 1989), which
442 includes several species of *Hibolithes*, with *H. argentinus*, and *Belemnopsis* species of
443 Indo-Pacific affinities as rare components. Tithonian Antarctic belemnite associations
444 also include *Duvalia/Produvalia* (Mutterlose, 1986; Crame and Howlett, 1988), a
445 strongly compressed genus with a dorsal median groove that is supposed to have
446 immigrated into the Indo-Pacific region from the Tethys during the Tithonian (Stevens,
447 1965) or as early as the Callovian (Challinor and Hikuroa, 2007).

448 Concerning Argentinean Jurassic successions, Doyle et al. (1996) recognized two
449 successive belemnite faunas for the Upper Jurassic–Lower Cretaceous namely early
450 *Belemnopsis* (Oxfordian) and *Hibolithes* (Tithonian–Berriasian), which was named after
451 *H. argentinus*. In that contribution, the Lower Cretaceous occurrences of *B.*
452 *patagoniensis* were treated separately. Meanwhile, *B. launzeloti* is an upper Tithonian–
453 Hauterivian species that defines an assemblage biozone in Antarctica (Howlett, 1989)
454 and has been recorded from the lower Valanginian of Madagascar (Besairie, 1936;
455 Combémoré, 1988), the upper Tithonian of the Austral Basin in the Lago Argentino
456 area (*cf.* Leanza, A., 1967), and with reservations from the Valanginian–Hauterivian of
457 southern Chile (Stinnesbeck et al., 2014). So, *Belemnopsis* records described in this
458 contribution indicate that this genus should also be considered in the Neuquén Basin as

459 a component of the well-established Upper Jurassic–Lower Cretaceous *Hibolithes* fauna
460 of Gondwana (Doyle et al., 1996; Challinor, 1991).

461 Whenever possible, the correlation of the Neuquén Basin belemnite records in
462 each locality with the current ammonoid Andean zonation (Aguirre-Urreta et al., 2019;
463 Leanza et al., 2020; Lescano et al., 2022) is summarized in Figure 7. *Hibolithes* sp. from
464 the Arroyo La Manga area was found by Gerth (1925) in beds assigned to the
465 *Argentiniceras noduliferum* Biozone, just above an interval bearing *Lytrochoplites*
466 *burckhardti* (Mayer-Eymar) and *Spiticeras acutum* Gerth among other ammonoids of
467 the *Substeueroceras koeneni* Biozone. *Hibolithes* sp. reported from Pampa Tril were
468 retrieved from the *Virgatosphinctes andesensis/Pseudolissoceras zitteli*,
469 *Windhauseniceras internispinosum*, and *Corongoceras alternans* Biozones, according
470 to Alberti et al. (2020). *Belemnopsis* cf. *B. patagoniensis* from Cerro Domuyo were
471 found in two positions within the *W. internispinosum* Biozone, near its base (bearing *W.*
472 *internispinosum* and *Corongoceras lotenoense* Spath) and loosed near its top. From the
473 upper portion of the *W. internispinosum* Biozone as well, in Las Loicas, were recorded
474 three successive beds with fragmentary *Hibolithes argentinus?* rostra associated with
475 *Catutosphinctes rafaeli* Leanza, H. and Zeiss. In Las Alcantarillas, *Belemnopsis* cf. *B.*
476 *patagoniensis* was retrieved from a bed with *Himalayites concurrens* Leanza, A. and
477 *Micracanthoceras?* sp. at the base of the *Corongoceras alternans* Biozone. From the
478 same locality was recorded a poorly preserved *Hibolithes* sp. from the *A. noduliferum*
479 Biozone, just below a bed with *Andiceras fallax* (Steuer) and *Substeueroceras*
480 *disputabile* Leanza, A. Finally, at Cañada de Leiva, *Belemnopsis* cf. *B. launzeloti* was
481 recorded from the top of the *C. alternans* Biozone, from an interval sandwiched
482 between "*Berriasella*" *bardensis* Krantz (= *Blanfordiceras bardense* in Parent et al.,
483 2011), "*Thurmanniceras discoidalis*" (Gerth) (= *Parodontoceras calistoides* in Parent et

484 al. 2011) and *Substeueroceras striolatissimum* (Steuer) occurrences, being the later an
485 index species of the *S. koeneni* Biozone across the basin (e.g., Groeber et al., 1952;
486 Vennari and Aguirre-Urreta, 2019).

487 FIGURE 7 AROUND HERE

488 **5.3. Paleoecological, paleogeographical and paleoenvironmental considerations.**

489 In stark contrast with the fossil record of the Upper Jurassic and, particularly, the Lower
490 Cretaceous successions of the Austral Basin, belemnite occurrences in time-equivalent
491 units of the Neuquén Basin are quite infrequent or even surprisingly absent from the
492 Valanginian onwards, both in shallow and deep facies (Aguirre-Urreta et al., 2008).
493 Here we discuss paleogeographical, paleoenvironmental, and paleoecological factors
494 concerning belemnite occurrences in the Neuquén Basin.

495 Episodes of transgressive or high sea level could have facilitated the access of
496 active swimming belemnites (or aid in the dispersing of their early ontogenetic stages)
497 to the Neuquén Basin. Belemnite records from Cerro Domuyo, Las Alcantarillas, and
498 Las Loicas were found in middle outer ramp to basinal deposits of a transgressive
499 system tract of the second-order Second Composite Depositional sequence, involving
500 the lower portion of the *W. internispinosum* Biozone to the lower portion of the *S.*
501 *koeneni* Biozone (Kietzmann et al., 2014, 2021). Meanwhile, belemnites from Cañada
502 de Leiva come from a storm-dominated middle ramp setting, corresponding to a
503 highstand system tract (Kietzmann et al., 2014, 2018). However, other factors than a
504 high local sea level alone would have influenced belemnite distribution.

505 Although some authors have modeled the existence of counter-oceanic currents at
506 the latitude of the Neuquén Basin (Parrish, 1992), up to the authors' best knowledge,
507 there were no physical geographic barriers between the Austral and the Neuquén Basin
508 that prevented a free interchange of nektonic fauna as both were retro-arc basins open to

509 the Paleo-Pacific. Indeed, both basins share several ammonoid genera (Riccardi, 1991),
510 although comparatively poorly preserved and less abundant in the case of the former
511 (Riccardi, 2008). Nevertheless, two alternative paleogeographic models for the western-
512 margin configuration of the Neuquén Basin were proposed, either enabling a direct
513 connection with the Paleopacific Ocean through a discontinuous arc of volcanic islands
514 (e.g., Howell et al., 2005) or a more restricted seaway in the north-western sector: the
515 Curepto Strait (Vicente et al., 2005). Nevertheless, the high diversity and Tethyan
516 affinities of several Neuquén Basin ammonoids and herpetofauna militate against the
517 possibility of a confined basin (Gasparini and Fernández, 2005; Riccardi, 2015).
518 Besides, Alberti et al. (2020) recently demonstrated, over Sr/Ca ratios and their
519 comparison with the global curve, the existence of water exchange with the open ocean
520 in several localities of the Neuquén Province.

521 Regarding average seawater temperature and salinity, Alberti et al. (2020) found
522 some strong influence of rainfall and/or a continental freshwater influx in their
523 northernmost sampled sites, and interpreted a water temperature of 20–25 °C from $\delta^{18}\text{O}$
524 data from oyster and brachiopod shells. That temperature range agrees well with
525 previous estimations by Gómez Dacal (2018) and Gómez Dacal et al. (2019) of a
526 medium water temperature of 23 °C from oysters of the Neuquén Basin, and 24 °C from
527 belemnites (*Belemnopsis* sp.) of the Austral Basin. Temperatures obtained in both cases
528 are in agreement with the supposed range of 10–30 °C that belemnites inhabited by
529 comparison with modern cephalopod preferences (Hoffmann and Stevens, 2020),
530 though we do not know to which extent minor temperature changes affected belemnite
531 distribution since it is considered the main parameter controlling their migration routes
532 at species and genus levels (e.g., Mutterlose, 1988; Mutterlose et al., 2022; Stevens,
533 1973). Other paleoecologic requirements listed by Hoffmann and Stevens (2020), such

534 as well-oxygenated waters, with a normal salinity of 27–37 psu and a depth range of no
535 more than 200 m would characterize the Vaca Muerta Formation Upper Jurassic
536 carbonate ramp (*e.g.*, Kietzmann et al., 2014). Even if, locally, some of the parameters
537 would deviate from the optimum (*e.g.*, by the influx of freshwater), belemnites could
538 have just actively swum away from such conflicting areas.

539 In most recent revisions of the group, belemnites are considered medium-sized
540 predators of Mesozoic food webs, most likely feeding upon small vertebrates, mollusks,
541 and crustaceans (Hoffman and Stevens, 2020 and references therein). All these food
542 sources were present at the Neuquén Basin, and possibly some v-shaped scars over
543 ammonoid shells were caused by non-lethal belemnite attacks (Vennari, 2011). In time,
544 belemnites were occasionally preyed upon by marine reptiles such as ichthyosaurs and
545 metriorhynchids (Fernández et al., 2019), which might have gotten into the basin mostly
546 for reproductive purposes (Gasparini and Fernández, 2005). However, most probably,
547 belemnite hatchlings were consumed by pachycormids, a stem-teleost clade of
548 suspensivorous-feeding fishes that were quite successful from the Early Jurassic to the
549 Late Cretaceous (Friedman et al., 2010). Pachycormids, which are well-known from the
550 Neuquén Basin (Gouiric-Cavalli, 2017; Gouiric-Cavalli and Arratia, 2022), to date have
551 not been recorded in the Austral Basin, though they have cited from Chile (Arratia,
552 2015). Although pachycormid gill rakers spacing was suitable to filter out both
553 hatchlings and juvenile ammonites (Tajika et al., 2018), the different reproductive
554 strategies of both cephalopod groups may have turned the odds in favor of ammonite
555 survival. On the one hand, ammonites, either semelparous or iteroparous, are assumed
556 to have been extreme r-strategists, with high fecundity rates and the possibility, for
557 medium-sized macroconchs, to lay more than 10,000 eggs (De Baets et al., 2015;
558 Mironenko and Rogov, 2015; Tajika et al., 2018; Greif et al., 2022). Even more,

559 ovoviviparity has been proven for some ammonites (Mironenko and Rogov, 2015), thus
560 incrementing the chances of post-hatching survival. Nevertheless, direct evidence of this
561 predation relation should be proven through bromalite analyses (*i.e.*, oral or anal
562 depositions or intestinal content) as conducted in other basins (Zatón and Salamon,
563 2008; Bigurrarena et al., 2019; Lukeneder et al., 2020). Belemnites, on the other hand,
564 are assumed to have been r-strategists (e.g., Doguzhaeva et al. 2013) and to have
565 depicted high fecundity rates as well, but of several magnitude order lower than that of
566 ammonites due to a smaller size difference between adults and embryos (Tajika et al.,
567 2018). In consequence, ammonite surviving into adulthood was more likely than for
568 belemnites.

569 To sum up, marine connections, seawater physical parameters, or food sources
570 availability seem not to have constrained, on their own, the presence of belemnites in
571 the Neuquén Basin in favor of ammonites. However, they were likely affected by the
572 predation pressure exerted by filter-feeding organisms on hatchlings, combined with
573 unsuitable paleoecological conditions over shallow marine settings relatively close to
574 the coast where belemnites were assumed to inhabit (Naef, 1922; Doguzhaeva et al.,
575 2013, Mutterlose, 2022).

576 **6. Concluding remarks**

577 We presented in this contribution new records of Upper Jurassic belemnites from the
578 Vaca Muerta Formation, Neuquén Basin, west-central Argentina. The studied rostra
579 were assigned to *Belemnopsis* cf. *B. patagoniensis*, *Belemnopsis* cf. *B. launceloti*, and
580 *Hibolithes argentinus?*. These are the first thoroughly described belemnites from the
581 Neuquén Basin. By their association with ammonoid faunas, the new belemnite records
582 can be referred to the Tithonian *Windhausenicerias internispinosum* to *Corongoceras*
583 *alternans* Biozones. The biostratigraphic framework suggests that *Belemnopsis* should

584 be contemplated as a component of the widely-known *Hibolithes* fauna recorded across
585 the Late Jurassic–Early Cretaceous of Gondwana. Affinities of the described fauna seem
586 close to Madagascan–Eastern African belemnite associations of the Ethiopian
587 Subprovince, thus reinforcing the hypothesis of a Trans-Gondwana seaway as an open
588 marine connection at least since the Tithonian.

589 The baffling scarcity of belemnites from the Late Jurassic in the Neuquén Basin and
590 their absence from the Early Cretaceous upwards is in utter contrast with their abundant
591 record in the Austral Basin. Though still unresolved, it is suggested that a combination
592 of factors, such as local variations in seawater salinity/temperature in shallow
593 environments and, particularly, the predation pressure of filter-feeding fishes on
594 belemnite hatchlings, may have affected the presence of the group in the basin. Also,
595 the possible existence of counter-currents at the latitude of the Neuquén Basin may have
596 limited the extension to lower latitudes of the *Belemnopsis/Hibolithes* Gondwanan
597 faunas.

598 **Acknowledgments**

599 The authors are grateful to all field assistants, to Leonardo Pazo (Fundación Azara) for
600 preparing the specimens, and to the editors and reviewers that greatly improved this
601 contribution. This work was partially financed by the ANPCyT PICT 1016-2016 and
602 PICT 2020-02067 (V.V.V.); Sepkoski Grant 2019 (V.V.V.); PIP CONICET
603 11220170101117CO (B.A.U.) and 11220200102844CO (V.V.V.); CNRS Grants
604 SYSTER-TELLUS “ACJC” in 2019 and “GC-JC” in 2021; and MITI AAP Temps
605 “AstroCarb” (M.M. and P.P.). This is contribution R-445 of the Instituto de Estudios
606 Andinos “Don Pablo Groeber”.

607 **References**

- 608 Aguirre-Urreta, B., Naipauer, M., Lescano, M., López-Martínez, R., Pujana, I., Vennari,
609 V., Concheyro, A., De Lena, L.F., Ramos, V. 2019. The Tithonian chrono-
610 biostratigraphy of the Neuquén Basin and related Andean areas: a review and
611 update. *Journal of South American Earth Sciences*, 92, 350–367.
612 <https://doi.org/10.1016/j.jsames.2019.03.019>
- 613 Aguirre-Urreta, M.B., Casadío, S., Cichowolski, M., Lazo, D.G., Rodriguez, D.L.,
614 2008. Afinidades paleobiogeográficas de los invertebrados cretácicos de la
615 Cuenca Neuquina. *Ameghiniana*, 45, 593–613.
- 616 Aguirre-Urreta, M.B., Suárez, M., 1985. Belemnites de una secuencia turbidítica
617 volcanoclástica de la Formación Yaghán, Titoniano–Cretácico Inferior del
618 extremo sur de Chile. 4° Congreso Geológico Chileno, Antofagasta, Actas 1, 1–
619 16.
- 620 Alberti, M., Parent, H., Garrido, A.C., Andersen, N., Garbe-Schönberg, D., Danise, S.,
621 2020. Stable isotopes ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$) and element ratios (Mg/Ca, Sr/Ca) of Jurassic
622 belemnites, bivalves and brachiopods from the Neuquén Basin (Argentina):
623 challenges and opportunities for palaeoenvironmental reconstructions. *Journal of*
624 *the Geological Society*, 178, JGS2020-163. <https://doi.org/10.1144/jgs2020-163>
- 625 Arratia, G. 2015. Los peces osteíctios fósiles de Chile y su importancia en los contextos
626 paleobiogeográfico y evolutivo. *Publicación Ocasional del Museo de Historia*
627 *Natural, Chile*, 63, 35–83.
- 628 Bayle, E., 1878. Atlas. Première partie. Fossiles principaux des terrains. –Explication de
629 la Carte géologique de la France, Paris, 4, part 1, 158 plates.
- 630 Besairie, H., 1930. Recherches géologiques a Madagascar. *Bulletin de la Société d'*
631 *Histoire naturelle de Toulouse*, 60, 345–642.

- 632 Besairie, H., 1936. Recherches géologiques a Madagascar. Première suite. La géologie
633 du Nord-Ouest. *Memoires de L'Académie Malgache*, 21, 145–148.
- 634 Bigurrarena Ojeda, M.A., Gouiric-Cavalli, S., Pérez, L.M., Burlaille, L.A., Moly, J.J.,
635 Reguero, M., Santillana, S. 2019. Las bromalitas de la Formación Ameghino
636 (Jurásico Tardío) de la Península Antártica. *Publicación Electrónica de la*
637 *Asociación Paleontológica Argentina. Suplemento Resúmenes*, R17.
- 638 Blainville, H.D., 1827. *Memoire sur les belemnites, considerees zoologiquement et*
639 *geologiquement*. F. G. Levrault, Paris, 136 pp.
- 640 Bonarelli, G., Nágera, J.J., 1961. Observaciones geológicas en las inmediaciones del
641 Lago San Martín. *Boletín de la Dirección Nacional de Minería y Geología*, B27,
642 1–39.
- 643 Caminos, R., Nullo, F., 1979. Descripción geológica de la Hoja 67e, Isla de los Estados.
644 *Servicio Geológico Nacional, Boletín*, 175, 5–52.
- 645 Campos, L., Fernández, M.S., Herrera, Y., Talevi, M., Concheyro, A., Gouiric-Cavalli,
646 S., O'Gorman, J.P., Santillana, S.N., Acosta-Burlaille, L., Moly, J.J., Reguero,
647 M.A., 2020. Bridging the southern gap: First definitive evidence of Late Jurassic
648 ichthyosaurs from Antarctica and their dispersions routes. *Journal of South*
649 *American Earth Sciences*, 109, 103259.
650 <https://doi.org/10.1016/j.jsames.2021.103259>
- 651 Cecca, F., 1999. Palaeobiogeography of Tethyan ammonites during the Tithonian (latest
652 Jurassic). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 147, 1–37.
653 [https://doi.org/10.1016/S0031-0182\(98\)00149-7](https://doi.org/10.1016/S0031-0182(98)00149-7)

- 654 Challinor, A.B., 1991. Belemnite successions and faunal provinces in the southwest
655 Pacific, and the belemnites of Gondwana. *BMR Journal of Australian Geology*
656 *and Geophysics*, 12, 301–325.
- 657 Challinor, A.B., 1999. Belemnite biostratigraphy of the New Zealand Late Jurassic
658 Mangaoran (Early Puarooan) Substage and the Puarooan Stage revisited, New
659 Zealand *Journal of Geology and Geophysics*, 42, 369–393.
660 <https://doi.org/10.1080/00288306.1999.9514852>
- 661 Challinor, A.B., 2001. Stratigraphy of Tithonian (Ohauan-Puarooan) marine beds near
662 Port Waikato, New Zealand, and a redescription of *Belemnopsis aucklandica*
663 (Hochstetter). *New Zealand Journal of Geology and Geophysics*, 44, 219–242.
664 <https://doi.org/10.1080/00288306.2001.9514936>
- 665 Challinor, A.B., 2003. Synonymy and stratigraphic ranges of *Belemnopsis* in the
666 Heterian and Ohauan Stages (Callovian-Tithonian), southwest Auckland, New
667 Zealand, *New Zealand Journal of Geology and Geophysics*, 46, 79–94.
668 <https://doi.org/10.1080/00288306.2003.9514997>
- 669 Challinor, A.B., Hikuroa, D.C.H., 2007. New Middle and Upper Jurassic Belemnite
670 Assemblages from West Antarctica (Latady Group, Ellsworth Land): Taxonomy
671 and Paleobiogeography. *Palaeontologia Electronica*, 10, 1–29. [http://palaeo-](http://palaeo-electronica.org/paleo/2007_1/assemblage/index.html)
672 [electronica.org/paleo/2007_1/assemblage/index.html](http://palaeo-electronica.org/paleo/2007_1/assemblage/index.html)
- 673 Christensen, W.K., 1990. Upper Cretaceous belemnite biostratigraphy of Europe.
674 *Cretaceous Research*, 11, 371–386. [https://doi.org/10.1016/S0195-](https://doi.org/10.1016/S0195-6671(05)80048-5)
675 [6671\(05\)80048-5](https://doi.org/10.1016/S0195-6671(05)80048-5)
- 676 Combémoré, R., 1988. Les bélemnites de Madagascar. *Documents de la Laboratoire de*
677 *Géologie de la Faculté des Sciences de Lyon*, 104, 1–239.

- 678 Crame, J.A, Howlett, P.J., 1988. Late Jurassic and Early Cretaceous biostratigraphy of
679 the Fossil Bluff Formation, Alexander Island. *British Antarctic Survey Bulletin*,
680 78, 1–35.
- 681 Dana, J.D., 1848. *Geology*. In *U.S. Exploring Expedition 1839–1842*, under Ch.
682 Wilkes. Chapters 15–16. New York Public Library, New York.
- 683 De Baets, K., Landman, N.H., Tanabe, K. 2015. Ammonoid embryonic development.
684 In: Klug, C., Korn, D., De Baets, K., Kruta, I. & Mapes, R.H. (eds), *Ammonoid*
685 *paleobiology: from anatomy to paleoecology*. 113–205 Dordrecht: Springer.
- 686 Dessanti, R.N., 1973. Descripción geológica de la Hoja 29b, Bardas Blancas, provincia
687 de Mendoza. *Servicio Nacional de Minería y Geología, Boletín*, 139, 1–70.
- 688 Doguzhaeva, L.A., Weis, R., Delsate, D., Mariotti, N. 2013. Embryonic shell structure
689 of Early–Middle Jurassic belemnites, and its significance for belemnite expansion
690 and diversification in the Jurassic. *Lethaia*, 47, 49–65.
- 691 Doyle, P., 1987. Lower Jurassic–Lower Cretaceous belemnite biogeography and the
692 origin of the Mesozoic Boreal Realm. *Palaeogeography, Palaeoclimatology,*
693 *Palaeoecology*, 61, 237–254.
- 694 Doyle, P., Bennet, M.R., 1995. Belemnites in biostratigraphy. *Palaeontology*, 38, 815–
695 829.
- 696 Doyle, P., Howlett, P., 1989. Antarctic belemnites biogeography and the break-up of
697 Gondwana. In: J.A. Crame (ed.), *Origins and Evolution of the Antarctic Biota*,
698 *Geological Society Special Publication*, 47, pp. 167–182.
699 <https://doi.org/10.1144/GSL.SP.1989.047.01.13>
- 700 Doyle, P., Kelly, S.R.A., 1988. The Jurassic and Cretaceous belemnites of Kong Karls
701 land, Svalbard. *Norsk Polarinstitut Skrifter*, 189, 1–77.

- 702 Doyle, P., Kelly, S.R.A., Pirrie, D., Riccardi, A.C., 1997. Jurassic belemnite distribution
703 patterns: implications of new data from Antarctica and Argentina. *Alcheringa: An*
704 *Australasian Journal of Paleontology*, 21, 219–228.
705 <http://dx.doi.org/10.1080/03115519708619175>
- 706 Doyle, P., Kelly, S.R.A., Pirrie, D., Riccardi, A.C., Olivero, E.B., 1996. Jurassic
707 belemnite biostratigraphy of the Southern Hemisphere: a comparative study from
708 Antarctica and Argentina. *Revista de la Asociación Geológica Argentina*, 51,
709 331–338.
- 710 Doyle, P., Pirrie, D., 1999. Belemnite distribution patterns. In F. Olóriz, F.J.,
711 Rodríguez-Tovar (eds), *Advancing Research on Living and Fossil Cephalopods*
712 pp. 419–436. Kluwer Academic-Plenum Publishers, New York.
- 713 Favre, F., 1908. Die Ammoniten der unteres Kreide Patagonien. *Neues Jahrbuch für*
714 *Mineralogie, Geologie und Paläontologie*, 25, 601–647.
- 715 Fernández, M., Herrera, Y., Vennari, V., Campos, L., de la Fuente, M., Talevi, M.,
716 Aguirre-Urreta, M.B. 2019. Marine reptiles from the Jurassic/Cretaceous
717 transition at the High Andes, Mendoza, Argentina. *Journal of South American*
718 *Earth Sciences*, 92, 658–673. <https://doi.org/10.1016/j.jsames.2019.03.013>
- 719 Feruglio, E., 1936. *Paleontographica Patagónica*. Memoria del Instituto Geológico de
720 Padova, 11, 1–384.
- 721 Feruglio, E., 1949. *Geología de la Patagonia*. Vol. 1, pp. 1–323. Dirección General de
722 Y.P.F., Buenos Aires.
- 723 Friedman, M., Shimada, K., Martin, L.D., Everhart, M.J., Liston, J., Maltese, A.,
724 Triebold, M. 2010. 100-million year dynasty of giant planktivorous bony fishes in
725 the Mesozoic seas. *Science* 327: 990-993.

- 726 Fuchs, D., Iba, Y., Ifrim, C., Nishimura, T., Kennedy, W.J., Keupp, H., Stinnesbeck,
727 W., Tanabe, K., 2013. *Longibelus* gen. nov., a new Cretaceous coleoid genus
728 linking Belemnnoidea and early Decabrachia. *Palaeontology*, 56, 1081–1106.
729 <https://doi.org/10.1111/pala.12036>
- 730 Gasparini, Z., Fernández, M.S., 2005. Jurassic marine reptiles of the Neuquén Basin:
731 records, faunas and their palaeobiogeographic significance. In: G.D. Veiga, L.A.
732 Spalletti, J.A. Howell, E. Schwarz (eds.), *The Neuquén Basin: A case study in*
733 *sequence stratigraphy and basin dynamics*, Geological Society, London, Special
734 Publication 252, 279–294. <https://doi.org/10.1144/GSL.SP.2005.252.01.14>
- 735 Gerth, H., 1925. Contribuciones a la estratigrafía y la paleontología de los Andes
736 Argentinos II. La Fauna Neocomiana de la Cordillera Argentina en la parte
737 meridional de la provincia de Mendoza: Academia Nacional de Ciencias,
738 Córdoba, Actas, 9, 57–132.
- 739 Gómez Dacal, A.R., 2018. Petrología, geoquímica y diagénesis de sedimentitas y fósiles
740 carbonáticos del Jurásico Superior–Cretácico Inferior de las cuencas Neuquina y
741 Austral, Argentina. PhD Thesis (unpublished). Universidad de La Plata, La Plata,
742 Argentina.
- 743 Gómez Dacal, A.R., Richiano, S.M., Gómez Peral, L.E., Spalletti, L.A., Sial, A.N.,
744 Poiré, D.G., 2019. Evidence of Warm Seas in High Latitudes of Southern South
745 America During the Early Cretaceous. *Cretaceous Research*, 95, 8–20.
746 <https://doi.org/10.1016/j.cretres.2018.10.021>.
- 747 Gouiric-Cavalli, S., 2017. Large and mainly unnoticed: the first lower Tithonian record
748 of a suspension-feeding pachycormid from Southern Gondwana. *Ameghiniana*,
749 54, 283–289. <https://doi.org/10.5710/AMGH.16.11.2016.3029>

- 750 Gouiric-Cavalli, S., Arratia G., 2022. A new Pachycormiformes (Actinopterygii) from
751 the Upper Jurassic of Gondwana sheds light on the evolutionary history of the
752 group, *Journal of Systematic Palaeontology*, DOI:
753 [10.1080/14772019.2022.2049382](https://doi.org/10.1080/14772019.2022.2049382)
- 754 Grant-Mackie, J.A., Aita, Y., Balme, B.E., Campbell, H.J., Challinor, A.B., MacFarlan,
755 D.A.B., Molnar, R.E., Stevens, G.R., Thulborn, R.A., 2000. Jurassic
756 palaeobiogeography of Australasia. *Memoir of the Association of Australasian*
757 *Palaeontologists*, 23, 311–353.
- 758 Greif, M., Nebelsick, J.H., and Klug, C. 2022. Extreme abundance of ammonoids in
759 mass accumulations from the Late Devonian of the Moroccan Anti-Atlas. *Acta*
760 *Palaeontologica Polonica* 67: 667–684. <https://doi.org/10.4202/app.00935.2021>
- 761 Groeber, P., Stipanovic, P.N., Mingramm, A., 1952. Jurásico. In: *Geografía de la*
762 *República Argentina*, tomo 2. Sociedad Argentina de Estudios Geográficos,
763 Buenos Aires, pp. 143–347.
- 764 Gulisano, C.A., Gutiérrez-Pleimling, A.R., 1994. The Jurassic of the Neuquén Basin.
765 *Field Guide, Mendoza Province. Asociación Geológica Argentina, Serie E*, 3,
766 *Secretaría de Minería de la Nación, Publicación*, 159, 1-103.
- 767 Harrington, H.H., 1943. Observaciones geológicas en la Isla de los Estados. *Anales del*
768 *Museo Argentino de Ciencias Naturales*, 49, *Geología* 29, 29–52.
- 769 Hoffmann, R., Richter, D.K., Neuser, R.D., Jöns, N., Linzmeier, B.J., Lemanis, R.E.,
770 Füsseis, F., Xiao, X., Immenhauser, A., 2016. Evidence for a composite organic-
771 inorganic fabric of belemnite rostra: implications for palaeoceanography and
772 palaeoecology. *Sedimentary Geology*, 341, 203–215.
773 <https://doi.org/10.1016/j.sedgeo.2016.06.001>

- 774 Hoffmann, R., Stevens, K., 2020. The palaeobiology of belemnites –foundation for the
775 interpretation of rostrum geochemistry. *Biological Reviews*, 95, 94–123.
776 <https://doi.org/10.1111/brv.12557>
- 777 Howell, J.A., Schwarz, E., Spalletti, L.A., Veiga, G.D., 2005. The Neuquén Basin: an
778 overview. In: G.D. Veiga, L.A. Spalletti, J.A. Howell, E. Schwarz (eds.), *The*
779 *Neuquén Basin: A case study in sequence stratigraphy and basin dynamics*,
780 Geological Society, London, Special Publication, 252, 1–14.
781 <https://doi.org/10.1144/GSL.SP.2005.252.01.01>
- 782 Howlett, P.J., 1989, Late Jurassic–Early Cretaceous cephalopods of eastern Alexander
783 Island, Antarctica. *Special Papers in Palaeontology*, 41, 5–72.
- 784 Iba, Y., Sano, S.I., Mutterlose, J., 2014. The early evolutionary history of belemnites:
785 new data from Japan. *PLoSOne*, 9, e95632.
786 <https://doi.org/10.1371/journal.pone.0095632>
- 787 Ippolitov, A.P., Desai, B., Arkadiev, V.V., 2015. First find of *Parabelemnopsis*, the
788 alien belemnite from the Southern Hemisphere, in the Upper Berriasian of Central
789 Crimea. *The International Scientific Conference on the Jurassic/Cretaceous*
790 *Boundary, Proceedings volume, Samara*, 52–59.
- 791 Jeletzky, J.A., 1965. Taxonomy and phylogeny of fossil Coleoidea (= Dibranchiata).
792 Geological Survey of Canada, paper 65-2, 72–76.
- 793 Jeletzky, J.A., 1946. Zur Kenntnis der oberkretazischen Belemniten. *Geologiska*
794 *Föreningens i Stockholm Förhandlingar*, 68, 87–105.
- 795 Jeletzky, J.A., 1966. *Comparative morphology, phylogeny, and classification of fossil*
796 *coleoidea*. *The University of Kansas Paleontological Contributions, Mollusca*
797 *Article 7*, pp. 1–162. University of Kansas Paleontological Institute, Lawrence.

- 798 Jeletzky, J.A., 1983. Macroinvertebrate paleontology, biochronology, and
799 paleoenvironments of Lower Cretaceous and Upper Jurassic rocks, Deep Sea
800 Drilling Project Hole 511, Eastern Falkland Plateau. DSDP report 35, 951–975.
801 <https://doi.org/10.2973/dsdp.proc.71.135.1983>
- 802 Kietzmann, D.A., Vennari, V.V., 2013. Sedimentología y estratigrafía de la Formación
803 Vaca Muerta (Tithoniano-Berriasiano) en el área del cerro Domuyo, norte de
804 Neuquén, Argentina. *Andean Geology*, 40, 41–65.
805 <http://dx.doi.org/10.5027/andgeoV40n1-a02>
- 806 Kietzmann, D.A., Iglesia Llanos, M.P., Palacio, J.P., Sturlesi, M.A., 2021. Facies
807 analysis and stratigraphy across the Jurassic-Cretaceous boundary in a new
808 basinal Tithonian–Berriasian section of the Vaca Muerta Formation, Las
809 Tapaderas, Southern Mendoza Andes, Argentina. *Journal of South American
810 Earth Sciences*, 109, 103267. <https://doi.org/10.1016/j.jsames.2021.103267>
- 811 Kietzmann, D.A., Iglesia-Llanos, M.P., Martínez, M.K., 2018. Astronomical calibration
812 of the Tithonian–Berriasian in the Neuquén Basin, Argentina: A contribution from
813 the Southern Hemisphere to the Geologic Time Scale. In: M. Montenari (ed.),
814 *Stratigraphy & Timescales 3*, Elsevier, 328–355.
815 <https://doi.org/10.1016/bs.sats.2018.07.003>
- 816 Kietzmann, D.A., Palma, R.M., 2011. Las tempestitas peloidales de la Formación Vaca
817 Muerta (Tithoniano-Valanginiano) en el sector surmendocino de la Cuenca
818 Neuquina, Argentina. *Latin American Journal of Sedimentology and Basin
819 Analysis*, 18, 121–149.
- 820 Kietzmann, D.A., Palma, R.M., Riccardi, A.C., Martín-Chivelet, J., López-Gómez, J.,
821 2014. Sedimentology and sequence stratigraphy of a Tithonian–Valanginian
822 carbonate ramp (Vaca Muerta Formation): a misunderstood exceptional source

- 823 rock in the southern Mendoza area of the Neuquén basin, Argentina. *Sedimentary*
824 *Geology*, 302, 64–86. <http://dx.doi.org/10.1016/j.sedgeo.2014.01.002>
- 825 Krantz, F., 1928. La Fauna del Tithono superior y medio en la parte meridional de la
826 provincia de Mendoza. Academia Nacional de Ciencias, Córdoba, *Actas*, 10, 9–
827 57.
- 828 Leanza, A.F., 1967. Anotaciones sobre los fósiles Jurásico-Cretácicos de Patagonia
829 Austral (Colección Feruglio) conservados en la Universidad de Bologna. *Acta*
830 *Geológica Lilloana*, 9, 121–186.
- 831 Leanza, H.A., Vennari, V.V., Aguirre-Urreta, B., Concheyro, A., Lescano, M., Ivanova,
832 D., Kietzmann, D.A., López-Martínez, R., Martz, P.A., Paolillo, M.A., Guler, V.,
833 Pujana, I., Paz, M., 2020. Relevant marine paleobiological markers of the Vaca
834 Muerta Formation. *AAPG Memoir*, 121, 61–98. DOI:
835 [10.1306/13682224M1203830](https://doi.org/10.1306/13682224M1203830)
- 836 Legarreta, L., Uliana, M.A., 1991. Jurassic–Cretaceous marine oscillations and
837 geometry of back-arc basin fill, central Argentine Andes. *International*
838 *Association of Sedimentologists, Special Publication*, 12, 429–450.
839 <https://doi.org/10.1002/9781444303896.ch23>.
- 840 Lena, L., López-Martínez, R., Lescano, M., Aguirre-Urreta, B., Concheyro, A., Vennari,
841 V., Naipauer, M., Samankassou, E., Pimentel, M., Ramos, V.A., Schaltegger, U.,
842 2019. High-precision U-Pb ages in the early Tithonian to early Berriasian and
843 implications for the numerical age of the Jurassic–Cretaceous boundary. *Solid*
844 *Earth*, 10, 1–14. <https://doi.org/10.5194/se-10-1-2019>
- 845 Lescano, M., Vennari, V., López-Martínez, R., Concheyro, A., Aguirre-Urreta, B.,
846 Pellenard, P., Martínez, M. 2022. New Tithonian biostratigraphic data from the

- 847 Neuquén Basin, Argentine Andes: a contribution from the Southern Hemisphere.
848 11^o International Congress on the Jurassic System. Budapest, p. 68.
- 849 Lukeneder, A., Surmik, D., Gorzelak, P., Niedźwiedzki, P., Brachaniec, T., Salamon,
850 M. 2020. Bromalites from the Upper Triassic Polzberg section (Austria); insights
851 into trophic interactions and food chains of the Polzberg palaeobiota. Nature
852 Scientific Reports 10, 20545. <https://doi.org/10.1038/s41598-020-77017-x>
- 853 Medina, F.A., Riccardi, A.C., 2005. Desmoceratidae, Silesitidae and Kossmaticeratidae
854 (Ammonitida) from the Upper Aptian–Albian of Patagonia (Argentina). Revue de
855 Paléobiologie, Genève, 24, 251–286.
- 856 Mironenko, A.A., Rogov, M.A. 2015. First direct evidence of ammonoid ovoviviparity.
857 Lethaia, 49: 245–260. <https://doi.org/10.1111/let.12143>.
- 858 Mitchell, F.S., 2015. A reassessment of the validity and affinities of *Belemnites sulcatus*
859 Miller, 1826, *Belemnopsis* Edwards in Gray, 1849, and *Belemnopsis* Bayle, 1878.
860 Carnet de Géologie, 15, 31–39. <https://doi.org/10.4267/2042/56399>
- 861 Monks, N., Hardwick, J.D., Gale, A.S., 1996. The function of the belemnite guard.
862 Paläontologische Zeitschrift, 70, 425–431.
- 863 Montfort, D. De, 1808. Conchyliologie systematique et classification methodique des
864 Coquilles. Coquilles univalves, cloisonnees. Tome I. Chez F. Schoell, Paris, 410
865 pp.
- 866 Mutterlose, J., 1986. Upper Jurassic Belemnites from the Orville Coast, Western
867 Antarctica, and their palaeobiogeographical significance. British Antarctic Survey
868 Bulletin, 70, 1–22.

- 869 Mutterlose, J., 1988. Migration and evolution patterns in Upper Jurassic and Lower
870 Cretaceous belemnites. In Wiedmann, J., Kullmann, J. (eds) *Cephalopods: Present*
871 *and past*. Schweizerbart, Stuttgart, pp. 525–537.
- 872 Mutterlose, J., 1990. A belemnite scale for the Lower Cretaceous. *Cretaceous Research*,
873 11, 1–15. [https://doi.org/10.1016/S0195-6671\(05\)80039-4](https://doi.org/10.1016/S0195-6671(05)80039-4)
- 874 Mutterlose, J., 1992. Early Cretaceous belemnites from the East Indian Ocean and their
875 paleobiogeographic implications. *Proceedings of the Ocean Drilling Program*,
876 *Scientific Results*, 123, 443–450.
877 <https://doi.org/10.2973/odp.proc.sr.123.123.1992>
- 878 Mutterlose, J., Picollier, M.-C., Dzyuba, O. 2022. The first belemnite of boreal ancestry
879 from the Early Cretaceous (Valanginian) of the western Tethys: implications for
880 belemnite ecology. *Papers in Paleontology*, e1455.
- 881 Naef, A., 1922. *Die fossilen Tintenfische*. Gustav Fischer, Jena, 322 pp.
- 882 Naipauer, M., Comerio, M., Lescano, M., Vennari, V., Aguirre-Urreta, M.B., Pimentel,
883 M., Ramos, V.A., 2020. The Huncal Member of the Vaca Muerta Formation,
884 Neuquén Basin of Argentina: insight into biostratigraphy, structure, U-Pb detrital
885 zircon, ages and provenance. *Journal of South American Earth Sciences*, 100,
886 102567. <https://doi.org/10.1016/j.jsames.2020.102567>
- 887 Olivero, E.B., 1987. Cefalópodos y bivalvos titonianos y hauterivianos de la Formación
888 Lago La Plata, Chubut. *Ameghiniana* 24, 181–202.
- 889 Parent, H., Garrido, A.C., Scherzinger, A., Schweigert, G., Fözy, I., 2015. The
890 Tithonian–Lower Valanginian stratigraphy and ammonite fauna of the Vaca
891 Muerta Formation in Pampa Tril, Neuquén Basin, Argentina. *Boletín del Instituto*
892 *de Fisiografía y Geología*, 86, 1–96.

- 893 Parent, H., Scherzinger, A., Schweigert, G., 2011. The Tithonian–Berriasian ammonite
894 fauna and stratigraphy of Arroyo Cieneguita, Neuquén- Mendoza Basin,
895 Argentina. *Boletín del Instituto de Fisiografía y Geología*, 79–81, 21–94.
- 896 Parent, H., Schweigert, G., Scherzinger, A., Garrido, A.C., 2017. Additional Tithonian
897 and Berriasian ammonites of the Vaca Muerta Formation in Pampa Tril, Neuquén
898 Basin, Argentina. *Volumina Jurassica*, 15, 139–154.
899 <https://doi.org/10.5604/01.3001.0010.6576>
- 900 Parrish, J.T., 1992. Jurassic climate and oceanography of the Pacific region. In: G.E.G.
901 Westermann (ed.), *The Jurassic of the Circum-Pacific*, Cambridge University
902 Press, Cambridge, pp. 365–379.
- 903 Riccardi, A.C., 1971. Estratigrafía en el Oriente de la Bahía de La Lancha, Lago San
904 Martín, Santa Cruz, Argentina. *Revista del Museo de la Plata, Sección Geología*,
905 7, 245–318.
- 906 Riccardi, A.C., 1976. Paleontología y edad de la Formación Springhill. 1° Congreso
907 Geológico Chileno, Santiago de Chile, Actas 1, 41–56.
- 908 Riccardi, A.C., 1977. Berriasian invertebrate faunas from the Springhill Formation in
909 southern Patagonia. *Neues Jahrbuch für Geologie und Paläontologie*, 155, 216–
910 152.
- 911 Riccardi, A.C., 1988. The Cretaceous System of southern South America. *Geological*
912 *Society of America Memoir*, 168, 1–161.
- 913 Riccardi, A.C., 1991. Jurassic and Cretaceous marine connections between the
914 Southeast Pacific and Tethys. *Palaeogeography, Palaeoclimatology,*
915 *Palaeoecology*, 87, 155–189. [https://doi.org/10.1016/0031-0182\(91\)90134-D](https://doi.org/10.1016/0031-0182(91)90134-D)

- 916 Riccardi, A.C., 2008. El Jurásico de la Argentina y sus amonites. *Revista de la*
917 *Asociación Geológica Argentina*, 63, 625–643.
- 918 Riccardi, A.C., 2015. Remarks on the Tithonian–Berriasian ammonite biostratigraphy
919 of west-central Argentina. *Volumina Jurassica*, 13, 23–52.
- 920 Riccardi, A.C., Damborenea, S.E., Manceñido, M.O., Leanza, H.A., 2011.
921 Megainvertebrados jurásicos y su importancia geobiológica. In: H.A. Leanza, C.
922 Arregui, O. Carbone, J.C. Danieli, J.M. Vallés (eds.), *Geología y Recursos*
923 *Naturales de la Provincia del Neuquén*, Asociación Geológica Argentina, Buenos
924 Aires, p. 441–464.
- 925 Riegraf, W., 1980. Revision der Belemniten der Schwäbischen Jura. Part 7. *Palaeonto-*
926 *graphica Abteilung A*, 169, 128–206.
- 927 Riegraf, W., 1999. Taxonomic status of the belemnite genus *Belemnopsis* Bayle 1878.
928 *Paläontologische Zeitschrift*, 73, 59–76.
- 929 Riley, T.R., Crame, J.A., Thomson, M.R.A., Cantrill, D.J., 1997. Late Jurassic
930 (Kimmeridgian–Tithonian) macrofossil assemblage from Jason Peninsula, Graham
931 Land: evidence for a significant northward extension of the Latady Formation.
932 *Antarctic Science*, 9, 434–442.
- 933 Smith, A.G., Smith, D.G., and Funnell, B.M., 1994. *Atlas of Mesozoic and Cenozoic*
934 *Coastlines*. Cambridge, Cambridge University Press, 99 p.
- 935 Stevens, G.R., 1965. The Jurassic and Cretaceous belemnites of New Zealand, and a
936 review of the Jurassic and Cretaceous belemnites of the Indo-Pacific region.
937 *Geological Survey of New Zealand, Palaeontological Bulletin*, 36, 1–233.

- 938 Stevens, G.R., 1973. Jurassic belemnites. In: A. Hallam (ed.), Atlas of
939 Palaeobiogeography, Elsevier, New York, pp. 259–274.
- 940 Stevens, K., in press. Belemnite phylogeny and decline during the mid-Cretaceous.
941 BioRxiv. <https://doi.org/10.1101/2021.10.11.463885>
- 942 Stinnesbeck, W., Frey, E., Rivas, L., Pardo Pérez, J., Leppe Cartes, M., Salazar Soto,
943 C., Zambrano Lobos, P., 2014. A Lower Cretaceous ichthyosaur graveyard in
944 deep marine slope channel deposits at Torres del Paine National Park, southern
945 Chile. *Geology Society of America Bulletin*, 126, 1317–1339.
946 <https://doi.org/10.1130/B30964.1>
- 947 Stolley, E., 1911. Beiträge zur Kenntnis der Cephalopoden der Norddeutschen Unteren
948 Kreide 1: Die Belemniten der Norddeutschen Unteren Kreide. *Geologische und*
949 *paläontologische Abhandlungen: Neue Folge*, 10: 201–272.
- 950 Stolley, E., 1928. *Heliceras Dana als Erhaltungszustand von Belemnopsis*. *Neues*
951 *Järbuch für Mineralogie, Geologie und Paläontologie, Beilage Band B*, 60, 315–
952 324.
- 953 Tajika, A., Nützel, A., Klug, C., 2018. The old and the new plankton: ecological
954 replacement of associations of mollusc plankton and giant filter feeders after the
955 Cretaceous? *PeerJ* 6: e4219. <https://doi.org/10.7717/peerj.4219>
- 956 Tate, R., 1867. On some secondary fossils from South Africa. *Quarterly Journal of the*
957 *Geological Society of London*, 23, 139–175.
- 958 Vennari, V., 2011. Estudio preliminar del registro de heridas letales y subletales en
959 amonoides del Jurásico Tardío (Tithoniano) de Mendoza. 18° Congreso
960 Geológico Argentino, Neuquén, Actas, 1494-1495.

- 961 Vennari, V.V., Aguirre-Urreta, M.B., 2019. Intraspecific variability, biostratigraphy and
 962 paleobiological significance of the Southern Gondwana ammonoid genus
 963 *Lytohoplites* Spath. *Journal of Paleontology*, 93, 702–726.
 964 <https://doi.org/10.1017/jpa.2019.1>.
- 965 Vennari, V.V., Lescano, M.A., Naipauer, M., Aguirre-Urreta, M.B., Concheyro, A.,
 966 Shaltegger, U., Armstrong, R., Pimentel, M., Ramos, V.A., 2014. New constraints
 967 in the Jurassic/Cretaceous boundary in the High Andes using high precision U-Pb
 968 data. *Gondwana Research*, 26, 374–385. <https://doi.org/10.1016/j.gr.2013.07.005>
- 969 Vicente, J.C., 2005. Dynamic paleogeography of the Jurassic Andean Basin: patterns of
 970 transgression and localization of main straits through the magmatic arc. *Revista de*
 971 *la Asociación Geológica Argentina*, 60, 221–250.
- 972 Weaver, C., 1931. Paleontology of the Jurassic and Cretaceous of West Central
 973 Argentina. *Memoirs of the University of Washington*, 1, 1–496.
- 974 Willey, I.E., 1973. Belemnites from South-Eastern Alexander Island. II, The occurrence
 975 of the Family Belemnopseidae in the Upper Jurassic and Lower Cretaceous.
 976 *Bulletin of the British Antarctic Survey*, 36, 33–59.
- 977 Zátón, M., Salamon, M. 2008. Durophagus predation on middle Jurassic Molluscs, as
 978 evidenced from shell fragmentation. *Palaeontology* 51, 1, 63–70.
 979 <https://doi.org/10.1111/j.1475-4983.2007.00736.x>

980 **Figure captions**

981 **Figure 1. 1.1.** Upper Jurassic–Lower Cretaceous outcrops in the Neuquén and Austral
 982 Basin, Argentina, with referred (in bold) and studied (circled) belemnite bearing
 983 localities: **BN**, Bahía Nassau; **CL**, Cañada de Leiva; **CM**, Chacay Melehue; **DO**, Cerro
 984 Domuyo; **IE**, Isla de los Estados; **LA**, Las Alcantarillas; **LAR**, Lago Argentino; **LB**,

985 Lago Belgrano; **LL**, Las Loicas; **LO**, Cerro Lotena; **LSM**, Lago San Martín; **1.2–1.4**.
 986 Detailed maps of the studied belemnite-bearing localities, **1.2**. Cañada de Leiva; **1.3**.
 987 Las Loicas and Las Alcantarillas; **1.4**. Cerro Domuyo.

988 **Figure 2.** Log sections of the belemnite-bearing localities indicating the stratigraphic
 989 position of belemnite records and selected ammonoid taxa.


990 **Figure 3.** Hummocky cross-bedding bioclastic packstone (**CLBe**) thin section, views of
 991 the densely packed poorly sorted and oriented packstone with micrite and organic
 992 matter. **1**, In the center, star-shaped crinoid columnals (**O**) in between other highly
 993 fragmented echinoderm fragments with micrite cement, and bone fragment (**B**) with
 994 silica replacement. **2**, Note orientated fragmented bioclasts, probably bivalve shells
 995 (white arrows), bone fragment (**B**) and echinoderm stems. **3**, Different sections of stems
 996 (**S**) in a micritic matrix (**M**). **4**, Bone fragments (**B**), small gastropods and, (**G**) pyrite
 997 with abundant stems fragments (**S**). Scales bars= 2 mm (**1, 2**), 1 mm (**3**); 0.5 mm (**4**).


998 **Figure 4.** Belemnites from the Vaca Muerta Formation, Neuquén Basin. **1–6**,
 999 *Belemnopsis* cf. *B. patagoniensis* (Favre, 1908). **1–3**, MCNAM-PI 24881, **1**, ventral
 1000 view, **2**, right lateral view, **3**, cross-section; **4–6**, CPBA 22924.6, **4**, ventral view, **5**, left
 1001 lateral view, **6**, cross-section. **7–13**, *Belemnopsis* cf. *B. launzeloti* Howlett, 1989. **7–9**,
 1002 MCNAM-PI 24647.2, **7**, right lateral view, **8**, ventral view, **9**, cross-section; **10–13**,
 1003 MCNAM-PI 24647.1, **10**, ventral view, **11**, left lateral view, **12**, upper stem cross-
 1004 section, **13**, lower stem cross-section. **14–16**, *Hibolithes argentinus?* CPBA 23224, **14**,
 1005 outline view, **15**, lateral view, **16**, cross-section. All specimens were covered with
 1006 ammonium chloride. Scale bars= 10 mm.





1007 **Figure 5.** *Belemnopsis* rostra sketch diagrams. **1–2**, *Belemnopsis* cf. *B. patagoniensis*
 1008 (Favre, 1908), MCNAM-PI 24881, **1**, outline; **2**, right profile; **3–4**, *Belemnopsis* cf. *B.*
 1009 *launzeloti* Howlett, 1989, MCNAM-PI 24647.1, **3**, left profile, **4**, outline.


1010 **Figure 6.** Paleobiogeographic distribution of *Belemnopsis* and *Hibolithes* during the
 1011 upper Tithonian–lower Berriasian. Data from Favre (1908), Besairie (1930, 1936),
 1012 Feruglio (1936-1937, 1949), Harrington (1943), Stevens (1965, and references therein),
 1013 Willey (1973), Riccardi (1977), Jeletzki (1983), Combémorrel (1988), Crame and
 1014 Howlett (1988), Howlett (1989), Doyle and Howlett (1989, and references therein),
 1015 Aguirre-Urreta and Suárez (1985, and references therein), Mutterlose (1986, and
 1016 references therein), Olivero (1987), Challinor (1991, and references therein), Challinor
 1017 and Hikuroa (2007), Crame *et al.* (2000), and Stinnesbeck *et al.* (2014). Map based on
 1018 Smith *et al.* (1994). **1**, Neuquén Basin; **2**, Lago San Martín, Austral Basin; **3**, Lago
 1019 Argentino, Austral Basin; **4**, Torres del Paine, Chile; **5**, Tierra del Fuego Archipelago);
 1020 **6**, Alexander Island, Antarctic Peninsula; **7**, Malvinas/Falkland Plateau; **8**, Orville Coast
 1021 and Ellsworth Land, Western Antarctica; **9**, Eastern Wedell Sea; **10**, Madagascar; **11**,
 1022 Kutchh; **12**, Northwestern Pakistan; **13**, Spiti Shales; **14**, Australia; **15**, New Guinea; **16**,
 1023 Moluccas and Sula Islands; **17**, Misool, Indonesia; **18**, New Zealand.


1024 **Figure 7.** Summary of occurrences of the Tithonian–Berriasian Neuquén Basin
 1025 belemnite records according to the current Andean ammonoid-based biostratigraphic
 1026 scheme (modified from Aguirre-Urreta *et al.*, 2019; Leanza *et al.*, 2020, Lescano *et al.*,
 1027 *in press*). **White stars**, *Hibolithes* sp. records (Gerth, 1925; Alberti *et al.*, 2020); **grey**
 1028 **stars**, *Hibolithes argentinus* and *H. argentinus?* records (this work); **black stars**,
 1029 *Belemnopsis* cf. *B. patagoniensis* records (this work). **DO**, Cerro Domuyo; **CL**, Cañada
 1030 de Leiva; **LA**, Las Alcantarillas; **LL**, Las Loicas; **LM**, Arroyo La Manga; **PT**, Pampa
 1031 Tril.














AGE		BIOZONE	OCCURRENCES					
			LM	CL	LL	LA	DO	PT
Berriasian	Late	?						
	Mid.	<i>Argentinerias noduliferum</i>	☆			☆		
	Early	?						
Tithonian	Late	<i>Substeueroceras koeneni</i>		★				
		<i>Corongoceras alternans</i>						☆
		?			★	★		
		<i>W. internispinosum</i>			★		★	
		?						☆
		<i>A. proximus</i>						
	Early	<i>Pseudolissoceras zitteli</i>						
	<i>Virgatosphinctes andesensis</i>							
	?							

Highlights

- New Tithonian belemnite data from the Neuquén Basin, western Argentina.
- 21 Belemnites are assigned to the genera *Belemnopsis* and *Hibolithes*.
- The belemnite association is close to Madagascan-Eastern African faunas.
- Generic composition and affinity of the belemnite fauna add new evidence supporting an open Trans-Gondwana seaway from the Tithonian onwards.

Authors statement

Verónica V. Vennari: Conceptualization, methodology, data acquisition, investigation, writing, supervision, and funding acquisition. **Beatriz Aguirre-Urreta:** investigation, data acquisition, writing, and funding acquisition. **Luciana S. Marin:** data acquisition, methodology and writing. **Pierre Pellenard:** data acquisition, writing, and funding acquisition. **Mathieu Martinez:** data acquisition, writing, and funding acquisition. **Maisa Tunik:** data acquisition and writing.

Journal Pre-proof

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof