



Improved near-infrared high-resolution solar spectrum from ACS NIR onboard TGO

Karim Gizatullin, Alexander Trokhimovskiy, Anna Fedorova, Oleg Koralev,
Franck Montmessin, Daria Betsis, Jean-Loup Bertaux, Monique Spite

► To cite this version:

Karim Gizatullin, Alexander Trokhimovskiy, Anna Fedorova, Oleg Koralev, Franck Montmessin, et al.. Improved near-infrared high-resolution solar spectrum from ACS NIR onboard TGO. European Planetary Science Congress 2018, Sep 2018, Berlin (Germany), Germany. pp.id.EPSC2018-565. insu-04001187

HAL Id: insu-04001187

<https://insu.hal.science/insu-04001187>

Submitted on 4 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Improved near-infrared high-resolution solar spectrum from ACS NIR onboard TGO

Karim Gizatullin (1,2), Alexander Trokhimovskiy (1), Anna Fedorova (1) Oleg Koralev (1), Franck Montmessin (3), Daria Betsis(1), Jean-Loup Bertaux (1,3), Monique Spite (4) , and the ACS Team

(1) Space Research Institute (IKI), Moscow, Russia, (2) Moscow Institute of Physics and Technology (MIPT), Dolgoprudny, Russia, (3) LATMOS/IPSL, UVSQ Université Paris-Saclay, UPMC Univ. Paris 06, CNRS, Guyancourt, France, (4) GEPI Observatoire de Paris, CNRS, Guyancourt, France

1 Introduction

The Atmospheric Chemistry Suite (ACS) is Russian contribution to ESA-Roscosmos ExoMars 2016 Trace Gas Orbiter (TGO) mission [1], [2]. It arrived to Mars in October 2016. ACS is a package of three high sensitive infrared spectrometers with high resolve power ($>10,000$) and cover from 0.7 to $17\mu\text{m}$ — the visible to thermal infrared range [1].

In this work we present first results for high resolution solar spectra observed by ACS NIR [3] instrument in near infrared range.

2 Observations

The ACS NIR is a near infrared spectrometer, extension of SPICAM-IR instrument family [4], the main concept of which relies on the combination of an acousto-optic tunable filter (AOTF) and an echelle diffraction grating [1]. This combination gives resolving power $\frac{\lambda}{\Delta\lambda} \approx 25,000$ in spectral range of $0.73 - 1.65\mu\text{m}$ that corresponds to the echelle diffraction orders from 48 through 105. This is the first instrument that can measure with such high resolution in near-IR range outside the atmosphere. Here we present results that were obtained in June 2016 during Mid Cruise Checkout TGO Payload (MCC) observations of Sun. The NIR solar spectrum shows some undiscovered lines and differences from the known solar spectra in the range of $1.3 - 1.5\mu\text{m}$ where the strong water absorption in the Earth atmosphere prevents from getting pure solar spectrum. The comparison with the Continuum Absorption at Visible and Infrared Wavelengths and its Atmospheric Relevance (CAVIAR) [5] solar spectrum and theoretical solar spectrum will be presented here.

References

- [1] O. Koralev, F. Montmessin, A. Trokhimovskiy, A. A. Fedorova, A. V. Shakun, A. V. Grigoriev, B. E. Moshkin, N. I. Ignatiev, F. Forget, F. Lefèvre, K. Anufreychik, I. Dzuban, Y. S. Ivanov, Y. K. Kalinnikov, T. O. Kozlova, A. Kungurov, V. Makarov, F. Martynovich, I. Maslov, D. Merzlyakov, P. P. Moiseev, Y. Nikolskiy, A. Patrakeev, D. Patsaev, A. Santos-Skipko, O. Sazonov, N. Semena, A. Semenov, V. Shashkin, A. Sidorov, A. V. Stepanov, I. Stupin, D. Timonin, A. Y. Titov, A. Viktorov, A. Zharkov, F. Altieri, G. Arnold, D. A. Belyaev, J. L. Bertaux, D. S. Betsis, N. Duxbury, T. Encrénaz, T. Fouchet, J.-C. Gérard, D. Grassi, S. Guerlet, P. Hartogh, Y. Kasaba, I. Khatuntsev, V. A. Krasnopolsky, R. O. Kuzmin, E. Lellouch, M. A. Lopez-Valverde, M. Luginin, A. Määttänen, E. Marcq, J. Martin Torres, A. S. Medvedev, E. Millour, K. S. Olsen, M. R. Patel, C. Quantin-Nataf, A. V. Rodin, V. I. Shematovich, I. Thomas, N. Thomas, L. Vazquez, M. Vincendon, V. Wilquet, C. F. Wilson, L. V. Zasova, L. M. Zelenyi, and M. P. Zorzano, “The atmospheric chemistry suite (acs) of three spectrometers for the exomars 2016 trace gas orbiter,” *Space Science Reviews*, vol. 214, p. 7, Nov 2017.
- [2] O. Koralev, A. Trokhimovsky, A. V. Grigoriev, A. Shakun, Y. S. Ivanov, B. Moshkin, K. Anufreychik, D. N. Timonin, I. Dziuban, Y. K. Kalinnikov, *et al.*, “Three infrared spectrometers, an atmospheric chemistry suite for the exomars 2016 trace gas orbiter,” *Journal of Applied Remote Sensing*, vol. 8, no. 1, p. 084983, 2014.
- [3] A. Trokhimovskiy, O. Koralev, Y. K. Kalinnikov, A. Fedorova, A. V. Stepanov, A. Y. Titov, I. Dziuban, A. Patrakeev, and F. Montmessin, “Near-infrared echelle-aotf spectrometer acs-nir for the exomars trace gas orbiter,” in *Infrared Remote Sensing and Instrumentation XXIII*, vol. 9608, p. 960809, International Society for Optics and Photonics, 2015.
- [4] O. Koralev, J.-L. Bertaux, A. Fedorova, D. Fonteyn, A. Stepanov, Y. Kalinnikov, A. Kiselev, A. Grigoriev,

- V. Jegoulev, S. Perrier, *et al.*, “Spicam ir acousto-optic spectrometer experiment on mars express,” *Journal of Geophysical Research: Planets*, vol. 111, no. E9, 2006.
- [5] K. P. Menang, M. D. Coleman, T. D. Gardiner, I. V. Ptashnik, and K. P. Shine, “A high-resolution near-infrared extraterrestrial solar spectrum derived from ground-based fourier transform spectrometer measurements,” *Journal of Geophysical Research: Atmospheres*, vol. 118, no. 11, pp. 5319–5331, 2013.