

The Carboniferous (Serpukhovian) macroflora from the "Coteaux du Pont Barré", Beaulieu-sur-Layon (Maine-et-Loire), South of the Armorican Massif, France

Christine Strullu-Derrien, Michel Ballèvre, Alan R.T. Spencer, Erwan Guillou, Jean-Paul Gislard, Hubert Lardeux, Christopher J Cleal

▶ To cite this version:

Christine Strullu-Derrien, Michel Ballèvre, Alan R.T. Spencer, Erwan Guillou, Jean-Paul Gislard, et al.. The Carboniferous (Serpukhovian) macroflora from the "Coteaux du Pont Barré", Beaulieusur-Layon (Maine-et-Loire), South of the Armorican Massif, France. Botany Letters, 2023, 170 (2), pp.183-193. 10.1080/23818107.2023.2183899 . insu-04023063

HAL Id: insu-04023063 https://insu.hal.science/insu-04023063

Submitted on 10 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Botany Letters



ISSN: (Print) (Online) Journal homepage: https://www.tandfonline.com/loi/tabg21

The Carboniferous (Serpukhovian) macroflora from the "Coteaux du Pont Barré", Beaulieu-sur-Layon (Maine-et-Loire), South of the Armorican Massif, France

Christine Strullu-Derrien, Michel Ballèvre, Alan R.T. Spencer, Erwan Guillou, Jean-Paul Gislard, Hubert Lardeux & Christopher J. Cleal


To cite this article: Christine Strullu-Derrien, Michel Ballèvre, Alan R.T. Spencer, Erwan Guillou, Jean-Paul Gislard, Hubert Lardeux & Christopher J. Cleal (2023): The Carboniferous (Serpukhovian) macroflora from the "Coteaux du Pont Barré", Beaulieusur-Layon (Maine-et-Loire), South of the Armorican Massif, France, Botany Letters, DOI: 10.1080/23818107.2023.2183899

To link to this article: https://doi.org/10.1080/23818107.2023.2183899





ARTICLE



The Carboniferous (Serpukhovian) macroflora from the "Coteaux du Pont Barré", Beaulieu-sur-Layon (Maine-et-Loire), South of the Armorican Massif, France

Christine Strullu-Derrien (Da,b), Michel Ballèvre (Dc, Alan R.T. Spencer (Db,d), Erwan Guilloue, Jean-Paul Gislardf, Hubert Lardeux⁹ and Christopher J. Cleal (1)^h

alnstitut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, Paris, France; bScience Group, The Natural History Museum, London, UK; Géosciences Rennes, Bât.15 campus Beaulieu, Université de Rennes, Rennes, France; ^aDepartment of Earth Science & Engineering, Imperial College London, London, UK; ^aChemin d'Orléans, Brissac-Loire-Aubance, France; frue Édouard Vaillant, Angers, France; Le Verger-Beaucé, Melesse, France; School of Earth Sciences, University of Bristol, Life Sciences Building, Bristol, UK

ABSTRACT

The "Coteaux du Pont Barré" in Beaulieu-sur-Layon is a Regional Natural Reserve which is home to exceptional flora and fauna. Recently fossil plants have been discovered at the site adding to its natural heritage significance. The exposure which is part of the "Sillon Houiller de la Basse-Loire" contains Serpukhovian-age (330-320 Ma) remains of sphenophytes, ferns, and lyginopteridopsid and cycadopsid pteridosperms, which are described for the first time as well as the geology of the site. Most Carboniferous macroflora previously described from the Maine-et-Loire has been based on ex situ specimens from now abandoned and inaccessible coal mines. The newly described macroflora shows evidence of in situ remains and differs somewhat in composition from those reported from the coal workings, especially in the total absence of lycopsids and Calymmotheca pteridosperms, and instead having abundant medullosalean foliage (Neuralethopteris). The co-occurrence of Sphenopteris elegans and Neuralethopteris densifolia suggests a slightly younger age compared to the macrofloras documented from the coal-bearing deposits. However, this could be an ecological consequence of the flora growing in better-drained substrate habitats.

ARTICLE HISTORY Received 21 January 2023

Accepted 16 February 2023 **KEYWORDS** Carboniferous:

Serpukhovian; coal swamps; macrofloras; Maine-et-Loire; Sillon houiller de la Basse-

1. Introduction

The coal-swamps that extended over large areas of palaeotropical Euramerica during the Pennsylvanian and Cisuralian (late Carboniferous and early Permian) was one of the most important Palaeozoic terrestrial biomes (e.g. DiMichele et al. 2001; Greb et al. 2006; Cleal et al. 2012; Cleal 2021a, b), providing the habitat in which many biotic groups diversified including insects, arachnids, reptiles and gymnosperms, and having a major impact on atmospheric composition and climate through enhanced transpiration and carbon sequestration. As a result, this biome has attracted considerable interest from geologists and palaeontologists, to try to determine its long-term impact on Earth history.

However, the earliest evolution of the coal swamp biome in the latest Mississippian (Serpukhovian) is less well understood. Palaeobotanical remains (hereafter referred to as macrofloras - see Pardoe et al. 2021) of the Serpukhovian proto-coal swamps are well preserved in the Pays-de-Loire region of northwestern France and were documented in detail notably by Bureau (1913-1914) and more recently by StrulluDerrien et al. (2021, 2022). Most of this evidence was derived from coal workings that are no longer accessible and, although many of the fossils are still available in museum geoheritage collections, it is difficult to interpret the depositional context of the macrofloras.

In this paper, we document a macroflora that was recently collected from an exposure, the "Coteaux du Pont Barré", along the Layon River in Beaulieu-sur-Layon, 20 km south of Angers (Maine-et-Loire). The site is well known, especially for its extant flora long studied by the botanist Robert Corillion (1908–1997), who first established the protection of the area (Corillion 1957). The macroflora was discovered some years ago by the winemaker who owned the site and his collaborators. Nowadays, most of the site belongs to the "Domaine Belargus", a leading biodynamic winery in the Loire Valley, and is managed for its protection by the "Ligue de Protection des Oiseaux (LPO) Anjou".

This macroflora differs in composition from those reported earlier from the coal workings in this area, which we have interpreted as representing

CONTACT Christine Strullu-Derrien 🖾 c.strullu-derrien@nhm.ac.uk 🖸 Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, UA, CP39, 57 rue Cuvier, Paris 75 005, France; Christopher J. Cleal 🔯 chris.cleal@bristol.ac.com

built upon in any way.

vegetation that grew in comparatively less welldrained habitats.

2. Geological setting

The studied outcrop is part of the "Sillon houiller de la Basse-Loire" (Bureau 1913-1914), a narrow (0.5 to 2

km) zone of Carboniferous coal-bearing outcrops extending over a length of about 100 km. This zone is bounded between two major domains, the Saint-Georges-sur-Loire Unit to the north, and the Mauges Unit to the south (Figure 1a, b; e.g. Cartier et al. 2001; Cartier and Faure 2004; Ballèvre et al. 2009; Lin et al. 2019). The deposition setting for these coal-bearing strata is interpreted as narrow pull-apart basins, deve-




Figure 1. (a) Geological map showing the location of the studied area (Beaulieu-sur-Layon). (b) Location of the area (rectangle) within the Armorican Massif (NASZ: North Armorican Shear Zone; SASZ(N) and SASZ(S): respectively, the northern and southern branches of the South Armorican Shear Zone; NSEF: Nort-sur-Erdre Fault). (c) Overview of the "Coteaux du Pont Barré" on the right bank of the Layon River (river visible on the right side). (d) the "Coteau des Treilles" vineyard. (e-g) Different views of the outcrop. Mesocalamites (cast of stem) (g), along the bedding surface of a conglomerate.

loped during dextral displacement along the crustalscale Nort-sur-Erdre Fault (Ballèvre et al. 2009).

The outcrop is located on the right bank of the Layon River, about 500 m north-west of Pont-Barré (Figures 1c, 2a). It is situated in a Regional Natural Reserve, with the majority of the outcrop located along the middle part of the slope dominating the Layon River, and above the renowned vineyard, the "Coteau des Treilles" (Figure 1d). Two main formations are seen (Blaise et al. 1986):

- (1) To the north-east, mafic schists of the Saint-Georges-sur-Loire Unit display a strongly dipping foliation to the NNE. They consist of mafic volcanics, in most cases fine-grained tuffs and breccias, sometimes containing flattened limestone clasts, up to 10-12 cm long. No fossils have been identified in these fine-grained, fully recrystallized limestones.
- (2) To the south-west of the schists are found steeply-dipping, alternating layers Carboniferous conglomerates, sandstones sometimes grading to siltstones, and less frequently shaly layers. Several layers of conglomerates may be seen. Most conglomerates grade from clast- to matrix-supported, the coarser varieties are very poorly sorted while finer layers are much better sorted. They display well-rounded, abundant pebbles of various lithologies, namely milky quartz (derived from quartz veins), rhyolites and tuffs, schists and quartzites. The size of the pebbles is smaller in the southerly layers compared to those close to the contact with the St Georges schists, where an outcrop contains pebbles up to 30 cm. The size of the pebbles, and their nature (with large amounts of phtanites, i.e. siliceous shales) suggest a rather proximal deposit, mainly derived from the northerly situated St-Georges-sur-Loire Unit, consistent with previous observations (Blaise et al. 1986). Fossil plants have been mainly found within reddish coloured sandy-to -silty layers interbedded within conglomerates (Figures. 1e-g; red star in Figure 2). These are observed in situ in the upper part of the vine-"Coteau yard slope des Treilles". A Mesocalamites (cast of stem) occurs in situ (Figure 1g) along the bedding surface of a polygenic conglomerate. This indicates transport and deposition of the plant remains by fast-flowing water (e.g. rivers).

The contact between the St-Georges-sur-Loire and the Carboniferous sediments is not well exposed in the Pont-Barré area. At a regional scale, this contact is generally considered as a major, crustal-scale, fault, as shown on the geological map of Thouarcé (Blaise et al. 1986). This fault has been called the Nort-sur-Erdre (or Layon) Fault, and displays a long, polyphase Variscan history (Cartier and Faure 2004; Ballèvre et al. 2009; Lin et al. 2019). In addition, it has been reactivated during several periods during the Mesozoic and the Cenozoic, as indicated in the studied area by the difference in altitude of the base of the transgression by Cenomanian sediments (Figure 2; Poirier 1935; Louail 1984; Blaise et al. 1986; Sellier 2020)

3. Material and methods

Specimens collected on the outcrops were borrowed from Erwan Guillou of the "Ligue de Protection des Oiseaux (LPO) Anjou" where they are currently stored. Later, they will be on exhibit at the "Domaine Belargus" winery located at Saint-Lambert-du-Lattay, Maine-et-Loire, France. Specimens under the names PalB 001 to PalB017 were photographed under natural light, using a Nikon D500 camera with a 180 mm macro lens.

4. Descriptions

In all cases, genera and species referred to here are fossil-taxa (sensu Cleal et al. 2010; Cleal and Thomas 2021), whereas suprageneric taxa are whole-plant taxa. Following Agosti et al. (2022), full details of the authorship of all taxa are given.

Class: EQUISETOPSIDA Agardh, 1825


- Genus: Mesocalamites Hirmer (1927) emend. Remy and Remy (1978)


Mesocalamites cistiiformis (Stur) Hirmer (1927) (Figures 1g, 3a-c, e)

(≡ *Calamites cistiiformis* Stur 1877)

Description: Linear, unbranched axes up to 20 mm wide, preserved for up to 150 mm long. Fine longitudinal ribs with bluntly acuminate ends, ca. 15 ribs per cm across axis. Clearly marked nodes across axis, with longitudinal ribs sometimes passing directly across or sometimes off-set. Internodes at least 100 mm long, as no preserved stem shows more than one node. No branch scars preserved.

Remarks: These axes represent calamite stems, some definitely as pit-casts, but others as possible stem compressions. They are characterised by long internodes, very fine longitudinal ribs that only sometimes alternate at the nodes, and an absence of branch scars, and are very similar to the specimens figured by Crookall (1969, pl. 111, Figures 1, 2), Purkynová (1970, pl. 8, Figure 4) and Josten (1983, pl. 2, Figures 1, 2) as M. cistiiformis.








Figure 2. Geological setting of the studied site, indicated by a red star. (a) Simplified geological map of the studied area, modified from Blaise et al. (1986). The Nort-sur-Erdre Fault (NSEF) separates the Saint-Georges-sur-Loire and the Ardenay Units, to the North and South, respectively. A narrow band of Carboniferous sediments ("Sillon houiller de la Basse-Loire") is located along the southern contact with the Nort-sur-Erdre Fault. Palaeozoic rocks are unconformably covered with a thin sheet of Cenomanian sands and clays. (b, c) Schematic cross-sections of the studied area. Note that the plant-bearing beds are observed in sandy to silty layers interbedded within conglomerates.

Jongmans (1911) combined this species with *Calamites cistii* Brongniart (1828), which also has axes with fine longitudinal ribs, long internodes and few branch scars, but the ribs in the latter species consistently alternate at

the nodes (Crookall 1969). There is also a comparison with *Mesocalamites taitianus* (Jongmans and Kidston) Hirmer (1927), but the latter has rather wider ribs, shorter internodes and large, oval branch scars.




Figure 3. (a-c, e) Mesocalamites cistiiformis (Stur) Hirmer (PalB 001–003, PalB 005); (a, b) pith casts; (c) ?cast of stem; (e) cast of stem showing leaf-attachment scars. (d) Annularia cf. radiata (Brongniart) Sternberg (PalB 004), leaf whorl. (f) Palaeostachya cf. mostensis Purkynová (PalB 006), part of strobilus showing bracts and axillary ?sporangiophores. Scale bars: 0.42 cm (a); 0.55 cm (b); 0.47 cm (c); 0.84 cm (d); 0.50 cm (e); 0.30 cm (f).

Figure 4. (a, b) Pecopteris plumosa (Artis) Brongniart (PalB 007, 008), slender, obliquely attached pinnules. (c, e) ?Mariopteris sp. (PalB 010, 011), linguaeform, pinnatifid pinnules typical of Mariopteris. (d) Sphenopteris elegans (Brongniart) Sternberg (PalB 009), ultimate pinna with typical incised pinnules. Scale bars: 0.37 cm (a), 0.40 cm (b, c); 0.55 cm (d); 0.15 cm (e).

Specimens PalB 001-004, PalB 005.

- Genus: Annularia Sternberg KM von (1821) Annularia cf. radiata (Brongniart) Sternberg KM von (1825) (Figure 3d)

Description: A single specimen of a stem ca. 5 mm wide with 25-20 leaves attached at a node (the number of leaves in the incompletely preserved leaf whorl is estimated). Leaves are basally fused to form a 1-mm-wide collar around the stem. Leaves ca. 20 mm long, 2 mm wide, linear-lanceolate, widest in middle, with acute apex.

Remarks: Since this single specimen represents the foliage on a relatively wide stem, it is difficult to definitely ascertain the shape of the leaf whorl. However, the general shape of the leaves and the presence of a basal collar makes it most comparable to A. radiata and Annularia ramosa Stur, (1887). These species of sphenophyte foliage are poorly constrained, and we have used A. radiata as the comparison simply because it is the more widely used name.

- Genus: Palaeostachya Weiss (1876) Palaeostachya cf. mostensis Purkynová (1970) (Figure 3f)

Description. Strobilus slightly tapered, maximum (proximal) width 9 mm, preserved for a length of 41 mm. Strobilus axis ca. 1 mm wide with whorls of at least 4 bracts at intervals of 4–5 mm. Bracts initially directed proximally, straight for the first half of length and then curve so their distal part is ca 45° to strobilus axis. Large carbonaceous mass (?sporangiophore) attached to axil of bract and strobilus axis.

Remarks. This strobilus clearly shows what are presumably the sporangiophores in an axillary indicating that it belongs Palaeostachya. Most sphenophyte strobili from Namurian macrofloras belong to Calamostachys (e.g. Stockmans and Willière 1953; Josten 1983), but Purkynová (1970) described Palaeostachya strobili from the upper Arnsbergian (uppermost Serpukhovian) of Upper Silesia. Although we have only one specimen of this strobilus, it is of generally similar size and shape and so we tentatively compare it with Purkynová's specimens.

Specimen PalB 006

Class: FILICOPSIDA Brongniart (1822)

- Genus: Pecopteris (Brongniart) Sternberg KM von (1825) emend. Cleal (2015)

(≡ *Filicites (Pecopteris)* Brongniart (1822)

Pecopteris plumosa (Artis) Brongniart (1836) (Figure 4a, b)

(≡ *Filicites plumosus* Artis 1825)

Description: Bipinnate pinna fragments up to 40 mm long. Penultimate rachis straight and slender (1 mm wide). Ultimate rachises slender (ca. 0.5 mm wide), alternately arranged and oblique to penultimate rachis. Ultimate pinnae up to 15 mm long, spaced 5 mm apart, gradually tapered along most of length. Pinnules alternate, oblique, subtriangular with rounded to acute apex, ca 2 mm long, 1 mm wide. The venation is unclear, but each pinnule appears to have a prominent midvein along most of its length.

Remarks: This specimen is very similar to the P. plumosa figured by Dalinval (1960, pl. 11, Figure 1; pl. 12, Figure 2) and Josten (1983, pl. 46, Figure 3), and corresponds to the forma delicatula of that species.

Specimens PalB 007-008

Class: LYGINOPTERIDOPSIDA Novák (1961)

- Genus: Sphenopteris (Brongniart) Sternberg KM von (1825)

(≡ *Filicites (Sphenopteris)* Brongniart (1822) Sphenopteris elegans (Brongniart) Sternberg KM von (1825) (Figure 4d)

(≡ *Filicites* (*Sphenopteris*) *elegans* Brongniart (1822)

Description: Once-pinnate fragments of fronds up to 55 mm long, 12 mm wide, gradually tapered to a deeply incised terminal segment. Rachis slender (<1 mm wide), straight or slightly deflected. Lateral pinnules alternate, obliquely inserted at 45° or less in the distal part of pinna, gradually changing to 60-70° in more proximal positions. Pinnules rhomboidal, up to 6 mm long, 5 mm wide, deeply incised into up to eight segments, each segment especially in more proximal parts of pinnule further divided into up to three subsidiary segments; segments/subsegments narrowly lanceolate with bluntly acuminate apex. Thin, basally decurrent midvein extends for most of the pinnule length and producing thin lateral veins, one for each ultimate segment. Pinnule lamina well developed so that there are only narrow gaps between each segment.

Remarks: These pinna fragments with deeply incised, somewhat obliquely attached pinnules, with bluntly acuminate ultimate segments, compare well with S. elegans as illustrated by Kidston (1923) and Purkynová (1970). Although the pinnule shape is typical for this species, their size is rather small, but this is probably just a function of them coming from a relatively distal position in the pinna. There are some similarities with the Westphalian Palmatopteris furcata (Brongniart) Potonié (1893), but the pinnules of the latter are not so oblique to the rachis and have more bluntly terminated pinnule lobes. S. elegans typically ranges between the upper Cardioneuropteris antecedens and Calymmotheca larischii Zones (Wagner 1984; Opluštil et al. 2021)

Specimen PalB 009

- Genus: Mariopteris Zeiller (1878)
- -? Mariopteris sp. (Figure 4c, e)

Description. Linguaeform, pinnatifid pinnules with rounded to bluntly acuminate apex and partly constricted base, 10-15 mm long, 3-5 mm wide. Thin, strongly decurrent midvein extends for much of the pinnule length. Lateral veins alternate and produce a bunch of final veins in each of the pinnule lobes (details of the veining pattern not preserved).

Remarks. These fragments compare with remains of Mariopteris fronds from Pennsylvanian macrofloras such as Mariopteris muricata (Sternberg) Zeiller (1878), but are too small to make a definite species assignment.

Specimens PalB 010, 011

Class: CYCADOPSIDA Brongniart, 1843

- Genus: Neuralethopteris Cremer (1893) emend. Laveine (1967)

Neuralethopteris densifolia Josten (1983) (Figure 5) Description. Linguaeform, more-or-less bilaterally

symmetrical pinnules with parallel margins, broadly rounded apex and constricted base, up to 9 mm wide;



Figure 5. Neuralethopteris densifolia Josten (PalB 012–017), with pinnules of very variable length-to-breadth ratio, but consistent venation, as is typical for this fossil-genus. Scale bars: 0.47 cm (a); 0,50 cm (b, c); 0.40 cm (d); 0.70 cm (e); 0.30 cm (f).

largest preserved pinnule 40 mm long. Most pinnules are isolated, but the few that are attached are at 80–90° to the rachis. Prominent midvein extends for most of the pinnule length. Lateral veins alternately arise from midvein and narrow angle, rapidly arched in a somewhat angular pattern and then extend to the pinnule margin, which it meets at 70–90°. Lateral veins fork up to four times depending on pinnule width, producing a vein density of 45–50 per cm on pinnule margin.

Remarks. Although these pinnate fragments are small, the basally constricted, bilaterally symmetrical pinnules with a strong midvein clearly places them within the fossil-genus Neuralethopteris, and the often elongate pinnules with a dense, slightly angular venation indicates N. densifolia. This is a rare species that has only previously been recorded from the Ruhr in north Germany (Josten 1983; Jenchen and Schultka 1993), where it occurs in the Pecopteris aspera



Zone of Kinderscoutian to Yeadonian (Wagner 1984; Opluštil et al. 2021).

Specimens PalB 012-017.

5. Interpretation of the macroflora

Compared with many of the Maine-et-Loire macrofloras described by Bureau (1913-1914) and Strullu-Derrien et al. (2021), which originated from grey, coalbearing sequences, the Pont Barré macroflora is relatively low in diversity. It also differs in the total absence of lycopsids and of Calymmotheca pteridosperms and instead has abundant medullosalean foliage (Neuralethopteris). Similar fern - pteridosperm macrofloras of about the same age from Upper Silesia are interpreted as the remains of vegetation from better-drained substrates and are termed "flöz-fern" macrofloras (Havlena 1961; Gastaldo et al. 2009a, b). This ecological interpretation appears to be supported by the Pont Barré macroflora being preserved in coloured, reddish clastic sediment, which could indicate well-drained conditions and a low water table (see, for instance, comments by Besly and Turner 1983; Besly and Fielding 1989). As the Pont Barré fossils are relatively fragmentary and preserved in a high-energy water deposited stratigraphical sequence, they are clearly allochthonous. However, since there is no evidence of taxa such as Adiantites, Pseudadiantites and Rhacopteridium that have been interpreted as allochthonous remains of hinterland vegetation (e.g. Wagner 2001), it is likely that this macroflora was derived from vegetation growing in well-drained alluvial habitats within the basin (see discussion by Thomas and Cleal 2017).

It is difficult to place biostratigraphically such a low diversity macroflora and the co-occurrence of Sphenopteris elegans and Neuralethopteris densifolia appears to be in conflict as their previously known ranges do not overlap. However, the lower limit of the range of N. densifolia is not well constrained, being based on the less well-documented floras of the lower Namurian of the Ruhr (Josten 1983), whereas the upper limit of the range of S. elegans is better constrained to the lower Chokierian such as in Upper Silesia (Purkynová 1970). In our view, the Pont Barré macroflora can probably be assigned to the Calymmotheca larischii Zone indicating Arnsbergian or early Chokierian age in the Heerlen Regional Chronostratigraphy, making it a little younger than the macrofloras documented from the coal-bearing deposits of this area (Strullu-Derrien et al. 2021); this would make a closer comparison with the macrofloras from the Poruba Formation of Upper Silesia (e.g. Purkynová 1970), which also includes Neuralethopteris.

6. Conclusions

The "Coteaux du Pont Barré" at Beaulieu-sur-Layon (Maine-et-Loire, France) has yielded a newly discovered macroflora of late Mississippian age. The allochthonous fossils occur in coloured, reddish clastic deposits suggesting relatively well-drained conditions, and reflect a vegetation dominated by medullosalean pteridosperms, tedeleacean ferns and sphenophytes. It is markedly different from most of the other macrofloras from the "Sillon houiller de la Basse-Loire" (e.g. Strullu-Derrien et al. 2021), which are more diverse and tend to be dominated by lycopsids (e.g. Lepidodendron) and Calymmotheca (pteridosperm) fronds, and are probably the remains of vegetation from swampier, poorer-drained habitats. The Pont Barré macrofloras therefore provides a valuable, contrasting insight into the vegetation that formed the origins of the coal swamp biome, which in early Westphalian (early Pennsylvanian) times expanded to cover swathes of tropical Euramerica.

Acknowledgments

We thank Ivan Massonnat (owner) and Jo Pithon (previous owner) of the "Domaine Belargus" vineyard and winery, Valérie Chalumeau from the LPO Anjou and Julien Le Berre (for image 3e). CS-D is supported for her work by the Fondation ARS Cuttoli-Paul Appell-Fondation de France.

Author contributions

CS-D designed the research program; CJC and CS-D contributed to the palaeobotany; MB, HL, ARTS contributed to the geology; J-PG photographed the fossil plants (except 3e) and CS-D photographed the site and the outcrop; CJC, MB, CS-D wrote the first version of the manuscript; all authors contributed to the final version of the manuscript.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The work has been partly supported by a grant from the Fondation ARS Cuttoli-Paul Appell to C S-D.

ORCID

Christine Strullu-Derrien http://orcid.org/0000-0003-1131-9546

Michel Ballèvre (b) http://orcid.org/0000-0001-6431-9051 Alan R.T. Spencer http://orcid.org/0000-0001-6590-

Christopher J. Cleal http://orcid.org/0000-0002-8750-

References

- Agosti, D, L Bénichou, W Addink, C Arvanitidis, T Catapano, G Cochrane, M Dillen, M Döring, T Georgiev, I Gérard. 2022. Recommendations for use of annotations and persistent identifiers in taxonomy and biodiversity publishing. Research Ideas and Outcomes. 8:
- Artis, ET. 1825. Antediluvian phytology, illustrated by a collection of the fossil remains of plants peculiar to the coal formations of Great Britain. London: Cumberland and others.
- Ballèvre, M, V Bosse, C Ducassou, P Pitra. 2009. Palaeozoic history of the Armorican Massif: Models for the tectonic evolution of the suture zones. Comptes Rendus Geoscience. 341(2-3):174-201. doi:10.1016/j.crte.2008. 11.009.
- Besly, BM, CR Fielding. 1989. Palaeosols in Westphalian coal-bearing and red-bed sequences, central and northern England. Palaeogeog Palaeoclimat Palaeoecol. 70 (4):303-330. doi:10.1016/0031-0182(89)90110-7.
- Besly, BM, P Turner. 1983. Origin of red beds in a moist tropical climate (Etruria Formation, Upper Carboniferous, UK). In R Wilson, editor. Residual deposits. London: Geological Society (Spec. Publ., 11); pp. 131-147.
- Blaise, J, A Arnaud, R Brossé, P Cavet, J Depagne, M Gruet, H Lardeux, O Limasset. 1986. Notice explicative de la carte géologique de France, feuille Thouarcé (484). Orléans: BRGM.
- Brongniart, A. 1822. Sur la classification et la distribution des végétaux fossiles en général, et sur ceux des terrains de sédiment supérieur en particulier (1). Mém Mus Hist Nat Paris. 8:203-240.
- Brongniart, A. 1828. Histoire des Végétaux fossiles. Vol. I (2), Paris: Dufour and d'Ocagne.
- Brongniart, A. 1836. Histoire des Végétaux fossiles. Vol. I (10), Paris: Dufour and d'Ocagne.
- Bureau, É. 1913. Bassin Houiller de la Basse Loire. Fascicule II. Description des Flores Fossiles. Paris: Imprimerie Nationale (Études des Gîtes Minéraux de la France). -1914Atlas published 1913, Text published 1914.
- Cartier, C, M Faure. 2004. The Saint-Georges-sur-Loire olistostrome, a key zone to understand the Gondwana-Armorica boundary in the Variscan belt (Southern Brittany, France). Int J Earth Sci. 93 (6):945-958. doi:10.1007/s00531-004-0398-3.
- Cartier, C, M Faure, H Lardeux. 2001. The Hercynian orogeny in the South Armorican Massif (Saint-Geroges-sur-Loire Unit, Ligerian Domain, France): rifting and welding of continental stripes. Terra Nova. 13(2):143-149. doi:10. 1046/j.1365-3121.2001.00334.x.
- Cleal, CJ. 2015. The generic taxonomy of Pennsylvanian age marattialean fern frond adpressions. Palaeontogr Abt B. 292(1-3):1-21. doi:10.1127/palb/292/2015/1.
- Cleal, CJ. 2021a. Paleozoic plants. In D Alderton S Elias, editors. Encyclopedia of Geology. 2nd ed. Vol. 3. London: Academic Press; pp. 461-475.
- Cleal, CJ. 2021b. Tiering on land trees and forests (Late Palaeozoic). eLS. 2:158-169.
- Cleal, CJ, BA Thomas. 2021. Naming of parts: the use of fossil-taxa in palaeobotany. Fossil Imprint. 77(1):166-186. doi:10.37520/fi.2021.013.
- Cleal, CJ, BA Thomas, J McNeill, AE Orchard, JC David. 2010. Botanical nomenclature and plant fossils. Taxon. 59 (1):261-268. doi:10.1002/tax.591024.

- Cleal, CJ, D Uhl, B Cascales-Miñana, BA Thomas, AR Bashforth, SC King, EL Zodrow. 2012. Plant biodiversity changes in Carboniferous tropical wetlands. Earth-Science Reviews. 114(1-2):124-155. doi:10.1016/j. earscirev.2012.05.004.
- Corillion, R. 1957. Sur le caractère méridional et la richesse de la flore de Pont-Barré. Bull Soc Bot Fr. 104(Suppl. 2):60-66. doi:10.1080/00378941.1957.10835174.
- Cremer, L. 1893. Über die Fossilen Farne des Westfälischen Carbons und ihre Bedeutung für eine Gliederung des letzeren. Mitt Geol Mus Berggewerkschaftskasse. 1:1-49.
- Crookall, R. 1969. Fossil plants of the Carboniferous rocks of Great Britain [Second Section]. Part 5. Mem Geol Surv G B Palaeont. 4:573-792.
- Dalinval, A. 1960. Contribution à l'étude des Pécoptéridées. Les Pecopteris du bassin houiller du Nord de la France. Études Géol Atlas Topogr Souterr. 1(3):1-222.
- DiMichele, WA, HW Pfefferkorn, RA Gastaldo. 2001. Response of late Carboniferous and Early Permian plant communities to climate change. Ann Rev Earth Planet Sci. 29(1):461-487. doi:10.1146/annurev.earth.29.1.461.
- Gastaldo, RA, E Purkyňová, Z Šimůnek. 2009a. Megafloral perturbation across the Enna marine zone in the Upper Silesian basin attests to late Mississippian (Serpukhovian) deglaciation and climate change. Palaios. 24(6):351-366. doi:10.2110/palo.2007.p07-027r.
- Gastaldo, RA, E Purkyňová, Z Šimůnek, MD Schmitz. 2009b. Ecological persistence in the late Mississippian (Serpukhovian, Namurian A) megafloral record of the Upper Silesia Basin, Czech Republic. Palaios. 24 (6):336-350. doi:10.2110/palo.2008.p08-084r.
- Greb, SF, WA DiMichele, RA Gastaldo. 2006. Evolution and importance of wetlands in earth history. In S Greb, and W DiMichele, editors. Wetlands Through Time. Boulder CO: Geological Society of America (Spec. Pap. 399); p.
- Havlena, V. 1961. Die flöznahe und flözferne Flora des oberschlesischen Namurs A und B. Palaeontogr Abt B.
- Hirmer, M. 1927. Handbuch der Paläobotanik. Oldenbourg: Munich and Berlin.
- Jenchen, U, S Schultka. 1993. Die ehemalige Ziegeleitongrube Voßacker und die Abgrabung am Küchenberg, zwei fossilführende Aufschlüsse im tiefen Oberkarbon. Geol.Paläont.Westf. 22:41-85.
- Jongmans, WJ. 1911. Anleitung zur Bestimmung der Karbonpflanzen West-Europas mit besonderer Berücksichtigung der in den Niederlanden und den benachbarten Länderen gefundenen oder noch zu erwartenden Arten. I Band Thallophytae, Equisetales, Sphenophyllales Mededeelingen van de Rijksopsporing van Delfstoffen. 3:1-482.
- Josten, K-H. 1983. Die fossilen Floren im Namur des Ruhrkarbons. Fortschr Geol Rheinland Westfalen. 31:1–327.
- Kidston, R. 1923. Fossil plants of the Carboniferous rocks of Great Britain. Parts 1-4. Mem Geol Surv Gt Brit Mem, Paleont. 2:1-376.
- Laveine, J-P. 1967. Contribution a l'étude de la flore du terrain houiller. Les Neuroptéridées du Nord de la France. Études Géol Atlas Topogr Souterr. 1(5):1–344.
- Lin, W, M Faure, X-H Li, W Ji. 2019. Pre-Variscan tectonic setting of the south margin of Armorica: insights from detrital zircon ages distribution and Hf isotopic composition of the St-Georges-sur-Loire Unit (S. Armorican



- Massif, France). Tectonophys. 766:340-378. doi:10.1016/ j.tecto.2019.06.015.
- Louail, J. 1984. La transgression crétacée au Sud du Massif Armoricain. Cénomanien de l'Anjou et du Poitou. Crétacé supérieur de Vendée. Mém Soc Géol Minéral Bretagne. 29:1-333.
- Novák, FA. 1961. Vyšší rostliny. Tracheophyta. Prague: Czechoslovak Acad. Sciences.
- Opluštil, S, CJ Cleal, J Wang, M Wang. 2021. Carboniferous macrofloral biostratigraphy: an overview. In S Lucas, J Schneider, X Wang, and S Nikolaeva, editors. The Carboniferous Timescale. London: Geological Society (Spec. Publ. 512); p. 813-863.
- Pardoe, HS, CJ Cleal, CM Berry, B Cascales-Miñana, B Davis, JB Diez, MV Filipova-Marinova, T Giesecke, J Hilton, D Ivanov, et al. 2021. Palaeobotanical experiences of plant diversity in deep time. 2: how to measure and analyse past plant biodiversity. Palaeogeog Palaeoclimat Palaeoecol. 580:110618. doi:10.1016/j. palaeo.2021.110618.
- Poirier, L. 1935. Essai sur la morphologie de l'Anjou méridional (Mauges et Saumurois). Ann Géog. 44(251):474-491. doi:10.3406/geo.1935.11159.
- Potonié, H. 1893. Ueber einige Carbonfarne. III. Theil. Jahrb. Kön. Preuss. Geol. Landesanst. Bergakad. Berlin. 12:1-36.
- Purkynová, E. 1970. Die Unternamurflora des Beckens von Horní Slezsko (CSSR). Paläontol Abh, Abt B. 3:129-268.
- Remy, W, R Remy. 1978. Calamitopsis n. gen. und die Nomenklatur und Taxonomie von Calamites Brongniart 1828. Argumenta Palaeobot. 5:1-10.
- Sellier, D. 2020. Essai sur la notion de coteau en Pays de la Loire. Les Cahiers Nantais. 2020:127-147.
- Sternberg KM von. 1821. Versuch einer geognostisch-botanischen Darstellung der Flora der Vorwelt. Vol. 1(2), Leipzig und Prag: F. Fleischer.
- Sternberg KM von. 1825. Versuch einer geognostisch-botanischen Darstellung der Flora der orwelt. Vol. 1(4), Regensburg: E. Brenck's Wittwe.
- Stockmans, F, Y Willière 1953. Végétaux namuriens de la Belgique (Texte). Brussels Association pour l'Étude de la

- Paléontologie et de la Stratigraphie Houillères (Publication No. 13) Atlas published 1952.
- Strullu-Derrien, C, CJ Cleal, C Ducassou, ART Spencer, E Stolle, VO Leshyk. 2021. A rare late Mississippian flora from Northwestern Europe (Maine-et-Loire Coalfield, Pays de la Loire, France). Rev Palaeobot Palynol. 285:104359. doi:10.1016/j.revpalbo.2020. 104359.
- Strullu-Derrien, C, AR Spencer, CJ Cleal, VO Leshyk. 2022. The 330-320 million-year-old tranchée des Malécots (Chaudefonds-sur-Layon, South of the Armorican Massif, France): a rare geoheritage site containing in situ palaeobotanical remains. Geoheritage. 14(2):1-17. doi:10.1007/s12371-022-00673-0.
- Stur, D. 1877. Die Culm-Flora der Ostrauer und Waldenburger Schichten. Abh Kaiserlich-Königlichen Geol Reichsanstalt. 8(2):107-472.
- Stur, D. 1887. Die Carbon-Flora der Schatzlarer Schichten. Abtheilung 2: Die Calamarien der Carbon-flora der Schatzlarer Schichten. Abh Kaiserlich-Königlichen Geol Reichsanstalt. 11(2):1-240.
- BA, CJ Cleal. 2017. Distinguishing Pennsylvanian-age lowland, extra-basinal and upland vegetation. Palaeobiodiversity and Palaeoenvironments. 97(2):273-293. doi:10.1007/s12549-017-0277-0.
- Wagner, RH. 1984. Megafloral zones of the Carboniferous. C R 9e Congr Int Strat Géol Carbon. 2:109-134. Washington, 1979.
- Wagner, RH. 2001. The extrabasinal elements in lower Pennsylvanian floras of the Maritime Provinces, Canada: of Adiantites, Pseudadiantites Rhacopteridium. Pseudadiantites and Rhacopteridium Rev Esp Paleont. 16(2):187–207. doi:10.7203/sjp.16.2.21599.
- Weiss, CE. 1876. Beiträge zur fossilen Flora. Steinkohlen-Calamarien mit besonderer Berücksichtigung ihrer Fructificationen. Abh geol Spez-Kt Preuss, Thüring Staaten. 2(1):1-149.
- Zeiller, R. 1878. Végétaux fossiles du terrain houiller. Atlas Explic Carte Géol Fr. 4:159-176.