

On the cyclotron absorption line and evidence of the spectral transition in SMC X-2 during 2022 giant outburst

G. K. Jaisawal¹,¹★ G. Vasilopoulos^{2,3}, S. Naik⁴, C. Maitra⁵, C. Malacaria⁶, B. Chhotaray^{4,7}, K. C. Gendreau⁸, S. Guillot⁹, M. Ng¹⁰ and A. Sanna¹¹

¹*DTU Space, Technical University of Denmark, Elektrovej 327-328, DK-2800 Lyngby, Denmark*

²*Université de Strasbourg, CNRS, Observatoire Astronomique de Strasbourg, UMR 7550, F-67000 Strasbourg, France*

³*Department of Physics, National and Kapodistrian University of Athens, University Campus Zografos, GR-15783, Athens, Greece*

⁴*Astronomy and Astrophysics Division, Physical Research Laboratory, Navrangpura, Ahmedabad, 380009, Gujarat, India*

⁵*Max-Planck-Institut für Extraterrestrische Physik, Gießenbachstraße 1, D-85748 Garching, Germany*

⁶*International Space Science Institute (ISSI), Hallerstrasse 6, CH-3012 Bern, Switzerland*

⁷*Indian Institute of Technology Gandhinagar, Palaj, Gandhinagar, 382055, Gujarat, India*

⁸*Astrophysics Science Division, NASA's Goddard Space Flight Center, Greenbelt, MD 20771, USA*

⁹*Institut de Recherche en Astrophysique et Planétologie, UPS-OMP, CNRS, CNES, 9 Avenue du Colonel Roche, BP 44346, F-31028 Toulouse Cedex 4, France*

¹⁰*MIT Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

¹¹*Dipartimento di Fisica, Università degli Studi di Cagliari, SP Monserrato-Sestu km 0.7, I-09042 Monserrato, Italy*

Accepted 2023 March 10. Received 2023 March 10; in original form 2023 February 13

ABSTRACT

We report comprehensive spectral and temporal properties of the Be/X-ray binary pulsar SMC X-2 using X-ray observations during the 2015 and 2022 outbursts. The pulse profile of the pulsar is unique and strongly luminosity dependent. It evolves from a broad-humped into a double-peaked profile above luminosity $3 \times 10^{38} \text{ erg s}^{-1}$. The pulse fraction of the pulsar is found to be a linear function of luminosity as well as energy. We also studied the spectral evolution of the source during the latest 2022 outburst with *NICER*. The observed photon index shows a negative and positive correlation below and above the critical luminosity, respectively, suggesting evidence of spectral transition from the sub-critical to supercritical regime. The broad-band spectroscopy of four sets of *NuSTAR* and *XRT/NICER* data from both outbursts can be described using a cut-off power-law model with a blackbody component. In addition to the 6.4 keV iron fluorescence line, an absorption-like feature is clearly detected in the spectra. The cyclotron line energy observed during the 2015 outburst is below 29.5 keV, however latest estimates in the 2022 outburst suggest a value of 31.5 keV. Moreover, an increase of 3.4 keV is detected in the cyclotron line energy at equal levels of luminosity observed in 2022 with respect to 2015. The observed cyclotron line energy variation is explored in terms of accretion induced screening mechanism or geometrical variation in line forming region.

Key words: stars: neutron – pulsars: individual: SMC X-2 – X-rays: stars.

1 INTRODUCTION

Be/X-ray binaries (BeXRBs) represent two-thirds of the population of high mass X-ray binaries. These systems consist of a massive ($> 10 M_{\odot}$) optical companion and a compact object (usually a neutron star) in a binary system. The optical companion in BeXRBs is a non-supergiant OB spectral type star that shows Balmer series emission lines and infrared excess at a point in its life (Reig 2011). The above characteristics originate from an equatorial circumstellar disk that is formed around the Be star due to its rapid rotation at velocities of more than 75 per cent of Keplerian limit (Porter & Rivinius 2003). The compact object in the system, on the other hand, accretes directly from the Be-circumstellar disk. Two kinds of X-ray outbursts are observed from BeXRBs. First, Type-I outbursts are short (only a few weeks long), (quasi-)periodic events that reach a peak luminosity of

$\leq 10^{37} \text{ erg s}^{-1}$ and occur close to the periastron passage of the binary system. The second category of outbursts is giant in nature, where the peak luminosity reaches $\geq 10^{37} - 10^{38} \text{ erg s}^{-1}$. The latter usually lasts for a multiple or significant portion of the orbit and does not follow any orbital dependencies.

A significant population of BeXRBs is found in the Milky Way and the Magellanic Clouds with luminosities in the range of $10^{34} - 10^{38} \text{ erg s}^{-1}$ (Liu, van Paradijs & van den Heuvel 2006; Reig 2011). Our Galaxy hosts about 60–70 such systems (Reig 2011; Walter et al. 2015). The Small Magellanic Cloud (SMC; a neighbouring irregular dwarf galaxy), however, contains about 68 pulsars (and a total of 122 sources including candidates) despite being only a few per cent of the mass of the Milky Way Galaxy (Reig 2011; Coe & Kirk 2015; Walter et al. 2015; Haberl & Sturm 2016; Yang et al. 2017; Vinciguerra et al. 2020).

SMC X-2 is a 2.37 s pulsating source in a BeXRB system located inside SMC (Schurch et al. 2011) at a distance of 62 kpc (Hilditch, Howarth & Harries 2005; Graczyk et al. 2014). It was the second

* E-mail: gaurava@space.dtu.dk




Figure 1. The light curves of SMC X-2 during its 2015 (top) and 2022 (bottom) X-ray outbursts with *MAXI* (2–20 keV; red dot), *NICER* (0.5–10 keV; blue square), and *Swift/XRT* (0.3–10 keV; grey ellipse). The vertical dotted lines show the start date of the *NuSTAR* observations. In the bottom panel, the XRT light curve is scaled by a factor of seven to compare with the *NICER* light curve.

brightest X-ray source in the galaxy when *SAS 3* discovered it in October 1977. The recorded luminosity was $8.4 \times 10^{37} \text{ erg s}^{-1}$ (assuming a distance of 65 kpc) in the 2–11 keV energy band (Clark et al. 1978; Clark, Li & van Paradijs 1979). Later observations with the *HEAO*, *Einstein*, and *ROSAT* missions demonstrated the X-ray transient nature of the source (Marshall et al. 1979; Seward & Mitchell 1981; Kahabka & Pietsch 1996). Thanks to *RXTE* (*Rossi X-ray Timing Explorer*) and *ASCA* (*Advanced Satellite for Cosmology and Astrophysics*), coherent X-ray pulsations from SMC X-2 were detected for the first time during the 2000 January–April outburst (Corbet et al. 2001). This discovery made it possible to identify the accreting object as a neutron star. The optical observations of the system, on the other hand, suggested two possible early-type stars at an angular separation of 2.5 arcsec as the potential counterparts of the pulsar. The Optical Gravitational Lensing Experiment *I*-band photometric studies found variability at a period of 18.62 ± 0.02 d from one of the stars (northern star) (Schurch et al. 2011). This periodicity closely matched the orbital period of ≈ 18.4 d measured by studying the X-ray pulse-period modulation during the 2002 and 2015 outbursts (Townsend et al. 2011; Li et al. 2016). The optical counterpart (northern star) is identified to be an O9.5 III–V emission star (McBride et al. 2008).

The system displayed major outbursts during 1977 October (Marshall et al. 1979; Seward & Mitchell 1981), 2002 January–April (Corbet et al. 2001), 2015 August–November (Kennea et al. 2015; Jaisawal & Naik 2016), and 2022 June–August (Coe et al. 2022). Detailed studies with *XMM-Newton*, *Swift/XRT* (X-Ray Telescope), and *NuSTAR* were performed during the 2015 outburst when the pulsar was accreting in the super-Eddington regime with a peak

outburst luminosity of $6 \times 10^{38} \text{ erg s}^{-1}$ (Kennea et al. 2015; Negoro et al. 2015; Jaisawal & Naik 2016). Ionized emission lines from N, O, Ne, Si, and Fe were detected in the data from the *XMM-Newton* observation (La Palombara et al. 2016). Moreover, with broad-band coverage of *NuSTAR* in the range of 3–79 keV, the neutron star magnetic field was estimated for the first time to be $\approx 3 \times 10^{12}$ G based on the discovery of a cyclotron resonance scattering feature (CRSF) at around 27 keV (Jaisawal & Naik 2016). A negative correlation between the luminosity and cyclotron line energy was also reported in the study. In addition to this, the pulsar is known to show luminosity-dependent pulse profiles evolving from single to double-peaked (Jaisawal & Naik 2016; Li et al. 2016; Roy et al. 2022). The onset of the propeller regime observed in late 2015 also provided a measurement of the magnetic field to be $\approx 3 \times 10^{12}$ G (Lutovinov et al. 2017), in agreement with the estimation through CRSF detection.

After 7 yr of X-ray quiescence, SMC X-2 became active on 2022 June (Kennea, Coe & Evans 2022). The pulsar reached a peak luminosity of $1 \times 10^{38} \text{ erg s}^{-1}$, 3 weeks after its detection in the soft X-ray band (Coe et al. 2022). Unlike the 2015 outburst, the *MAXI/GSC* (Matsuoka et al. 2009) did not detect any significant outburst peak in 2–20 keV band due to the relatively moderate nature of the outburst in addition to the observational coverage gap (see e.g. Fig. 1). We monitored the target closely with *NICER* and *NuSTAR* to understand the spectral and temporal evolution during the 2022 outburst. We also analysed three *NuSTAR* observations from the 2015 outburst to examine the cyclotron line and its evolution and also studied the broad-band temporal and spectral properties of the source. Section 2 reports the details of the observations and data analysis methods. The timing and spectral results are presented in Section 3, followed by a discussion and conclusion in Section 4.

2 OBSERVATIONS AND DATA ANALYSIS

2.1 *NICER*

Launched in 2017 June, the *NICER* X-ray timing instrument (XTI, Gendreau, Arzoumanian & Okajima 2012; Gendreau et al. 2016) is a soft XRT (0.2–12 keV) attached to the International Space Station. The XTI consists of a total of 56 co-aligned concentrator optics. Each optics collimates the soft X-ray photons on to a silicon drift detector at the focus (Prigozhin et al. 2012). A high time resolution of ~ 100 ns (rms) and spectral resolution of ≈ 85 eV at 1 keV are achieved by *NICER*. The effective area of the instrument is $\approx 1900 \text{ cm}^2$ at 1.5 keV, with 52 active detectors. The total field of view of *NICER* is $\approx 30 \text{ arcmin}^2$.

NICER monitored SMC X-2 after the onset of the 2022 outburst in June. We used a total of 37 observations under the ObsIDs 52028301xx with a net exposure time of ≈ 39.6 ks for studying the pulsar emission during the outburst between 2022 June 20 and August 17. We reduced the *NICER* data using the *nicer12*¹ script available under HEASOFT version 6.30. The reprocessing is done in the presence of the gain and calibration database files of version 20200722. The cleaned events obtained from the pipeline are used in our further analysis. In the beginning, the events from the South Atlantic Anomaly region were filtered out. We further applied standard filtering criteria based on the elevation angle from the Earth limb, pointing offset, and the offset from the bright

¹https://heasarc.gsfc.nasa.gov/docs/nicer/analysis_threads/nicer12/.

Table 1. Log of simultaneous observations of SMC X-2 with *NuSTAR* and *Swift*/XRT or *NICER* during its 2015 and 2022 X-ray outbursts.


Observatory/ Instrument	ObsID	Start date	Exposure (ks)
<i>NuSTAR</i>	90102014002	2015-09-25T21:51:08	24.5
<i>Swift</i> /XRT	00034073002	2015-09-25T22:32:58	1.8
<i>Swift</i> /XRT	00081771002	2015-10-12T21:30:58	1.5
<i>NuSTAR</i>	90102014004	2015-10-12T21:41:08	23
<i>Swift</i> /XRT	00034073042	2015-10-21T14:08:58	4
<i>NuSTAR</i>	90101017002	2015-10-21T21:31:08	26.7
<i>NICER</i>	5202830119	2022-07-13T07:50:21	1.1
<i>NuSTAR</i>	90801319002	2022-07-13T20:18:43	42.7

Earth. Good time intervals were produced using the *nimaketime* task. Final spectra and light curves were created with the XSELECT package of FTOOLS. The spectral background corresponding to each observation is generated using the *nibackgen3C50*² tool (Remillard et al. 2022). The response matrix and ancillary response files of version 20200722 are applied in our spectral analysis.

2.2 NuSTAR

NuSTAR is the first hard X-ray focusing observatory launched in 2012 June (Harrison et al. 2013). It consists of two co-aligned grazing angle incidence telescopes. The mirrors in each optic are coated with multilayers of Pt/SiC and W/Si that reflect the soft to hard X-ray photons. A Cadmium–Zinc–Telluride detector at the focal point of each unit is sensitive to the 3–79 keV photons. Following the recent outburst of SMC X-2, we requested a *NuSTAR* target of opportunity observation. The target was observed for an effective exposure of 42.7 ks on 2022 July 13–14 (Table 1). Standard analysis procedures were followed for data reduction with NUSTARDAS 1.9.7 software. Unfiltered events were processed in the presence of the CALDB of version 20220802 using *nupipeline* task. Source products are extracted from a circular region of 120 arcsec radius around the central coordinates on each detector using the *nuproducts* task. The background products are also accumulated in a similar manner from a source-free circular region of 120 arcsec radius. The background-subtracted light curves from both detector modules of *NuSTAR* were combined for our timing studies.

We also reduced the data from the *NuSTAR* observations of SMC X-2 during its 2015 major outburst (Table 1) by following the above procedure. The first observation was close to the peak of the outburst in 2015 September, whereas the other two were in the declining phase. Corresponding *Swift*/XRT observations are analysed to perform broad-band spectroscopy (see also Jaisawal & Naik 2016; Lutovinov et al. 2017 for XRT data analysis and descriptions; Evans et al. 2009). All four *NuSTAR* observations in the 2015–2022 timeline are simply referred to as Obs-I, Obs-II, Obs-III, and Obs-IV in the paper. For spectroscopy, we have grouped each *NICER*, *NuSTAR*, and XRT spectra for a minimum of 32 counts per channel bin to achieve a good signal-to-noise ratio using *grppha*.

**Figure 2.** The 3–79 keV pulse profiles of SMC X-2, observed with *NuSTAR*, arranged in order of luminosity. The top two (Obs-I and Obs-II) and bottom (Obs-III) panels are from the 2015 outburst. The third panel (Obs-IV) represents the pulse profile from the recent 2022 outburst. L_{38} stands for 0.5–100 keV unabsorbed luminosity in $10^{38} \text{ erg s}^{-1}$ unit.

3 RESULTS

3.1 Timing analysis

We searched for X-ray pulsations in the 3–79 keV *NuSTAR* light curves of SMC X-2 using the χ^2 -maximization technique (Leahy 1987). The barycentric corrected pulse period of the neutron star was estimated to be 2.37197(2), 2.37141(2), 2.37257(2), and 2.37283(1) s during Obs-I, Obs-II, Obs-III, and Obs-IV, respectively. The light curves from all the *NuSTAR* observations are folded with the corresponding pulse period to obtain pulse profiles in the 3–79 keV band, which are shown in Fig. 2. A double-peaked profile is clearly evident at a luminosity³ of $6 \times 10^{38} \text{ erg s}^{-1}$ (top panel of Fig. 2). The shape of the pulse profile changes as luminosity decreases to $< 3 \times 10^{38} \text{ erg s}^{-1}$ where a hump-like structure appears in the profile between 0.5 and 1.5 pulse phase. The pulse profiles exhibit similar morphology at an equal level of luminosities observed during the 2015 (Obs-II) and 2022 (Obs-IV) outbursts.

³The 0.5–100 keV unabsorbed luminosity is calculated at a distance of 62 kpc after broad-band spectroscopy that is presented in Section 3.2.

²https://heasarc.gsfc.nasa.gov/docs/nicer/toolsnicer_bkg_est_tools.html.




Figure 3. The energy resolved pulse profiles of SMC X-2 from the *NuSTAR* observations during its 2015 (Obs-I, Obs-II, and Obs-III) and 2022 (Obs-IV) outbursts. L_{38} stands for 0.5–100 keV unabsorbed luminosity in units of $10^{38} \text{ erg s}^{-1}$.

We adopted the following definition to evaluate the pulse fraction (PF) of the pulsar, i.e. the relative amplitude of pulsed modulation:

$$\text{PF} = \frac{F_{\text{max}} - F_{\text{min}}}{F_{\text{max}} + F_{\text{min}}}, \quad (1)$$

where, F_{max} and F_{min} are the maximum and minimum intensities observed in the pulse profile, respectively. The estimated PFs during Obs-I, Obs-II, Obs-III, and Obs-IV are $\approx 45.7 \pm 0.4$ percent, 29.2 ± 0.4 percent, 17.6 ± 0.2 percent, and 26.2 ± 0.3 percent, respectively. The observed PF is correlated with luminosity. This means that the pulsed emission from the accretion column or the hot spot contributes more than the unpulsed radiation (originating from accretion flow or from the surface) when the source luminosity increases.

To examine the evolution of pulsed beam geometry of the source, we folded the energy-resolved light curves in 3–7, 7–15, 15–25, 25–40, and 40–79 keV from the *NuSTAR* observations (Fig. 3). Doubled peaked profiles are observed throughout soft to hard X-ray energy ranges during Obs-I. The emission from both poles is clearly apparent at this stage. As the luminosity decreases to a level of $\sim 3 \times 10^{38} \text{ erg s}^{-1}$, the pulse profiles below 15 keV (Obs-II and Obs-IV) appear with a hump-like structure that further evolves into a double-peak at higher energies. The Obs-III is at around the Eddington limit of a classical neutron star. At this point, the soft to hard pulse profiles are single-peaked with a hump in the middle. We further studied the energy evolution of the PF from the pulse profiles. Fig. 4 shows positive dependencies between the energy and the PF. An extremely low PF of about 11 percent is found during Obs-III in the 3–7 keV band.

3.2 Spin evolution

We used the barycentric corrected *NICER* data to investigate the evolution of the temporal properties with the outburst and in particular, the spin evolution. To search for a periodic signal, we followed the same methodology adopted for *NICER* monitoring of SXP 15.6




Figure 4. The evolution of pulse fraction with energy, obtained from the energy resolved pulse profiles of the pulsar.

(see Vasilopoulos et al. 2022). First, we adopted the epoch folding Z-search test implemented through *HENDRICS* and *Stingray* (Huppenkothen et al. 2019). Then, we refined the temporal solution and its uncertainty based on the ToA (time of arrival) of individual pulses by using *PINT*⁴ (Luo et al. 2021). For some snapshot phases connecting the ToAs was challenging to impossible due to multiple peaks in the periodogram with similar intensity, a problem often encountered in slow pulsars observed with gaps (e.g. see appendix of

⁴<https://github.com/nanograv/pint/>.




Figure 5. Period evolution of SMC X-2 during the 2022 outburst using *NICER* and *NuSTAR* observations in top panel. The bottom panel shows the pulse period evolution observed by *RXTE* during the 2000 January–May outburst.

Zolotukhin et al. 2017; Vasilopoulos et al. 2018). The resulting period measurements are shown in Fig. 5. In the same plot, we mark the *NuSTAR* measurement. We note that *NICER* observations where a defined period was not obtained are marked as ‘bad’ and no error is given as their uncertainty is similar to the range of the plot.

In the spin evolution, we see clear signatures of orbital motion, consistent with the 18.38 d period reported in the literature based on *RXTE* data (Townsend et al. 2011), whereas a similar periodicity is also seen in optical data of the system (e.g. Roy et al. 2022). Modelling of the archival *RXTE* data has revealed an almost circular orbit with an eccentricity of about 0.07. The sampling of the *RXTE* data was sparse and in similar orbital phases. In contrast, the new *NICER* data cover a smaller baseline but better sample the orbital cycle near the peak of the outburst. Thus, their study could help improve the old orbital solution. To model the orbital evolution, we used a model composed of the orbital signature and the first derivative of the frequency similar to Townsend et al. (2011). Moreover, we followed a Bayesian approach to effectively sample the whole parameter space of the underline model (Karaferias et al. 2023). We applied the same method to the *NICER* data, and the *RXTE* measurements.⁵ Given the issues with the *NICER* period determination at the later stages of the 2022 monitoring, we constrained our fit within MJD 59750–59790. We find that the 2022 monitoring data yield a somehow eccentric orbit ($e = 0.27$), while the period is better constrained by the *RXTE* data. Interestingly, if we maintain no priors for the eccentricity the

⁵Data points were extracted from Townsend et al. (2011).




Figure 6. Three representative *NICER* spectra in 0.5–10 keV fitted with an absorbed cut-off power-law model. Black, red, and green correspond to an unabsorbed luminosity of 4.4×10^{37} , 6.8×10^{37} , and 1.3×10^{38} erg s^{−1} in the 0.5–10 keV range, respectively.

fit to the *RXTE* data yields a lower eccentricity consistent with the one reported by Townsend et al. (2011). For comparison, we plotted both solutions to the newly obtained and archival data shown in Fig. 5.

3.3 The continuum emission during 2022 outburst

The soft X-ray energy spectrum of SMC X-2 is studied to understand the emission at different phases of the 2022 outburst. We used *NICER* observations at multiple epochs of the outburst in this study. Each 0.5–10 keV *NICER* spectrum can be fitted statistically well with an absorbed cut-off power-law model in XSPEC (Arnaud 1996). Fig. 6 shows the energy spectra from these observations together. We considered TBabs model at Wilm abundance (Wilms, Allen & McCray 2000) and Verner cross-section (Verner et al. 1996) to account for the local and interstellar medium absorptions along the source direction in our spectral fit. We found that the equivalent hydrogen column density (N_H) varies in the range of $1\text{--}3 \times 10^{21}$ atoms cm^{−2} during these observations. To reduce the spectral fitting degeneracy in the soft X-ray band, the column density is fixed at an average value of 1.43×10^{21} cm^{−2}, which is consistent with the value observed with *XMM-Newton* in 2015 within error bars (La Palombara et al. 2016). We then fitted each individual spectrum with an absorbed cut-off power-law model at the above-fixed column density. No signature of iron fluorescence line is detected in the *NICER* spectra. The 0.5–10 keV unabsorbed flux is estimated using the cflux convolution model. The parameter errors are estimated for the 90 per cent confidence interval in this paper.

The evolution of the spectral parameters during the outburst is shown in Fig. 7. The power-law photon index and cut-off energy are variable across the rise (blue) and decay (red) phases of the outburst. The peak luminosity in the 0.5–10 keV range is found to be 1.3×10^{38} erg s^{−1} (for a source distance of 62 kpc) around the outburst peak. To examine the spectral changes and associated spectral transition, the parameters are studied with respect to the 0.5–10 keV unabsorbed luminosity in Fig. 8. The photon index and cut-off energy follow a positive correlation (with a Pearson correlation coefficient of 0.61 and 0.66, respectively) with luminosity above 9×10^{37} erg s^{−1}. The




Figure 7. Spectral parameters of SMC X-2 after fitting each *NICER* energy spectrum with an absorbed cut-off power-law model. The absorption column density is fixed at $1.43 \times 10^{21} \text{ cm}^{-2}$. The unabsorbed luminosity is estimated in 0.5–10 keV range for a source distance of 62 kpc. The outburst peaked around 2022 July 10 with a luminosity of $1.3 \times 10^{38} \text{ erg s}^{-1}$ in the *NICER* band. The square (blue) and squashbox (red) symbols correspond to the rising and declining parts of the 2022 outburst, respectively.

index stays close to a value of 0.35 at luminosity in the range of $(6\text{--}9) \times 10^{37} \text{ erg s}^{-1}$. This could be identified as a critical limit of the pulsar where the spectral shape changes. Below this limit, the source shows a negative correlation (with a Pearson correlation coefficient of -0.57) between luminosity and the photon index. Similar behaviour is observed between luminosity and the cut-off energy (with a Pearson correlation coefficient of -0.54).

We also fitted each *NICER* spectrum with two absorption components together with a cut-off power-law model. One of the column densities was fixed at a foreground value $4 \times 10^{20} \text{ cm}^{-2}$ (Dickey & Lockman 1990) at the Solar abundance. Another column density was free in our study at the SMC abundance of 0.2 Solar (Russell & Dopita 1992) to incorporate the local absorption. The SMC column density changes in the range $0.5\text{--}3 \times 10^{21} \text{ cm}^{-2}$ during the outburst. All the fitted parameters, such as SMC column density, photon index, and the cut-off energy, however, were found to decrease with luminosity. A gradual decline may be because of spectral degeneracy among these parameters in the soft X-ray band. This is similar to the findings discussed above in this section. Though we cannot totally disentangle the variation of the local absorption during the outburst therefore to reduce the degeneracy among the parameters, we fixed the SMC




Figure 8. The evolution of photon index and cut-off energy with the 0.5–10 keV luminosity of the pulsar. Square (blue) and squashbox (red) symbols correspond to the rising and declining parts of the 2022 outburst, respectively.

column density to an average value of $1.25 \times 10^{21} \text{ cm}^{-2}$. This resulted in similar kinds of parameter variations as reported in Figs 7 and 8.

3.4 Broad-band spectroscopy of the pulsar during 2015 and 2022 giant outbursts

We further performed broad-band spectroscopy of SMC X-2 in the 1–70 keV range to investigate the continuum as well as the nature of the cyclotron absorption line and its evolution. Three sets of XRT and *NuSTAR* quasi-simultaneous data from the 2015 outburst are used along with the 2022 July *NICER* and *NuSTAR* pointed observations (Fig. 9). These spectra are additionally binned for representation purposes using `setplot rebin` command of XSPEC. The pulsar spectrum can be described by standard models such as the Fermi–Dirac cut-off power-law, the negative and positive power-law with exponential cut-off (NPEX; Makishima et al. 1999), and the cut-off power-law with a blackbody component (see also Jaisawal & Naik 2016). In our study, we used an absorbed cut-off power-law model with a blackbody component (`bbbodyrad` in XSPEC) to study all four data sets. The column density was pegged at a lower limit of $1 \times 10^{20} \text{ cm}^{-2}$ in the fit. A 6.4 keV Gaussian component for an iron fluorescence line is also needed. In addition to the emission, a cyclotron absorption feature appears in the data and is fitted with a Gaussian absorption line. The cyclotron lines are detected at $<30 \text{ keV}$ and at 31.5 keV in the 2015 and 2022 data sets, respectively (Table 2). We used the Markov chain Monte Carlo model in XSPEC




Figure 9. The broad-band energy spectra of the pulsar during its 2015 and 2022 outbursts are fitted with an absorbed cut-off power-law model along with a blackbody, and an iron emission line as well as a Gaussian absorption feature for a cyclotron line (top). The middle and bottom panels of each figure correspond to the spectral residuals with and without an iron line and cyclotron feature component to the best-fitting spectra, respectively. (a)–(d) denote the sets of *NuSTAR* and *Swift/XRT* (green) or *NICER* (magenta) data in Table 1, respectively.

for error estimation on the spectral parameters. The Goodman Ware algorithm with 20 walkers and a total length of 200 000 is considered in our study. The spectral parameters at a 90 per cent confidence interval are presented in Table 2. We also show the 68 per cent, 90 per cent, 95 per cent, and 99.7 per cent confidence level contour maps for cyclotron line energy and width for second and fourth *NuSTAR* observations in Fig. 10.

The cyclotron line energy, width, and line strength decreased with the luminosity during the 2015 outburst (Fig. 11). The second and fourth *NuSTAR* observations (see Table 2) are at almost the same luminosity level. However, the cyclotron line parameters, especially the cyclotron line energies, are found to be marginally higher (within 90 per cent level) in 2022 July. The magnetic field of the neutron star can be estimated to be 3.5×10^{12} G based on this recent detection.

Moreover, to study the model-independency of the cyclotron line, we show the parameter evolution using an absorbed NPEX model with a GABS component. The cyclotron parameters are consistent between both models (Fig. 11).

Following the broad-band fit, we found that the 0.5–10 keV luminosity represents about 40 per cent of the total pulsar emission in a 0.5–100 keV band. Therefore, an average bolometric correction of 2.5 can be applied to the outburst emission measured by *NICER*. This is consistent with the typical estimates where the soft band mostly contributes approximately 30–40 per cent of the pulsar emission (Anastasopoulou et al. 2022; Vasilopoulos et al. 2022). The 2022 outburst of SMC X-2 peaked at around MJD 59769.4 with a bolometric corrected luminosity of 3.2×10^{38} erg s^{−1}. This is close to half of the peak value the pulsar attained in the 2015 outburst.

Table 2. Best-fitting spectral parameters (with 90 % errors) of SMC X-2 obtained from four simultaneous *NuSTAR* and *Swift*/XRT or *NICER* observations. The fitted model consists of an absorbed cut-off power-law continuum with a blackbody and a Gaussian function for the iron emission line and cyclotron absorption line components.

Parameters	Obs-I + XRT	Obs-II + XRT	Obs-III + XRT	Obs-IV + <i>NICER</i>
N_H^a	$0.01^{+0.01}_{-0.01}$	$0.01^{+0.08}_{-0.01}$	$0.01^{+0.04}_{-0.01}$	$0.01^{+0.6}_{-0.01}$
Photon index	-0.16 ± 0.06	-1 ± 0.1	-1.33 ± 0.1	-0.77 ± 0.1
E_{cut} (keV)	6.4 ± 0.2	5.2 ± 0.3	5 ± 0.3	5.9 ± 0.3
kT_{bb} (keV)	1.1 ± 0.1	0.95 ± 0.04	0.95 ± 0.04	0.85 ± 0.02
Norm _{bb}	5.5 ± 1.5	9.1 ± 1.1	5.8 ± 0.7	12.4 ± 0.7
Radius _{bb} (km)	14.5 ± 2	18.7 ± 1.1	14.9 ± 0.9	21.8 ± 0.6
Fe line energy (keV)	6.39 ± 0.09	6.32 ± 0.07	6.37 ± 0.07	6.3 ± 0.1
Fe line eq. width (eV)	71 ± 10	72 ± 14	88 ± 18	83 ± 18
Cyclotron line energy (E_c) (keV)	26.8 ± 0.8	$28.1^{+1.7}_{-1.4}$	$29.5^{+2.7}_{-1.7}$	$31.5^{+1.3}_{-1.1}$
Cyclotron line width (σ_c) (keV)	5.5 ± 0.8	$5.5^{+1.5}_{-1.1}$	$6.6^{+2.3}_{-1.4}$	$6.5^{+0.9}_{-0.8}$
Cyclotron line strength (τ_c)	$3.9^{+1.1}_{-0.8}$	$4.4^{+2.9}_{-1.6}$	$6.2^{+6.9}_{-2.5}$	$8.7^{+3.1}_{-2.2}$
Luminosity ^b (0.5–10 keV)	2.75 ± 0.02	1.13 ± 0.01	0.7 ± 0.01	1.10 ± 0.01
Luminosity ^b (0.5–100 keV)	6.08 ± 0.02	2.87 ± 0.02	1.97 ± 0.01	2.82 ± 0.01
χ^2_ν (ν)	1.04 (1276)	0.96 (1125)	1.06 (602)	1.09 (1549)

^a Equivalent hydrogen column density (in 10^{22} atoms cm^{-2}).

^b Luminosity in 10^{38} erg s^{-1} units, assuming the source distance of 62 kpc (Hilditch et al. 2005; Haschke, Grebel & Duffau 2012; Graczyk et al. 2014).

Other parameters in Table 2 such as the blackbody temperature was found to be in a narrow range of 0.85–1.1 keV during both outbursts. We also measured the corresponding emission radius (Radius_{bb}) to examine the possible origin and location of thermal emission. The observed Radius_{bb} is estimated to be in the range of 14–22 km (see Table 2), which is close to the size of a neutron star. This suggests that the thermal component may originate from the neutron star surface or from the accretion column during the 2015 and 2022 outbursts.

4 DISCUSSION AND CONCLUSIONS

The broad-band emission from the accreting X-ray pulsars is understood to be due to thermal and bulk Comptonizations of seed photons from the hot spots within the accretion column mounted on the magnetic poles of the neutron star (Becker & Wolff 2007). The underlying physical phenomena of these sources can be understood by studying the pulse emission geometry and the spectral evolution of the pulsar during X-ray outbursts. At lower luminosities such as 10^{34} – 10^{35} erg s^{-1} , the accreted material falls freely until it gets halted and settled by Coulomb interactions in a hydrodynamical shock near the surface (Basko & Sunyaev 1975). This leads to a simplistic pulsed emission geometry in the form of a pencil beam where the X-ray photons propagate along the magnetic field lines. At this stage, the radiation is mostly dominated by bulk Comptonization (Becker & Wolff 2005, 2007). When the accretion rate increases, a radiation-dominated shock is expected to be formed in the accretion column at a critical luminosity that leads to geometrical changes or spectral transition in the emission (Basko & Sunyaev 1976; Becker & Wolff 2007). The infalling material interacts closely with the shock and gets decelerated through it before settling down on to the neutron star. The photons below the shock region diffuse through the column side-wall in the form of a fan beam pattern. The anticipated emission geometry is a mixture of fan and pencil beam patterns at a luminosity of $\sim 10^{36}$ – 10^{37} erg s^{-1} . The thermal as well as bulk Comptonization processes contribute significantly to the observed broad-band emission at this stage. In the supercritical luminosity regime ($> 10^{37}$ erg s^{-1}), a pure fan beam pattern is expected because of a strong radiative shock that makes the accretion column optically thick for photons propagating along the magnetic field lines.

SMC X-2 was observed with *NuSTAR* at luminosities above the Eddington limit of a classical neutron star during the 2015 and 2022 outbursts. The *NuSTAR* pulse profiles are found to be luminosity dependent. At a luminosity of 6.1×10^{38} erg s^{-1} (Obs-I), a doubled peaked profile appears throughout soft to hard X-rays suggesting the emission from both the poles of the neutron star as the observed peaks are separated by 0.5 phase in the pulse profile. The emission geometry, however, evolves below this luminosity, where the soft X-ray profiles are mainly broad or contain a hump-like structure. The emission from both poles is clearly apparent in the hard X-rays in the case of Obs-II and IV. The pulse fraction of the pulsar increases with energy clearly up to 40 keV, indicating an increase in the pulsating photons contributing to emission at higher luminosities. This suggests the rising height of the accretion column or mound. The profile changes with luminosity can be understood in terms of evolving emission geometry, where the significant contribution of the fan beam is anticipated in the supercritical regime.

We also detected a spectral transition in SMC X-2 during the 2022 outburst with *NICER*. The photon index shows a positive and negative dependency on luminosity below and above a range of value in the range of 6 – 8×10^{37} erg s^{-1} in 0.5–10 keV range. Considering a bolometric correction factor of 2.5, we expect a critical luminosity of 1.5 – 2×10^{38} erg s^{-1} at which the emission pattern is expected to change. According to Becker et al. (2012), the critical luminosity depends on the magnetic field assuming disk accretion on to a classical neutron star with a $1.4 M_\odot$ mass and a 10 km radius.

$$L_{\text{crit}} = 1.49 \times 10^{37} \left(\frac{B}{10^{12} \text{ G}} \right)^{16/15} \text{ erg s}^{-1} \quad (2)$$

Using equation (2), the magnetic field B can be estimated to be in the range of 8.7 – 11.4×10^{12} G for the above range of luminosity values. The estimated magnetic field differs by a factor of 3–4 from the measurement based on the detection of a cyclotron line in the spectrum (Jaisawal & Naik 2016). In case no bolometric correction is applied, the magnetic field from equation (2) turns out to be 3.7 – 4.8×10^{12} G, i.e. much closer to the estimates based on cyclotron line energy.




Figure 10. The 68 per cent, 90 per cent, 95 per cent, and 99.7 per cent (from innermost to outermost) confidence intervals contour maps between cyclotron line energy and width at an equivalent level of luminosity from 2015 (upper) and 2022 (bottom) outbursts. (b) and (d) have the usual meaning as denoted in Fig. 9. The ‘+’ sign corresponds to the best-fitting values for both parameters. The colour bars denote the parameter confidence range from 68 per cent to 99.7 per cent on a scale of zero to one, respectively.

4.1 The evolution of cyclotron absorption line and its implication on the magnetic field

Both the 2015 and 2022 X-ray outbursts of SMC X-2 lasted for more than 2 months with peak bolometric luminosities of 6×10^{38} erg s $^{-1}$ and 3×10^{38} erg s $^{-1}$, respectively (see Fig. 1). We report broad-band spectral properties of the pulsar during the latest giant outburst using data from *NICER* and *NuSTAR* observations. The 2015 *NuSTAR* and *Swift*/XRT observations are also used to study the cyclotron absorption line and its long-term evolution. We found relatively higher cyclotron line energy (with an increase of 3.4 keV) during the current outburst in comparison to the 2015 measurement made around a similar luminosity. In addition, the line energy is anticorrelated with luminosity in the 2015 outburst. The cyclotron line is a unique absorption-like feature that is observed in the 10–100 keV spectrum of X-ray pulsars (Meszaros 1992). They appear due to a resonance scattering of photons with electrons in quantized Landau energy states in the presence of a strong magnetic field. The field strength can be estimated directly using a 12-B-12 rule




Figure 11. Change in the cyclotron line parameters such as line energy, width, and its strength (τ_{cyc}) with 0.5–100 keV unabsorbed luminosity. The measurements from the 2015 and 2022 outbursts are shown in solid circles and squares, respectively. The red and blue colours correspond to line parameters obtained after an absorbed cut-off power-law model along with blackbody and GABS, and an absorbed NPEX model with a GABS model, respectively.

depending on the line energy since it is expressed as $E_{\text{cyc}} = 11.6 B_{12} \times (1+z)^{-1}$ keV, where B_{12} is the magnetic field in 10^{12} G unit, and z is the gravitational red-shift.

The centroid energy of the cyclotron line in principle measures the field strength at the line-forming region. A change in line energy with luminosity is observed in several accretion-powered X-ray pulsars during outbursts (Nakajima et al. 2006; Tsygankov, Lutovinov & Serber 2010; Jaisawal, Naik & Epili 2016; Staubert et al. 2019). In some cases like Her X-1 (Staubert et al. 2007), the cyclotron energy and luminosity are positively correlated, whereas there are pulsars such as 4U 0115+63 (Nakajima et al. 2006; Tsygankov et al. 2010) and V 0332+53 (Tsygankov et al. 2010) that show a negative correlation between these parameters (Becker et al. 2012). The positive correlation is expected in the sub-critical region (below the critical luminosity) when the pressure of the accreting plasma pushes the hydrodynamical shock or line-forming region closer to the neutron star surface with an increasing accretion rate (luminosity). On the other hand, a formation of radiation-dominated shock is expected in the accretion column at higher accretion rates. The shock drifts upwards in the column due to strong radiation pressure with increasing mass accretion rate in the supercritical regime (Becker et al. 2012). Therefore, a negative correlation between the cyclotron line energy and luminosity is expected at this phase. This is observed in SMC X-2 when the pulsar was accreting in the supercritical regime during the 2015 outburst. Alternatively, the formation of a cyclotron line and its observed negative correlation can be understood by the reflection of accretion column photons on to the neutron star surface

(Poutanen et al. 2013). The hypothesis suggests that the cyclotron line gets formed on the surface rather than in the accretion column where the gradient is high to explain the observed changes in the line energy. As the accretion rate increases, a larger accretion column covers a substantial portion of the atmosphere from the pole to the equator (high to low magnetic field regions). This can also explain the origin of the observed negative correlation between the cyclotron energy and luminosity in SMC X-2 resulted due to a change in illumination pattern.

It is important to note that the cyclotron line energy of SMC X-2 is different at a given luminosity observed at Obs-II and Obs-IV during different outbursts. The *MAXI*/GSC light curves in Fig. 1 show a clear rise in the source intensity after MJD 57259 and 59738 during the 2015 and 2022 outbursts, respectively. The source reached a luminosity of $2\text{--}3 \times 10^{38} \text{ erg s}^{-1}$ (Obs-II and IV) after 49 and 36 d since the onset of respective outbursts. Our measurements suggest an increase of 3.4 keV in the cyclotron line energy between these data sets at comparable luminosities. This difference in the cyclotron line energy corresponds to a change in the magnetic field of 3.8×10^{11} G, assuming a classical neutron star and gravitational redshift of 0.3. Such an enhancement in the cyclotron energy can be understood by an accretion-induced screening of the magnetic field or geometrical changes in the line-forming region. Theoretical studies suggest that the accretion flow distribution from poles to lower latitudes may drag the field lines (Choudhuri & Konar 2002 and references therein). This mechanism could effectively lower down the original field strength even on a shorter time-scale in the top layer of the distributed material, considering no effect of magnetic buoyancy. However, in reality, the screening mechanism depends on the ratio of time-scales for field burial by the advection mechanism and the re-emerging of the field through magnetic buoyancy and Ohmic rediffusion (Choudhuri & Konar 2002). An increase of 3.4 keV in cyclotron line energy in the latest modest outburst suggests that the screening mechanism is not effective as it is in the case of the 2015 giant outburst, where a similar luminosity is achieved in the decay phase almost 49 d after the onset. The neutron star by this point would accumulate a factor of two or more mass with an integrated value of $3.5 \times 10^{24} \text{ g}$ in 2015.

A similar possibility is discussed for V 0332+53 during its 2004/05 outburst where the cyclotron line energy at a given luminosity in the decay got reduced by $\approx 1.5 \text{ keV}$ with respect to the rising phase of the outburst (Cusumano et al. 2016). However, Doroshenko et al. (2017) argued that the expected time-scales for field burring/reemerging are significantly shorter than those observed in the case of Vela X-1 and Her X-1. They also noticed discrepancies between the spin rates in rising and declining parts of the outburst that could explain the difference in the magnetic field based on the accretion-torque theory. In addition, they suggested that the change in the accretion disk and effective magnetospheric radius in different phases of the outburst lead to a variable column height, resulting in an evolving cyclotron line through X-ray reflection on the surface (Doroshenko et al. 2017). In the case of SMC X-2, however, we observe the increase in cyclotron line energy in the decay phase of two outbursts at a comparable luminosity. This is important to recall that the pulse profile observed during Obs-II and IV appears identical, suggesting similar accretion geometry of the pulsar. The accretion-induced screening mechanism, thus, seems a possible explanation for the observed behaviour over the geometrical change in the line-forming region.

The long term evolution of the cyclotron line is well studied for two sources, Her X-1 and Vela X-1 (La Parola et al. 2016; Staubert et al. 2019). The cyclotron centroid energy of Her X-1 has shown a secular decay of $\approx 5 \text{ keV}$ in 20 yr before stabilizing between 2012 and

2019. There was also a period in 1993–1996 when the line energy was increased by a few keV. The time evolution of the cyclotron line energy is still a matter of study. It is believed that the observed decay/increase in line energy represents a geometrical change within the emission region or in the configuration of the local magnetic field rather than a change in the global dipolar magnetic field of the neutron star (Mukherjee, Bhattacharya & Mignone 2013; Staubert et al. 2019). This could be further understood in terms of a marginal imbalance between the accretion rate and the losing or settling rate of the accreted material on the accretion mound. If the settling rate is slightly higher than the leakage of the material from the bottom of the mound, this may lead to an increase in the column height as well as a change in the local magnetic field configuration of the line forming region. A line decay is expected naturally in this case. Over time, the excessive accumulation of the material and its pressure on the mound may result in a forced redistribution of the material across the surface. This would adjust the accretion mount to its unperturbed configuration where higher line energy would be expected. The time-scale for such redistribution is unclear. But, it may happen after continuous accumulation of matter over a long period or (multiple) strong outbursts like those observed in 2002 and 2015 of SMC X-2. Future observations of the source during outbursts will allow understanding the behaviour of the cyclotron line better.

ACKNOWLEDGEMENTS

We thank the referee for constructive suggestions on the paper. This research has made use of data obtained through HEASARC Online Service, provided by the NASA/GSFC, in support of NASA High Energy Astrophysics Programs. This work used the NuSTAR Data Analysis Software (NUSTARDAS) jointly developed by the ASI Science Data Center (ASDC, Italy) and the California Institute of Technology (USA). SG acknowledges the support of the CNES. GV acknowledges support by H.F.R.I. through the project ASTRAPE (Project ID 7802).

DATA AVAILABILITY

We used archival data of *NICER*, *NuSTAR*, *Swift*, and *MAXI* observatories for this work.

REFERENCES

- Anastasopoulou K., Zezas A., Steiner J. F., Reig P., 2022, *MNRAS*, 513, 1400
- Arnaud K. A., 1996, in Jacoby G. H., Barnes J., eds, ASP Conf. Ser. Vol. 101, Astronomical Data Analysis Software and Systems V, Astron. Soc. Pac., San Francisco, p. 17
- Basko M. M., Sunyaev R. A., 1975, *A&A*, 42, 311
- Basko M. M., Sunyaev R. A., 1976, *MNRAS*, 175, 395
- Becker P. A., Wolff M. T., 2005, *ApJ*, 630, 465
- Becker P. A., Wolff M. T., 2007, *ApJ*, 654, 435
- Becker P. A. et al., 2012, *A&A*, 544, A123
- Choudhuri A. R., Konar S., 2002, *MNRAS*, 332, 933
- Clark G., Doxsey R., Li F., Jernigan J. G., van Paradijs J., 1978, *ApJ*, 221, L37
- Clark G., Li F., van Paradijs J., 1979, *ApJ*, 227, 54
- Coe M. J., Kirk J., 2015, *MNRAS*, 452, 969
- Coe M. J., Kennea J. A., Evans P. A., Buckley D. A. H., Townsend L. J., Monageng I., 2022, *Astron. Telegram*, 15500, 1
- Corbet R. H. D., Marshall F. E., Coe M. J., Laycock S., Handler G., 2001, *ApJ*, 548, L41
- Cusumano G., La Parola V., D’Ai A., Segreto A., Tagliaferri G., Barthelmy S. D., Gehrels N., 2016, *MNRAS*, 460, L99

- Dickey J. M., Lockman F. J., 1990, *ARA&A*, 28, 215
- Doroshenko V., Tsygankov S. S., Mushtukov A. A., Lutovinov A. A., Santangelo A., Suleimanov V. F., Poutanen J., 2017, *MNRAS*, 466, 2143
- Evans P. A. et al., 2009, *MNRAS*, 397, 1177
- Gendreau K. C., Arzoumanian Z., Okajima T., 2012, in Proc. SPIE Conf. Ser., p. 13,
- Gendreau K. C., et al., 2016, in Space Telescopes and Instrumentation 2016: Ultraviolet to Gamma Ray. p. 99051H
- Graczyk D. et al., 2014, *ApJ*, 780, 59
- Haberl F., Sturm R., 2016, *A&A*, 586, A81
- Harrison F. A. et al., 2013, *ApJ*, 770, 103
- Haschke R., Grebel E. K., Duffau S., 2012, *AJ*, 144, 107
- Hilditch R. W., Howarth I. D., Harries T. J., 2005, *MNRAS*, 357, 304
- Huppenkothen D. et al., 2019, *ApJ*, 881, 39
- Jaisawal G. K., Naik S., 2016, *MNRAS*, 461, L97
- Jaisawal G. K., Naik S., Epili P., 2016, *MNRAS*, 457, 2749
- Kahabka P., Pietsch W., 1996, *A&A*, 312, 919
- Karaferias A. S., Vasilopoulos G., Petropoulou M., Jenke P. A., Wilson-Hodge C. A., Malacaria C., 2023, *MNRAS*, 520, 281
- Kennea J. A. et al., 2015, *Astron. Telegram*, 8091, 1
- Kennea J. A., Coe M. J., Evans P. A., 2022, *Astron. Telegram*, 15434, 1
- La Palombara N., Sidoli L., Pintore F., Esposito P., Mereghetti S., Tiengo A., 2016, *MNRAS*, 458, L74
- La Parola V., Cusumano G., Segreto A., D’Aì A., 2016, *MNRAS*, 463, 185
- Leahy D. A., 1987, *A&A*, 180, 275
- Li K. L., Hu C. P., Lin L. C. C., Kong A. K. H., 2016, *ApJ*, 828, 74
- Liu Q. Z., van Paradijs J., van den Heuvel E. P. J., 2006, *A&A*, 455, 1165
- Luo J. et al., 2021, *ApJ*, 911, 45
- Lutovinov A. A., Tsygankov S. S., Krivonos R. A., Molkov S. V., Poutanen J., 2017, *ApJ*, 834, 209
- McBride V. A., Coe M. J., Negueruela I., Schurch M. P. E., McGowan K. E., 2008, *MNRAS*, 388, 1198
- Makishima K., Mihara T., Nagase F., Tanaka Y., 1999, *ApJ*, 525, 978
- Marshall F. E., Boldt E. A., Holt S. S., Mushotzky R. F., Pravdo S. H., Rothschild R. E., Serlemitsos P. J., 1979, *ApJS*, 40, 657
- Matsuoka M. et al., 2009, *PASJ*, 61, 999
- Meszaros P., 1992, *Theoretical Astrophysics*, University of Chicago Press, Chicago
- Mukherjee D., Bhattacharya D., Mignone A., 2013, *MNRAS*, 430, 1976
- Nakajima M., Mihara T., Makishima K., Niko H., 2006, *ApJ*, 646, 1125
- Negoro H. et al., 2015, *Astron. Telegram*, 8088, 1
- Porter J. M., Rivinius T., 2003, *PASP*, 115, 1153
- Poutanen J., Mushtukov A. A., Suleimanov V. F., Tsygankov S. S., Nagirner D. I., Doroshenko V., Lutovinov A. A., 2013, *ApJ*, 777, 115
- Prigozhin G. et al., 2012, in High Energy, Optical, and Infrared Detectors for Astronomy V. p. 845318
- Reig P., 2011, *Ap&SS*, 332, 1
- Remillard R. A. et al., 2022, *AJ*, 163, 130
- Roy A., Cappallo R., Laycock S. G. T., Christodoulou D. M., Vasilopoulos G., Bhattacharya S., 2022, *ApJ*, 936, 90
- Russell S. C., Dopita M. A., 1992, *ApJ*, 384, 508
- Schurch M. P. E., Coe M. J., McBride V. A., Townsend L. J., Udalski A., Haberl F., Corbet R. H. D., 2011, *MNRAS*, 412, 391
- Seward F. D., Mitchell M., 1981, *ApJ*, 243, 736
- Staubert R., Shakura N. I., Postnov K., Wilms J., Rothschild R. E., Coburn W., Rodina L., Klochkov D., 2007, *A&A*, 465, L25
- Staubert R. et al., 2019, *A&A*, 622, A61
- Townsend L. J., Coe M. J., Corbet R. H. D., Hill A. B., 2011, *MNRAS*, 416, 1556
- Tsygankov S. S., Lutovinov A. A., Serber A. V., 2010, *MNRAS*, 401, 1628
- Vasilopoulos G., Haberl F., Carpano S., Maitra C., 2018, *A&A*, 620, L12
- Vasilopoulos G., Jaisawal G. K., Maitra C., Haberl F., Maggi P., Karaferias A. S., 2022, *A&A*, 664, A194
- Verner D. A., Ferland G. J., Korista K. T., Yakovlev D. G., 1996, *ApJ*, 465, 487
- Vinciguerra S. et al., 2020, *MNRAS*, 498, 4705
- Walter R., Lutovinov A. A., Bozzo E., Tsygankov S. S., 2015, *A&A Rev.*, 23, 2
- Wilms J., Allen A., McCray R., 2000, *ApJ*, 542, 914
- Yang J., Laycock S. G. T., Christodoulou D. M., Fingerman S., Coe M. J., Drake J. J., 2017, *ApJ*, 839, 119
- Zolotukhin I. Y., Bachetti M., Sartore N., Chilingarian I. V., Webb N. A., 2017, *ApJ*, 839, 125

This paper has been typeset from a $\mathrm{\LaTeX}$ file prepared by the author.