



**HAL**  
open science

## How to create the highest manganese ore grade in the world? A geodynamic story in the Atlas of Morocco.

Augustin Dekoninck, Jocelyn Barbarand, Gilles Ruffet, Yves Missenard, Nadine Mattielli, Rémi Leprêtre, Abdellah Mouttaqi, Omar Saddiqi, Michèle Verhaert, Johan Yans

### ► To cite this version:

Augustin Dekoninck, Jocelyn Barbarand, Gilles Ruffet, Yves Missenard, Nadine Mattielli, et al.. How to create the highest manganese ore grade in the world? A geodynamic story in the Atlas of Morocco.. EGU General Assembly 2023, Apr 2023, Vienna, Austria. pp. EGU23-5478, 2023, 10.5194/egusphere-egu23-5478 . insu-04083146

**HAL Id: insu-04083146**

**<https://insu.hal.science/insu-04083146>**

Submitted on 27 Apr 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# How to create the highest manganese ore grade in the world?

## A geodynamic story in the Atlas of Morocco




Augustin DEKONINCK<sup>(1,2)</sup>, Jocelyn BARBARAND<sup>(3)</sup>, Gilles RUFFET<sup>(4,5)</sup>, Yves MISSENARD<sup>(3)</sup>, Nadine MATTIELLI<sup>(1)</sup>, Rémi LEPRETRE<sup>(3,6)</sup>, Abdellah MOUTTAQI<sup>(7)</sup>, Omar SADDIQI<sup>(8)</sup>, Michèle VERHAERT<sup>(2)</sup>, Johan YANS<sup>(2)</sup>

### Introduction

The Imini mining district hosts the **largest Mn resources of North Africa** due to two laterally extensive ore bodies (~25-30 km) of **nearly pure pyrolusite** (MnO<sub>2</sub>). A third medium-grade coronadite-rich (K,Ba,PbMn<sub>8</sub>O<sub>16</sub>) Mn ore occurs discontinuously over a wider area of ~100 km. The two ore types are hosted in a poorly deformed 10-15m-thick Cenomanian-Turonian dolostone unit located in the southern foreland of the High Atlas of Morocco. Until now, the **source of the metals and the timing** of ore deposition remains unclear for this unconventional karst-hosted deposit.

### Objectives and methods

We examine the **Pb isotope** composition of the Mn ores and the most representative country rocks to track down potential sources of the metals. We also study the age of the K-bearing Mn oxides present in the Mn ore by <sup>40</sup>Ar/<sup>39</sup>Ar. By combining these two methods we are able to produce a consistent model of the ore deposition.



### Key interpretations

- **Atlas tectonics** is the first-order control of the Mn mineralization.
- Periods with regionally high elevations modified the climate to wetter conditions that brought rather **continuous supplies of meteoric water** to feed temporary aquifers.
- **Relief building created the required hydraulic head** to sustain (1) fluid-rock interaction between O<sub>2</sub>-poor acidic fluids and the Triassic series, (2) migration of the metal-rich fluid and (3) overpressure fluid in the Imini depositional site.
- The disappearance of Triassic series in front of the Imini anticline forced these low-temperature (<100°C) fluids to **mix with oxygenated ground and alkaline waters** resident in the karst system and precipitate Mn oxides.
- The N70°-oriented **Atlas tectonic structure controls the orientation** of the Mn deposits.
- Post-Cretaceous uplifts generated **additional supplies and/or in-situ remobilizations** of the primary late Cretaceous medium-grade ore to form the high-grade pyrolusite-rich ore in Imini.

### Results

The new Pb isotopic ratios show that **Triassic series (basalts and Fe-rich sandstones) are likely the source** of the metals. The <sup>40</sup>Ar/<sup>39</sup>Ar dating of these K-Mn oxides shows that the Mn-rich ore bodies formed during at least **three periods**:

- late Cretaceous to late Paleocene (>58 Ma)
- late Eocene (c. 36.3 Ma)
- early Burdigalian to early Serravalian (c. 19–20 Ma and c. 13 Ma)

These periods **coincide with three known Atlas geodynamic deformation** events during late Cretaceous, late Eocene and the Early(?)–Middle Miocene.

