



The source of electrons at a weakly outgassing comet

Peter Stephenson, Kathrin Altwegg, Arnaud Beth, Jim Burch, Christopher Carr, Jan Deca, Anders Eriksson, Marina Galand, Karl-Heinz Glassmeier, Charlotte Goetz, et al.

► To cite this version:

Peter Stephenson, Kathrin Altwegg, Arnaud Beth, Jim Burch, Christopher Carr, et al.. The source of electrons at a weakly outgassing comet. 16th Europlanet Science Congress 2022, 2022, à renseigner, Unknown Region. 10.5194/epsc2022-1047 . insu-04089892

HAL Id: insu-04089892

<https://insu.hal.science/insu-04089892>

Submitted on 5 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



The source of electrons at a weakly outgassing comet

Peter Stephenson¹, Kathrin Altwegg², Arnaud Beth³, Jim Burch⁴, Christopher Carr¹, Jan Deca⁵, Anders Eriksson⁶, Marina Galand¹, Karl-Heinz Glassmeier⁷, Charlotte Goetz⁸, Pierre Henri^{9,10}, Kevin Heritier¹, Fredrik Johansson⁶, Zoe Lewis¹, Hans Nilsson¹¹, and Martin Rubin²

¹Imperial College London, Physics, London, United Kingdom of Great Britain – England, Scotland, Wales

(p.stephenson18@imperial.ac.uk)

²Physikalisches Institut, University of Bern, Bern, Switzerland

³Department of Physics, Umeå University, Umeå, Sweden

⁴South West Research Institute, San Antonio, Texas

⁵Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, Colorado

⁶Swedish Institute of Space Physics, Uppsala, Sweden

⁷Institut für Geophysik und extraterrestrische Physik, TU Braunschweig, Braunschweig, Germany

⁸ESTEC, European Space Agency, Noordwijk, the Netherlands

⁹Lagrange, OCA, CNRS, UCA, Nice, France

¹⁰LPC2E, CNRS, Orleans, France

¹¹Swedish Institute of Space Physics, Kiruna, Sweden

The Rosetta spacecraft escorted comet 67P/Churyumov-Gerasimenko for two years along its orbit, from Aug 2014 to Sep 2016, observing the evolution of the comet from a close perspective. The Rosetta Plasma Consortium (RPC) monitored the plasma environment at the spacecraft throughout the escort phase.

Cometary electrons are produced by ionization of the neutral gas coma. This occurs through photoionization by extreme ultraviolet photons, and through electron-impact ionization (EII) by collisions of energetic electrons with the coma. Far from perihelion, EII is, at times, more prevalent than photoionization (Galand et al., 2016; Heritier et al., 2018), but the EII frequency has not been assessed across the whole mission. The source of the cometary electrons, and the origin of the ionizing electrons is still unclear.

We have calculated the electron impact ionization (EII) frequency throughout the Rosetta mission and at its location from measurements of RPC's Ion and Electron Sensor (RPC/IES). EII ionization is confirmed as the dominant source of cometary electrons and ions when far from perihelion but is much more variable than photoionization. We compare the EII frequency with properties of the neutral coma and cometary plasma to identify key drivers of the energetic electron population. The EII frequency is structured by outgassing rate and magnetic field strength.

The first 3D collision model of electrons at a comet (Stephenson et al. 2022) is also utilised to assess the origin of electrons within the coma. The model uses self-consistently calculated electric and magnetic fields from a fully-kinetic and collisionless Particle-in-Cell model (Deca et al. 2017, 2019) as an input. The modelling approach confirms cometary electrons are produced by impacts of energetic electrons at low outgassing. The ionizing electrons are identified as solar wind electrons

that have undergone acceleration in an ambipolar potential well, confirming the results of the data analysis.

Deca J, Divin A, Henri P, Eriksson A, Markidis S, et al. 2017. *Physical Review Letters*. 118(20):205101

Deca J, Henri P, Divin A, Eriksson A, Galand M, et al. 2019. *Physical Review Letters*. 123:55101

Galand M, H  ritier K L., Odelstad E, Henri P, Broiles TW., et al. 2016. *Monthly Notices of the Royal Astronomical Society*. 462:S331–51

Heritier KL, Galand M, Henri P, Johansson FL, Beth A, et al. 2018. *A&A*, p. 618:A77–A77

Stephenson P, Galand M, Deca J, Henri P, Carnielli G. 2022. *Monthly Notices of the Royal Astronomical Society*. 511(3):4090–4108