



HAL
open science

Cold electrons at a weakly outgassing comet

Peter Stephenson, Marina Galand, Jan Deca, P. Henri

► **To cite this version:**

Peter Stephenson, Marina Galand, Jan Deca, P. Henri. Cold electrons at a weakly outgassing comet. 16th Europlanet Science Congress 2022, 2022, à renseigner, Unknown Region. 10.5194/epsc2022-972 . insu-04089893

HAL Id: insu-04089893

<https://insu.hal.science/insu-04089893>

Submitted on 5 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Cold electrons at a weakly outgassing comet

Peter Stephenson¹, Marina Galand¹, Jan Deca², and Pierre Henri^{3,4}

¹Imperial College London, Physics, London, United Kingdom of Great Britain – England, Scotland, Wales

(p.stephenson18@imperial.ac.uk)

²Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, Colorado

³Lagrange, OCA, CNRS, UCA, Nice, France

⁴LPC2E, CNRS, Orleans, France

The Rosetta spacecraft rendezvoused with comet 67P/Churyumov-Gerasimenko in August 2014 and escorted it for two years until September 2016. The plasma surrounding the comet was probed by the Rosetta Plasma Consortium (RPC), comprising five instruments. The Langmuir Probe (RPC/LAP; Engelhardt et al., 2018) and Mutual Impedance Probe (RPC/MIP; Wattieaux et al, 2020; Gilet et al., 2020) measured a cold population of electrons (< 1 eV) within the cometary environment, that was persistent throughout the mission.

Cometary electrons are typically produced at 10 eV through ionization of the neutral gas coma. This is either through photoionization, by absorption of an extreme ultraviolet (EUV) photon, or electron-impact ionization, by collisions of energetic electrons with the coma. Cold electrons are formed by cooling the warm, newly born electrons, through inelastic collisions with the cometary neutrals. Assuming they flow radially away from the nucleus, the electron cooling should only be significant for the high outgassing rates found around perihelion ($Q > 3 \times 10^{27} \text{ s}^{-1}$). However, the cold population was observed at large heliocentric distances (> 3.5 au) and very weak outgassing rates ($Q < 10^{26} \text{ s}^{-1}$).

We have developed the first 3D collisional model of electrons at a comet, featuring a spherically symmetric coma of pure water. Electric and magnetic fields are self-consistently calculated using a fully-kinetic, collisionless Particle-in-Cell simulation model (Deca et al., 2017; 2019), which are used as an input for the test particle model. Electron-neutral collisions are treated as a stochastic process. The collision processes include elastic scattering, inelastic excitations, and electron-impact ionization.

With the test particle model, we demonstrate that electrons are trapped in an ambipolar potential well around the nucleus. This greatly increases the efficiency of collisional processes compared to the case of radial outflow. A cold electron population is formed at weak outgassing rates ($Q = 10^{26} \text{ s}^{-1}$), far below the threshold for cooling under radial outflow.

We quantify the increased efficiency of electron collisions when using the complex electric and magnetic fields, compared to radial outflow. The location of the electron cooling exobase is calculated accounting for electron trapping and compared with the cold electron measurements from Rosetta. This explains the cold electron observations at low outgassing rates.

Deca J, Divin A, Henri P, Eriksson A, Markidis S, et al. 2017. *Physical Review Letters*. 118(20):205101

Deca J, Henri P, Divin A, Eriksson A, Galand M, et al. 2019. *Phys. Rev. Lett.* 123:55101

Engelhardt IAD, Eriksson AI, Vigren E, Valières X, Rubin M, et al. 2018. 15

Gilet N, Henri P, Wattieaux G, Traoré N, Eriksson AI, et al. 2020. *A&A*. 640:A110–A110

Wattieaux G, Henri P, Gilet N, Vallières X, Deca J. 2020. *A&A*. 638:A124–A124