



HAL
open science

Current and projected patterns of warming and marine heatwaves in the Southern Indian Ocean

Clara Azarian, Laurent Bopp, Alice Pietri, Jean-Baptiste Sallée, Francesco d'Ovidio

► **To cite this version:**

Clara Azarian, Laurent Bopp, Alice Pietri, Jean-Baptiste Sallée, Francesco d'Ovidio. Current and projected patterns of warming and marine heatwaves in the Southern Indian Ocean. *Progress in Oceanography*, 2023, 215, 10.1016/j.pocean.2023.103036 . insu-04139547

HAL Id: insu-04139547

<https://insu.hal.science/insu-04139547>

Submitted on 10 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Current and projected patterns of warming and marine heatwaves in the**

2 **Southern Indian Ocean**

3
4
5
6
7
8 Clara Azarian^{1,2}, Laurent Bopp³, Alice Pietri⁴, Jean-Baptiste Sallée¹, Francesco d'Ovidio¹

9
10
11 1 Sorbonne Université, CNRS, IRD, MNHN, Laboratoire d'Océanographie et du Climat:
12 7 Expérimentations et Approches Numériques (LOCEAN-IPSL), Paris, France.

13 8 2 Ecole Nationale des Ponts et Chaussées (ENPC), Champs-sur-Marne, France.

14 9 3 Ecole Normale Supérieure / Université PSL, CNRS, Ecole Polytechnique, Sorbonne
15 10 Université, Paris, PSL University, Laboratoire de Météorologie Dynamique (LMD-IPSL) Paris,
16 11 France.

17 12 4 Instituto del Mar del Peru (IMARPE), Callao, Peru.

18
19
20
21
22 15 Corresponding author: C. Azarian; clara.azarian@locean.ipsl.fr
23 16
24
25
26
27 17
28
29
30
31
32 18
33
34
35
36 19
37
38
39
40 20
41
42
43
44 21
45
46
47
48
49 22
50
51
52
53 23
54
55
56
57 24
58
59
60
61
62
63
64
65

25 Abstract

26

27 The Southern Indian Ocean (20-120°E, 70-30°S) hosts an exceptional biodiversity that contributed to
28 the inscription of the French and Australian natural reserves on the UNESCO World Heritage List. This
29 region is a “hot spot” for ocean heat uptake and already experiences intense marine heat waves (MHW),
30 as evidenced in 2011/2012 over the Kerguelen Plateau. In the coming decades, this region is also
31 expected to face supplemental anthropogenic warming, depending on future greenhouse gas emissions,
32 with unknown consequences for its marine ecosystems. Here, we present a regional analysis of ocean
33 warming and MHW based on the analyses of historical observations and Coupled Model
34 Intercomparison Project Phase 6 (CMIP6) climate projections. Consistent with observations over the
35 last decades, we find an intensification through the 21st century of surface warming and MHW over a
36 band located between 40°S and 55°S within the Antarctic Circumpolar Current region. CMIP6 models
37 also project much faster climate velocities (i.e. the speed and direction at which isotherms drift in the
38 wake of climate change) in the mesopelagic (200-1000m) than at the surface (0-200m). Lastly, a
39 comparison between the two Shared Socioeconomic Pathways (SSP1-2.6 and SSP2-4.5) analysed in
40 this study shows much larger changes in the second half of the 21st century for the higher emission
41 scenario. These results suggest that the subantarctic islands will probably be mostly affected by
42 warming and MHW under both scenarios, although committing to SSP1-2.6 could substantially
43 alleviate the pressure on ecosystems in the long term. This study also highlights the need to consider a
44 tri-dimensional environment that may evolve at different paces when designing efficient conservation
45 measures.

46

47 Keywords

48

49 global climate models, ocean warming, ocean extremes, Southern Indian Ocean

50

51 1 Introduction

52

53 **Ocean circulation.** The Southern Indian Ocean (20-120°E, 70-30°S) is characterized by dynamic jets
54 and a system of fronts associated with the Antarctic Circumpolar Current (ACC). The ACC is a strong
55 eastward current (Donohue et al., 2016), driven by strong westerly winds (Rintoul and Naveira
56 Garabato, 2013). As the ACC encounters the Kerguelen Plateau, it divides in several branches, with
57 most of the ACC transport passing north of the Plateau (Park et al., 1991,1993, 2009). A strong jet is
58 also observed through the Fawn Trough, a deep passage across the Plateau at 56°S (Park et al., 2008;
59 Roquet et al., 2009; Van Wijk et al., 2010; Vivier et al., 2015; Figure 1). Northwest of the Southern
60 Indian Ocean, the Agulhas Current, a western boundary current, flows southward along the east coast
61 of Africa. At around 40°S, this flow is split with the Agulhas Return Current transporting water eastward
62 (Lutjharms et al., 2006) and the Agulhas leakage transporting water to the South Atlantic (Schmidt et
63 al., 2021). Generally, the frontal system and the circulation in the Southern Indian Ocean is an important
64 biogeographical driver of phytoplankton and zooplankton communities which are at the basis of rich
65 ecosystems (Hunt et al., 2001; Matsuno et al., 2020; Mishra et al., 2020; Venkataramana et al., 2020;
66 Cotte et al., 2022).

67

68 **Conservation efforts to protect a rich biodiversity.** The Southern Indian Ocean region hosts several
69 endemic species, including large populations of megafauna of high patrimonial and economical value
70 like king penguins, yellow-nosed albatrosses, southern elephant seals, krill, toothfish, and many others
71 (Delord et al., 2014). To protect the rich biodiversity of the Southern Indian Ocean, conservation
72 measures have been implemented both in the exclusive economic zones (EEZs) of the subantarctic
73 islands and in the areas under the Commission for the Conservation of Antarctic Marine Living
74 Resources (CCAMLR). Part of EEZs of the following territories, Prince Edward, Crozet, Kerguelen

75 and Heard and McDonald's islands, are managed through Marine Protected Areas (MPAs). The French
76 national natural reserve (i.e. managed area with zones under strong or integral protection) within Crozet,
77 Kerguelen and Saint-Paul and Amsterdam EEZ was created in 2006 and extended in 2016. The South
78 African MPA around Prince Edward Islands have been officially implemented in 2013 and according
79 to CCAMLR it contributes to a representative system of MPAs, to be used as a scientific reference for
80 future management plans, to foster the recovery of Patagonian toothfish populations and to reduce the
81 bycatch of toothfish fishery on marine seabirds. The Australian MPA around Heard and McDonald
82 Islands was created in 2002 and has been extended in 2014. In total, 5.8% of the Southern Indian Ocean
83 is covered by MPAs and only within EEZs. The new frontier for conservation is now the vast open
84 ocean region that extends beyond EEZs (Della Penna et al. 2017), known as the "High Seas". A
85 workshop held in 2019 organized by the CCAMLR Scientific Committee developed a scientific work
86 program for pelagic spatial planning in the eastern subantarctic region to investigate the relevance of
87 adding new subantarctic spatial conservation tools in the High Seas of the CCAMLR area (Makhado et
88 al., 2019).

89
90 **Conservation and climate change.** One key step in the design of a conservation plan is the evaluation
91 of threats, including climate change (Meredith et al., 2019). The sector directly northward and within
92 the Antarctic Circumpolar Current (ACC), corresponding to the latitudinal band 30°-50°S, in the
93 Southern Indian Ocean is one of the regions of the world that has experienced the largest increase in
94 ocean heat content in recent decades (Llovel & Terray, 2016; Roemmich et al., 2015; Sallée, 2018; Fox-
95 Kemper et al., 2021). This large heat content increase has been explained by an increased heat uptake
96 in the subpolar region transported northward by Ekman transport and subducted within and north of the
97 ACC (Armour et al., 2016; Frölicher et al., 2015; Morrison et al., 2016; Sallée, 2018). In addition to
98 large-scale processes (> 100 km), finer scale processes (1-100 km), which are key for the Southern
99 Indian Ocean circulation (Kostianoy et al., 2004) and can be expected to intensify (Martinez-Moreno
100 et al., 2019; Martinez-Moreno et al., 2021), could also potentially modulate climate change signal
101 locally, maybe inducing important spatial heterogeneity in the temperature trends (e.g. anomalous

102 vertical heat transport driven by submesoscale fronts altering oceanic heat uptake, Siegelman et al.,
103 2020).

104

105 In the Southern Indian Ocean, species distributions are strongly shaped by the large meridional
106 temperature gradient characterizing the region, which covers a transition zone from the subtropic to the
107 Antarctic. Some species are even living close to their thermal tolerance limit (e.g., icefish, Kock &
108 Everson, 2003; Collins et al., 2010) and it has been observed that warming could have a negative impact
109 on the abundance of subantarctic krill, flagellates and Notothenioid fish (Constable et al. 2014) or even
110 potentially on the juvenile recruitment of Patagonian toothfish (Belchier and Collins, 2008). Climate
111 velocity, that is, the speed and direction at which isotherms drift in the wake of climate change (Loarie
112 et al., 2009), can be a useful tool to study biodiversity redistributions due to climate change (Garcia
113 Molinos et al., 2016). Although there is no specific studies on the Southern Ocean using this tool, there
114 are global studies that tend to show that the Southern Ocean may be a generally « slow-moving » basin
115 compared to other regions (Burrows et al., 2014) but also that there may be an important spatial
116 heterogeneity of this indicator (Brito-Morales et al., 2020).

117

118 Given the rich biodiversity and the observed and projected warming trends in the Southern Indian Ocean
119 (Fox-Kemper et al., 2021), there is an urgent need to integrate climate change in biodiversity
120 conservation and management. However, if climate change is increasingly presented as one of the
121 threats on biodiversity in conservation plans, such a threat is rarely quantified and no associated actions
122 are indicated. This issue will also likely be raised in line with the momentum fostered by the
123 negotiations of a treaty on the conservation and sustainable use of marine biodiversity in areas beyond
124 national jurisdiction (BBNJ), notably for the development of MPAs in the High Seas (Ban et al., 2014;
125 Maxwell et al., 2020; Crespo et al., 2020). Despite an increasing awareness that climate change might
126 hinder conservation measures efficiency and despite increasing literature on the need to develop new
127 dynamical tools (Tittensor et al., 2019; Crespo et al., 2020), conservation policies seem to remain only
128 focused on managing direct anthropogenic pressures as long as policy-makers have no concrete and
129 operational courses of action to take into account the impacts of climate change (Wilson et al., 2020).

130


1
2 131 **Current lack of relevant knowledge on climate change for policy-makers.** Levers for potential
3
4 132 action remain today unclear at the regional level and require a better understanding of climate change
5
6 133 related hazards at relevant spatial and temporal scales for conservation (Wiens and Bachelet, 2010 ;
7
8 134 Carr et al., 2011; Butt et al., 2016; Jones et al., 2016). Today, a gap remains in a systematic description
9
10
11 135 of local observed and future changes over the entire Southern Indian Ocean.

12
13
14 136 In this study, we aim at addressing this shortcoming, by presenting a regional analysis of past and future
15
16 137 ocean warming based on historical observations and climate projections under different scenarios. Here,
17
18 138 we use a combination of decadal observations and CMIP6 simulations over the entire Southern Indian
19
20 139 Ocean to estimate current trends and analyse projections to help identifying areas that are most likely
21
22 140 to undergo significant temperature changes under two future emission scenarios : Shared
23
24 141 Socioeconomic Pathways (SSPs) 1-2.6 (high mitigation scenario) and 2-4.5 (modest mitigation
25
26 142 scenario; projected to lead to a global warming in 2100 similar to modelled pathways assuming
27
28 143 continuation of policies implemented by the end of 2020; IPCC, 2022). We estimate the average
29
30 144 temperature trends and extreme events, both listed as “climatic impact-drivers” of marine ecosystems
31
32 145 according to the IPCC Assessment Report 6 (Chen et al., 2021; Cooley et al., 2022). We document
33
34 146 average temperature trends in terms of warming patterns (°C/decade) as well as climate velocities
35
36 147 (km/decade). This analysis is performed for the ocean surface (0-200 m) and for the mesopelagic (200-
37
38 148 1000 m) regions. Extreme events are characterized in terms of marine heatwaves (MHW, Hobday et
39
40
41 149 al., 2016; Frölicher et al., 2018). Su and co-authors (2021) characterized MHW but focusing only on
42
43 150 the northern part of the Kerguelen Plateau. Our study extends the MHW analysis to the Southern Indian
44
45 151 Ocean and uses CMIP6 models to project the evolution of MHW throughout the 21st century.

46
47
48
49
50
51 152
52
53
54 153 Figure 1: Ocean circulation in the Southern Indian Ocean. The mean geostrophic velocity between 1993
55
56 154 and 2020 using AVISO product (<https://doi.org/10.48670/moi-00148>) is shown (grey arrows). Some
57
58 155 specific currents are indicated in blue and the Antarctic Circumpolar Current (ACC) fronts, the Northern
59
60
61
62
63
64
65

156 Boundary (NB), Subantarctic Front (SAF), Polar Front (PF), Southern ACC front (SACCF) and the
1 Southern Boundary (SB) are shown in brown (constructed from mean dynamic topography, source:
2 157
3
4 158 Park and Durand, 2019, Park et al., 2019).

5
6
7
8 159



9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39 160

42 161 2 Material and Methods

45 162 2.1 Data

47 163 2.1.1 Observations

48
49
50
51 164

52
53 165 The Operational Sea Surface Temperature and Ice Analysis (OSTIA) system run by the UK's Met
54
55 166 Office (Good et al., 2020) provides daily sea surface temperature free of diurnal variability at a 0.05deg.
56
57 167 x 0.05deg. horizontal grid resolution between 1982 and 2019 (available at

168 <https://marine.copernicus.eu/>). This product combines satellite measurements from both infrared and
169 microwave radiometers with in-situ measures from ships, and drifting and moored buoys.

170

171 The National Oceanic and Atmospheric Administration Daily Optimum Interpolation Sea Surface
172 Temperature (NOAA OI SST V2 High Resolution) dataset, provided by the NOAA PSL (available
173 from their website at <https://psl.noaa.gov>), combines sea surface temperature observations (SST at 0.2
174 m) from different platforms (satellites, ships, buoys and Argo floats) with a 0.25deg x 0.25deg
175 horizontal grid resolution between 1982 and 2019 (Reynolds et al. 2007; Banzon et al. 2016). Contrary
176 to OSTIA dataset, OISST does not include satellite measurements from microwave radiometers as SST
177 inputs (Yang et al., 2021).

178

179 2.1.2 Global climate models – CMIP6

180

181 Models from the Coupled Model Intercomparison Project 6 (CMIP6; Eyring et al., 2016) are used in
182 this study (Table 1). These are coupled ocean-atmosphere models developed by 49 different climate
183 modelling groups¹ that have been used to carry out historical and projection simulations notably to
184 investigate how the Earth system responds to forcing (Eyring et al., 2016).

¹ https://wcrp-cmip.github.io/CMIP6_CVs/docs/CMIP6_institution_id.html

1
2
3 186 Table 1: CMIP6 models used, using yearly data over multiple depths and daily data at the surface and
4
5 187 scenarios associated. CMIP6 models outputs are available at : [https://esgf-](https://esgf-node.llnl.gov/projects/cmip6/)
6
7 188 [node.llnl.gov/projects/cmip6/](https://esgf-node.llnl.gov/projects/cmip6/).

	Yearly data – multiple depth levels	Daily data - surface	Ocean horizontal resolution (before interpolation)	References
ACCESS-CM2	Historical,ssp1 26,ssp245	Historical, ssp126, ssp245	1°	Bi et al., 2020
ACCESS-ESM1-5	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Ziehn et al., 2020
BCC-CSM2-MR	Historical, ssp126,ssp245	Historical, ssp126, ssp245	1°	Wu et al., 2019
CAMS-CSM1-0	Historical, ssp126,ssp245	/	1°	Rong et al., 2019
CanESM5	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Swart et al., 2019; Christian et al., 2021
CanESM5-CanOE	Historical, ssp126, ssp245	/	1°	Swart et al., 2019; Christian et al., 2021
CESM2	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Danabasoglu et al., 2020

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

CESM2-WACCM	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Danabasoglu et al., 2020
CMCC-CM2-SR5	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Cherchi et al., 2019
CMCC-ESM2	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Lovato et al., 2022
CNRM-CM6-1	Historical, ssp126, ssp245	Historical,ssp12 6,ssp245	1°	Voldoire et al., 2019
CNRM-CM6-1-HR	/	Historical,ssp12 6,ssp245	0.25°	Voldoire et al., 2019
CNRM-ESM2-1	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Séférian et al., 2019
GFDL-CM4	Historical, ssp245	Historical, ssp245	0.25°	Held et al., 2019; Dunne et al., 2020
GFDL-ESM4	Historical, ssp126, ssp245	/	0.5°	Dunne et al., 2020
HadGEM3-GC31-LL	Historical, ssp126,ssp245	/	1°	Kuhlbrodt et al., 2018; Andrews et al., 2020
EC-Earth3	/	Historical, ssp126, ssp245	1°	Döscher et al., 2021

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

EC-Earth3-CC	Historical, ssp245	/	1°	Döscher et al., 2021
EC-Earth3-Veg	Historical, ssp126, ssp245	/	1°	Döscher et al., 2021
IPSL-CM6A-LR	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Boucher et al., 2020
MIROC-6	/	Historical, ssp126, ssp245	1°	Tatebe et al., 2019
MIROC-ES2L	Historical, ssp126, ssp245	/	1°	Hajima et al., 2020
MPI-ESM1-2-HR	Historical, ssp126, ssp245	Historical, ssp126, ssp245	0.4°	Müller et al., 2018; Mauritsen et al., 2019
MPI-ESM1-2-LR	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1.5°	Mauritsen et al., 2019
MRI-ESM2-0	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°x0.5°	Yukimoto et al., 2019
NESM3	Historical, ssp126, ssp245	Historical, ssp126, ssp245	1°	Cao et al., 2021
UKESM1-0-LL	Historical, ssp126, ssp245	/	1°	Sellar et al., 2019

189

190

191 Historical and projection simulations from the ScenarioMIP experiments are used. Historical
192 simulations cover the period 1850-2014 and use historical forcing, mostly based on observations, that
193 includes among others greenhouse gas, aerosol concentrations and solar forcing (Eyring et al., 2016).
194 Projection simulations from the ScenarioMIP experiments cover 2015-2100 and over multiple emission
195 trajectories, described through so-called Shared Socioeconomic Pathways (SSPs). More specifically,
196 SSPs describe development pathways according to different economic and political strategies (O'Neill
197 et al., 2017), and subsequent forcing levels similar to the ones used in CMIP5, that are the
198 Representative Concentration Pathways (RCPs). However, SSPs represent an improvement in the sense
199 that those scenarios include updated data on recent emission trends (O'Neill et al., 2016) and allow to
200 explore the implication of different climate change mitigation policies (Riahi et al., 2017). In this study,
201 for all analyses, two projection scenarios are considered: the SSPs 1-2.6 and 2-4.5. The SSP1-2.6
202 describes a world that shifts to a more sustainable path leading to an estimated warming of 1.8 °C by
203 the end of the century (as compared to pre-industrial) whereas SSP2-4.5 represents a path in which
204 socio-economic trends do not shift significantly from the historical patterns, with an estimated warming
205 of 2.7 °C by the end of the century. The focus on those two scenarios allows the comparison between
206 two possible futures whose occurrence will depend on countries' commitments to reduce their net
207 greenhouse gas emissions (Hausfather & Peters, 2020).

208 Depending on the analysis, the variable considered is either *thetao* (sea temperature across different
209 depth levels) or *tos* (sea surface temperature) as only *tos* was available daily. Each model output is
210 regridded to the same regular 1°-1° horizontal grid using distance weighted average remapping (using
211 climate data operators « cdo » remapdis) as in Kwiatkowski et al., 2020. Each model output is also
212 regridded vertically following the World Ocean Atlas standard discretization (33 vertical intervals from
213 the surface (0 m) to the abyssal seafloor (5500 m)).

214 There are several ways to combine the results from multiple models. The traditional approach used by
215 the IPCC, which is the main method used here, is to use a model ensemble mean, all models being
216 weighted equally, and then to study the projections for fixed future periods. For changes in temperature
217 and projected MHW characteristics, future periods of 20 years are considered: the near term (2021-

218 2040), the mid term (2041-2060) and the long term (2081-2100) as in the IPCC Assessment Report 6
1
2 219 (Chen et al., 2021). An alternative method is also considered to further support the projected trends
3
4 220 obtained with the traditional method: the time-shift approach (Herger et al., 2015; Chen et al. 2021; Lee
5
6 221 et al., 2021). Instead of specifying a fixed time period, this approach specifies a fixed global warming
7
8 222 value compared to pre-industrial, typically 1.5°C, 2°C and 3°C (hereafter referred to as « global
9
10 223 warming levels » or GWL 1.5°C, 2°C and 3°C, respectively). These GWLs are in general reached at
11
12 224 different time periods depending on the models and on the scenarios. This method allows to account for
13
14 225 the different climate sensitivities of the models and facilitates the comparison between global warming
15
16 226 and regional spatial patterns. For each model and under each SSP, warming levels are defined as 20-
17
18 227 year running means of globally averaged atmospheric surface temperature anomaly compared to the
19
20 228 pre-industrial period (1850-1900). Then the change in a given metric (temperature, MHW indicator) is
21
22 229 estimated using a climatology of 20 years centred on the first year for which the warming level exceeded
23
24 230 a given threshold (e.g., 1.5°C), relative to the 1850-1900 reference value. These results are then
25
26 231 averaged over the model ensemble and over the SSPs considered, all simulations being weighted
27
28 232 equally. This method is applied to estimate the spatial patterns in warming and in MHW intensity for
29
30 233 GWL 1.5°C, GWL 2°C and GWL 3°C. To be able to investigate GWL 3°C, given that not all models
31
32 234 for SSP1-2.6 and SSP2-4.5 may reach this threshold, outputs from SSP5-8.5 are also used
33
34 235 (Supplementary Figure S1).
35
36
37
38
39
40

41 236
42
43
44 237 Models are also confronted with observations over the historical period for evaluation purposes. More
45
46 238 precisely, we evaluate how well the spatial patterns and mean trends are reproduced by the models and
47
48 239 compared to observations. This comparison is used as an indicator of what processes affecting trends
49
50 240 may be or may not be reproduced by the model ensemble. Such information contributes to the
51
52 241 interpretation of projections and uncertainties. Models and observations warming rates and MHW mean
53
54 242 intensity are compared respectively over 1982-2019 (to account for natural interannual variability) and
55
56 243 1984-2014 (MHW analysis requiring 30 years of data). CMIP6 historical outputs are thus completed
57
58 244 with SSP2-4.5 outputs until 2019, since SSP2-4.5 is most representative of today's emission pathway
59
60
61
62
63
64
65

245 (Fricko et al., 2017; Hausfather and Peters, 2020). The temperature trends significance is determined
1
2 246 using the coefficient of determination (R^2) of the linear regression as well as a signal-to-noise ratio
3
4 247 (SNR) comparing the trend to the interannual variability of the anomaly between the expected (from
5
6 248 the linear regression) and observed temperature, as defined in Auger et al., 2021.
7
8

9
10 249 To quantify the impact of natural variability on the projected trends, we also exploit additional
11
12 250 simulations using the IPSL-CM6A-LR model, which exhibits an important centennial climate
13
14 251 variability (Bonnet et al., 2021), through the analysis of the temperature trends of 11 members (r1i1p1fl,
15
16 252 r2i1p1fl, r3i1p1fl, r4i1p1fl, r5i1p1fl, r6i1p1fl, r10i1p1fl, r11i1p1fl, r14i1p1fl, r22i1p1fl,
17
18 253 r25i1p1fl). These members were all initialized in 1850, but with distinct initial conditions, mostly
19
20 254 covering different states regarding the initial trend of the Atlantic Meridional Overturning Circulation
21
22 255 (Bonnet et al., 2021). First, the temperature trends over the historical period (1975-2015) of the different
23
24 256 members are compared. Second, projected changes in surface temperature for the near, mid and long
25
26 257 term periods are compared. Finally, Pearson correlation coefficients between spatial patterns of changes
27
28 258 in surface temperature for each member simulation relative to the r1i1p1fl simulation over each year
29
30 259 between 2015 and 2100 in SSP2-4.5 are estimated. The aim is to determine whether the initial
31
32 260 conditions of the simulations and natural variability still play an important role in determining the
33
34 261 regional characteristics of the projected trends compared to the radiative forcing.
35
36
37
38
39

40 262

41 42 43 263 **2.2 Climate velocity**

44
45 264

46
47
48 265 For climate velocity we use the classical definition as the ratio between the warming rate and the local
49
50 266 spatial gradient of a variable as in Loarie et al., 2009 and Brito-Morales et al., 2018. The intensity of
51
52 267 horizontal climate velocity v for temperature (T) is thus here defined as
53
54

55
56 268
$$v = \frac{\frac{\partial T}{\partial t}}{||\nabla T||}$$

269

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

270 By convention, v is positive in case of a warming ($\frac{\partial T}{\partial t} > 0$) and negative in case of a cooling.

271 The climate velocity vector \vec{v} is then defined as

$$\vec{v} = \frac{-\partial T}{\partial t} \left[\frac{\frac{\partial T}{\partial x}}{\|\nabla T\|^2} \cdot \vec{u}_x + \frac{\frac{\partial T}{\partial y}}{\|\nabla T\|^2} \cdot \vec{u}_y \right].$$

273 By convention, \vec{u}_x (longitudinal unit vector) is positive eastward and \vec{u}_y (latitudinal unit vector) is
274 positive northward. \vec{v} points to the direction to follow to remain at the same temperature.

275

276 For CMIP6 outputs, climate velocity is estimated over different historical periods of 50 years (1850-
277 1900, 1950-2000, 1955-2005 and 1965-2015) to check the sensitivity of the pre-industrial or historical
278 climate velocity that could be used as a baseline for comparison with the projected climate velocities.
279 Then climate velocity is estimated over 2015-2065 and 2050-2100 for SSP1-2.6 and SSP2-4.5,
280 respectively. Climate velocity is estimated for different depth zones: surface (0-200 m) and mesopelagic
281 (200-1000 m). The average temperature weighted by the thickness of each standard depth layer is used
282 to calculate a horizontal climate velocity. The climate velocities obtained from each model are then
283 averaged to get the mean-ensemble results.

284

285 **2.3 Marine heatwaves**

286

287 The definition of a marine heatwave (MHW) as a "discrete prolonged anomalously warm water event"
288 was introduced in Hobday et al., 2016 to allow for comparison with other studies increasingly adopting
289 this standardized definition. In this work, a MHW is detected when the temperature is above a given
290 threshold for at least 5 days. This threshold is defined as the 90th (for model evaluation) or the 99th
291 percentile (for projections, to focus on the most intense events) of the data distribution. The climatology

292 and threshold are estimated for each grid point of the region of interest and for each day of the year
293 using an 11-day window. The threshold is also smoothed by applying a 30-day moving average. The
294 climatology is estimated over at least 30 years of data, in order to smooth out climate mode of variability
295 (e.g., ENSO, AMO; Scannell et al., 2016).

296 MHW can be quantified by different metrics. The definition of MHW intensity varies and can be
297 determined relative to the seasonal climatology or to a chosen threshold (Hobday et al., 2016). Here we
298 define MHW intensity relative to the threshold (90th percentile for model evaluation, 99th percentile
299 for projections) and the mean intensity is weighted by the number of days affected. In addition, the
300 mean number of days affected by MHW per year, also called mean annual MHW days, has been found
301 strongly correlated to some observed ecological performance in the marine environment (e.g., seagrass
302 density, Smale et al., 2019). The two metrics, mean MHW intensity and mean MHW annual days, are
303 therefore used here to characterize MHW as potential climatic impact-drivers of marine ecosystems.

304

305 **3 Results**

306 **3.1. Historical warming trends and marine heatwaves**

307 *3.1.1. Warming trends*

308

309 The mean warming rate observed over the Southern Indian Ocean (20°-120°E, 70°-30°S) between 1982
310 and 2019 is 0.03°C/decade ($\pm 0.08^\circ\text{C}/\text{decade}$, spatial standard deviation), as estimated by a linear
311 regression of observation-based sea surface temperature from the OSTIA product, and 0.07°C/decade
312 ($\pm 0.09^\circ\text{C}/\text{decade}$) using OISST dataset (See Section 2.1.1). Such regional averaged temperature trend
313 hides important east/west contrasts (Figure 2, also for OISST as shown in Supplementary Figure S2)
314 with warming trends north of the area and east of the Kerguelen Plateau between 0.1 and 0.3 °C/decade
315 that are found significant (R^2 between 0.4 and 0.5, $\text{SNR} > 1$). The cooling pattern west of the Kerguelen

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

316 Plateau is not found significant, but has also been obtained on different observational datasets (Yang et
317 al., 2021), suggesting high temporal variability in this area.


318 In comparison, the 24-single-member-CMIP6-models produces an ensemble mean warming rate from
319 1982 to 2019 close to the one obtained with OISST dataset (0.085 ± 0.053 °C/decade, spatial standard
320 deviation). A majority of CMIP6 models (17 over 24) simulate higher mean warming rates than the
321 observed one with OISST dataset but none shows an averaged cooling in the region (Figure 2,
322 Supplementary Figure S3). The simulated spatial patterns of temperature trends are quite different
323 between different model simulations, but also between different ensemble members of the same model
324 (IPSL-CM6A-LR, Supplementary Figure S5A). The initial conditions of the simulations as well as
325 decadal and/or multidecadal variability might thus play an important role in the regionalisation of
326 temperature trends over the historical period.

327 Historical warming rates from the multimodel ensemble are more homogeneous than from any
328 individual models or observations, despite greater warming rates being found between 40° and 50°S
329 north and west of Prince Edward, Marion and Crozet islands (Figure 2). The multimodel ensemble mean
330 underestimates warming trends north of the ACC and east of the Kerguelen Plateau. It shows a general
331 warming throughout the area. A lower signal and/or lower agreement between models is found south
332 of the Kerguelen Plateau near the Antarctic shelves.

333

334 Figure 2: Warming rates between 1982 and 2019 in the Southern Indian Ocean, using linear regression
335 on surface temperature, using OSTIA observations (A) and a 24-single-member-CMIP6-model
336 ensemble (B). Hatching indicates areas where the intermodel standard deviation of the warming rate is
337 greater than the mean value, suggesting a low change or a low robustness of the output. For each model
338 and for the 24-model ensemble, the mean warming rate over the area is also calculated and compared
339 to the mean warming rates estimated from the OSTIA and OISST datasets (C). Bars colored in red
340 indicate a higher warming rate than the one derived from OISST (gray bar) and in blue if it is a lower
341 rate.

342



343

344 *3.1.2. Climate velocity*

345

346 The mean climate velocity over the Southern Indian Ocean is 4.9 km/decade (± 27 km/decade, spatial
 347 standard deviation) using OSTIA dataset and 10.64 km/decade (± 34 km/decade) using OISST dataset
 348 between 1982 and 2019 (see Sections 2.1.1 and 2.2). An east/west contrast is observed similar to what
 349 is found in section 3.1.1. East of the Southern Indian Ocean (70° - 120° E, 70° - 30° S), the mean climate
 350 velocity is around 12 km/decade using OSTIA dataset and around 18 km/decade using OISST dataset.

351

352 *3.1.3. MHW intensity*

353


354 The regional mean MHW intensity observed over the Southern Indian Ocean between 1984 and 2014
 355 is 0.35°C above the 90th percentile of the local temperature distribution using the OSTIA dataset
 356 (Figure 3). Detection of MHW from the OSTIA dataset highlights a pattern of greater MHW intensity

357 north of the Kerguelen Plateau, in the region of influence of the ACC (Figure 3, pattern also identified
1
2 358 in Su et al., 2021). The 19-model ensemble, reproduced the spatial pattern with greater mean MHW
3
4 359 intensity north of the Plateau, notably over the Crozet Plateau, and the less intense MHW west of the
5
6 360 Kerguelen Plateau (Figure 3, Supplementary Figure S4).
7
8

9 361 However, the multimodel mean underestimates the intensity of MHW north of the ACC compared to
10
11 362 observations. The mean MHW intensity between 40°S and 30°S obtained from the multimodel mean is
12
13
14 363 26% lower than the one obtained from observations.
15
16

17 364

18
19
20 365 Figure 3: Mean Marine Heatwaves (MHW) intensity as defined by the anomaly above the threshold
21
22 366 (90th percentile) between 1984 and 2014 in the Southern Indian Ocean using OSTIA observations (A)
23
24 367 and a 19-single-member-CMIP6-model ensemble (B). Hatching indicates areas where the intermodel
25
26 368 standard deviation of the MHW intensity is greater than the mean value, suggesting a low robustness
27
28
29 369 of the output.
30
31
32
33
34
35
36



51
52 370

53
54
55 371

56 372 **3.2 Projected warming trends and marine heatwaves**

57 373 *3.2.1 The impact of natural variability on the projected temperature trends*

374

1
2 375 To quantify the impact of natural variability on the projected temperature trends, we first focus our
3
4 376 analysis on one model (IPSL-CM6A-LR), for which we use an ensemble of simulations and investigate
5
6 377 the similarity between members of the simulated patterns for historical and projected warming (see
7
8
9 378 Section 2.1.2).

10
11
12 379 The mean warming rate over the Southern Indian Ocean varies from 0.031°C/decade to 0.11°C/decade
13
14 380 over 1975-2015 between the 11 members of IPSL-CM6A-LR (Supplementary Figure S5). Important
15
16 381 differences regarding the spatial patterns are found, with members producing a general warming
17
18 382 throughout the region (e.g. r6i1p1f1), others showing a cooling north of the area (e.g. r3i1p1f1,
19
20 383 r4i1p1f1, r22i1p1f1) or a cooling east of the Kerguelen Plateau (e.g. r2i1p1f1, r14i1p1f1, r25i1p1f1).

21
22
23 384 Across the ensemble members, the mean change in surface temperature over the Southern Indian Ocean
24
25 385 in SSP2-4.5 ranges from 0.15°C to 0.44°C in the near term (2021-2040), from 0.43 to 0.66°C in the mid
26
27 386 term (2041-2060) and from 0.83°C to 1.06°C in the long term (2081-2100), as compared to the historical
28
29 387 period (1995-2014, Supplementary Figure 5B). The timeseries of Pearson correlation coefficient
30
31 388 relative to r1i1p1f1 member pinpoints the increased similarity between the members' patterns, notably
32
33 389 from 2050 onward (correlation coefficient mainly over 0.5, Supplementary Figure 5C). It could be that
34
35 390 the remaining differences between the members' projections is to be attributed to a non-deterministic
36
37 391 component in ocean response, notably intrinsic ocean variability (Dijkstra, 2016).

38
39
40
41
42 392 This suggests that over the historical period the natural variability is strong and drives a large spread of
43
44 393 temperature trends between ensemble members. With increasing radiative forcing, this spread
45
46 394 decreases, suggesting that the projected long term spatial patterns are not dependent on natural
47
48 395 variability.

49
50
51 396


52 53 397 *3.2.2 Projected warming*

54
55 398

56
57
58 399 Over the Southern Indian Ocean and for SSP2-4.5, the projected warming is 0.29°C ($\pm 0.06^\circ\text{C}$,
59
60 400 intermodel variability) in the near term (2021-2040), 0.58°C ($\pm 0.09^\circ\text{C}$) in the mid term (2041-2060)

1 401 and 1.05°C ($\pm 0.15^\circ\text{C}$) in the long term (2081-2100), compared to the historical period (1995-2014).
2 402 While the mean surface temperature change is similar in both scenarios in the near term, it is 1.2 and
3
4 403 1.7 times higher for SSP2-4.5 than for SSP1-2.6, in the mid and long term, respectively (Figure 7).
5
6
7 404 Using the 24-model ensemble, a larger warming is simulated between 40°S and 50°S and up to 55°S,
8
9 405 thus in regions including the Indian Ocean subantarctic Exclusive Economic Zones (EEZs, see Figure
10
11 406 4). This latitudinal band of large warming is located in the ACC and expands southward with time and
12
13
14 407 in SSP2-4.5.
15
16
17 408 Between 35° and 40°S, warming is lower than in the latitudinal band encompassing the subantarctic
18
19 409 island's EEZs (warming of 0.3-0.4°C in SSP2-4.5 mid term). South of 55°S, surface temperature
20
21 410 changes remain below 1°C even in the long term in SSP2-4.5. The region south of the ACC has been
22
23 411 shown to experience lower warming due to the local circulation and stratification change (Armour et
24
25 412 al., 2016; Haumann et al., 2020).
26
27
28 413
29
30
31 414 Figure 4: Surface temperature change from the historical period (1995-2014) for the mid-term (A and
32
33 415 B) and long term (C and D) projected periods under SSP1-2.6 (A and C) and SSP2-4.5 (B and D) using
34
35 416 CMIP6 multimodel mean. Contour lines of similar temperature change are indicated in cyan. Hatched
36
37 417 areas indicate where the intermodel standard deviation of the temperature change is higher than the
38
39 418 mean value, suggesting a less robust output.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



3.2.3 Projected climate velocity

For the Southern Indian and in SSP2-4.5, the average climate velocities over 2015-2065 is 38.53 southward (± 5.8 , intermodel variability) and 47.43 (± 8.1) km/decade at the surface and in the mesopelagic, respectively. When averaged over 2050-2100, it is 31.62 (± 5.6) and 57.46 (± 9.8) km/decade, at the surface and in the mesopelagic, respectively. At the end of the century, surface (mesopelagic) climate velocities are expected to be 5.3 times (1.8 times) faster in SSP2-4.5 as compared to SSP1-2.6 (Figure 7). Mesopelagic climate velocities under both scenarios are also about 5 times faster for the period 2015-2065 compared to the historical period 1955-2005.

Negative surface climate velocities (i.e. northward) over 2050-2100, are found around 35°S in SSP1-2.6 (Figure 5), meaning that this area is expected to cool towards the end of the 21st century. This

1 431 cooling is only found at the surface and not at depth under SSP1-2.6, and is not projected under SSP2-
2 432 4.5.


3
4
5 433 Important differences are found between surface and mesopelagic climate velocities. Mesopelagic
6
7 434 conditions might undergo faster climate velocities than at the surface and that, already over 2015-2050
8
9 435 (Figure 7). For the two projected periods and for both scenarios, mesopelagic climate velocities
10
11 436 averaged over the Southern Indian Ocean are between one and four times faster than surface climate
12
13 437 velocities. In contrast, for different periods of 50 years between 1850 and 2015, surface climate
14
15 438 velocities are found between 1.5 and 3.1 faster than mesopelagic climate velocities. Those differences
16
17 439 between climate velocities in the mesopelagic and at the surface are accentuated in some areas. Those
18
19 440 differences can reach up to 160 km/decade southwest of Australia, 120 km/decade west of Kerguelen
20
21 441 Plateau or vary between 40 and 70 km/decade north of Crozet Plateau in SSP2-4.5. Faster climate
22
23 442 velocities in the mesopelagic compared to the surface for a same location (Figure 5) can be due to the
24
25 443 weaker spatial temperature gradients at depth as it is for instance the case west of the Australian coast..
26
27
28
29

30 444

31
32
33 445 Figure 5: Surface (0-200m, A and B) and mesopelagic (200-1000m, C and D) climate velocities, defined
34
35 446 as the ratio between the warming rate and the temperature spatial gradient between 2050 and 2100 using
36
37 447 multimodel means under SSP1-2.6 (A and C) and SSP2-4.5 (B and D). It can be interpreted as the
38
39 448 velocity an individual should adopt to remain at the same temperature. By convention, positive climate
40
41 449 velocities correspond to warming trends and negative ones to cooling trends. Black arrows indicate the
42
43 450 direction of the climate velocity.
44
45
46

47 451
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



452

3.2.4 Projected marine heatwaves

453


454

455 In the Southern Indian Ocean, the projected mean MHW intensity is 0.28°C (above 99th percentile
 456 threshold) in the near term for both scenarios and 0.39°C or 0.63°C in the long term, respectively in
 457 SSP1-2.6 or SSP2-4.5. The areas most affected by this intensification in the long term are located
 458 between 40°S and 55°S and in particular over Prince Edward, Marion, Crozet, Kerguelen and Heard
 459 and McDonald Islands (Figure 6). The spatial pattern of mean MHW intensity in SSP2-4.5 for the near
 460 term is similar to the historical pattern (Figure 3). North of the ACC, mean MHW intensity, for this
 461 scenario and time horizon, varies between 0.3 and 0.4°C. In the long term under SSP2-4.5, a maximum
 462 MHW intensity of 0.97°C is found over Crozet Plateau and the zonal maximum MHW intensity is found
 463 to be 0.81°C at 48°S. The pattern obtained is similar to the pattern of surface temperature change (Figure

464 4), suggesting that the detected MHW might be caused by a mean shift of the temperature distribution
1
2 465 rather than a change in the spread of the temperature distribution (as in Oliver et al., 2019a).
3
4
5 466 Additional analyses, investigating the daily temperature distribution shift in SSP2-4.5 relative to the
6
7 467 historical period, show that the relative shift in the mean of the temperature distribution south of 50°S
8
9 468 is much larger than the changes in the spread of the distribution, whereas north of 50°S similar orders
10
11 469 of magnitude of relative change are found in terms of increased mean temperatures and increased
12
13 470 distributions spread (Supplementary Figure S6). In particular, the area between 35° and 40°S shows a
14
15 471 larger relative change in the spread of the distribution than in the mean of the distribution, with increases
16
17 472 of respectively 9.98% and 7.60% in the long term compared to the historical period.
18
19
20
21 473 In the Southern Indian Ocean, the projected mean annual MHW days is 46.1 days/year in the near term
22
23 474 and 119.3 days/year in the long term in SSP1-2.6. These values increase to 42.8 days/year and 206
24
25 475 days/year respectively in SSP2-4.5 (Figure 6). Annual MHW days under both scenarios are similar in
26
27 476 the near term but it is 23% higher in the mid term and 73% higher in the long term in SSP2-4.5 compared
28
29 477 to SSP1-2.6 (Figure 7). Similar as for MHW intensity, the most impacted area is generally the latitudinal
30
31 478 zone between 40°S and 55°S in the ACC region. In SSP2-4.5, in the near term, the zonal number of
32
33 479 annual MHW days is maximum between 40°S and 55°S varying between 44.5 and 62.9 days, the
34
35 480 maximum being reached at 50°S. In the long term, the zonal number of annual MHW days is maximum
36
37 481 between 40°S and 57°S varying between 194 and 290 days, the maximum being reached at 51°S. A
38
39 482 permanent state of MHW is almost reached over the Kerguelen Plateau (312 days west of the Plateau).
40
41 483 When considering the relationship between changes in surface temperatures and MHW annual days for
42
43 484 the three projected time periods, it appears that over the whole area a permanent state of MHW could
44
45 485 be reached for a 2°C regional mean surface temperature change relative to the historical period
46
47 486 (Supplementary Figure S7).
48
49
50
51
52
53 487
54
55
56 488 Figure 6: Projected mean MHW intensity (as defined by the anomaly above the threshold of 99th
57
58 489 percentile relatively, A and C) and projected mean number of days per year affected by MHW (B and
59
60
61
62
63
64
65

490 D) under SSP2-4.5 using a 19-single-member-CMIP6-model ensemble for both 2021-2040 (A and B)
 1
 2 491 and 2081-2100 (C and D). MHW were detected relative to a historical seasonally-varying climatology
 3
 4 492 (1984-2014). Contour lines of similar metric are indicated in cyan. Hatching indicates areas where the
 5
 6 493 intermodel standard deviation of the metric is greater than the mean value, suggesting a low robustness
 7
 8
 9 494 of the output.


10
 11
 12 495



13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45 496

497 Figure 7: Synthesis of the different climate impact drivers metrics over the Southern Indian Ocean (20°-
 498 120°E 70°-30°S) : change in surface temperature (A, relative to 1995-2014); the annual number of day
 499 affected by MHW as defined using the 99th percentile threshold (C); average surface (B) and
 500 mesopelagic (D) climate velocities. The error bars correspond to the intermodel standard deviation for
 501 each metric.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



3.3 Patterns of warming-related climatic impact-drivers for GWL 1.5°C, 2°C and 3°C

Patterns of warming-related climatic impact-drivers are also investigated using a timeshift approach (see Section 2.1.2). Not only is this method used to confirm the robustness of the spatial patterns obtained through the traditional approach, but this method also contributes to describing future changes in accordance with targets defined in the Paris Agreement², regardless of the mitigation pathway (Chen et al., 2021).

² https://unfccc.int/sites/default/files/english_paris_agreement.pdf


1 512 The mean sea surface temperature change over the Southern Indian Ocean is 0.6°C ($\pm 0.2^{\circ}\text{C}$, intermodel
2 513 variability), 0.7°C ($\pm 0.2^{\circ}\text{C}$) and 1.1°C ($\pm 0.3^{\circ}\text{C}$) greater than in 1850-1900 for global warming levels
3
4 514 (GWLs, see Section 2.1.2) of 1.5°C , 2°C and 3°C , respectively, as computed from the CMIP6 model
5
6 515 ensemble (Figure 8; we remind that GWLs are defined in respect to pre-industrial temperatures and not
7
8 516 in respect to the present). In the three cases, the patterns are similar, only the amplitude increases with
9
10 517 increasing global warming. Surface temperature changes are warmer than 0.8°C between 40° and 50°S
11
12 518 for GWLs 1.5°C and 2°C and warmer than 1.2°C for GWL 3°C at the same latitudes. Surface
13
14 519 temperature changes stay below 0.4°C and 0.6°C near the Antarctic, respectively for GWLs 2°C and
15
16 520 3°C .

17
18
19
20 521 The mean MHW intensity is 0.37°C ($\pm 0.037^{\circ}\text{C}$, intermodel variability), 0.43°C ($\pm 0.050^{\circ}\text{C}$) and
21
22 522 0.43°C ($\pm 0.050^{\circ}\text{C}$), for GWLs 1.5°C , 2°C and 3°C , respectively (Figure 8). For GWL 1.5°C , the most
23
24 523 intense MHW are found in the subtropical region and north of the ACC (40° - 45°S , 20° - 90°E) varying
25
26 524 between 0.5°C and 0.6°C . For GWLs 2°C and 3°C , mean MHW intensity between 40° and 50°S can be
27
28 525 warmer than respectively 0.5°C and 0.8°C . For both warming trends and MHW characteristics, the
29
30 526 patterns obtained here using GWLs are similar to the ones obtained using projections on fixed future
31
32 527 periods (Section 3.2).

33
34 528
35
36
37
38
39
40 529 Figure 8: Change in surface temperature for global warming levels (GWLs) 1.5°C (A), 2°C (B) and 3°C
41
42 530 (C) and mean MHW intensity for GWLs 1.5°C (D), 2°C (E) and 3°C (F) warming world relative to the
43
44 531 pre-industrial period (1850-1900) using a time-shift approach and CMIP6 mean ensembles. The number
45
46 532 of simulations used (using SSP1-2.6, SSP2-4.5 and SSP5-8.5) in each case is indicated in
47
48 533 parentheses. Contour lines of similar temperature change or MHW intensity are indicated in cyan.
49
50 534 Stapling indicates areas where the intermodel standard deviation is higher than the mean value,
51
52 535 suggesting a less robust output.

53
54
55
56 536
57
58
59
60
61
62
63
64
65

Change in surface temperature



537

538 4 Discussion

539 4.1. Regional characteristics of temperature-related climatic impact-drivers

540

541 This study presents a regional analysis of current and projected trends of temperature-related climatic
542 impact-drivers in the Southern Indian Ocean: temperature change, temperature-driven climate
543 velocities and marine heatwaves (MHW). Our main findings can be summarized as follows.

544

545 **Warming patterns are spatially heterogeneous in the region.** Global ocean surface warming rates
546 have been estimated at around 0.08°C/decade between 1880 and 2020 (NOAA, 2022) and the IPCC
547 Assessment Report 6 has indicated a 0.60°C increase in global SST from 1980 to 2020 (Fox-Kemper et
548 al., 2021). In the Southern Indian Ocean, a greater warming north of the Antarctic Circumpolar Current

549 (ACC) of around 0.11°C/decade has been observed between 1950 and 2012, while cooling patterns
1
2 550 have been observed near the Antarctic (Armour et al., 2016; Sallée et al., 2018). The mean warming
3
4 551 trend here in the Southern Indian Ocean between 1982 and 2019 is lower, between 0.03°C/decade and
5
6 552 0.07°C/decade (Figure 2, Supplementary Figure S2). Observed warming trends are significant in the
7
8
9 553 northern part of the region and east of Kerguelen Plateau and could vary between 0.1 and 0.3°C/decade.

10
11
12 554

15 555 **Climate velocities are faster over Kerguelen Plateau as compared to global ocean averages.** The
16
17 556 global ocean median climate velocity has been estimated around 13 km/decade over 1900-2010 (Gupta
18
19 557 et al., 2015) or 12 km/decade over 1955-2005 (Brito-Morales et al., 2020) at the surface and around 6.3
20
21 558 km/decade in the mesopelagic layer over 1955-2005 (200 - 1000 m). The mean surface climate velocity
22
23
24 559 over the Southern Indian Ocean is found between 4.9 and 10.6 km/decade between 1982 and 2019 using
25
26 560 OSTIA and OISST datasets. In particular, surface climate velocities are particularly fast over Heard and
27
28
29 561 McDonalds Islands EEZ (20.3 km/decade, using OSTIA dataset ; 44.0 km/decade, using OISST
30
31 562 dataset), Saint-Paul and Amsterdam EEZ (12.9 km/decade ; 24.4 km/decade) and Kerguelen EEZ (11.8
32
33 563 km/decade ; 27.4 km/decade) while they are slower than global estimates or more variable over Crozet
34
35 564 EEZ (-2.6 km/decade ; 18.2 km/decade) and Prince Edward Islands EEZ (-5.8 km/decade ; 5.9). Crozet
36
37 565 and Prince Edward Islands EEZ are located in the area which seems to undergo a local cooling that
38
39
40 566 could be associated with high temporal variability (Figure 2, Supplementary Figure S2). However, the
41
42 567 analyses using IPSL-CM6A-LR members (see Sections 2.1.2 and 3.2.1) suggest that in the long term,
43
44 568 radiative forcing will likely dominate warming trends compared to internal variability, and an important
45
46 569 warming as well as fast climate velocities are projected over the Crozet Plateau (Section 3.2).

47
48
49
50 570

53 571 **MHW events are particularly intense at the ACC northern boundary between 20°E and 95°E**
54
55 572 **(north of the Kerguelen Plateau).** The patterns of MHW mean intensity between 1984 and 2014
56
57 573 obtained with the OSTIA dataset (Figure 3) are consistent with past estimates (Su et al., 2021).
58
59
60 574 However, zooming out from the northern part of the Kerguelen Plateau shows not only that the MHW

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

575 are more intense at the northern boundary of the ACC, but also that MHW are generally more intense
576 northward up to the subtropical zone. The global ocean mean intensity of MHW estimated from NOAA
577 observations and defined as the anomaly of temperature compared to the 90th percentile threshold is
578 around 0.35 °C (Plecha & Soares, 2020) when it can reach 1.2°C north of Crozet EEZ.

579
580 Over the historical period, no significant spatial contrasts in mean annual MHW days are
581 observed in the Southern Indian Ocean north of 60°S. The global ocean mean annual number of
582 days affected by MHW has been estimated from around 30 days/year in 1900 to around 60 days/year
583 in 2020 (Holbrook et al., 2020, using observations and models). Between 1984 and 2014 using OSTIA
584 dataset, a mean number of around 24 days per year affected by MHW (90th percentile) is observed
585 north of 60°S, with in average 23 annual MHW days over Kerguelen and Crozet EEZ (data not shown).
586 South of 60°S, the mean annual MHW days is lower (16 days/year).


587
588 Five contrasted zones emerge. When comparing the projected regional characteristics of both
589 warming and MHW characteristics (climatic impact-drivers), five zones (from the subtropics to the
590 Antarctic continent) associated with specific dynamical features stand out relative to the level of change
591 expected to occur (Figure 9). The first (north of 35°S, subtropical region) and fourth (between 50°S and
592 55°S, southern part of the ACC) zones are affected by intermediate amplitudes of warming-related
593 climatic impact-drivers. The second (between 35°S and 40°S, under the influence of the Agulhas Return
594 Current) and fifth (south of 55°S, Antarctic colder water) zones have weaker warming and weaker
595 changes in mean MHW intensity and in mean annual MHW days. Finally, the third zone (north of the
596 ACC and covering subantarctic islands) is the one with the most important magnitudes of warming-
597 related climatic impact-drivers, especially over topographic features: the Crozet Plateau, Kerguelen and
598 the Southeastern ridge.

599 **The projected mean MHW intensity might be underestimated in CMIP6.** It can be noted that the
1
2 600 projected mean MHW intensity might be underestimated, as it was found over the historical period
3
4 601 relative to observations (Figure 3) and in global studies using CMIP6 models (Pilo et al., 2019; Plecha
5
6 602 et al., 2020). Since the observations suggest that the most intense MHW are related to mesoscale activity
7
8
9 603 and that this mesoscale activity is not always well reproduced by climate models (Pilo et al., 2019; Su
10
11 604 et al., 2021), but could be expected to intensify (Hogg et al., 2015; Patara et al., 2016; Martinez-Moreno
12
13 605 et al., 2019; Martinez-Moreno et al., 2021), projections in terms of MHW intensity may be
14
15 606 underestimated.
16
17
18
19 607

21
22 608 Figure 9: Representation of the 5 zones identified in this study in terms of climatic impact-drivers'
23
24 609 trends (warming and marine heatwaves) along with the identification of areas for which surface
25
26 610 temperature increases by more than 1°C at the end of the century relative to 1995-2014 under SSP1-2.6
27
28
29 611 and SSP2-4.5; and areas for which mean MHW annual days are higher than 250 day/year at the end of
30
31 612 the century under SSP2-4.5 using CMIP6 models. Areas of chlorophyll a concentration greater than
32
33 613 0.25 mg.m-3 (using Copernicus-GlobColour product provided by ACRI-ST company, between 1997
34
35 614 and 2020) are identified in green to allow for comparison between spatialised climate trends and
36
37 615 spatialised biological characteristics.
38
39
40

41 616
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



4.2 Implications of climate mitigation

This study also shows how regional patterns of temperature change and marine heatwaves can differ between SSP1-2.6 and SSP2-4.5. Not only can the magnitude of changes be mitigated in SSP1-2.6 but some patterns may be reversed at the end of the century, for instance through the surface cooling obtained around 35°S. However, the major differences between SSP1-2.6 and SSP2-4.5 mainly occurred in the long term.

The consequences of following one SSP instead of another can also be identified at the EEZ scale, both in terms of emerging trends and of magnitude of change. The projected changes in surface temperature over Saint-Paul and Amsterdam EEZ in SSP1-2.6 remain in the range of the observed interannual variability at that location, even in the long term, which is not the case for the other EEZs. Already in the mid term in SSP1-2.6 Crozet, Heard and McDonalds Islands, Kerguelen and Prince Edward Islands


631 EEZ changes in surface temperature are found to be respectively 2.6, 2.5, 2.8 and 3.1 times higher than
1
2 632 the observed (using OSTIA) interannual variability in sea surface temperature (Figure 10). This
3
4 633 indicates that emerging warming trends will also occur in a high mitigation scenario (SSP1-2.6) for
5
6 634 subantarctic islands.
7

8
9
10 635 At the EEZ level, major differences between the two SSPs used in this study mainly occur in the long
11
12 636 term. In the long term , the change in surface temperature is 0.62°C , 0.49°C , 0.58°C , 0.65°C and
13
14 637 0.43°C warmer in SSP2-4.5 compared to SSP1-2.6 for Crozet, Heard and McDonalds Islands,
15
16 638 Kerguelen, Prince Edward Islands and Saint-Paul and Amsterdam EEZ, respectively. In the long term,
17
18 639 mean MHW intensity is 0.39°C , 0.33°C, 0.40°C, 0.40°C and 0.23°C warmer in SSP2-4.5 compared to
19
20 640 SSP1-2.6 for respectively Crozet, Heard and McDonalds Islands, Kerguelen, Prince Edward Islands
21
22 641 and Saint-Paul and Amsterdam EEZ (Figure 10).
23
24

25
26 642 This study therefore shows that the choice of a mitigation strategy and socio-economic development
27
28 643 pathway can have direct regional consequences on climate change patterns in the long term, even at the
29
30 644 scale of EEZs.
31
32

33
34 645

35
36
37
38 646 Figure 10: Map of subtropical and subantarctic islands' exclusive economic zones (EEZs) in the
39
40 647 Southern Indian Ocean (A). Change in surface temperature relative to 1995-2014 (B), mean MHW
41
42 648 intensity (D) and surface (C) and mesopelagic (E) climate velocities averaged over the Southern Indian
43
44 649 Ocean (SIO) and different EEZs : Crozet, Heard and Mc Donald Islands (HMI), Kerguelen (KP), Prince
45
46 650 Edward Islands (PEI) and Saint-Paul and Amsterdam (SPA), for SSP1-2.6 and SSP2-4.5. Changes in
47
48 651 surface temperature and mean MHW intensity are estimated over 2021-2040 (NT), 2041-2060 (MT)
49
50 652 and 2081-2100 (LT). Climate velocities were estimated over 2015-2065 (MT) and 2050-2100 (LT).
51
52 653 Change in surface temperature results are compared to the observed interannual variability in sea
53
54 654 surface temperature in each location calculated from the OSTIA dataset between 1982 and 2019.
55
56
57
58
59
60
61
62
63
64
65



655
656
657 **4.3 Coherence with known physical mechanisms and remaining questions**

658 The aim here is to better understand the projected intensities and patterns of warming and marine
 659 heatwaves in the Southern Indian Ocean. This requires an evaluation of the models and a mechanistic
 660 understanding of the potential heat uptake and redistribution processes.

661 It can be firstly noted that CMIP6 models have improved compared to CMIP5 models for some key
1
2 662 metrics in the Southern Ocean, such as the Antarctic Circumpolar Current (ACC) strength or the
3
4 663 representation of differences in density across latitudes of the ACC (Beadling et al., 2020). In addition,
5
6 664 the spatial patterns and interannual variability of MHW are better represented in CMIP6 compared to
7
8
9 665 CMIP5 models, and more generally tend to be better represented in higher resolution climate models
10
11 666 (Qiu et al., 2021).

12
13
14 667 There are at least three ways to evaluate a climate model: 1) to compare the model outputs to
15
16 668 observation-based products, 2) to verify the agreement between multiple models and/or models'
17
18 669 versions (this is done here by using an ensemble of models) and 3) to confront the projections to known
19
20
21 670 physical mechanisms (Baumberger et al., 2017). Given the important role of natural variability for
22
23 671 shaping the spatial patterns of historical warming trends, it is difficult to evaluate the quality of the
24
25 672 CMIP6 ensemble in simulating warming trends in the Southern Indian Ocean by simply comparing
26
27 673 them to past observations (Raäisaänen, 2007; Baumberger et al., 2017; Gopika et al., 2020). We
28
29
30 674 therefore focus here on determining whether the projected patterns of warming and marine heatwaves
31
32 675 are coherent with known or potential heat redistribution processes.

33
34
35 676 The intensity of the Southern Indian Ocean warming is not directly correlated to the intensity of global
36
37 677 warming. The models used in our study have a wide range of Equilibrium Climate Sensitivities (ECS;
38
39 678 defined as the long term global warming obtained after a doubling of the atmospheric CO_2 above its
40
41
42 679 pre-industrial concentration; Charney et al., 1979; Forster et al., 2021). The ECS of the model ensemble
43
44 680 we used in this study vary from 2.29 to 5.62 °C (Forster et al., 2021). However, no correlation was
45
46 681 found between the mean ocean warming in the region and the model ECS, that is, models displaying
47
48 682 greater warming are not necessarily the ones with greater ECS (data not shown), suggesting that the
49
50
51 683 regional warming in the Southern Indian Ocean is sensitive to local processes, reproduced differently
52
53 684 in different models. It is therefore particularly important to investigate whether the projected warming
54
55 685 and marine heatwaves patterns in the Southern Indian Ocean can be explained by known physical
56
57
58 686 processes active in the region.

687 The coherence of the projected patterns of warming-related climatic impact-drivers with known or
688 potential heat redistribution processes is discussed here to both assess the adequacy of the models
689 (Baumberger et al., 2017) and understand their main physical drivers.

690

691 *4.3.1 The impact of winds and local circulation on warming-related climatic*
692 *impact-drivers' patterns*

693

694 CMIP6 projections, either through the traditional or the time-shift approaches (i.e. using GWLs, see
695 Section 2.1.2), suggest an intensification of MHW and warming mostly in the region of influence of the
696 ACC, in the subantarctic latitudinal band between 40° and 55°S. Such result is consistent with the
697 observed increased heat uptake in the subpolar region transported northward by Ekman transport and
698 subducted within and north of the ACC (Armour et al., 2016; Frölicher et al., 2015; Morrison et al.,
699 2016; Sallée, 2018; Huguenin et al., 2022). The strength of this northward Ekman transport is modulated
700 by the position and strength of the westerlies which is particularly strong over the Indian sector of the
701 Southern Ocean (Lin et al., 2018). The westerlies are expected to intensify and to shift poleward (Lee
702 et al., 2021) and thus the associated northward transport of heat content anomaly (anomaly caused by
703 warming) may be strengthened, accumulating heat over the subantarctic (Swart et al., 2018; Fox-
704 Kemper et al., 2021; Silvy et al., 2022). In addition, southward heat transport also occur through eddy
705 advection, partly compensating the Ekman northward heat transport (Farneti et al., 2010; Saenko et al.,
706 2018), but there are still uncertainties on the degree of this compensation (Fox-Kemper et al., 2021).

707 Another pattern that stands out in this study is the lesser intensification of warming and marine
708 heatwaves between 35° and 40°S. This latitudinal band is mostly under the influence of the Agulhas
709 Current system, which consists in the Agulhas Current, a western boundary current flowing southward
710 along the African coast, which brings warm waters into the Indian Ocean at around 40°S through the
711 Agulhas Return Current up to 70°E (Lutjharms et al., 2006). The remaining waters are transported to

1 712 the South Atlantic between 32°S and 42°S and this westerward inflow of relatively warm and salty
2 713 water exiting the Indian sector is called the Agulhas leakage (Schmidt et al., 2021). Projections from
3
4 714 global climate models suggest an increase in Agulhas leakage (Rouault et al., 2009; Backeberg et al.,
5
6 715 2012; Biastoch & Böning, 2013), and a decrease of Agulhas Current transport by 11 to 23% at the end
7
8 716 of the century (Stellema et al., 2019; Ma et al., 2020; Sen Gupta et al., 2021). If we assume that the
9
10 717 decreasing Agulhas Current feeds both the increasing Agulhas leakage and the Agulhas Return Current,
11
12 718 and that the Agulhas Current is the main source of water for the Agulhas Return Current, then the
13
14 719 Agulhas Return Current would have to decrease. A decreasing Agulhas Return Current would slow
15
16 720 down a large heat transport source in the northern part of the Southern Indian Ocean, which could
17
18 721 potentially explain the lesser warming projected in this area.
19
20
21
22

23 722 The Agulhas Current system transport is also modulated by the position and strength of the westerlies.
24
25 723 Indeed, an equatorward shift of the westerlies (as observed in summer in SSP1-2.6, Bracegirdle et al.,
26
27 724 2020) can be associated with an increase of the Agulhas leakage (Durgadoo et al., 2013).
28
29
30

31 725 Latitudinal differences in projected warming trends and climate velocities seem therefore consistent
32
33 726 with known mechanisms in the region. In particular, the position and strength of the westerlies can have
34
35 727 a different effect on a specific zone, depending on how westerlies impact the local circulation.
36
37
38

39 728

40
41
42 729 *4.3.2 Remaining questions: on the role of unresolved spatial scales and*
43
44 730 *modes of variability*
45
46

47 731

48
49
50
51 732 The projected trends in warming and marine heatwaves therefore seem consistent with potential
52
53 733 mechanisms of heat uptake and transport but some key questions remain. Uncertainties remain
54
55 734 concerning the role of the ocean mesoscale circulation and the role of the decadal and longer variability.
56
57

58 735
59
60
61
62
63
64
65

736 Finer scale processes (meso- and submesoscale) have the potential to impact warming trends. Eddies
1
2 737 can play a significant role in meridional heat transport (Morrison et al., 2016) and can for instance
3
4 738 contribute to modulate heat uptake through submesoscale ventilation (Dove et al., 2021). Increased
5
6 739 upward heat transport at submesoscale fronts can also alter surface heat uptake capacity (Siegelmann
7
8 740 et al., 2020) but the impact of such a process on warming trends locally is yet to be studied. In a changing
9
10 741 climate with intensifying winds, it is expected that mesoscale activity might change (Martinez-Moreno
11
12 742 et al., 2019; Martinez-Moreno et al., 2021). However, it remains unclear how such changes might also
13
14 743 impact the projected warming trends. To investigate whether resolution could play a role in determining
15
16 744 the patterns of projected temperature changes, we focus on two models, GFDL-CM4 and MPI-ESM1-
17
18 745 2-HR with a resolution of 0.25° and 0.4° respectively. As for the model ensemble, a pattern of greater
19
20 746 surface warming over the subantarctic is observed (Supplementary Figure S8). This suggests that the
21
22 747 influence of smaller scale processes on the warming patterns may remain limited, at least for the larger
23
24 748 eddies represented in higher resolution CMIP6 models. The net impact of eddies on the heat budget and
25
26 749 how it will evolve under climate change remain an active field of research (Hewitt et al., 2022).

30
31
32 750

33
34
35 751 Internal variability is not negligible in the Southern Indian ocean between 1982 and 2019 but it is
36
37 752 challenging to evaluate whether the representation of decadal and multidecadal variability modes and
38
39 753 their interaction with the local circulation in the models is accurate. The Southern Indian Ocean is
40
41 754 indeed under the influence of various modes of natural variability but depending on the time scale of
42
43 755 interest, for instance for policy-makers, different modes of natural variability are to be considered. The
44
45 756 Indian Ocean Subtropical Dipole (IOSD) and the El-Niño-Southern Oscillation (ENSO) can influence
46
47 757 patterns of SST anomaly (Behera and Yamagata, 2001; Huang and Shuckla, 2008; Morioka et al., 2010)
48
49 758 on decadal timescales and the Southern Annular Mode (SAM) on even shorter timescales (Sallée et al.,
50
51 759 2010). As the above-mentioned climatic modes of variability may not be in the same phase in reality
52
53 760 and in the climate models, this increases the uncertainty on near-term climate projections (Hurrell et al.,
54
55 761 2010; Chen et al., 2021).

56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

762 Multidecadal variability may also not be negligible in the Southern Indian Ocean and could potentially
763 impact projected trends on longer timescales (Zhang et al., 2018). Indeed, the natural variability
764 observed in this study is important over a 37-year period. The east/west contrast obtained in the
765 observations (Section 3.1.1) could result from the response to the positive-to-negative phase transition
766 of the Atlantic Multidecadal Variability (AMV) but the teleconnexion processes associated may not be
767 well represented in climate models (Chung et al., 2022). There are therefore uncertainties on how
768 natural modes of variability and their interactions with the local circulation will evolve, and whether
769 they will enhance, buffer or mask the projected changes.

771 **4.4 Potential impacts on ecosystems**

772
773 Using the warming-related climatic impact-drivers regional characteristics, it is then important to make
774 the link with the regional ecological characteristics. Here, based on the observed and projected trends
775 in warming and MHW and based on knowledge on the ecology of the Southern Indian Ocean, examples
776 of potential impacts are discussed.

777
778 Climate change might induce a shift in habitats for various species, thus potentially affecting the
779 distribution of species with conservation and/or economic values (Reisinger et al., 2022). The region
780 studied here covers three biogeochemical provinces (i.e. oceanographically and ecologically relatively
781 homogeneous regions) as defined by Reygondeau et al., 2020, distributed latitudinally from north to
782 south: the South Subtropical convergence, the Subantarctic water ring and the Antarctic (Reygondeau
783 et al., 2020). Our results suggest that, as far as temperature interactions are concerned, a poleward shift
784 of these provinces is expected and this shift is shown to be faster for the mesopelagic layer compared
785 to the surface with potential ecological impacts for species living across multiple depth layers
786 throughout their life cycle. Such shift between surface and mesopelagic conditions is particularly

1 787 relevant for the Saint-Paul and Amsterdam region which is currently located at the transition between
2 788 different bioregions: the South subtropical convergence in the north and the Subantarctic water ring in
3
4 789 the south. By the end of the century, the temperature conditions south of Saint-Paul and Amsterdam
5
6 790 EEZ will probably be similar to the ones in the south subtropical convergence province. For the north
7
8 791 of the EEZ, the mesopelagic layer temperature conditions will shift faster and could be similar to the
9
10 792 ones in the Indian South Subtropical Gyre province. Our study helps identifying other areas that could
11
12 793 be undergoing an important shift between surface and mesopelagic conditions and therefore where long
13
14 794 term biophysical monitoring could be of special interest: the north-western side of Kerguelen Plateau,
15
16 795 slightly up to the southern part of the Crozet Plateau which has been identified as a foraging zone for
17
18 796 top predators (e.g. Pütz et al., 2002).

19
20
21
22
23 797 The consequences of the long term shift of thermal conditions in top predators' foraging zones are
24
25 798 anticipated by extreme events in the near term (or in the past). The 1997 MHW that occurred in the
26
27 799 northern part of the Southern Indian Ocean has been associated with a southward shift of the Polar Front
28
29 800 south of the Crozet Plateau where king penguins (*Aptenodytes patagonicus*) from Crozet islands usually
30
31 801 forage. The temperature anomaly was so intense and associated with such a major shift of the foraging
32
33 802 zones that these zones were barely accessible to the penguins leading to important mortality of those
34
35 803 populations (Bost et al., 2015). Over the whole region, the mean MHW intensity of this event remained
36
37 804 greater than 0.5°C for 36 days (obtained using OSTIA dataset). Given models projections it appears
38
39 805 that such a drastic event may no longer be an extreme phenomenon and if associated with a significant
40
41 806 shift in the Polar Front position, it could lead to systematic mass mortality events for top predators.

42
43
44
45
46 807 Intermediate levels of the food webs can also be affected by warming and MHW. Indeed, some
47
48 808 populations of fish are already living at the edge of their thermal tolerance. Low-antarctic (i.e. just south
49
50 809 of the Antarctic Polar Front) species can live in conditions above their upper optimal range of
51
52 810 temperature (1°C to 1.5°C) but further increases in temperature may become too physiologically
53
54 811 demanding, especially as the intensification projected appears to occur over their distribution areas
55
56 812 (Kock and Everson, 2003).

57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

813 Besides, a whole range of MHW potential impacts in the Southern Indian Ocean has not yet been
814 covered. Such impacts could include primary production changes (Hayashida et al., 2020), changes in
815 species growth and abundance (Oliver, 2019b; Smale & Wernberg, 2013), changes in population
816 structure (Smale et al., 2017), behavioral changes, notably for reproduction and foraging (Fromant et
817 al., 2021), changes in geographic distributions of species (Cure et al., 2017; Smale & Wernberg, 2013)
818 or even genetic changes (e.g., Coleman et al., 2020). These impacts are yet to be studied in the Southern
819 Indian Ocean.

820 821 **4.5 Climate change and conservation**

822
823 A regionalisation of temperature-related climatic impact-drivers, associated with further research on
824 climate change impacts on ecosystems, should contribute to the further consideration of climate change
825 impacts in conservation design and management. Increasing literature aims at addressing this issue and
826 suggests management tools to help integrate climate resilience into MPA design (Tittensor et al., 2019;
827 Crespo et al., 2020). However, efforts are needed to identify the risks posed by climate change at the
828 regional scale. Our study of warming-related climatic impact-drivers mostly addresses the hazards
829 dimension of a risk assessment (which would also include an analysis of the vulnerability and exposure
830 of ecosystems, Chen et al., 2021) and points out the need to include the vertical analysis in the eco
831 regionalisation process, since different layer depths may be impacted differently. In particular, this
832 study shows that the relative difference between mesopelagic and surface climate velocity (a « climatic
833 vertical shear ») is highest over Kerguelen Plateau and Saint-Paul and Amsterdam EEZ under both
834 scenarios while it is the lowest over Heard and McDonalds Islands and Prince Edward Islands EEZ
835 (Figure 10). Yet, mesopelagic climate velocities over Heard and McDonalds Islands and Prince Edward
836 Islands EEZ are still between 20 and 30% faster than surface climate velocities in SSP2-4.5; and
837 respectively between 50% and 60% and between 35% and 66% faster in SSP1-2.6 (Figure 10).

1 838 The 3D information provided by climate velocity could be further used to project potential changes in
2 839 marine biodiversity distribution and help inform conservation management (Arafeh-Dalmau et al.,
3
4 840 2021). Through this type of methodology, other concepts related to climate velocity can be used to
5
6 841 understand the impact of climate change on species distribution, notably i) climate residence time
7
8 842 (Loarie et al., 2009) and ii) climate refugia (Burrows et al., 2014; Brito-Morales et al., 2018). Indeed, it
9
10 843 can be useful to know not only if certain environmental conditions will be found and where but also
11
12 844 how long these conditions will last in a specific area (climate residence time, Loarie et al., 2009). The
13
14 845 concept of climate refugia aims to identify areas that are relatively less impacted by climate change,
15
16 846 meaning low climate velocity and/or long climate residence time (Brito-Morales et al., 2018), but it can
17
18 847 also be based on the time of emergence of the climate change signal to identify temporary refugia
19
20 848 (Bruno et al., 2018). This concept could guide the development of surveillance programs to follow the
21
22 849 evolution of environmental conditions on specific zones that are particularly affected by climatic
23
24 850 impact-drivers. Climate velocity could also be used to anticipate future expansions of species and to
25
26 851 check if current MPA networks cover these expansions (Arafeh-Dalmau et al., 2021). Note that, to be
27
28 852 even more biologically meaningful, climate velocity should be combined with other constraints of
29
30 853 dispersion such as habitat permeability or connectivity (Brito-Morales et al., 2018). It may also be
31
32 854 important to consider multiple variables such as pH (acidification), primary production or zooplankton
33
34 855 abundance, since the interaction between multiple variables changes can result in multi-directional
35
36 856 distribution shifts (e.g., VanDerWal et al., 2013).

37
38 857 From the characterisation of warming and marine heatwaves patterns presented here, it is difficult to
39
40 858 identify potential climate refugia. However, this study pinpoints areas that are scientifically interesting
41
42 859 to further study the impacts of climate change on ecosystems, for instance the area south of Crozet near
43
44 860 the Antarctic Polar Front and Saint-Paul and Amsterdam EEZ. Indeed, the area south of Crozet, in the
45
46 861 main zone of intensification identified in this study, is an important foraging zone for top predators (e.g.
47
48 862 Pütz et al., 2002) and is expected to undergo intense warming and increases in MHW intensity and
49
50 863 duration but it is also an area that can be subject to important natural variability (as seen on Figure 2).
51
52 864 It can therefore be very interesting to study whether this variability in the long term will be enough to
53
54
55
56
57
58
59
60
61
62
63
64
65

1 865 counteract warming trends and whether ecosystems in this area compared to similar ecosystems east of
2 866 the Kerguelen Plateau (at the same latitudes) will develop different features as a response to different
3
4 867 changing conditions. The Saint-Paul and Amsterdam area can also be interesting to protect for further
5
6 868 scientific investigation on how the difference between surface and mesopelagic temperature changes
7
8 869 might affect endemic ecosystems.
9

10
11
12 870

13
14
15 871

16
17
18 872 **5. Conclusion**
19

20
21
22 873 The aim of this work is to provide a systematic description of observed and future temperature changes
23
24 874 and marine heatwaves over the entire Southern Indian Ocean and to compare the projections over this
25
26 875 region under two scenarios that reflect two possible socio-economic pathways.
27

28
29
30 876 This analysis shows the spatial heterogeneity of temperature trends and marine heatwaves
31
32 877 characteristics at the regional scale, highlighting also some limits. The projected warming trends appear
33
34 878 to be consistent with warming mechanisms identified in the literature, notably related to the dynamic
35
36 879 of the ACC and the intensification and shifts of westerly winds. Concerning MHW intensity pattern, a
37
38 880 better spatial fit between the observations and the models was obtained, although intensities may be
39
40 881 underestimated in the models, mainly because of resolution limits, and current spatial patterns of MHW
41
42 882 intensity are expected to intensify.
43

44
45
46 883 This intensification of both warming and marine heatwaves characteristics is expected to occur mostly
47
48 884 over the subantarctic islands, with consequences on endemic species and ecosystems that are still to be
49
50 885 further studied. It can be noted that the difference between SSP1-2.6 and SSP2-4.5 projections are
51
52 886 mostly significant in the long term, with changes in both scenarios that could be important for
53
54 887 ecosystems already in the near and mid term, highlighting the need to anticipate adaptation measures.
55
56 888 Such measures would need to also consider that surface and deeper conditions may evolve at different
57
58 889 pace.
59
60

61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

890 Through a regional example, this study also reaffirms the need to globally commit to strong mitigation
891 strategies and to follow a sustainable socio-economic development pathway to alleviate the potential
892 impacts of warming and MHW.

893

894 Acknowledgements

895

896 We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling,
897 that is responsible for CMIP. The authors also thank the IPSL modelling group for the software
898 infrastructure, which facilitated CMIP analysis. We acknowledge funding support from CNES and from
899 the program LEFE led by CNRS-INSU (projects KERTREND and KERTREND-SAT). C.A. was
900 financially supported by the French Ministry of Ecological Transition. A.P. acknowledges the financial
901 support of the Project Concytec - World Bank "Characterization and forecast of extreme events in the
902 Peruvian sea using an operational system of oceanic information", through its executing unit Fondo
903 Nacional de Desarrollo Científico, Tecnológico y de Innovación Tecnológica (Fondecyt). L.B.
904 acknowledges funding from the European Union's Horizon 2020 research and innovation program
905 under grant agreement no. 820989 (project COMFORT).

906

907 Declaration of Competing Interest

908

909 The authors declare that they have no known competing financial interests or personal relationships that
910 could have appeared to influence the work reported in this paper.

911

1
2
3
4 912 Appendix A : Supplementary material

5 913

6
7
8 914 References

9
10
11 915

12
13
14
15 916 Ab, M. (2019). Expert workshop on pelagic spatial planning for the Eastern Subantarctic region (domains
16
17 917 4, 5 and 6). 72.

18
19
20
21 918 Andrews, M. B., Ridley, J. K., Wood, R. A., Andrews, T., Blockley, E. W., Booth, B., Burke, E., Dittus,
22
23 919 A. J., Florek, P., Gray, L. J., Haddad, S., Hardiman, S. C., Hermanson, L., Hodson, D., Hogan, E., Jones,
24
25 920 G. S., Knight, J. R., Kuhlbrodt, T., Misios, S., ... Sutton, R. T. (2020). Historical Simulations With
26
27 921 HadGEM3-GC3.1 for CMIP6. *Journal of Advances in Modeling Earth Systems*, 12(6),
28
29 922 e2019MS001995. <https://doi.org/10.1029/2019MS001995>

30
31
32
33 923 Arafteh-Dalmau, N., Brito-Morales, I., Schoeman, D. S., Possingham, H. P., Klein, C. J., & Richardson,
34
35 924 A. J. (2021). Incorporating climate velocity into the design of climate-smart networks of marine
36
37 925 protected areas. *Methods in Ecology and Evolution*, n/a(n/a). <https://doi.org/10.1111/2041-210X.13675>

38
39
40
41 926 Armour, K. C., Marshall, J., Scott, J. R., Donohoe, A., & Newsom, E. R. (2016). Southern Ocean
42
43 927 warming delayed by circumpolar upwelling and equatorward transport. *Nature Geoscience*, 9(7),
44
45 928 549- 554. <https://doi.org/10.1038/ngeo2731>

46
47
48
49 929 Auger, M., Morrow, R., Kestenare, E., Sallée, J.-B., & Cowley, R. (2021). Southern Ocean in-situ
50
51 930 temperature trends over 25 years emerge from interannual variability. *Nature Communications*, 12(1),
52
53 931 514. <https://doi.org/10.1038/s41467-020-20781-1>

54
55
56 932 Ban, N. C., Maxwell, S. M., Dunn, D. C., Hobday, A. J., Bax, N. J., Ardron, J., Gjerde, K. M., Game,
57
58 933 E. T., Devillers, R., Kaplan, D. M., Dunstan, P. K., Halpin, P. N., & Pressey, R. L. (2014). Better

- 934 integration of sectoral planning and management approaches for the interlinked ecology of the open
1
2 935 oceans. *Marine Policy*, 49, 127- 136. <https://doi.org/10.1016/j.marpol.2013.11.024>
3
4
5 936 Banzon, V., Smith, T. M., Chin, T. M., Liu, C., & Hankins, W. (2016). A long-term record of blended
6
7 937 satellite and in situ sea-surface temperature for climate monitoring, modeling and environmental
8
9
10 938 studies. *Earth System Science Data*, 8(1), 165- 176. <https://doi.org/10.5194/essd-8-165-2016>
11
12
13 939 Baumberger, C., Knutti, R., & Hirsch Hadorn, G. (2017). Building confidence in climate model
14
15 940 projections: An analysis of inferences from fit. *WIREs Climate Change*, 8(3), e454.
16
17 941 <https://doi.org/10.1002/wcc.454>
18
19
20
21 942 Beadling, R. L., Russell, J. L., Stouffer, R. J., Mazloff, M., Talley, L. D., Goodman, P. J., Sallée, J. B.,
22
23 943 Hewitt, H. T., Hyder, P., & Pandde, A. (2020). Representation of Southern Ocean Properties across
24
25 944 Coupled Model Intercomparison Project Generations: CMIP3 to CMIP6. *Journal of Climate*, 33(15),
26
27 945 6555- 6581. <https://doi.org/10.1175/JCLI-D-19-0970.1>
28
29
30
31 946 Behera, S. K., & Yamagata, T. (2001). Subtropical SST dipole events in the southern Indian Ocean.
32
33 947 *Geophysical Research Letters*, 28(2), 327- 330. <https://doi.org/10.1029/2000GL011451>
34
35
36 948 Belchier, M., & Collins, M. A. (2008). Recruitment and body size in relation to temperature in juvenile
37
38 949 Patagonian toothfish (*Dissostichus eleginoides*) at South Georgia. *Marine Biology*, 155, 493- 503.
39
40
41 950 <https://doi.org/10.1007/s00227-008-1047-3>
42
43
44 951 Bi, D., Dix, M., Marsland, S., O'Farrell, S., Sullivan, A., Bodman, R., Law, R., Harman, I., Srbinovsky,
45
46 952 J., Rashid, H. A., Dobrohotoff, P., Mackallah, C., Yan, H., Hirst, A., Savita, A., Dias, F. B., Woodhouse,
47
48 953 M., Fiedler, R., Heerdegen, A., ... Heerdegen, A. (2020). Configuration and spin-up of ACCESS-CM2,
49
50
51 954 the new generation Australian Community Climate and Earth System Simulator Coupled Model.
52
53 955 *Journal of Southern Hemisphere Earth Systems Science*, 70(1), 225- 251.
54
55 956 <https://doi.org/10.1071/ES19040>
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 957 Bost, C. A., Cotté, C., Terray, P., Barbraud, C., Bon, C., Delord, K., Gimenez, O., Handrich, Y., Naito,
958 Y., Guinet, C., & Weimerskirch, H. (2015). Large-scale climatic anomalies affect marine predator
959 foraging behaviour and demography. *Nature Communications*, 6(1), 8220.
960 <https://doi.org/10.1038/ncomms9220>
- 961 Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet,
962 R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F.,
963 Cozic, A., Cugnet, D., D'Andrea, F., ... Vuichard, N. (2020). Presentation and Evaluation of the IPSL-
964 CM6A-LR Climate Model. *Journal of Advances in Modeling Earth Systems*, 12(7), e2019MS002010.
965 <https://doi.org/10.1029/2019MS002010>
- 966 Bracegirdle, T.J., Krinner, G., Tonelli, M., Haumann, F. A., Naughten, K. A., Rackow, T., Roach, L.
967 A. and Wainer, I. (2020). Twenty first century changes in Antarctic and Southern Ocean surface climate
968 in CMIP6. *Atmos Sci Lett.*, 21:e984. <https://doi.org/10.1002/asl.984>
- 969 Brito-Morales, I., García Molinos, J., Schoeman, D. S., Burrows, M. T., Poloczanska, E. S., Brown, C.
970 J., Ferrier, S., Harwood, T. D., Klein, C. J., McDonald-Madden, E., Moore, P. J., Pandolfi, J. M.,
971 Watson, J. E. M., Wenger, A. S., & Richardson, A. J. (2018). Climate Velocity Can Inform
972 Conservation in a Warming World. *Trends in Ecology & Evolution*, 33(6), 441- 457.
973 <https://doi.org/10.1016/j.tree.2018.03.009>
- 974 Brito-Morales, I., Schoeman, D. S., Molinos, J. G., Burrows, M. T., Klein, C. J., Arafteh-Dalmau, N.,
975 Kaschner, K., Garilao, C., Kesner-Reyes, K., & Richardson, A. J. (2020). Climate velocity reveals
976 increasing exposure of deep-ocean biodiversity to future warming. *Nature Climate Change*, 10(6),
977 576- 581. <https://doi.org/10.1038/s41558-020-0773-5>
- 978 Bruno, J. F., Bates, A. E., Cacciapaglia, C., Pike, E. P., Amstrup, S. C., van Hooidek, R., Henson, S.
979 A., & Aronson, R. B. (2018). Climate change threatens the world's marine protected areas. *Nature*
980 *Climate Change*, 8(6), 499- 503. <https://doi.org/10.1038/s41558-018-0149-2>

- 1 981 Burrows, M. T., Schoeman, D. S., Richardson, A. J., Molinos, J. G., Hoffmann, A., Buckley, L. B.,
2 982 Moore, P. J., Brown, C. J., Bruno, J. F., Duarte, C. M., Halpern, B. S., Hoegh-Guldberg, O., Kappel, C.
3
4 983 V., Kiessling, W., O'Connor, M. I., Pandolfi, J. M., Parmesan, C., Sydeman, W. J., Ferrier, S., ...
5
6 984 Poloczanska, E. S. (2014). Geographical limits to species-range shifts are suggested by climate velocity.
7
8 985 Nature, 507(7493), 492- 495. <https://doi.org/10.1038/nature12976>
9
10
11
12 986 Butt, N., Possingham, H. P., De Los Rios, C., Maggini, R., Fuller, R. A., Maxwell, S. L., & Watson, J.
13
14 987 E. M. (2016). Challenges in assessing the vulnerability of species to climate change to inform
15
16 988 conservation actions. Biological Conservation, 199, 10- 15.
17
18 989 <https://doi.org/10.1016/j.biocon.2016.04.020>
19
20
21
22 990 Cai, W., Cowan, T., Godfrey, S., & Wijffels, S. (2010). Simulations of Processes Associated with the
23
24 991 Fast Warming Rate of the Southern Midlatitude Ocean. Journal of Climate, 23(1), 197- 206.
25
26 992 <https://doi.org/10.1175/2009JCLI3081.1>
27
28
29
30 993 Cao, J., Ma, L., Liu, F., Chai, J., Zhao, H., He, Q., Wang, B., Bao, Y., Li, J., Yang, Y., Deng, H., &
31
32 994 Wang, B. (2021). NUIST ESM v3 Data Submission to CMIP6. Advances in Atmospheric Sciences,
33
34 995 38(2), 268- 284. <https://doi.org/10.1007/s00376-020-0173-9>
35
36
37 996 Carr, M. H., Woodson, C. B., Cheriton, O. M., Malone, D., McManus, M. A., & Raimondi, P. T. (2011).
38
39 997 Knowledge through partnerships : Integrating marine protected area monitoring and ocean observing
40
41 998 systems. Frontiers in Ecology and the Environment, 9(6), 342- 350. <https://doi.org/10.1890/090096>
42
43
44
45 999 Cavole, L. M., Demko, A. M., Diner, R. E., Giddings, A., Koester, I., Pagniello, C. M. L. S., Paulsen,
46
47 1000 M.-L., Ramirez-Valdez, A., Schwenck, S. M., Yen, N. K., Zill, M. E., & Franks, P. J. S. (2016).
48
49 1001 Biological Impacts of the 2013–2015 Warm-Water Anomaly in the Northeast Pacific. Oceanography,
50
51 1002 29(2), 273- 285. JSTOR.
52
53
54
55 1003 Charney, J. G., Arakawa, A., Baker, D. J., Bolin, B., Dickinson, R. E., Goody, R. M., Leith, C. E.,
56
57 1004 Stommel, H. M., & Wunsch, C. I. (1979). Carbon dioxide and climate: a scientific assessment. Climate
58
59 1005 Research Board, 13.
60
61
62
63
64
65

- 1006 Chen, D., M. Rojas, B.H. Samset, K. Cobb, A. Diongue Niang, P. Edwards, S. Emori, S.H. Faria, E.
1
21007 Hawkins, P. Hope, P.Huybrechts, M. Meinshausen, S.K. Mustafa, G.-K. Plattner, and A.-M. Tréguier,
3
41008 2021: Framing, Context, and Methods. In *Climate Change 2021: The Physical Science Basis*.
5
61009 Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on
7
81010 Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud,
9
10
111011 Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock,
12
131012 T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United
14
151013 Kingdom and New York, NY, USA, pp. 147–286, doi:10.1017/9781009157896.003.
16
17
181014 Cherchi, A., Fogli, P. G., Lovato, T., Peano, D., Iovino, D., Gualdi, S., Masina, S., Scoccimarro, E.,
19
20
211015 Materia, S., Bellucci, A., & Navarra, A. (2019). Global Mean Climate and Main Patterns of Variability
22
231016 in the CMCC-CM2 Coupled Model. *Journal of Advances in Modeling Earth Systems*, 11(1), 185- 209.
24
251017 <https://doi.org/10.1029/2018MS001369>
26
27
281018 Christian, J. R., Denman, K. L., Hayashida, H., Holdsworth, A. M., Lee, W. G., Riche, O. G. J., Shao,
29
30
311019 A. E., Steiner, N., & Swart, N. C. (2021). Ocean biogeochemistry in the Canadian Earth System Model
32
331020 version 5.0.3: CanESM5 and CanESM5-CanOE. *Geoscientific Model Development Discussions*,
34
351021 1- 68. <https://doi.org/10.5194/gmd-2021-327>
36
37
381022 Chung, E.-S., Kim, S.-J., Timmermann, A., Ha, K.-J., Lee, S.-K., Stuecker, M. F., Rodgers, K. B., Lee,
39
40
411023 S.-S., & Huang, L. (2022). Antarctic sea-ice expansion and Southern Ocean cooling linked to tropical
42
431024 variability. *Nature Climate Change*, 12(5), 461- 468. <https://doi.org/10.1038/s41558-022-01339-z>
44
45
461025 Civel-Mazens, M., Crosta, X., Cortese, G., Michel, E., Mazaud, A., Ther, O., Ikehara, M., & Itaki, T.
47
48
491026 (2021). Impact of the Agulhas Return Current on the oceanography of the Kerguelen Plateau region,
50
511027 Southern Ocean, over the last 40 kyrs. *Quaternary Science Reviews*, 251, 106711.
52
531028 <https://doi.org/10.1016/j.quascirev.2020.106711>
54
55
561029 Coleman, M. A., Minne, A. J. P., Vranken, S., & Wernberg, T. (2020). Genetic tropicalisation following
57
58
591030 a marine heatwave. *Scientific Reports*, 10(1), 12726. <https://doi.org/10.1038/s41598-020-69665-w>
60
61
62
63
64
65

- 1031 Collins, M. A., Brickle, P., Brown, J., & Belchier, M. (2010). Chapter Four - The Patagonian Toothfish :
1
21032 Biology, Ecology and Fishery. In M. Lesser (Éd.), *Advances in Marine Biology* (Vol. 58, p. 227- 300).
3
41033 Academic Press. <https://doi.org/10.1016/B978-0-12-381015-1.00004-6>
5
6
7
81034 Constable, A. J., Melbourne-Thomas, J., Corney, S. P., Arrigo, K. R., Barbraud, C., Barnes, D. K. A.,
9
101035 Bindoff, N. L., Boyd, P. W., Brandt, A., Costa, D. P., Davidson, A. T., Ducklow, H. W., Emmerson,
11
121036 L., Fukuchi, M., Gutt, J., Hindell, M. A., Hofmann, E. E., Hosie, G. W., Iida, T., ... Ziegler, P. (2014).
13
141037 Climate change and Southern Ocean ecosystems I: How changes in physical habitats directly affect
15
161038 marine biota. *Global Change Biology*, 20(10), 3004- 3025. <https://doi.org/10.1111/gcb.12623>
17
18
19
201039 Cooley, S., D. Schoeman, L. Bopp, P. Boyd, S. Donner, D.Y. Ghebrehiwet, S.-I. Ito, W. Kiessling, P.
21
221040 Martinetto, E. Ojea, M.-F. Racault, B. Rost, and M. Skern-Mauritzen (2022). *Oceans and Coastal*
23
241041 *Ecosystems and Their Services*. In: *Climate Change 2022: Impacts, Adaptation and Vulnerability*.
25
261042 Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on
27
281043 Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría,
29
301044 M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press,
31
32
331045 Cambridge, UK and New York, NY, USA, pp. 379–550, doi:10.1017/9781009325844.005.
34
35
361046 Cotté, C., Ariza, A., Berne, A., Habasque, J., Lebourges-Dhaussy, A., Roudaut, G., Espinasse, B., Hunt,
37
381047 B.P.V., Pakhomov, E.A., Henschke, N., Péron, C., Conchon, A., Koedooder, C., Iazard, L., Cherel, Y.
39
40
411048 (2022). Macrozooplankton and micronekton diversity and associated carbon vertical patterns and fluxes
42
431049 under distinct productive conditions around the Kerguelen Islands. *Journal of Marine Systems*, 226,
44
451050 471-492. <https://doi.org/10.1016/B978-0-12-391851-2.00018-0>.
46
47
48
491051 Crespo, G. O., Mossop, J., Dunn, D., Gjerde, K., Hazen, E., Reygondeau, G., Warner, R., Tittensor, D.,
50
511052 & Halpin. (s. d.). *Beyond static spatial management : Scientific and legal considerations for dynamic*
52
531053 *management in the high seas*.
54
55
561054 Cure, K., Hobbs, J.-P. A., Langlois, T. J., Abdo, D. A., Bennett, S., & Harvey, E. S. (2017).
57
58
591055 *Distributional responses to marine heat waves : Insights from length frequencies across the geographic*
60
61
62
63
64
65

- 1056 range of the endemic reef fish *Choerodon rubescens*. *Marine Biology*, 165(1), 1.
1
21057 <https://doi.org/10.1007/s00227-017-3259-x>
3
4
- 51058 Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons,
6
71059 L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P.
8
9
101060 H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., ... Strand, W. G.
11
121061 (2020). The Community Earth System Model Version 2 (CESM2). *Journal of Advances in Modeling*
13
141062 *Earth Systems*, 12(2), e2019MS001916. <https://doi.org/10.1029/2019MS001916>
15
16
- 171063 Della Penna, A., Koubbi, P., Cotté, C., Bon, C., Bost, C.-A., & d'Ovidio, F. (2017). Lagrangian analysis
18
191064 of multi-satellite data in support of open ocean Marine Protected Area design. *Future of oceanic animals*
20
211065 in a changing ocean, 140, 212- 221. <https://doi.org/10.1016/j.dsr2.2016.12.014>
22
23
24
- 251066 Delord, K., Barbraud, C., Bost, C.-A., Cherel, Y., Guinet, C., & Weimerskirch, H. (2014). Atlas of top
26
271067 predators from French Southern Territories in the Southern Indian Ocean (p.
28
291068 http://www.cebc.cnrs.fr/ecommm/Fr_ecomm/ecommm_ecor_OI1.html) [Research Report]. CNRS.
30
311069 https://doi.org/10.15474/AtlasTopPredatorsOI_CEBC.CNRS_FrenchSouthernTerritories
32
33
34
- 351070 Dijkstra, H. (2016). A Normal Mode Perspective of Intrinsic Ocean-Climate Variability. *Annual*
36
371071 *Review of Fluid Mechanics*, 48(1), 341- 363. <https://doi.org/10.1146/annurev-fluid-122414-034506>
38
39
- 40
411072 Donohue, K. A., Tracey, K. L., Watts, D. R., Chidichimo, M. P., and Chereskin, T. K. (2016). Mean
42
431073 Antarctic Circumpolar Current transport measured in Drake Passage. *Geophys. Res. Lett.*, 43, 11760–
44
451074 11767. doi:10.1002/2016GL070319
46
47
- 48
491075 Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arneth, A., Arsouze, T., Bergmann, T.,
50
511076 Bernadello, R., Bousetta, S., Caron, L.-P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini,
52
531077 P., Dekker, E., Doblas-Reyes, F. J., Docquier, D., Echevarria, P., ... Zhang, Q. (2021). The EC-Earth3
54
551078 Earth System Model for the Climate ModelIntercomparison Project 6 [Preprint]. *Climate and Earth*
56
571079 *system modeling*. <https://doi.org/10.5194/gmd-2020-446>
58
59
60
61
62
63
64
65

- 1080 Dove, L. A., Thompson, A. F., Balwada, D., & Gray, A. R. (2021). Observational Evidence of
1
21081 Ventilation Hotspots in the Southern Ocean. *Journal of Geophysical Research: Oceans*, 126(7),
3
41082 e2021JC017178. <https://doi.org/10.1029/2021JC017178>
5
6
7
81083 Dunne, J. P., Horowitz, L. W., Adcroft, A. J., Ginoux, P., Held, I. M., John, J. G., Krasting, J. P.,
9
101084 Malyshev, S., Naik, V., Paulot, F., Shevliakova, E., Stock, C. A., Zadeh, N., Balaji, V., Blanton, C.,
11
121085 Dunne, K. A., Dupuis, C., Durachta, J., Dussin, R., ... Zhao, M. (2020). The GFDL Earth System Model
13
141086 Version 4.1 (GFDL-ESM 4.1): Overall Coupled Model Description and Simulation Characteristics.
15
161087 *Journal of Advances in Modeling Earth Systems*, 12(11), e2019MS002015.
17
18
191088 <https://doi.org/10.1029/2019MS002015>
20
21
221089 Durgadoo, J. V., Loveday, B. R., Reason, C. J. C., Penven, P., & Biastoch, A. (2013). Agulhas Leakage
23
241090 Predominantly Responds to the Southern Hemisphere Westerlies. *Journal of Physical Oceanography*,
25
261091 43(10), 2113- 2131. <https://doi.org/10.1175/JPO-D-13-047.1>
27
28
29
301092 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016).
31
321093 Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and
33
341094 organization. *Geoscientific Model Development*, 9(5), 1937- 1958. [https://doi.org/10.5194/gmd-9-](https://doi.org/10.5194/gmd-9-1937-2016)
35
361095 [1937-2016](https://doi.org/10.5194/gmd-9-1937-2016)
37
38
39
401096 Farneti, R., Delworth, T. L., Rosati, A. J., Griffies, S. M., & Zeng, F. (2010). The Role of Mesoscale
41
421097 Eddies in the Rectification of the Southern Ocean Response to Climate Change. *Journal of Physical*
43
441098 *Oceanography*, 40(7), 1539- 1557. <https://doi.org/10.1175/2010JPO4353.1>
45
46
47
481099 Forster, P., T. Storelvmo, K. Armour, W. Collins, J.-L. Dufresne, D. Frame, D.J. Lunt, T. Mauritsen,
49
501100 M.D. Palmer, M. Watanabe, M. Wild, and H. Zhang (2021). The Earth's Energy Budget, Climate
51
521101 Feedbacks, and Climate Sensitivity. In *Climate Change 2021: The Physical Science Basis*. Contribution
53
541102 of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change
55
561103 [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L.
57
58
591104 Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T.
60
61
62
63
64
65

- 1105 Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United
1
21106 Kingdom and New York, NY, USA, pp. 923–1054, doi:10.1017/9781009157896.009
3
4
51107 Fox-Kemper, B., H.T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S.S. Drijfhout, T.L. Edwards, N.R. Golledge,
6
71108 M. Hemer, R.E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I.S. Nurhati, L. Ruiz, J.-B. Sallée,
8
9
101109 A.B.A. Slangen, and Y. Yu (2021). Ocean, Cryosphere and Sea Level Change. In Climate Change,
11
121110 2021. The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of
13
141111 the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L.
15
161112 Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E.
17
181113 Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)].
19
20
211114 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1211–1362,
22
231115 doi:10.1017/9781009157896.011
24
25
261116 Frölicher, T. L., Sarmiento, J. L., Paynter, D. J., Dunne, J. P., Krasting, J. P., & Winton, M. (2015).
27
281117 Dominance of the Southern Ocean in Anthropogenic Carbon and Heat Uptake in CMIP5 Models.
29
30
311118 Journal of Climate, 28(2), 862- 886. <https://doi.org/10.1175/JCLI-D-14-00117.1>
32
33
341119 Frölicher, T. L., Fischer, E. M., & Gruber, N. (2018). Marine heatwaves under global warming. Nature,
35
361120 560(7718), 360- 364. <https://doi.org/10.1038/s41586-018-0383-9>
37
38
39
401121 Fromant, A., Delord, K., Bost, C.-A., Eizenberg, Y. H., Botha, J. A., Cherel, Y., Bustamante, P.,
41
421122 Gardner, B. R., Brault-Favrou, M., Lec’hvien, A., & Arnould, J. P. Y. (2021). Impact of extreme
43
441123 environmental conditions : Foraging behaviour and trophic ecology responses of a diving seabird, the
45
461124 common diving petrel. Progress in Oceanography, 198, 102676.
47
48
491125 <https://doi.org/10.1016/j.pocean.2021.102676>
50
51
521126 García Molinos, J., Halpern, B., Schoeman, D. et al. (2016). Climate velocity and the future global
53
541127 redistribution of marine biodiversity. Nature Clim Change 6, 83–88.
55
561128 <https://doi.org/10.1038/nclimate2769>
57
58
59
60
61
62
63
64
65

- 1129 Good, S., Fiedler, E., Mao, C., Martin, M. J., Maycock, A., Reid, R., Roberts-Jones, J., Searle, T.,
1
21130 Waters, J., While, J., & Worsfold, M. (2020). The Current Configuration of the OSTIA System for
3
41131 Operational Production of Foundation Sea Surface Temperature and Ice Concentration Analyses.
5
61132 Remote Sensing, 12(4), 720. <https://doi.org/10.3390/rs12040720>
7
8
9
101133 Gopika, S., Izumo, T., Vialard, J., Lengaigne, M., Suresh, I., & Kumar, M. R. R. (2020). Aliasing of
11
121134 the Indian Ocean externally-forced warming spatial pattern by internal climate variability. Climate
13
141135 Dynamics, 54(1- 2), 1093- 1111. <https://doi.org/10.1007/s00382-019-05049-9>
15
16
171136 Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A.,
18
191137 Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., & Kawamiya, M. (2020).
20
211138 Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes
22
231139 and feedbacks. Geoscientific Model Development, 13(5), 2197- 2244. [https://doi.org/10.5194/gmd-13-](https://doi.org/10.5194/gmd-13-2197-2020)
24
25
261140 2197-2020
27
28
29
301141 Haumann, F. A., Gruber, N., & Münnich, M. (2020). Sea-Ice Induced Southern Ocean Subsurface
31
321142 Warming and Surface Cooling in a Warming Climate. AGU Advances, 1(2), e2019AV000132.
33
341143 <https://doi.org/10.1029/2019AV000132>
35
36
371144 Hausfather, Z., & Peters, G. P. (2020). Emissions – the ‘business as usual’ story is misleading. Nature,
38
391145 577(7792), 618- 620. <https://doi.org/10.1038/d41586-020-00177-3>
40
41
42
431146 Hayashida, H., Matear, R. J., Strutton, P. G., & Zhang, X. (2020). Insights into projected changes in
44
451147 marine heatwaves from a high-resolution ocean circulation model. Nature Communications, 11(1),
46
471148 4352. <https://doi.org/10.1038/s41467-020-18241-x>
48
49
50
511149 Held, I. M., Guo, H., Adcroft, A., Dunne, J. P., Horowitz, L. W., Krasting, J., Shevliakova, E., Winton,
52
531150 M., Zhao, M., Bushuk, M., Wittenberg, A. T., Wyman, B., Xiang, B., Zhang, R., Anderson, W., Balaji,
54
551151 V., Donner, L., Dunne, K., Durachta, J., ... Zadeh, N. (2019). Structure and Performance of GFDL’s
56
571152 CM4.0 Climate Model. Journal of Advances in Modeling Earth Systems, 11(11), 3691- 3727.
58
591153 <https://doi.org/10.1029/2019MS001829>
60
61
62
63
64
65

1154 Herger, N., Sanderson, B. M., and Knutti, R. (2015), Improved pattern scaling approaches for the use
1 in climate impact studies. *Geophys. Res. Lett.*, 42, 3486– 3494. doi: 10.1002/2015GL063569.
21155
3
4
51156 Hewitt, H., Fox-Kemper, B., Pearson, B., Roberts, M., and Klocke, D. (2022). The small scales of the
6
7 ocean may hold the key to surprises. *Nature Climate Change*, 12, 496–499. doi: 10.1038/s41558-022-
81157 01386-6
9
101158
11
12
131159 Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C. J., Benthuisen,
14
151160 J. A., Burrows, M. T., Donat, M. G., Feng, M., Holbrook, N. J., Moore, P. J., Scannell, H. A., Sen
16
171161 Gupta, A., & Wernberg, T. (2016). A hierarchical approach to defining marine heatwaves. *Progress in*
18
19 *Oceanography*, 141, 227- 238. <https://doi.org/10.1016/j.pocean.2015.12.014>
201162
21
22
231163 Hogg, A. McC., Meredith, M. P., Chambers, D. P., Abrahamsen, E. P., Hughes, C. W., & Morrison, A.
24
251164 K. (2015). Recent trends in the Southern Ocean eddy field. *Journal of Geophysical Research: Oceans*,
26
271165 120(1), 257- 267. <https://doi.org/10.1002/2014JC010470>
28
29
30
311166 Holbrook, N. J., Sen Gupta, A., Oliver, E. C. J., Hobday, A. J., Benthuisen, J. A., Scannell, H. A.,
32
331167 Smale, D. A., & Wernberg, T. (2020). Keeping pace with marine heatwaves. *Nature Reviews Earth &*
34
351168 *Environment*, 1(9), 482- 493. <https://doi.org/10.1038/s43017-020-0068-4>
36
37
38
391169 Huang, B., & Shukla, J. (2008). Interannual variability of the South Indian Ocean in observations and
40
411170 a coupled model. *INDIAN J. MAR. SCI.*, 37(1), 23.
42
43
441171 Huguenin, M.F., Holmes, R.M. & England, M.H. (2022). Drivers and distribution of global ocean heat
45
461172 uptake over the last half century. *Nat Commun* 13, 4921. <https://doi.org/10.1038/s41467-022-32540-5>
47
48
49
501173 Hunt, B. P. V., Pakhomov, E. A., and McQuaid, C. D. (2001). Short-term variation and long-term
51
521174 changes in the oceanographic environment and zooplankton community in the vicinity of a sub-
53
541175 Antarctic archipelago. *Marine Biology*, 138, 369–381. <https://doi.org/10.1007/s002270000467>
55
56
571176 IPCC (2022). Summary for Policymakers. In: *Climate Change 2022: Mitigation of Climate Change.*
58
591177 *Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on*
60
61
62
63
64
65

- 1178 Climate Change [P.R. Shukla, J. Skea, R. Slade, A. Al Khourdajie, R. van Diemen, D. McCollum, M.
1
21179 Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley, (eds.)].
3
41180 Cambridge University Press, Cambridge, UK and New York, NY, USA. doi:
5
61181 10.1017/9781009157926.001
7
8
9
101182 Jones, K. R., Watson, J. E. M., Possingham, H. P., & Klein, C. J. (2016). Incorporating climate change
11
121183 into spatial conservation prioritisation: A review. *Biological Conservation*, 194, 121- 130.
13
141184 <https://doi.org/10.1016/j.biocon.2015.12.008>
15
16
171185 Kock, K.-H., & Everson, I. (2003). Shedding new light on the life cycle of mackerel icefish in the
18
191186 Southern Ocean. *Journal of Fish Biology*, 63(1), 1- 21. <https://doi.org/10.1046/j.1095->
20
21
221187 8649.2003.00150.x
23
24
251188 Kostianoy, A. G., Ginzburg, A. I., Frankignoulle, M., & Delille, B. (2004). Fronts in the Southern Indian
26
271189 Ocean as inferred from satellite sea surface temperature data. *Journal of Marine Systems*, 45(1), 55- 73.
28
29
301190 <https://doi.org/10.1016/j.jmarsys.2003.09.004>
31
32
331191 Kravtsov, S., Grimm, C., & Gu, S. (2018). Global-scale multidecadal variability missing in state-of-
34
351192 the-art climate models. *Npj Climate and Atmospheric Science*, 1(1), 1- 10.
36
371193 <https://doi.org/10.1038/s41612-018-0044-6>
38
39
40
411194 Kuhlbrodt, T., Jones, C. G., Sellar, A., Storkey, D., Blockley, E., Stringer, M., Hill, R., Graham, T.,
42
431195 Ridley, J., Blaker, A., Calvert, D., Copsey, D., Ellis, R., Hewitt, H., Hyder, P., Ineson, S., Mulcahy, J.,
44
451196 Siahaan, A., & Walton, J. (2018). The Low-Resolution Version of HadGEM3 GC3.1 : Development
46
471197 and Evaluation for Global Climate. *Journal of Advances in Modeling Earth Systems*, 10(11),
48
491198 2865- 2888. <https://doi.org/10.1029/2018MS001370>
50
51
52
531199 Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P.,
54
551200 Gehlen, M., Ilyina, T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-
56
571201 Falcón, Y., Schwinger, J., Séférian, R., Stock, C. A., ... Ziehn, T. (2020). Twenty-first century ocean
58
591202 warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from
60
61
62
63
64
65

- 1203 CMIP6 model projections. *Biogeosciences*, 17(13), 3439- 3470. <https://doi.org/10.5194/bg-17-3439->
1
21204 2020
3
4
51205 Le Bars, D., Viebahn, J. P., & Dijkstra, H. A. (2016). A Southern Ocean mode of multidecadal
6
71206 variability. *Geophysical Research Letters*, 43(5), 2102- 2110. <https://doi.org/10.1002/2016GL068177>
8
9
10
111207 Lee, J.-Y., J. Marotzke, G. Bala, L. Cao, S. Corti, J.P. Dunne, F. Engelbrecht, E. Fischer, J.C. Fyfe, C.
12
131208 Jones, A. Maycock, J. Mutemi, O. Ndiaye, S. Panickal, and T. Zhou (2021). Future Global Climate:
14
151209 Scenario-Based Projections and NearTerm Information. In *Climate Change 2021: The Physical Science*
16
171210 Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel
18
191211 on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N.
20
211212 Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K.
22
231213 Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press,
24
251214 Cambridge, United Kingdom and New York, NY, USA, pp. 553–672,
26
271215 doi:10.1017/9781009157896.006.
28
29
30
31
321216 Lin, X., Zhai, X., Wang, Z., & Munday, D. R. (2018). Mean, Variability, and Trend of Southern Ocean
33
341217 Wind Stress: Role of Wind Fluctuations. *Journal of Climate*, 31(9), 3557- 3573.
35
361218 <https://doi.org/10.1175/JCLI-D-17-0481.1>
37
38
39
401219 Llovel, W., & Terray, L. (2016). Observed southern upper-ocean warming over 2005–2014 and
41
421220 associated mechanisms. *Environmental Research Letters*, 11(12), 124023.
43
441221 <https://doi.org/10.1088/1748-9326/11/12/124023>
45
46
47
481222 Loarie, S. R., Duffy, P. B., Hamilton, H., Asner, G. P., Field, C. B., & Ackerly, D. D. (2009). The
49
501223 velocity of climate change. *Nature*, 462(7276), 1052- 1055. <https://doi.org/10.1038/nature08649>
51
52
531224 Lovato, T., Peano, D., Butenschön, M., Materia, S., Iovino, D., Scoccimarro, E., Fogli, P. G., Cherchi,
54
551225 A., Bellucci, A., Gualdi, S., Masina, S., & Navarra, A. (2022). CMIP6 Simulations With the CMCC
56
571226 Earth System Model (CMCC-ESM2). *Journal of Advances in Modeling Earth Systems*, 14(3),
58
591227 e2021MS002814. <https://doi.org/10.1029/2021MS002814>
60
61
62
63
64
65

- 1228 Lutjeharms, J. R. E. (Éd.). (2006). The Agulhas return flow. In *The Agulhas Current* (p. 209- 231).
1
21229 Springer. https://doi.org/10.1007/3-540-37212-1_7
3
4
- 51230 Ma, J., Feng, M., Lan, J. and Hu, D., 2020. Projected future changes of meridional heat transport and
6
71231 heat balance of the Indian Ocean. *Geophysical Research Letters*, 47(4), p.e2019GL086803
8
9
- 10
111232 Makhado, A.B., Lowther, A., Koubbi, P., Ansorge, I., Brooks,C., Cotté, C., Crawford, R., Dlulisa, S.,
12
131233 d'Ovidio, F., Fawcet, S., Freeman, D., Grant, S., Huggett, J., Hindell, M., Hulley,P.A., Kirkman, S.,
14
151234 Lamont, T., Lombard, M., Masothla, M.J., Lea, M.-A., Oosthuizen,W.C., Orgeret, F., Reisinger, R.,
16
171235 Samaai, T., Sergi, S., Swadling, K., Somhlaba, S., Van de Putte, A., Von de Meden, C., and Yemane,
18
19
201236 D. (2019). Expert Workshop on Pelagic Spatial Planning for the eastern subantarctic region (Domains
21
221237 4, 5 and 6). SC-CAMLR-38/BG/29.
23
24
- 251238 Martínez-Moreno, J., Hogg, A. McC., Kiss, A. E., Constantinou, N. C., & Morrison, A. K. (2019).
26
271239 Kinetic Energy of Eddy-Like Features From Sea Surface Altimetry. *Journal of Advances in Modeling*
28
29
301240 Earth Systems, 11(10), 3090- 3105. <https://doi.org/10.1029/2019MS001769>
31
32
- 331241 Martínez-Moreno, J., Hogg, A. M., England, M. H., Constantinou, N. C., Kiss, A. E., & Morrison, A.
34
351242 K. (2021). Global changes in oceanic mesoscale currents over the satellite altimetry record. *Nature*
36
371243 *Climate Change*, 11(5), 397- 403. <https://doi.org/10.1038/s41558-021-01006-9>
38
39
- 40
411244 Matsuno, K., Wallis, J. R., Kawaguchi, S., Bestley, S. and Swadling, K. M. (2020). Zooplankton
42
431245 community structure and dominant copepod population structure on the southern Kerguelen Plateau
44
451246 during summer 2016. *Deep-Sea Research II*, 174, 104788. <https://doi.org/10.1016/j.dsr2.2020.104788>
46
47
- 48
491247 Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M.,
50
511248 Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak,
52
531249 H., Hagemann, S., Hedemann, C., ... Roeckner, E. (2019). Developments in the MPI-M Earth System
54
551250 Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO2. *Journal of Advances in*
56
571251 *Modeling Earth Systems*, 11(4), 998- 1038. <https://doi.org/10.1029/2018MS001400>
58
59
60
61
62
63
64
65

- 1252 Maxwell, S. M., Gjerde, K. M., Conners, M. G., & Crowder, L. B. (2020). Mobile protected areas for
1 biodiversity on the high seas. *Science*, 367(6475), 252- 254. <https://doi.org/10.1126/science.aaz9327>
2
3
4
- 51254 Meredith, M., M. Sommerkorn, S. Cassotta, C. Derksen, A. Ekaykin, A. Hollowed, G. Kofinas, A.
6
7
81255 Mackintosh, J. Melbourne-Thomas, M.M.C. Muelbert, G. Ottersen, H. Pritchard, and E.A.G. Schuur
9
101256 (2019). Polar Regions. In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate
11
121257 [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K.
13
141258 Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. Cambridge
15
161259 University Press, Cambridge, UK and New York, NY, USA, pp. 203–320.
17
18
191260 <https://doi.org/10.1017/9781009157964.005>.
20
21
- 221261 Mills, K. E., Pershing, A. J., Brown C. J., Chen, Y., Chiang, F.-S., Holland, D. S., Lehuta, S., Nye, J.
23
241262 A., Sun, J. C., Thomas, A. C., & Wahle, R. A. (2013). Fisheries Management in a Changing Climate :
25
261263 Lessons from the 2012 Ocean Heat Wave in the Northwest Atlantic. *Oceanography*, 26(2), 191- 195.
27
28
29
- 301264 Mishra, R. K., Naik, R. K., Venkataramana, V., Jena, B., AnilKumar, N., Soares, M. A., Sarkar, A. and
31
321265 Singh, A. (2020). Phytoplankton biomass and community composition in the front zones of Southern
33
341266 Ocean. *Deep-Sea Research II*, 178, 104799. <https://doi.org/10.1016/j.dsr2.2020.104799>
35
36
- 371267 Morrison, A. K., Griffies, S. M., Winton, M., Anderson, W. G., & Sarmiento, J. L. (2016). Mechanisms
38
391268 of Southern Ocean Heat Uptake and Transport in a Global Eddy Climate Model. *Journal of Climate*,
40
41
421269 29(6), 2059- 2075. <https://doi.org/10.1175/JCLI-D-15-0579.1>
43
44
- 451270 Müller, W. A., Jungclaus, J. H., Mauritsen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M.,
46
471271 Ghosh, R., Haak, H., Ilyina, T., Kleine, T., Kornblueh, L., Li, H., Modali, K., Notz, D., Pohlmann, H.,
48
49
501272 Roeckner, E., Stemmler, I., ... Marotzke, J. (2018). A Higher-resolution Version of the Max Planck
51
521273 Institute Earth System Model (MPI-ESM1.2-HR). *Journal of Advances in Modeling Earth Systems*,
53
541274 10(7), 1383- 1413. <https://doi.org/10.1029/2017MS001217>
55
56
- 571275 Oliver, E. C. J. (2019a). Mean warming not variability drives marine heatwave trends. *Climate*
58
59
601276 Dynamics, 53(3), 1653- 1659. <https://doi.org/10.1007/s00382-019-04707-2>
61
62
63
64
65

- 1277 Oliver, E. C. J., Burrows, M. T., Donat, M. G., Sen Gupta, A., Alexander, L. V., Perkins-Kirkpatrick,
1
21278 S. E., Benthuyzen, J. A., Hobday, A. J., Holbrook, N. J., Moore, P. J., Thomsen, M. S., Wernberg, T.,
3
41279 & Smale, D. A. (2019b). Projected Marine Heatwaves in the 21st Century and the Potential for
5
61280 Ecological Impact. *Frontiers in Marine Science*, 6, 734. <https://doi.org/10.3389/fmars.2019.00734>
7
8
9
101281 O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R.,
11
121282 Kriegler, E., Lamarque, J.-F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., & Sanderson, B. M. (2016).
13
141283 The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geoscientific Model*
15
161284 *Development*, 9(9), 3461- 3482. <https://doi.org/10.5194/gmd-9-3461-2016>
17
18
19
201285 O'Neill, B. C., Kriegler, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B.
21
221286 J., van Vuuren, D. P., Birkmann, J., Kok, K., Levy, M., & Solecki, W. (2017). The roads ahead :
23
241287 Narratives for shared socioeconomic pathways describing world futures in the 21st century. *Global*
25
261288 *Environmental Change*, 42, 169- 180. <https://doi.org/10.1016/j.gloenvcha.2015.01.004>
27
28
29
301289 Ortuño Crespo, G., Mossop, J., Dunn, D., Gjerde, K., Hazen, E., Reygondeau, G., Warner, R., Tittensor,
31
321290 D., & Halpin, P. (2020). Beyond static spatial management : Scientific and legal considerations for
33
341291 dynamic management in the high seas. *Marine Policy*, 122, 104102.
35
361292 <https://doi.org/10.1016/j.marpol.2020.104102>
37
38
39
401293 Park Y.-H., Gambéroni, L., and Charriaud, E. (1991). Frontal structure and transport of the Antarctic
41
421294 Circumpolar Current in the south Indian Ocean sector, 40-80°E. *Marine Chemistry*, 35, 45-62.
43
441295 [https://doi.org/10.1016/S0304-4203\(09\)90007-X](https://doi.org/10.1016/S0304-4203(09)90007-X)
45
46
47
481296 Park, Y.-H., Gamberoni, L., and Charriaud, E. (1993). Frontal structure, water masses, and circulation
49
501297 in the Crozet Basin. *J. Geophys. Res.*, 98(C7), 12361– 12385. doi:10.1029/93JC00938
51
52
531298 Park, Y.-H., Roquet, F., Durand, I. and Fuda, J.-L. (2008). Large-scale circulation over and around the
54
551299 Northern Kerguelen Plateau. *Deep Sea Research II*, 55(5-7), 566-581.
56
571300 <https://doi.org/10.1016/j.dsr2.2007.12.030>
58
59
60
61
62
63
64
65

- 1301 Park, Y.-H., Vivier, F., Roquet, F., and Kestenare, E. (2009). Direct observations of the ACC transport
1
21302 across the Kerguelen Plateau. *Geophys. Res. Lett.*, 36, L18603. doi:10.1029/2009GL039617.
3
4
- 51303 Park Y.-H., and Durand, I. (2019). Altimetry-derived Antarctic Circumpolar Current Fronts. *SEANOE*.
6
71304 <https://doi.org/10.17882/59800>
8
9
- 10
111305 Park Y.- H., Park T., Kim T.- W., Lee S.- H., Hong C.- S., Lee J.- H., Rio M.- H., Pujol M.- I.,
12
131306 Ballarotta M., Durand I., Provost C. (2019). Observations of the Antarctic Circumpolar Current over
14
151307 the Udintsev Fracture Zone, the narrowest choke point in the Southern Ocean. *Journal of Geophysical*
16
171308 *Research: Oceans*, 124, 4511-4528. <https://doi.org/10.1029/2019JC015024>
18
19
- 20
211309 Patara, L., Böning, C. W., & Biastoch, A. (2016). Variability and trends in Southern Ocean eddy activity
22
231310 in 1/12° ocean model simulations. *Geophysical Research Letters*, 43(9), 4517- 4523.
24
251311 <https://doi.org/10.1002/2016GL069026>
26
27
- 28
291312 Pietri, A., Colas, F., Mogollon, R., Tam, J., & Gutierrez, D. (2021). Marine heatwaves in the Humboldt
30
311313 current system : From 5-day localized warming to year-long El Niños. *Scientific Reports*, 11(1), 1- 12.
32
331314 <https://doi.org/10.1038/s41598-021-00340-4>
34
35
- 361315 Pilo, G. S., Holbrook, N. J., Kiss, A. E., & Hogg, A. M. (2019). Sensitivity of Marine Heatwave Metrics
37
381316 to Ocean Model Resolution. *Geophysical Research Letters*, 46(24), 14604- 14612.
39
40
411317 <https://doi.org/10.1029/2019GL084928>
42
43
- 441318 Plecha, S. M., & Soares, P. M. M. (2020). Global marine heatwave events using the new CMIP6 multi-
45
461319 model ensemble : From shortcomings in present climate to future projections. *Environmental Research*
47
481320 *Letters*, 15(12). <https://doi.org/10.1088/1748-9326/abc847>
49
50
- 51
521321 Pütz, K. (2002). Spatial and Temporal Variability in the Foraging Areas of Breeding King Penguins.
53
541322 *The Condor*, 104(3), 528- 538. <https://doi.org/10.1093/condor/104.3.528>
55
56
57
58
59
60
61
62
63
64
65

- 1323 Qiu, Z., Qiao, F., Jang, C. J., Zhang, L., & Song, Z. (2021). Evaluation and projection of global marine
1 heatwaves based on CMIP6 models. *Deep Sea Research Part II: Topical Studies in Oceanography*, 194,
2 1324 104998. <https://doi.org/10.1016/j.dsr2.2021.104998>
3
4 1325
5
6
7
8 1326 Raäisaänen, J. (2007). How reliable are climate models? *Tellus A: Dynamic Meteorology and*
9
10 1327 *Oceanography*, 59(1), 2- 29. <https://doi.org/10.1111/j.1600-0870.2006.00211.x>
11
12
13 1328 Reisinger, R. R., Corney, S., Raymond, B., Lombard, A. T., Bester, M. N., Crawford, R. J. M., Davies,
14
15 1329 D., de Bruyn, P. J. N., Dilley, B. J., Kirkman, S. P., Makhado, A. B., Ryan, P. G., Schoombie, S.,
16
17 1330 Stevens, K. L., Tosh, C. A., Wege, M., Whitehead, T. O., Sumner, M. D., Wotherspoon, S., ... Pistorius,
18
19 P. A. (2022). Habitat model forecasts suggest potential redistribution of marine predators in the southern
20 1331 Indian Ocean. *Diversity and Distributions*, 28(1), 142- 159. <https://doi.org/10.1111/ddi.13447>
21
22 1332
23
24
25 1333 Reygondeau, G., Cheung, W. W. L., Wabnitz, C. C. C., Lam, V. W. Y., Frölicher, T., & Maury, O.
26
27 1334 (2020). Climate Change-Induced Emergence of Novel Biogeochemical Provinces. *Frontiers in Marine*
28
29 *Science*, 7, 657. <https://doi.org/10.3389/fmars.2020.00657>
30 1335
31
32
33 1336 Reynolds, Richard W., Thomas M. Smith, Chunying Liu, Dudley B. Chelton, Kenneth S. Casey,
34
35 1337 Michael G. Schlax (2007). Daily High-Resolution-Blended Analyses for Sea Surface Temperature. *J.*
36
37 1338 *Climate*, 20, 5473-5496. Reynolds, Richard W., Thomas M. Smith, Chunying Liu, Dudley B. Chelton,
38
39 1339 Kenneth S. Casey, Michael G. Schlax, 2007: Daily High-Resolution-Blended Analyses for Sea Surface
40
41 1340 Temperature. *J. Climate*, 20, 5473-5496.
42
43
44
45 1341 Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin,
46
47 1342 K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., Kc, S., Leimbach, M., Jiang, L., Kram,
48
49 1343 T., Rao, S., Emmerling, J., ... Tavoni, M. (2017). The Shared Socioeconomic Pathways and their
50
51 energy, land use, and greenhouse gas emissions implications : An overview. *Global Environmental*
52 1344 *Change*, 42, 153- 168. <https://doi.org/10.1016/j.gloenvcha.2016.05.009>
53
54 1345
55
56
57 1346 Rintoul, S. R. and Naveira Garabato, A.C. (2013). Chapter 18 - Dynamics of the Southern Ocean
58
59 1347 circulation. In G. Siedler, S. M. Griffies, J. Gould, & J. A. Church (Eds.), *International geophysics*,
60
61
62
63
64
65

- 1348 ocean circulation and climate (Vol. 103, pp. 471– 492). Waltham, MA: Academic Press.
1
21349 <https://doi.org/10.1016/B978-0-12-391851-2.00018-0>
3
4
- 51350 Roemmich, D., Church, J., Gilson, J., Monselesan, D., Sutton, P., & Wijffels, S. (2015). Unabated
6
71351 planetary warming and its ocean structure since 2006. *Nature Climate Change*, 5(3), 240- 245.
8
9
101352 <https://doi.org/10.1038/nclimate2513>
11
12
- 131353 Rong, X.-Y., Li, J., Chen, H.-M., Xin, Y.-F., Su, J.-Z., Hua, L.-J., & Zhang, Z.-Q. (2019). Introduction
14
151354 of CAMS-CSM model and its participation in CMIP6. *Advances in Climate Change Research*, 15(5),
16
171355 540. <https://doi.org/10.12006/j.issn.1673-1719.2019.186>
18
19
- 20
211356 Roquet, F., Park, Y.-H., Guinet, C., Bailleul, F., and Charassin, J.-B. (2009). Observations of the Fawn
22
231357 Trough Current over the Kerguelen Plateau from instrumented elephant seals. *Journal of Marine*
24
251358 *Systems*, 78, 377-393. doi:10.1016/j.jmarsys.2008.11.017
26
27
- 28
291359 Saenko, O. A., Yang, D., & Gregory, J. M. (2018). Impact of Mesoscale Eddy Transfer on Heat Uptake
30
311360 in an Eddy-Parameterizing Ocean Model. *Journal of Climate*, 31(20), 8589- 8606.
32
331361 <https://doi.org/10.1175/JCLI-D-18-0186.1>
34
35
- 361362 Sallée, J. B., Speer, K. G., & Rintoul, S. R. (2010). Zonally asymmetric response of the Southern Ocean
37
381363 mixed-layer depth to the Southern Annular Mode. *Nature Geoscience*, 3(4), 273- 279.
39
40
411364 <https://doi.org/10.1038/ngeo812>
42
43
- 441365 Sallée, J.-B. (2018). Southern Ocean Warming. *Oceanography*, 31(2), 52- 62. JSTOR.
45
46
- 471366 Scannell, H. A., Pershing, A. J., Alexander, M. A., Thomas, A. C., & Mills, K. E. (2016). Frequency of
48
491367 marine heatwaves in the North Atlantic and North Pacific since 1950. *Geophysical Research Letters*,
50
51
521368 43(5), 2069- 2076. <https://doi.org/10.1002/2015GL067308>
53
54
- 551369 Schlegel, R. W., Oliver, E. C. J., Hobday, A. J., & Smit, A. J. (2019). Detecting Marine Heatwaves
56
571370 With Sub-Optimal Data. *Frontiers in Marine Science*, 6, 737. <https://doi.org/10.3389/fmars.2019.00737>
58
59
60
61
62
63
64
65

- 1371 Schmidt, C., Schwarzkopf, F. U., Rühls, S., & Biastoch, A. (2021). Characteristics and robustness of
1
21372 Agulhas leakage estimates : An inter-comparison study of Lagrangian methods. *Ocean Science*, 17(4),
3
41373 1067- 1080. <https://doi.org/10.5194/os-17-1067-2021>
5
6
7
81374 Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C.,
9
101375 Berthet, S., Chevallier, M., Sénési, S., Franchistéguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy,
11
121376 O., Guérémy, J.-F., Moine, M.-P., M'Sadek, R., Ribes, A., Rocher, M., Roehrig, R., Salas-y-Mélia, D.,
13
141377 Sanchez, E., Terray, L., Valcke, S., Waldman, R., Aumont, O., Bopp, L., Deshayes, J., Éthé, C., and
15
161378 Madec, G. (2019). Evaluation of CNRM Earth System model, CNRM-ESM2-1: role of Earth system
17
181379 processes in present-day and future climate. *J. Adv. Model. Earth Syst.*, 11, 4182–4227.
19
20
211380 <https://doi.org/10.1029/2019MS001791>
22
23
241381 Sellar, A. A., Jones, C. G., Mulcahy, J., Tang, Y., Yool, A., Wiltshire, A., O'connor, F. M., Stringer,
25
261382 M., Hill, R., Palmieri, J., Woodward, S., Mora, L., Kuhlbrodt, T., Rumbold, S., Kelley, D. I., Ellis, R.,
27
281383 Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou,
29
301384 S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths,
31
32
331385 P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J.,
34
351386 Liddi coat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahann, A., Smith, R. S.,
36
371387 Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and
38
391388 evaluation of the UK Earth System Model (2019). *J. Adv. Model. Earth Syst.*, 11, 4513–4558,
40
41
421389 <https://doi.org/10.1029/2019MS001739>
43
44
451390 Sen Gupta, A., Brown, J. N., Jourdain, N. C., van Sebille, E., Ganachaud, A., & Vergés, A. (2015).
46
471391 Episodic and non-uniform shifts of thermal habitats in a warming ocean. *Deep Sea Research Part II:*
48
491392 *Topical Studies in Oceanography*, 113, 59- 72. <https://doi.org/10.1016/j.dsr2.2013.12.002>
50
51
52
531393 Sen Gupta, A., Stellema, A., Pontes, G. M., Taschetto, A. S., Vergés, A., & Rossi, V. (2021). Future
54
551394 changes to the upper ocean Western Boundary Currents across two generations of climate models.
56
571395 *Scientific Reports*, 11(1), 9538. <https://doi.org/10.1038/s41598-021-88934-w>
58
59
60
61
62
63
64
65

- 1396 Siegelman, L., Klein, P., Rivière, P., Thompson, A. F., Torres, H. S., Flexas, M., & Menemenlis, D.
1
21397 (2020). Enhanced upward heat transport at deep submesoscale ocean fronts. *Nature Geoscience*, 13(1),
3
41398 50- 55. <https://doi.org/10.1038/s41561-019-0489-1>
5
6
7
81399 Smale, D. A., & Wernberg, T. (2013). Extreme climatic event drives range contraction of a habitat-
9
101400 forming species. *Proceedings of the Royal Society B: Biological Sciences*, 280(1754), 20122829.
11
121401 <https://doi.org/10.1098/rspb.2012.2829>
13
14
151402 Smale DA, Wernberg T, & Vanderklift MA. (2017). Regional-scale variability in the response of
16
171403 benthic macroinvertebrate assemblages to a marine heatwave. *Marine Ecology Progress Series*, 568,
18
191404 17- 30.
20
21
22
231405 Stellema, A., Sen Gupta, A., & Taschetto, A. S. (2019). Projected slow down of South Indian Ocean
24
251406 circulation. *Scientific Reports*, 9(1), 17705. <https://doi.org/10.1038/s41598-019-54092-3>
26
27
28
291407 Su, Z., Pilo, G. S., Corney, S., Holbrook, N. J., Mori, M., & Ziegler, P. (2021). Characterizing Marine
30
311408 Heatwaves in the Kerguelen Plateau Region. *Frontiers in Marine Science*, 7, 1119.
32
331409 <https://doi.org/10.3389/fmars.2020.531297>
34
35
361410 Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora,
37
381411 V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C.,
39
40
411412 Shao, A., Sigmond, M., Solheim, L., ... Winter, B. (2019). The Canadian Earth System Model version
42
431413 5 (CanESM5.0.3). *Geoscientific Model Development*, 12(11), 4823- 4873.
44
451414 <https://doi.org/10.5194/gmd-12-4823-2019>
46
47
48
491415 Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe,
50
511416 M., Saito, F., Chikira, M., Watanabe, S., Mori, M., Hirota, N., Kawatani, Y., Mochizuki, T., Yoshimura,
52
531417 K., Takata, K., O'ishi, R., ... Kimoto, M. (2019). Description and basic evaluation of simulated mean
54
551418 state, internal variability, and climate sensitivity in MIROC6. *Geoscientific Model Development*, 12(7),
56
571419 2727- 2765. <https://doi.org/10.5194/gmd-12-2727-2019>
58
59
60
61
62
63
64
65

- 1420 Tittensor, D. P., Beger, M., Boerder, K., Boyce, D. G., Cavanagh, R. D., Cosandey-Godin, A., Crespo,
1
21421 G. O., Dunn, D. C., Ghiffary, W., Grant, S. M., Hannah, L., Halpin, P. N., Harfoot, M., Heaslip, S. G.,
3
41422 Jeffery, N. W., Kingston, N., Lotze, H. K., McGowan, J., McLeod, E., ... Worm, B. (2019). Integrating
5
61423 climate adaptation and biodiversity conservation in the global ocean. *Science Advances*.
7
8
91424 <https://doi.org/10.1126/sciadv.aay9969>
- 10
11
121425 van Wijk, E. M., Rintoul, S. R., Ronai, B. M. and Williams, G. D. (2010). Regional circulation around
13
141426 Heard and McDonald Islands and through the Fawn Trough, central Kerguelen Plateau. *Deep-Sea*
15
161427 *Research I*, 57, 653-669. doi:10.1016/j.dsr.2010.03.001
- 17
18
19
201428 VanDerWal, J., Murphy, H. T., Kutt, A. S., Perkins, G. C., Bateman, B. L., Perry, J. J., & Reside, A. E.
21
221429 (2013). Focus on poleward shifts in species' distribution underestimates the fingerprint of climate
23
241430 change. *Nature Climate Change*, 3(3), 239- 243. <https://doi.org/10.1038/nclimate1688>
- 25
26
271431 Venkataramana, V., Anilkumar, N. Swadling, K., Mishra, R. K., Tripathy, S. C., Sarkar, A., Augusta,
28
291432 S. M., Sabu, P. and Pillai, H. (2020). Distribution of zooplankton in the Indian sector of the Southern
30
311433 Ocean. *Antarctic Science*, 32(3), 168-179. doi:10.1017/S0954102019000579
- 32
33
34
351434 Vivier, F., Park, Y-H., Sekma, H., and Le Sommer, J. (2015). Variability of the Antarctic Circumpolar
36
371435 Current transport through the Fawn Trough, Kerguelen Plateau. *Deep-Sea Research II*, 114, 12-26.
38
391436 <http://dx.doi.org/10.1016/j.dsr2.2014.01.017>
- 40
41
42
431437 Voldoire, A., Saint-Martin, D., S n si, S., Decharme, B., Alias, A., Chevallier, M., Colin, J., Gu r my,
44
451438 J.-F., Michou, M., Moine, M.-P., Nabat, P., Roehrig, R., Salas y M lia, D., S f rian, R., Valcke, S.,
46
471439 Beau, I., Belamari, S., Berthet, S., Cassou, C., ... Waldman, R. (2019). Evaluation of CMIP6 DECK
48
491440 Experiments With CNRM-CM6-1. *Journal of Advances in Modeling Earth Systems*, 11(7),
50
511441 2177- 2213. <https://doi.org/10.1029/2019MS001683>
- 52
53
54
551442 Wiens, J. A., & Bachelet, D. (2010). Matching the Multiple Scales of Conservation with the Multiple
56
571443 Scales of Climate Change. *Conservation Biology*, 24(1), 51- 62. <https://doi.org/10.1111/j.1523->
58
591444 [1739.2009.01409.x](https://doi.org/10.1111/j.1523-1739.2009.01409.x)
- 60
61
62
63
64
65


- 1445 Wilson, K. L., Tittensor, D. P., Worm, B., & Lotze, H. K. (2020). Incorporating climate change
1 adaptation into marine protected area planning. *Global Change Biology*, 26(6), 3251- 3267.
2
3
41447 <https://doi.org/10.1111/gcb.15094>
5
6
7
81448 Wu, T., Lu, Y., Fang, Y., Xin, X., Li, L., Li, W., Jie, W., Zhang, J., Liu, Y., Zhang, L., Zhang, F.,
9
101449 Zhang, Y., Wu, F., Li, J., Chu, M., Wang, Z., Shi, X., Liu, X., Wei, M., ... Liu, X. (2019). The Beijing
11
121450 Climate Center Climate System Model (BCC-CSM) : The main progress from CMIP5 to CMIP6.
13
141451 *Geoscientific Model Development*, 12(4), 1573- 1600. <https://doi.org/10.5194/gmd-12-1573-2019>
15
16
171452 Yang, C., Leonelli, F. E., Marullo, S., Artale, V., Beggs, H., Nardelli, B. B., Chin, T. M., Toma, V. D.,
18
191453 Good, S., Huang, B., Merchant, C. J., Sakurai, T., Santoleri, R., Vazquez-Cuervo, J., Zhang, H.-M., &
20
211454 Pisano, A. (2021). Sea Surface Temperature Intercomparison in the Framework of the Copernicus
22
231455 Climate Change Service (C3S). *Journal of Climate*, 34(13), 5257- 5283. [https://doi.org/10.2151/jmsj.2019- 051](https://doi.org/10.1175/JCLI-
24
25
261456 <u>D-20-0793.1</u>
27
28
29
301457 Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M.,
31
321458 Tanaka, T., Hosaka, M., Yabu, S., Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y., and
33
341459 Ishii, M. (2019). The Meteorological Research Institute Earth System Model version 2.0, MRI-ESM2.0:
35
361460 Description and basic evaluation of the physical component, <i>J. Meteorol. Soc. Jpn.</i>, 97, 931–965.
37
38
391461 <a href=)
40
41
421462 Zhang, Y., Feng, M., Du, Y., Phillips, H. E., Bindoff, N. L., & McPhaden, M. J. (2018). Strengthened
43
441463 Indonesian throughflow drives decadal warming in the Southern Indian Ocean. *Geophysical Research*
45
461464 *Letters*, 45(12), 6167- 6175. <https://doi.org/10.1029/2018GL078265>
47
48
49
501465 Ziehn, T., Chamberlain, M. A., Law, R. M., Lenton, A., Bodman, R. W., Dix, M., Stevens, L., Wang,
51
521466 Y.-P., Srbinovsky, J., Ziehn, T., Chamberlain, M. A., Law, R. M., Lenton, A., Bodman, R. W., Dix, M.,
53
541467 Stevens, L., Wang, Y.-P., & Srbinovsky, J. (2020). The Australian Earth System Model : ACCESS-
55
561468 ESM1.5. *Journal of Southern Hemisphere Earth Systems Science*, 70(1), 193- 214.
57
58
591469 <https://doi.org/10.1071/ES19035>
60
61
62
63
64
65

Appendix A: Supplementary Materials

Figure S1: List of models used for the timeshift approach analyses for both SSP 245 and SSP585.


For changes in temperature (yearly data) 22 models	For MHW intensity (daily data) 17 models
ACCESS-CM2 ACCESS-ESM1-5 BCC-CMS2-MR CAMS-CSM1-0 CanESM5 CanESM5-CanOE CESM2-WACCM CMCC-ESM2 CMCC-CM2-SR5 CNRM-CM6-1 GFDL-CM4 GFDL-ESM4 HadGEM3-GC31-LL EC-Earth3-CC EC-Earth3-Veg IPSL-CM6A-LR MIROC-ES2L MPI-ESM1-HR MPI-ESM1-LR MRI-ESM2-0 NESM3 UKESM1-0-LL	ACCESS-CM2 ACCESS-ESM1-5 BCC-CMS2-MR CanESM5 CESM2-WACCM CMCC-ESM2 CMCC-CM2-SR5 CNRM-CM6-1 CNRM-CM6-1-HR GFDL-CM4 EC-Earth3 IPSL-CM6A-LR MIROC6 MPI-ESM1-HR MPI-ESM1-LR MRI-ESM2-0 NESM3

Figure S2 : Temperature trends between 1982 and 2019 in the Southern Indian Ocean (top panel), using linear regression on sea surface temperature, using OISST (left) and OSTIA (right) datasets. Timeseries of sea surface temperature for two areas (bottom panel): one west of the Southern Indian Ocean, covering Prince Edward Islands and Crozet (orange) and one north of Saint-Paul and Amsterdam (green) using OISST (full) and OSTIA (dashed) datasets



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure S3: Warming rates between 1982-2019 for each of the 24 CMIP6 models studied.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure S4: MHW intensity over 1984-2014 (90th percentile) for each of the 19 CMIP6 models studied.




Figure S5 : (A) Warming trends between 1975 and 2015 of 10 IPSL-CM6A-LR members. (B) Change in surface temperature (2081-2100 minus 1995-2014) of the same 10 IPSL-CM6A-LR members under

SSP2-4.5. (C)Time series of mean Pearson correlation coefficient, as an indicator of spatial similarity,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

for each 10 IPSL-CM6A-LR members relative to r1i1p1f1 member, under SSP2-4.5.






Figure S6: Relative increase in the mean (full) or spread (hatched) of the daily temperature distribution under SSP2-4.5 relative to historical period (1995-2014) from 19 CMIP6 models for 2021-2040 (near term, NT), 2041-2060 (mid term, MT) and 2081-2100 (long term, LT).



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure S7: Relation between the change in surface temperature and MHW annual days for 20-year periods under SSP1-2.6 and SSP2-4.5 averaged over the area. Error bars correspond to intermodel variability.





Figure S8: Change in surface temperature for the near-, mid- and long-terms under SSP2-4.5 for two high resolution models: MIP-ESM1-2-HR and GFDL-CM4.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65