



HAL
open science

Arc volcano activity driven by small-scale metasomatism of the magma source

Ivan Vlastélic, N. Sainlot, Pablo Samaniego, B. Bernard, François Nauret, S. Hidalgo, Delphine Auclair, Abdelmouhcine Gannoun

► **To cite this version:**

Ivan Vlastélic, N. Sainlot, Pablo Samaniego, B. Bernard, François Nauret, et al.. Arc volcano activity driven by small-scale metasomatism of the magma source. *Nature Geoscience*, 2023, 16, pp.363-370. <10.1038/s41561-023-01143-0>. <insu-04155741>

HAL Id: insu-04155741

<https://insu.hal.science/insu-04155741v1>

Submitted on 19 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 **Arc volcano activity driven by small-scale metasomatism**
2 **of the magma source**

3
4
5 I. Vlastélic^{1*}, N. Sainlot¹, P. Samaniego^{1,2}, B. Bernard², F. Nauret¹, S. Hidalgo²,
6 D. Auclair¹, A. Gannoun¹

7
8 1 Université Clermont Auvergne, CNRS, IRD, OPGC, Laboratoire Magmas et
9 Volcans, F-63000 Clermont-Ferrand, France

10
11 2 Instituto Geofísico — Escuela Politécnica Nacional, Ladrón de Guevara E11-
12 253 y Andalucía, 6to piso ed. Ing. Civil, Quito, Ecuador

13
14 * Now at Observatoire Volcanologique et Sismologique de Guadeloupe, Institut
15 de Physique du Globe de Paris, UAR3454, Gourbeyre, Guadeloupe

16
17 Word count:

18 Main text plus introductory paragraph: 2891

19 Figure legends: 323

20 Methods: 1062

21
22 References:

23 Main text and figure legends: 53

24 Methods: 7

25
26 Figure : 4 (1.6 pages)

27 Fig. 1 (1/4 page) ; Fig. 2 (1/8 page); Fig. 3 (1/2 page) ; Fig. 4 (1/2 page) ; Fig. 5
28 (1/4 page).

29 Abstract

30

31 Arc volcanism arises from the release of fluids from descending slab, which lowers mantle
32 solidus and enables melting. Mantle metasomatism plays an important role in magma
33 production and arc volcanic activity at long spatial and temporal scales. However, the episodic
34 eruption of individual volcanoes is generally thought to be primarily regulated by the
35 evolution and recharge of crustal magma chambers, with no established link with mantle
36 processes. Using high-resolution isotope time-series we show here a link between the most
37 recent 1999-2016 eruptive activity of the Tungurahua volcano in Ecuador and the small-scale
38 metasomatic enrichment of the magma source. During the eruption the average rate of
39 magma production varies logarithmically with the degree of source enrichment. Such a
40 correlation points to a major control of source enrichment on magma supply and eruptive
41 activity, which is known so far only for basaltic shield volcanoes. The data also clearly show
42 that the 1999-2016 eruption sequence ended because of the exhaustion of the enriched
43 source during the very last weeks. Our findings suggest that the metasomatic fertilization of
44 the mantle wedge over short spatial and temporal scales regulates magma production, and
45 ultimately the last eruptive cycle of Tungurahua volcano. This is also a plausible mechanism
46 to produce long-lasting eruptions at andesitic volcanoes.

47

48

49 Volcanoes are fed by vertically extensive magmatic systems rooted at the crust-mantle
50 boundary¹. In arc setting, potentially large magma chambers may form atop these trans-
51 crustal systems and feed volcanic eruptions. Depth, temperature, and evolution of the shallow
52 magma reservoirs are reasonably well constrained from phase equilibria experiments²⁻⁴,
53 mineral chemistry⁵⁻⁷, and the temporal-compositional trends in the erupted magmas^{8,9}.
54 Conversely, owing to the shallow magma chamber filter, the deep roots of arc magmatic
55 systems are less well known. Thermal and physical models point these regions (the so-called
56 “deep crustal hot zones”) as the main place where arc magmas accumulate, differentiate and
57 segregate from their source¹⁰. Magma production in the mantle and deep transfer processes
58 are also increasingly thought to control the recharge and evolution of shallow magma
59 chambers, and ultimately volcanic activity^{11,12}. Here we take advantage of the intense activity
60 of Tungurahua stratovolcano (Ecuador) between 1999 and 2016, and the evidence from large
61 scale syn-eruptive uplift of the edifice for continuous replenishment of the shallow magma
62 chamber^{13,14} to constrain the compositional structure of the deep magma source by means of
63 isotope analysis. We use regularly emitted glassy ash samples which, unlike scarce lavas and
64 other eruptive products, provide a continuous record of magma composition and allow us to
65 reconstruct the small-scale distribution of melts in the volcano source and their progressive
66 extraction during the eruptive sequence. The high-resolution isotope record provides clues as
67 to how deep processes control the secular activity of the Tungurahua volcano.

68

69 **Tungurahua volcano**

70 Tungurahua is an andesitic stratovolcano located in Ecuador along the eastern Andean
71 Cordillera (Fig. 1a). Since 0.3 Ma the volcano produced mainly basaltic andesite and andesite,
72 and occasionally dacite, with no compositional evolution through time^{15,16}. Since 1532 the

73 volcano showed a frequent activity, with main eruptive phases in 1641-1646, 1773-1781,
74 1886-1888, 1916-1918 and 1999-2016^{17,18}. Extensive multi-disciplinary study of the last 1999-
75 2016 eruptive sequence allowed the identification of four stages (Fig. 2a): Stage 1 (1999-2005)
76 started quite intensely with VEI2-3 eruptions in 1999 and 2001¹⁹ and the activity generally
77 decreased through time²⁰. Magma transfer and accumulation resulted in large-scale inflation
78 of the volcano and passive SO₂ degassing^{21,13}. Between January and August 2006 (stage 2), the
79 injection of volatile-rich andesitic magma into a partially degassed magma chamber triggered
80 subplinian eruptions that produced Pyroclastic Density Currents (PDC) and lava flows²²⁻²⁴.
81 Following the 2006 paroxysm, the activity generally decreased between 2007 and mid 2009
82 (stage 3), especially after the February 2008 event that produced PDCs and small lava flows.
83 The activity became less continuous between 2010 and 2016 (stage 4), decreasing first until
84 2012, before increasing markedly from mid-2013 to mid-2014. Then the activity strongly
85 decreased before a last burst between December 2015 and March 2016. The eruptive activity
86 ended suddenly in March 2016 and only cold fumaroles occurred at the summit from 2017.
87 The less continuous activity after 2009 was interpreted as sequential plugging of the system
88 possibly due to lower magma feeding rate²⁵. Although few isotope data from 2006 lavas were
89 previously reported¹⁶, no study so far investigated isotope variations over the eruptive
90 sequence, and generally the role of magma source composition. In particular, the deep source
91 processes responsible for the onset and sudden end of this major eruptive episode remain
92 poorly known. Between November 1999 and March 2016, Tungurahua ash samples were
93 collected monthly to daily at the Choglontus station located below the main plume dispersion
94 axis (W-SW) and far from main roads (Figs. 1b and 2b). Ninety-eight samples were analyzed
95 for Pb and Sr isotope compositions (see Methods).

96

97 **Mass balance model**

98 Lead and strontium isotope compositions of ash samples are reported in supplementary Table
99 S1 and plotted against time in Fig. 3. The isotope compositions of ashes show rapid variations,
100 but no clear temporal trend over the 1999-2016 eruption sequence, beyond the decrease of
101 $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{206}\text{Pb}/^{204}\text{Pb}$, and the increase of $^{207}\text{Pb}/^{206}\text{Pb}$ during the last six months. The
102 variations of raw isotopic ratios have no simple interpretation because each isotope ratio is
103 influenced by more than two sources in Ecuador volcanism²⁶. However, the three isotope
104 dimensions of the Pb system can be used to disentangle the different magma sources. Three
105 main components contribute to the Pb isotopic composition of Tungurahua magmas¹⁶: (1)
106 Depleted mantle. (2) A chemically enriched component derived from slab dehydration and/or
107 melting. This component may occur as pyroxenite veins in the sub-arc mantle^{27,28}, or as later
108 products of melt-rock reactions, such as clinopyroxene-and/or amphibole-bearing cumulates
109 in the lower crust^{29,30}. This component will be referred to as enriched deep component given
110 the uncertainty on its exact nature and location. It will also be called “fertile” due to its lower
111 melting point. (3) Upper continental crust assimilated by mantle derived magmas. The later
112 crustal component must itself include a radiogenic and an unradiogenic end-member to
113 account for the large Pb isotope heterogeneity of basement rocks¹⁶. Although magma
114 components mix at the liquid state, mass balance calculation and results will refer to parental
115 rocks to circumvent the uncertainty related to melting. All compositions used in the model are
116 given in Table 1 (see Methods). The mass-balance equations were solved in three-dimension
117 ^{204}Pb -normalized isotope space using matrix formulation (see Methods).

118

119 **Magma source components versus eruptive activity**

120 The cumulative volume of magma and derived magma production rate (see Methods) are
121 plotted against time (Fig. 4a, b) together with the computed mass fractions of the four source
122 components (Fig. 4c-f). A major observation is that the depleted mantle and the enriched deep
123 component show antithetic temporal variations related to the main eruption stages. For
124 instance, the contribution of the enriched deep component peaked during the 2006 eruption
125 paroxysm (up to 70%), before decreasing between 2007 and mid-2009 (Stage 3), down to 23%
126 in June 2009, as the eruption activity decreased (Fig. 4d). The contribution of the enriched
127 deep component increased again during early stage 4, reaching 60% in August 2012 while the
128 activity had become again nearly continuous (between November 2011 and September 2012).
129 It stayed elevated (about 50%) during the following period of high activity (mid-2013 – mid-
130 2014) before decreasing at the end of the eruptive sequence, in particular during the last six
131 months (Fig. 4d). Remarkably, the fraction of the enriched deep component was the lowest
132 (12.6%) during the very last weeks of the sequence (February 27 to March 14, 2016), strongly
133 suggesting that the 1999-2016 eruption sequence of Tungurahua ended because of the
134 exhaustion of the enriched component in the mantle source. A corollary is that the whole
135 1999-2016 eruption sequence must result from the melting of a small fertile domain, whether
136 in the crust or in the mantle. On the other hand, the highly correlated ($r^2=0.88$) contributions
137 of the two crustal components do not show clear temporal trends, but generally more
138 variability during the first two years of the sequence, and between 2006 and 2009 (Fig. 4e, f).
139 To evaluate statistically the apparent relation between magma source composition and
140 eruption activity, we have analyzed the correlations between the deep source components
141 and the rate of deep magma supply. Figure 5 shows that magma supply rate correlates
142 positively ($r^2=0.61$) with the enriched deep component and negatively ($r^2=0.62$) with the
143 depleted mantle component. Moreover, it is important to note that linear variations of

144 magma source composition result in logarithmic variations of magma production rate,
145 demonstrating that even small variations of mantle fertility have a strong influence on magma
146 supply and eruptive activity. For example, over the course of the 1999-2016 eruption
147 sequence, a factor of 3 variation of the enriched component resulted in a factor of 300
148 variations of magma production rate (Fig. 5a). Although the correlations shown in Fig. 5 are
149 highly significant, they nevertheless show some scatter of the data. This scatter most likely
150 results from magma chamber and conduit processes occasionally uncoupling source and
151 surface processes. Such decoupling is especially apparent at the very beginning of the eruption
152 (1999-2001) (Fig. 4 b-d), where high magma production rate likely resulted from the
153 depressurization of the magma reservoir, whereas magma composition likely reflected mixing
154 between old unerupted magmas from the last 1916-1918 eruption stored in the shallow
155 reservoir and new magmas rising from the mantle. During stage 2 and 3, the tight coupling
156 between eruption activity and source composition are consistent with steady-state supply and
157 open-vent conditions. Conversely, during stage 4 the increase of the fertile component
158 predated the peak of surface activity (July 2013 – April 2014) possibly because the plugging of
159 the conduit²⁵ increased magma storage and mixing before eruption.

160

161 **Implications**

162 Arc volcanism arises from the input of slab fluids that changes mantle composition, lowers its
163 solidus and enables partial melting. In Ecuador, the variations of mantle metasomatism in
164 space and time are responsible for the compositional variations of magmas across^{26,30-32} and
165 along^{26,28} the arc, and the chemical evolution of long-lived volcanic complexes^{33,34}. Using high-
166 resolution Pb isotope time-series we show here for the first time a tight link between small
167 scale metasomatic enrichment of the magma source, and the short-term eruptive activity of

168 a volcano. This link is especially apparent after the early draining of the shallow magma
169 reservoir (1999-2001 magma surge) once the volcano is directly fed from the deep magma
170 source (Fig. 4). Such a link between magma source fertility and eruptive activity is known for
171 basaltic shield volcanoes, which are connected to their mantle sources by relatively simple
172 feeding systems^{35,36}, but is unexpected for Tungurahua stratovolcano that includes several
173 zones of magma storage between 1 and 15 km depth^{13,14,22,37,38}. In the context of continental
174 arc volcanism, the direct connection between mantle or lower-crust and eruption at the
175 surface has two major requirements. First, a process must convert the necessarily slow
176 melting processes into high melt flux to the shallow magma chamber. Reactive melt transport
177 has been shown to meet this requirement beneath mid-ocean ridges³⁹ as well as in arc
178 setting⁴⁰. This process is fundamentally based on the chemically unstable nature of primary
179 melts. In arc setting, primary melts with high silica and water contents dissolve olivine and
180 crystallize pyroxene⁴¹. Pyroxene is subsequently replaced by amphibole as the melt-rock
181 reaction proceeds near the mantle-crust boundary²⁹. The dissolution-precipitation reactions
182 increase locally the porosity and permeability of the mantle and lead to the formation of melt
183 channels. Keller and Katz⁴⁰ showed that dissolution of fertile material and magma transport
184 in such channels are enhanced by up to an order of magnitude. Very importantly, the
185 formation of melt channels is an unstable process that bridges the gap between the slow
186 melting of the mantle and the fast melt extraction required to feed sustained volcanic
187 eruptions. Rawson et al.⁴² suggested that reactive channels are likely to develop in the distal
188 parts of the mantle wedge where more stable thermal and rheological conditions prevail. The
189 location of Tungurahua volcano along the eastern Cordillera, 330 km away from the
190 subduction trench, could favor the development of reactive melt channels. The high magma
191 productivity of Tungurahua volcano might also be related to its location above a major fault

192 in the subducting plate (The Grijalva Fracture Zone) (Fig. 1a), because serpentinized
193 peridotites exhumed near fracture zones may deliver large amounts of volatiles to the sub-arc
194 mantle and enhance flux melting⁴³.

195 The second requirement is that the inflow of mantle derived magmas is not totally absorbed
196 by the magma storage system in the crust, so that part of the magma flow directly feeds the
197 eruptions. In support of such a process, large-scale uplift of the Tungurahua edifice between
198 2003 and 2014 indicates continuous replenishment of the magma reservoir during the
199 eruption sequence, with slightly more than half of the magma supplied to the reservoir
200 reaching the surface^{13,14}. Thus, Tungurahua seems to belong to the few stratovolcanoes from
201 the Central and South American arcs that are directly connected to their mantle source,
202 together with Irazu volcano in Costa Rica⁴⁴ and Ollagüe volcano in Chile⁴⁵. In such a situation,
203 extensive crustal contamination of Tungurahua magmas^{16,26} is at first glance paradoxical
204 because crustal assimilation is generally thought to be related to magma storage and
205 differentiation in the crust. The extensive contamination of Tungurahua andesitic magmas can
206 be reconciled with the evidence for short residence time in the crust if the magmas are
207 superheated during rapid ascent^{45,46}. In support of this hypothesis, the absence of amphibole
208 in Tungurahua andesites, together with thermometric and experimental constraints suggest
209 that the magmas evolved at temperatures higher than 950–1000 °C^{22,37}. The time scale of
210 magma ascent is not constrained here, but the residence time of magma in the reservoir must
211 be comparable to the time scale of geochemical fluctuations⁴⁷, on the order of years (Fig. 3).
212 The view that emerges for Tungurahua is that of a more dynamic magmatic system compared
213 to the deep crustal hot zones¹⁰ which require timescale of 10⁵–10⁶ years for thermal
214 incubation at depth, and 10–10³ years for magmatic differentiation from basalt to andesite.
215 One possible solution is that the trans-crustal plumbing system is a much more dynamic

216 environment than expected, with probably “active” zones with rapid magma ascent and other
217 zones dominated by magma accumulation and differentiation. In this frame, Tungurahua is a
218 good candidate to join the growing family of arc volcanoes, where seismic tomography⁴⁸,
219 volatiles diffusion along crystal embayments⁴⁹, and Ra-Th isotope disequilibria⁵⁰ indicate ultra-
220 fast magma ascent in well-developed melt channels. A major outcome of this study is that
221 eruption of the Tungurahua volcano requires at least 12% of fertile lithology in its source.
222 Below 12%, melt production is likely too low to sustain magma eruption at the surface, as
223 illustrated by the exhaustion of the fertile component a few weeks before the end of the 1999-
224 2016 eruption sequence. Conversely, when the fraction of the fertile lithology reaches 60-
225 70%, reactive channeling instabilities allow high melt supply to the volcano, and trigger
226 eruption paroxysm, such as in 2006. In support of such process, magma production increases
227 logarithmically with the degree of source enrichment (Fig. 5). The nature and location of the
228 fertile source lithology (i.e., the enriched deep component) remain uncertain, although its
229 isotopic signature requires a genetic link with the subducting oceanic crust. In any case, the
230 fact that the fractions of the enriched deep component and the depleted mantle are strictly
231 complementary (Fig. 4c,d) points to a mantle origin for the isotopic signal correlating with
232 volcanic activity. In such situation, our preferred interpretation is that metasomatic
233 fertilization of the mantle wedge over short spatial and temporal scales regulates magma
234 production rates and ultimately the secular activity of the volcano. Small-scale variations of
235 mantle metasomatism in southern Ecuador is independently evidenced by olivine-hosted melt
236 inclusions²⁸, and could be related to variations of the thermal structure and composition of
237 the subducting plate in the vicinity of the Grijalva Fracture Zone.

238 This work confidently resolves a link between the short-term activity of an arc volcano and
239 the small-scale heterogeneity of its magma source. This observation leaves some open

240 questions regarding the geometry of the melting source and how melting and melt extraction
241 occur. For instance, the episodic activity of arc volcanoes could reflect the melting of individual
242 fertile veins forming some kind of fractal tree in the mantle wedge. In such case, the spatial
243 distribution and the size of the veins, and the re-organization of melt channels driven by the
244 slow decompression of the mantle, will control melt production and volcano supply.
245 Alternatively, if each volcano is fed by a main melt channel in the mantle wedge⁵¹, short-term
246 variations in fluid delivery from the slab and related flux melting are expected to govern melt
247 production and volcano supply.

248 On a more practical point of view, the results of this work suggest that volcano monitoring
249 techniques should be designed to track the deeper process of the magmatic plumbing system.
250 For instance, monitoring the isotopic composition of juvenile ashes during a long-lived
251 eruption, or an eruption sequence, would provide information on how the fertility of the
252 magma source evolves as the eruption proceeds. Such information could be used to forecast
253 eruption paroxysm, major changes of activity, or eruption end. The volatility of Pb in magmatic
254 systems also raises the possibility to probe the hidden magmatic evolution of quiescent
255 volcanoes from the Pb isotope composition of volcanic gases⁵².

256

257 **Figure legends**

258

259 Fig. 1: Location maps. (a) Regional map showing the location of the Tungurahua volcano (star)
260 along the eastern Cordillera, other main volcanoes of the arc (triangles), the oceanic Grijalva
261 Fracture Zone and its projection on land (dashed line). (b) Map of Tungurahua volcano
262 showing the sampling location.

263

264 Fig. 2: History and sampling of the 1999-2016 eruption sequence of the Tungurahua volcano.
265 (a) Periods of eruption and inactivity are shown in black and white, respectively. The four main
266 eruption stages (S1-S4) are indicated^{25,53}. (b) Ash sampling during the sequence.

267

268 Fig. 3: Pb and Sr isotopic composition of ash samples plotted against time. Error bars (S.D.) are
269 within symbol size. The four main eruption stages (S1-S4) are indicated^{25,53}.

270

271 Fig. 4: Temporal evolution of eruptive activity and magma source components. The four main
272 eruption stages (S1-S4) are indicated^{25,53}. (a) Cumulative volume of magma. (b) Rate of deep
273 magma supply estimated by calculating the slope of the cumulative volume emission curve
274 over a moving two-year window (see Methods). The high magma production rate during the
275 first years (1999-2001) results from the depressurization of the plumbing system (shaded
276 area), whereas subsequent eruptive activity can be related to melt production at depth. (c-f)
277 Mass fractions of magma source components are plotted versus time: (c) Depleted mantle.
278 (d) Slab-derived enriched deep component, located either in the mantle or in the lower crust.
279 (e&f) Upper crustal components including Pb-poor radiogenic end-member and Pb-rich

280 unradiogenic end-member. Component compositions and model equations are given in the
281 methods.

282

283 Fig. 5: Rate of deep magma supply plotted against magma source components. The rate of
284 deep magma supply was estimated by calculating the slope of the cumulative volume emission
285 curve over a moving two-year window (see Methods). The rate of deep magma supply is
286 plotted against (a) the mass fraction of the enriched deep component, and (b) the mass
287 fraction of the depleted mantle component. Linear regressions and correlation coefficients at
288 95% confidence level are indicated.

289 **References**

290

- 291 1. Cashman, K. V., Sparks, R.S.J. & Blundy, J.D. Vertically extensive and unstable magmatic
292 systems: a unified view of igneous processes. *Science* **355**, eaag3055 (2017).
- 293 2. Rutherford, M.J., Sigurdsson, H., Carey, S. & Davis, A. The May 18, 1980, eruption of Mount
294 St. Helens .1. Melt composition and experimental phase equilibria. *J. Geophys. Res.* **90**,
295 2929-2947 (1985).
- 296 3. Moore, G. & Carmichael, I.S.E. The hydrous phase equilibria (to 3 kbar) of an andesite and
297 basaltic andesite from western Mexico: constraints on water content and conditions of
298 phenocryst growth. *Contrib. Mineral. Petrol.* **130**, 304-319 (1998).
- 299 4. Andujar, J., Scaillet, B., Pichavant, M. & Druitt, T.H. Differentiation conditions of a basaltic
300 magma from Santorini, and its bearing on the production of andesite in arc settings. *J.*
301 *Petrol.* **56**, 765–794 (2015).
- 302 5. Ginibre, C., Wörner , G. & Kronz, A. Crystal zoning as an archive for magma evolution.
303 *Elements* **3**, 261-266 (2007).
- 304 6. Morgan, D.J. et al. Time scales of crystal residence and magma chamber volume from
305 modelling of diffusion profiles in phenocrysts: Vesuvius 1944. *Earth Planet Sci. Lett.* **222**,
306 933-946 (2004).
- 307 7. Turner, S. & Costa, F. Measuring timescales of magmatic evolution. *Elements* **3**, 267-272
308 (2007).
- 309 8. Corsaro, R.A. & Pompilio, M. Magma dynamics in the shallow plumbing system of Mt. Etna
310 as recorded by compositional variations in volcanics of recent summit activity (1995–
311 1999). *J. Volcanol. Geotherm. Res.* **137**, 55-71 (2004).

- 312 9. Turner, S.J., Izbekov, P. & Langmuir, C. The magma plumbing system of Bezymianny
313 Volcano: Insights from a 54 year time series of trace element whole-rock geochemistry
314 and amphibole compositions. *J. Volcanol. Geotherm. Res.* **263**, 108–121 (2013).
- 315 10. Annen, C., Blundy, J.D. & Sparks, R.S.J. The genesis of intermediate and silicic magmas in
316 deep crustal hot zones. *J. Petrol.* **47**, 505–539 (2006).
- 317 11. Till, C.B. et al. The causes of spatiotemporal variations in erupted fluxes and compositions
318 along a volcanic arc. *Nat. Commun.* **10**, 1350 (2019).
- 319 12. Colucci, S. & Papale, P. Deep Magma Transport Control on the Size and Evolution of
320 Explosive Volcanic Eruptions. *Front. Earth Sci.* **9**, 681083 (2021).
- 321 13. Champenois, J. et al. Large-scale inflation of Tungurahua volcano (Ecuador) revealed by
322 Persistent Scatterers SAR interferometry. *Geophys. Res. Lett.* **41**, 5821–5828 (2014).
- 323 14. Muller, C. et al. Temporal evolution of the magmatic system at Tungurahua Volcano,
324 Ecuador, detected by geodetic observations. *J. Volcanol. Geotherm. Res.* **368**, 63-72
325 (2018).
- 326 15. Bablon, M. et al. Eruptive chronology of Tungurahua volcano (Ecuador) revisited based on
327 new K-Ar ages and geomorphological reconstructions. *J. Volcanol. Geotherm. Res.* **357**,
328 378–398 (2018).
- 329 16. Nauret, F. et al. The genetic relationship between andesites and dacites at Tungurahua
330 volcano, Ecuador. *J. Volcanol. Geotherm. Res.* **349**, 283-297 (2018).
- 331 17. Hall, M.L., Robin, C., Beate, B., Mothes, P. & Monzier, M. Tungurahua Volcano, Ecuador:
332 structure, eruptive history and hazards. *J. Volcanol. Geotherm. Res.* **91**, 1–21 (1999).
- 333 18. Le Pennec, J.-L. et al. The AD 1300–1700 eruptive periods at Tungurahua volcano, Ecuador,
334 revealed by historical narratives, stratigraphy and radiocarbon dating. *J. Volcanol.*
335 *Geotherm. Res.* **176**, 70–81 (2008).

- 336 19. Le Pennec, J.-L. et al. Impact of tephra falls on Andean communities: The influences of
337 eruption size and weather conditions during the 1999–2001 activity of Tungurahua
338 volcano, Ecuador. *J. Volcanol. Geotherm. Res.* **217-218**, 91-103 (2012).
- 339 20. Wright, H.M.N. et al. Estimating rates of decompression from textures of erupted ash
340 particles produced by 1999–2006 eruptions of Tungurahua volcano, Ecuador. *Geology*
341 **40**, 619–622 (2012).
- 342 21. Arellano, S.R. et al. Degassing patterns of Tungurahua volcano (Ecuador) during the 1999–
343 2006 eruptive period, inferred from remote spectroscopic measurements of SO₂
344 emissions. *J. Volcanol. Geotherm. Res.* **176**, 151–162 (2008).
- 345 22. Samaniego, P., Le Pennec, J.L., Robin, C. & Hidalgo, S. Petrological analysis of the pre-
346 eruptive magmatic process prior to the 2006 explosive eruptions at Tungurahua volcano
347 (Ecuador). *J. Volcanol. Geotherm. Res.* **199**, 69–84 (2011).
- 348 23. Eychenne, J., Le Pennec, J.L., Ramon, P. & Yepes, H. Dynamics of explosive paroxysms at
349 open-vent andesitic systems: High-resolution mass distribution analyses of the 2006
350 Tungurahua fall deposit (Ecuador). *Earth Planet. Sci. Lett.* **361**, 343-355 (2013).
- 351 24. Myers, M.L. et al. Replenishment of volatile-rich mafic magma into a degassed chamber
352 drives mixing and eruption of Tungurahua volcano. *Bull. Volcanol.* **76**, 872 (2014).
- 353 25. Hidalgo, S. et al. SO₂ degassing at Tungurahua volcano (Ecuador) between 2007 and 2013:
354 transition from continuous to episodic activity. *J. Volcanol. Geotherm. Res.* **298**, 1–14
355 (2015).
- 356 26. Ancellin, M.-A. et al. Across-arc versus along-arc Sr-Nd-Pb isotope variations in the
357 Ecuadorian volcanic arc. *Geochem. Geophys. Geosyst.* **18**, e2016GC006679 (2017).
- 358 27. Straub, S.M. et al. Formation of hybrid arc andesites beneath thick continental crust. *Earth*
359 *Planet Sci. Lett.* **303**, 337-347 (2011).

- 360 28. Narvaez, D., Rose-Koga, E., Samaniego, P., Koga, K. & Hidalgo, S. Constraining magma
361 sources using primitive olivine-hosted melt inclusions from Puñalica and Sangay
362 volcanoes (Ecuador). *Contrib. Mineral. Petrol.* **173**, 80 (2018).
- 363 29. Smith, D.J. Clinopyroxene precursors to amphibole sponge in arc crust. *Nat. Commun.* **5**,
364 4329 (2014).
- 365 30. Chiaradia, M., Müntener, O., Beate, B. & Fontignie, D. Adakite-like volcanism of Ecuador:
366 lower crust magmatic evolution and recycling. *Contrib. Mineral. Petrol.* **158**, 563–588
367 (2009).
- 368 31. Bourdon, E. et al. Magmatic response to early aseismic ridge subduction: the Ecuadorian
369 margin case (South America). *Earth Planet. Sci. Lett.* **205**, 123–138 (2003).
- 370 32. Le Voyer, M., Rose-Koga, E.F., Laubier, M. & Schiano, P. Petrogenesis of arc lavas from the
371 Rucu Pichincha and Pan de Azucar volcanoes (Ecuadorian arc): major, trace element,
372 and boron isotope evidences from olivine-hosted melt inclusions. *Geochem. Geophys.*
373 *Geosyst.* **9**, e2008GC002173 (2008).
- 374 33. Hidalgo, S. et al. Adakitic magmas in the Ecuadorian volcanic front: petrogenesis of the
375 Iliniza Volcanic Complex (Ecuador). *J. Volcanol. Geotherm. Res.* **159**, 366–392 (2007).
- 376 34. Samaniego, P., Robin, C., Chazot, G., Bourdon, E. & Cotten, J. Evolving metasomatic agent
377 in the Northern Andean subduction zone, deduced from magma composition of the
378 long-lived Pichincha volcanic complex (Ecuador). *Contrib. Mineral. Petrol.* **160**, 239–260
379 (2010).
- 380 35. Pietruszka, A.J., Rubin, K.H. & Garcia, M.O. ^{226}Ra - ^{230}Th - ^{238}U disequilibria of historical
381 Kilauea lavas (1790–1982) and the dynamics of mantle melting within the Hawaiian
382 plume. *Earth Planet Sci. Lett.* **186**, 15–31 (2001).

- 383 36. Vlastélic, I. et al. Control of source fertility on the eruptive activity of Piton de la Fournaise
384 volcano, La Réunion. *Sci. Rep.* **8**, 14478 (2018).
- 385 37. Andújar, J. et al. Structure of the plumbing system at Tungurahua volcano, Ecuador:
386 insights from phase equilibrium experiments on July–August 2006 eruption products. *J.*
387 *Petrol.* **58**, 1249–1278 (2017).
- 388 38. Koch, C.D. et al. Crustal thickness and magma storage beneath the Ecuadorian arc. *J. South*
389 *Amer. Earth Sci.* **110**, 103331 (2021).
- 390 39. Kelemen, P.B., Shimizu, N. & Salters, V.J.M. Extraction of mid-ocean-ridge basalt from the
391 upwelling mantle by focused flow of melt in dunite channels. *Nature* **375**, 747–753
392 (1995).
- 393 40. Keller, T. & Katz, R.F. The Role of Volatiles in Reactive Melt Transport in the Asthenosphere.
394 *J. Petrol.* **57**, 1073–1108 (2016).
- 395 41. Müntener, O., Kelemen, P.B. & Grove, T.L. The role of H₂O during crystallization of
396 primitive arc magmas under uppermost mantle conditions and genesis of igneous
397 pyroxenites: an experimental study. *Contrib. Mineral. Petrol.* **141**, 643-658 (2001).
- 398 42. Rawson, H. et al. Compositional variability in mafic arc magmas over short spatial and
399 temporal scales: Evidence for the signature of mantle reactive melt channels. *Earth*
400 *Planet Sci. Lett.* **456**, 66-77 (2016).
- 401 43. Paulatto, M. et al. Dehydration of subducting slow-spread oceanic lithosphere in the Lesser
402 Antilles. *Nat. Commun.* **8**, 15980 (2017).
- 403 44. Ruprecht, P. & Plank, T. Feeding andesitic eruptions with a high-speed connection from
404 the mantle. *Nature* **500**, 68-72 (2013).
- 405 45. Mattioli, M., Renzulli, A., Menna, M. & Holm, P.M. Rapid ascent and contamination of
406 magmas through the thick crust of the CVZ (Andes, Ollagüe region): Evidence from a

- 407 nearly aphyric high-K andesite with skeletal olivines. *J. Volcanol. Geotherm. Res.* **158**, 87-
408 105 (2006).
- 409 46. Marsh, B.D. On the cooling of ascending andesitic magma. *Phil. Trans. R. Soc. Lond. A.* **288**,
410 611-625 (1978).
- 411 47. Albarède, F. Residence time analysis of geochemical fluctuations in volcanic series.
412 *Geochim. Cosmochim. Acta* **57**, 615-621 (1993).
- 413 48. Koulakov, I. et al. Rapid changes in magma storage beneath the Klyuchevskoy group of
414 volcanoes inferred from time-dependent seismic tomography. *J. Volcanol. Geotherm.*
415 *Res.* **263**, 75-91 (2013).
- 416 49. Moussallam, Y. et al. Fast ascent rate during the 2017–2018 Plinian eruption of Ambae
417 (Aoba) volcano: a petrological investigation. *Contrib. Mineral. Petrol.* **174**, 90 (2019).
- 418 50. Turner, S., Evans, P. & Hawkesworth, C. Ultrafast Source-to-Surface Movement of Melt at
419 Island Arcs from ^{226}Ra - ^{230}Th Systematics. *Science* **292**, 1363-1366 (2001).
- 420 51. Tamura, Y., Tatsumi, Y., Zhao, D., Kido, Y. & Shukuno, H. Hot fingers in the mantle wedge:
421 new insights into magma genesis in subduction zones. *Earth. Planet. Sci. Lett.* **197**, 105–
422 116 (2002).
- 423 52. Vlastélic I. et al. Probing the hidden magmatic evolution of El Misti volcano (Peru) with the
424 Pb isotope composition of fumaroles. *Bull. Volcanol.* **84**, 17 (2022).
- 425 53. Bustillos, A.J., Romero, J.E., Guevara, C.A. & Díaz-Alvarado, J. Tephra fallout from the long-
426 lasting Tungurahua eruptive cycle (1999-2014): Variations through eruptive style
427 transition and deposition processes. *And. Geol.* **45**, 47-77 (2018).
- 428

429 **Corresponding author**

430 Correspondence and requests for materials should be addressed to I. Vlastelic

431 (vlastelic@ipgp.fr)

432

433 **Acknowledgements**

434 We thank J.E. Romero and two anonymous reviewers for their constructive comments and R.

435 Neely for handling the manuscript. This work is part of the “Laboratoire Mixte International”

436 program entitled “Seismes et Volcans dans les Andes du Nord”. The research was financed by

437 the Institut de Recherche pour le Développement, the French Government Laboratory of

438 Excellence initiative n° ANR-10-LABX-0006, the Region Auvergne and the European Regional

439 Development Fund. This is Laboratory of Excellence ClerVolc contribution number xxx.

440

441 **Author contribution**

442 I.V. designed the study, coordinated the analyses and wrote the paper

443 N.S. performed the isotopic analyses and wrote the paper

444 P.S. designed the study and contributed to data interpretation

445 B.B. collected samples and contributed to data interpretation

446 F.N. and S.H. contributed to the discussions and interpretation of the data

447 D.A. supervised Sr isotope measurements

448 M.G. supervised Pb isotope measurements

449

450 **Supplementary information**

451 Supplementary Table S1

452

453 **Competing interests**

454 The authors declare no competing interests.

455

456 **Methods**

457

458 **Samples.** Ash fallouts were sampled using ash collectors before 2010 and an ashmeter device
459 subsequently⁵⁴. The 1999-2016 ash samples are dominantly juvenile. They include black and
460 brown scoria, 100 to 600 μm in size, larger beige pumices (1000 to 5000 μm) and glass shards.
461 Ashes matrix is glassy to micro-crystallized and shows a uniform mineralogical assemblage
462 (plagioclases + orthopyroxenes + clinopyroxenes + Fe-Ti oxides \pm olivines). These observations
463 are consistent with previous descriptions^{22,23,53}.

464

465 **Analytical methods.** All chemical analyses were performed at the Laboratoire Magmas et
466 Volcans (Clermont-Ferrand, France) between 2018 and 2020. Lead and Sr isotope
467 compositions were measured on 5 mg of juvenile glassy fragments that were hand-picked
468 under a binocular microscope and washed in deionized water. Following sample dissolution
469 with HNO_3 -HF, Pb and Sr were separated using polyethylene columns filled with 50 μL of
470 Sr.Spec extraction chromatographic material⁵⁵. Full procedural blanks were 2.7 ± 1.1 ng for Sr
471 ($n= 13$) and 19 ± 15 pg for Pb ($n=9$), which are negligible compared to the amounts of Sr (2.1-
472 3.7 μg) and Pb (40-75 ng) extracted from the ash samples. Lead isotope compositions were
473 measured on a Thermo Fisher Scientific Neptune Plus MC-ICP-MS coupled to an Aridus II
474 desolvating nebulizer system. The Pb instrumental mass fractionation was corrected during
475 measurement by doping the Pb solutions with the Tl NBS997 standard. The NBS981 standard
476 was measured between every two samples to correct Pb ratios for instrumental drift and the
477 slight difference in mass fractionation between Pb and Tl. Data were normalized to the
478 NBS981 standard using the values of Todt et al.⁵⁶ ($^{206}\text{Pb}/^{204}\text{Pb}= 16.9356$, $^{207}\text{Pb}/^{204}\text{Pb}= 15.4891$,
479 $^{208}\text{Pb}/^{204}\text{Pb}= 36.7006$). Repeated analysis of the NBS981 standard every two samples yielded

480 average intra-session reproducibility of 115 ppm (2 S.D.) for the $^{206}\text{Pb}/^{204}\text{Pb}$ ratio. Strontium
 481 isotope compositions were measured on a Thermo Fisher Scientific Triton Thermal Ionization
 482 Mass Spectrometer (TIMS) and were corrected for mass fractionation using $^{86}\text{Sr}/^{88}\text{Sr}=0.1194$.
 483 Repeated measurement of the NBS SRM 987 standard during the course of the study yielded
 484 $^{87}\text{Sr}/^{86}\text{Sr}=0.710232 \pm 8$ (2 S.D.). Sample data were renormalized to the accepted value of
 485 0.710245. Repeated measurement (duplicate dissolution) of rock reference materials yielded
 486 $^{87}\text{Sr}/^{86}\text{Sr}= 0.705004 \pm 14$, $^{206}\text{Pb}/^{204}\text{Pb} = 18.756 \pm 0.005$, $^{207}\text{Pb}/^{204}\text{Pb} = 15.615 \pm 0.004$,
 487 $^{208}\text{Pb}/^{204}\text{Pb} = 38.708 \pm 0.026$ for BCR2 basalt (n=5), and $^{87}\text{Sr}/^{86}\text{Sr}= 0.703532 \pm 13$, $^{206}\text{Pb}/^{204}\text{Pb}$
 488 $= 18.301 \pm 0.005$, $^{207}\text{Pb}/^{204}\text{Pb} = 15.536 \pm 0.007$, $^{208}\text{Pb}/^{204}\text{Pb} = 38.249 \pm 0.024$ for JA-1 andesite
 489 (n=6).

490

491 **Mixing model.** The compositions of the four end-member components used in the mixing
 492 model are given in Table1.

Table 1

	Depleted mantle	Enriched deep component (arc pyroxenite)	High U/Pb Pb poor crust	Low U/Pb Pb rich crust
Pb (ppm)	0.018	0.06	9	17
$^{206}\text{Pb}/^{204}\text{Pb}$	18.40	19.10	19.20	18.79
$^{207}\text{Pb}/^{204}\text{Pb}$	15.50	15.56	15.72	15.66
$^{208}\text{Pb}/^{204}\text{Pb}$	38.00	38.70	39.22	38.60

493

494 The Pb concentration of the depleted mantle is from ref. ⁵⁷. The Pb content of the enriched
 495 deep component is assumed to be as in clinopyroxenes crystalizing early in arc magmatism²⁹.

496 Lead concentrations of basement rocks are from Nauret et al.¹⁶. Lead isotopic compositions
 497 of the four components are also from Nauret et al.¹⁶, with slight modification of the crustal
 498 compositions. This fine tuning is needed to model the most contaminated samples with high
 499 ²⁰⁷Pb/²⁰⁴Pb. The Andean continental crust being highly heterogeneous, it is possible that the
 500 crustal rocks assimilated during the 1999-2016 eruption of the Tungurahua volcano are not
 501 exactly the same as the five basement samples analyzed by Nauret et al.¹⁶. The mass fractions
 502 (x) of the four components are quantified in the ²⁰⁶Pb/²⁰⁴Pb —²⁰⁷Pb/²⁰⁴Pb—²⁰⁸Pb/²⁰⁴Pb
 503 isotope space by solving the mass balance equation system:

504

$$505 \text{ Pb}_1 \cdot x_1 \cdot [(64)_s - (64)_1] + \text{Pb}_2 \cdot x_2 \cdot [(64)_s - (64)_2] + \text{Pb}_3 \cdot x_3 \cdot [(64)_s - (64)_3] + \text{Pb}_4 \cdot x_4 \cdot [(64)_s - (64)_4] = 0$$

$$506 \text{ Pb}_1 \cdot x_1 \cdot [(74)_s - (74)_1] + \text{Pb}_2 \cdot x_2 \cdot [(74)_s - (74)_2] + \text{Pb}_3 \cdot x_3 \cdot [(74)_s - (74)_3] + \text{Pb}_4 \cdot x_4 \cdot [(74)_s - (74)_4] = 0$$

$$507 \text{ Pb}_1 \cdot x_1 \cdot [(84)_s - (84)_1] + \text{Pb}_2 \cdot x_2 \cdot [(84)_s - (84)_2] + \text{Pb}_3 \cdot x_3 \cdot [(84)_s - (84)_3] + \text{Pb}_4 \cdot x_4 \cdot [(84)_s - (84)_4] = 0$$

$$508 x_1 + x_2 + x_3 + x_4 = 1$$

509

510 where (64), (74) and (84) are for ²⁰⁶Pb/²⁰⁴Pb, ²⁰⁷Pb/²⁰⁴Pb and ²⁰⁸Pb/²⁰⁴Pb isotope ratios,
 511 respectively; Pb is for lead concentration, subscript S is for sample, and subscripts 1-4 are for
 512 the four mixing components reported in Table 1. The equation system can be written as the
 513 matrix product:

514

$$\begin{pmatrix} \text{Pb}_1 \cdot [(64)_s - (64)_1] & \text{Pb}_2 \cdot [(64)_s - (64)_2] & \text{Pb}_3 \cdot [(64)_s - (64)_3] & \text{Pb}_4 \cdot [(64)_s - (64)_4] \\ \text{Pb}_1 \cdot [(74)_s - (74)_1] & \text{Pb}_2 \cdot [(74)_s - (74)_2] & \text{Pb}_3 \cdot [(74)_s - (74)_3] & \text{Pb}_4 \cdot [(74)_s - (74)_4] \\ \text{Pb}_1 \cdot [(84)_s - (84)_1] & \text{Pb}_2 \cdot [(84)_s - (84)_2] & \text{Pb}_3 \cdot [(84)_s - (84)_3] & \text{Pb}_4 \cdot [(84)_s - (84)_4] \\ 1 & 1 & 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

515

516 or :

517

518 $A \cdot B = C$

519

520 the solution matrix B is obtained from

521

522 $B = A^{-1} \cdot C$

523

524 where A^{-1} is the inverse matrix of A.

525 We stress that the temporal trends inferred from the model (Fig. 4) do not depend on the

526 precise composition of the end-member components used (Table 1). In support of this idea,

527 consistent trends can be inferred from the raw $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{207}\text{Pb}/^{204}\text{Pb}$ ratios (Fig. 3b,d),

528 which dominantly record mantle composition and crustal processes, respectively^{16,26}.

529

530

531 **Rate of deep magma supply.** The cumulative volume of magma was determined using ash

532 fallout data from ref. ²⁰ for the 1999-2004 period, and ref. ⁵⁸ for the 2006-2014 period. Ash

533 volumes were converted into magma volumes assuming the density of magma is 2.5 times

534 that of ash deposit. For the August 2006 paroxysmal eruption, we used the complete mass

535 quantification of ref. ⁵⁹. For the 2014-2016 period, our ash fallout measurements correspond

536 to $9.4 \times 10^4 \text{ m}^3$ (09/03/2014-06/22/2015), $8.7 \times 10^4 \text{ m}^3$ (06/22/2015-09/05/2015), $8.7 \times 10^5 \text{ m}^3$

537 (09/05/2015-12/08/2015) and $4.2 \times 10^5 \text{ m}^3$ (02/26/2016-03/15/2016) of magma (B.B.

538 unpublished data and IG-EPN report available at:

539 <https://www.igepn.edu.ec/servicios/noticias/1463-erupcion-de-febrero-marzo-2016-del->

540 volcan). We also included in the calculation the 2010 and 2014 lava flows ($1 \times 10^6 \text{ m}^3$ each
541 according to ref. ⁶⁰ and B.B. unpublished data) and the accumulation of ballistic material in
542 the summit crater ($3.4 \times 10^6 \text{ m}^3$ between 2008 and 2012 and $0.8 \times 10^6 \text{ m}^3$ between 2012 and
543 2016, which translates into $2.1 \times 10^6 \text{ m}^3$ and $0.5 \times 10^6 \text{ m}^3$ of magma, respectively, according to
544 ref.⁵⁹). Summit crater data are from D. Yaguana 2020 engineering thesis
545 (<http://bibdigital.epn.edu.ec/handle/15000/21245>). The cumulative volume emission curve
546 was built with a time resolution of one point per month (Fig. 4a). The rate of deep magma
547 supply (Fig. 4b) was estimated by calculating the slope of the cumulative volume emission
548 curve over a moving two-year window (linear regression through 24 data points). A two-year
549 window was selected because it corresponds to the timescale of source component
550 fluctuations (Fig. 4c,d). Such a relatively wide time-window is expected to smooth monthly
551 variations of shallow magma supply, which has varied by a factor of 10^6 over the course of the
552 1999-2006 eruption²⁰. The calculated rate of magma supply was assigned to the middle of the
553 time window.

554

555 **Data availability**

556 All data generated during this study are included in this published article (and its
557 supplementary information files)

558

559 **Methods references**

560

561 54. Bernard, B. Homemade ashmeter: a low-cost, high-efficiency solution to improve tephra
562 field-data collection for contemporary explosive eruptions. *J. Applied Volcanol.* **2**, 1
563 (2013).

- 564 55. Deniel, C. & Pin, C. Single-stage method for the simultaneous isolation of lead and
565 strontium from silicate samples for isotopic measurements. *Anal. Chim. Acta* **426**, 95–
566 103 (2001).
- 567 56. Todt, W., Cliff, R.A., Hanser, A. & Hofmann, A.W. Evaluation of a ^{202}Pb – ^{205}Pb double spike
568 for high-precision lead isotope analysis. In: Basu, A. & Hart, S. (eds) *Earth Processes:
569 Reading the Isotopic Code. Am. Geophys. Union, Geophys. Mon.* **95**, 429–437 (1996).
- 570 57. Workman, R.K. & Hart, S.R. Major and trace element composition of the depleted MORB
571 mantle (DMM). *Earth Planet. Sci. Lett.* **231**, 53–72 (2005).
- 572 58. Bustillos, J., Romero, J., Troncoso, L., Guevara, A. Tephra fall at Tungurahua Volcano
573 (Ecuador)-1999-2014: An example of tephra accumulation from a long-lasting eruptive
574 cycle. *Geofis. Int.* **55**, 55-67. (2016)
- 575 59. Bernard, J., Eychenne, J., Le Pennec, J.-L. & Narvaez, D. Mass budget partitioning during
576 explosive eruptions: insights from the 2006 paroxysm of Tungurahua volcano, Ecuador,
577 *Geochem. Geophys. Geosyst.* **17**, 3224– 3240 (2016).
- 578 60. Kelfoun, K. & Vallejo, S. VolcFlow capabilities and potential development for the simulation
579 of lava. *Geol. Soc. London Spec. Publ.* **426**, 337-343 (2016).
- 580
- 581









