



HAL
open science

Transient response to changes in uplift rates in the northern Atlas-Meseta system (Morocco)

R. Clementucci, P. Ballato, L.L. Siame, C. Faccenna, S. Racano, G. Torreti,
R. Lanari, L. Leanni, V. Guillou

► **To cite this version:**

R. Clementucci, P. Ballato, L.L. Siame, C. Faccenna, S. Racano, et al.. Transient response to changes in uplift rates in the northern Atlas-Meseta system (Morocco). *Geomorphology*, 2023, 436, pp.108765. 10.1016/j.geomorph.2023.108765 . insu-04157642

HAL Id: insu-04157642

<https://insu.hal.science/insu-04157642v1>

Submitted on 10 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Transient response to changes in uplift rates in the northern Atlas-Meseta system**
2 **(Morocco)**

3 Clementucci R.^{1,2*}, Ballato P.¹, Siame L. L.², Faccenna C.^{1,3}, Racano S.⁴, Torreti G.¹, Lanari R.⁵,
4 Leanni L.², Guillou V.²

5

6 ¹ *Dipartimento di Scienze, Università Roma Tre, Largo San Leonardo Murialdo 1, 00146 Rome, Italy*

7 ² *Aix-Marseille Univ., CNRS, IRD, INRAE, Collège de France, CEREGE, Aix-en Provence, France*

8 ³ *GFZ-German Research Centre for Geosciences, Potsdam, Germany*

9 ⁴ *Institute for Geosciences, University of Potsdam, Potsdam, Germany*

10 ⁵ *Dipartimento di scienze, Università di Firenze, Italia*

11 **Corresponding author: rclementucci@ethz.ch*

12 *Current address: Department of Earth Sciences, ETH Zurich, Zurich, Switzerland.*

13 **Keywords:** landscape evolution, cosmogenic nuclides, transient topography, quantitative
14 geomorphology, Atlas Mountains

15

16 **Abstract**

17 Transient topography represents an opportunity for extracting information on the combined effect of
18 tectonics, mantle-driven processes, lithology and climate across different temporal and spatial scales.

19 The geomorphic signature of transient conditions can be used to unravel landscape evolution,
20 especially in areas devoid of stratigraphic constraints. The topography of the Western Moroccan

21 Meseta domain (WMM) is characterized by elevated non-lithological knickpoints, that delimit an
22 uplifted relict landscape, implying a transient response to a change in uplift rate that occurred during

23 the Cenozoic. Here, we determine denudation rates of selected watersheds and bedrock outcrops from
24 cosmogenic nuclides and perform stream profile, regional and basin-scale geomorphic analysis.

25 Denudation rates of the relict and the rejuvenated landscape range from 15 to 20 m/Myr and from 30
26 to 40 m/Myr, respectively. Rock uplift rates from river-profile inversions are 10-25 m/Myr from 45

27 to 22 Ma and 30-55 m/Myr from 22 to 10 Ma. Despite the different time scales, the inverted rates are

28 consistent with ^{10}Be averaged denudation rates (15-20 and 30-40 m/Myr) and river incision values
29 from Pleistocene lava flows (<10 and ~50 m/Myr) for the rejuvenated and relict regions of the WMM.
30 These results agree with geological data and indicate that the observed ~400 m of surface uplift in the
31 WMM started to develop possibly during the early Miocene (first phase). Given the wavelength of
32 the topographic swell forming the topography of the WMM, uplift is here interpreted to reflect
33 localized crustal thickening through magma addition or lithospheric thinning through mantle
34 delamination. More recently, the occurrence of late Miocene marine sediments at ~1200 m of
35 elevation indicates that the adjacent Folded Middle Atlas during the last 5-7 Ma experienced surface
36 uplift at ~170-220 m/Myr. Considering the cumulative amount of surface uplift that varies eastward
37 from 400 to 800 and 1200 m from the Meseta to the Tabular and the Folded Middle Atlas, as well as
38 the spatio-temporal pattern of alkaline volcanism (middle Miocene and Pliocene to Present), we
39 suggest that the most recent episode (second phase) of surface uplift was induced by a larger-scale
40 process that most likely included upwelling of asthenospheric mantle and to a lesser extent crustal
41 shortening in the Folded Middle Atlas.

42

43 **1. Introduction**

44 The competition between crustal deformation, mantle driven uplift and surface processes modulates
45 the growth and the decay of topography across diverse temporal and spatial scales (England and
46 Molnar, 1990; Faccenna and Becker, 2020; Babault et al., 2022). Analytical solutions and numerical
47 models suggest that topography should attain a steady state configuration when uplift is balanced by
48 surface erosion (Whipple, 2001; Willet and Brandon, 2002). Conversely, when such a balance is not
49 respected the landscape should be in disequilibrium conditions (*i.e.*, a transient landscape; *e.g.*,
50 Olivetti et al., 2016; Godard et al., 2019; Clementucci et al., 2022; 2023). Transient landscapes
51 represent the unique possibility to unravel the topographic evolution in areas devoid of stratigraphic
52 constraints since the landscape still contains geomorphological features relative to the pre-

53 perturbation stage. The transience results from different causes, such as base-level falls, drainage
54 reorganizations, climate changes and/or variations in uplift rates at crustal and deeper scales (*e.g.*,
55 Snyder et al., 2002; Roberts and White, 2010; Miller et al., 2013, Ballato et al., 2015; Sembroni et
56 al., 2017; Gallen, 2018; Fernandes et al., 2019; Faccenna and Becker, 2020; Racano et al., 2021;
57 Lanari et al., 2022). Therefore, the topography represents the superimposition of processes operating
58 at multiple timescales and acting at different rates, wavelengths, and amplitudes. Although diverse
59 approaches have been used to unpick these different signals, the combination of quantitative
60 geomorphology with denudation data from cosmogenic nuclides and river profile inversion represents
61 a powerful tool to infer the magnitude, timing, and rates of uplift through time (Miller et al., 2013;
62 Goren et al., 2014; Pastor et al., 2015). This comprehensive approach allows for exploration of the
63 uplift history of areas lacking stratigraphic constraints, with the potential to deconvolve the signals
64 of topographic growth operating at different spatial scales. The surface expression of large-scale uplift
65 and associated causes, in fact, remains often elusive mostly because the rates at which uplift occurs
66 is poorly understood. Accordingly, the integrated approach outlined above can provide fundamental
67 insights into geodynamic processes.

68 In this study, we focus on the Western Moroccan Meseta and the adjacent Tabular Middle Atlas, two
69 tectonically quiescent domains of the Atlas-Meseta system (Fig. 1b). Despite quiescence, these relief
70 features are characterized by elevated, extensive, low-gradient erosional surfaces bounded by steep
71 hillslopes that are adjusting through fluvial incision, to a relative base level drop. This landscape is
72 considered to represent a transient topography resulting from a Neogene phase of large-scale surface
73 uplift, most likely driven by mantle upwelling (*i.e.*, topographic rejuvenation, *e.g.*, Missenard et al.,
74 2006; Babault et al., 2008; Frizon de Lamotte et al., 2009; Miller and Becker, 2014; Clementucci et
75 al., 2023). The relict surfaces are characterized by a protracted phase of hillslope lowering,
76 documented by low temperature thermochronology. Data suggests slow hillslope exhumation rates
77 since the latest Cenozoic (< 28 m/Ma if we consider a maximum cooling of 50°C over the last 70 Ma
78 and a geothermal gradient of 25°C ; see thermal modelling in Barbero et al., 2010 and also the recent

79 review of Charton et al., 2020). These conditions provide a natural setting for investigating the
80 mechanisms of large-scale uplift in tectonically inactive (or slow deforming) areas and quantifying
81 rock uplift changes through time since the relict topography retains features of the pre-uplift event.

82 In this work, we combined denudation rates derived from in situ-produced cosmogenic nuclides
83 (^{10}Be) with regional and basin-scale geomorphic analysis and river profile inversions. Specifically,
84 we determined bedrock outcrops and basin-wide denudation rates from river-borne sediments to
85 gauge them with basin-averaged geomorphic metrics (slope, local relief, and normalized steepness
86 index). In addition, we estimated the magnitude of base-level fall by projecting the upper relict
87 channels toward the present-day river mouth (*e.g.*, Miller et al., 2013; Sembroni et al., 2020). This
88 allowed us to quantify the long-term magnitude of uplift of the Western Moroccan Meseta and the
89 surrounding Tabular and Folded Middle Atlas domains. Moreover, we reconstructed the Cenozoic
90 uplift history of the entire northern Atlas-Meseta system by combining the rock uplift evolution
91 through time inferred from fluvial modelling with available geological data. Our results show
92 consistent rates of denudation and rock uplift of the rejuvenated and the relict portions of the
93 landscape through time and highlight the transient adjustment of hillslope and channels in response
94 to a Cenozoic increase in rock uplift rates. Finally, we retraced back in time the uplift and the
95 topographic evolution of the northern sector of the Atlas-Meseta orographic system and discussed
96 possible driving mechanisms of the topographic rejuvenation.

97

98 **2. Geological setting**

99 The Atlas-Meseta of Morocco (NW Africa) includes mountain ranges (Anti-Atlas, High Atlas and
100 Middle Atlas) elevated tabular areas (Western Meseta, Tabular Middle Atlas and High Plateaus), and
101 sedimentary basins (Missour and Saiss among others; Fig. 1a). The WSW-ENE oriented, 4-km-high
102 High Atlas represents the highest mountain range of the Atlas-Meseta system. It consists mainly of
103 Mesozoic-Cenozoic syn- to post-rift deposits that were uplifted in the Eocene and Miocene (Arboleya

104 et al., 2004; Lanari et al., 2020a). The central sectors of the High Atlas merge towards the NE with
105 the SW-NE-striking Middle Atlas (Fig. 1a). This range is formed by the same deposits of the High
106 Atlas that were deformed most likely from the late Miocene, but with some angular unconformities
107 suggesting a more complex Meso-Cenozoic history of deformation and sedimentation (Charrière,
108 1984). Topographically, the Middle Atlas can be divided into two morpho-structural domains known
109 as the Tabular and Folded Middle Atlas (TMA and FMA; Fig. 1b), which are commonly interpreted
110 as the flank (footwall) and the depocenter (hanging wall) of a Mesozoic rift, respectively (Gomez et
111 al., 1996; 1998). The TMA domain ranges between 1500 and 2000 m of elevation and consists of
112 Mesozoic limestones that are slightly tilted westward. The FMA reaches 3000 m of elevation and
113 includes north-west- to south-east-verging folds mostly detached from the underlying basement along
114 Triassic evaporites (Charrière, 1984; Gomez et al., 1998). The folded range is bounded to the
115 northwest by the North Middle Atlas fault (NMAF) and to the southeast by the South Middle Atlas
116 Fault (SMAF; Fig. 1b). The elevated Western Moroccan Meseta (WMM) and High Plateau (HP) rises
117 to an elevation of 1500 m, without any evidence of Cenozoic tectonic deformation (Gomez et al.,
118 1999; Barbero et al., 2010). Particularly, the WMM is characterized by a complex assemblage of
119 Paleozoic rocks deformed during the Variscan orogeny (Hoepffner et al., 2005; Fig. 1b). This includes
120 Ordovician, Silurian, and Devonian schists and metasedimentary rocks, locally intruded by late
121 Carboniferous granitoids (320 and 270 Ma; Hoepffner et al., 2005 and references therein). This
122 domain is partially covered by undeformed Mesozoic and Eocene marine sedimentary rocks (*e.g.*,
123 Phosphatic series; Salvan, 1954; El Kiram et al., 2019; El Attmani et al., 2021). The Rif Mountains
124 to the north reaches peaks of up to 2500 m and represents a fold and thrust belt that experienced NW-
125 SE shortening due to the convergence of Africa and Eurasia (Jolivet and Faccenna, 2000; Abbassi et
126 al., 2020; Agharroud et al., 2021). There, contractional deformation migrated southward until 8 Ma
127 when the outermost tectonic units were emplaced onto the South Rifian Corridor (Krijgsman et al.,
128 1999; Capella et al., 2018; Abbassi et al., 2020; 2022).

129 The High and Middle Atlas domains are characterized by strain partitioning with a total amount of
130 shortening of ~24%, accommodated by both thrust and oblique/strike-slip faults (Gomez et al., 1999;
131 Lanari et al., 2020b). The amount of shortening across the thrust faults ranges between ~12% and 15
132 % (Missenard et al., 2006; Domenech et al., 2018; Lanari et al., 2020a; 2023) suggesting that crustal
133 thickening is insufficient to isostatically support the modern topography (Arboleya et al., 2004;
134 Missenard et al., 2006). Moreover, geophysical studies highlight the presence of an anomalously thin
135 lithosphere beneath the entire Atlas-Meseta system (Miller and Becker, 2014; Miller et al., 2015).
136 This is associated with continental intraplate volcanism that seems to be coeval with the main tectonic
137 events (Missenard et al., 2006; El Azzouzi et al., 2010; Lanari et al., 2020a). The lavas are well
138 exposed in the FMA, TMA, the WMM over a surface of about 960 km² and have a geochemical
139 affinity with the Canary mantle plume composition ranging from basanite to alkali basalt (Duggen et
140 al., 2009; El Azzouzi et al., 2010). The late Cenozoic phase of uplift is documented by the occurrence
141 of late Miocene (Messinian) shallow-water marine deposits in the Skoura basin at ~1200 m of
142 elevation (Babault et al., 2008) and the progressive shallowing upward trend of the Miocene marine
143 sediments of the Saïss and Guercif basin (Fig. 1a; Krijgsman et al., 1999).

144

145 **3. Methods**

146 **3.1. Topographic, stream analysis and long-term incision**

147 A Shuttle Radar Topography Mission Digital Elevation Model (SRTM DEM, pixel size of 30 m) was
148 used to extract the drainage network. Hillslope metrics, such as slope and local relief with a circular
149 sampling window of 2.5 km radius, as well as channel and basin metrics were extracted using ArcGIS
150 and the MATLAB software packages TopoToolbox (Schwanghart and Scherler, 2014) and
151 Topographic Analysis Kit (TAK, Forte and Whipple, 2019), together with additional MATLAB
152 functions (*e.g.*, Gallen, 2017). The relationships between river channel steepness and erosion/uplift
153 rates have been demonstrated in various geological settings (Kirby and Whipple, 2012). In

154 detachment-limited conditions, these relationships are modulated by the stream-power model, which
155 can be integrated, assuming spatially invariant uplift and erodibility, to estimate the integral quantity
156 χ (details in the section S1; Perron and Royden, 2013). This allows transforming river longitudinal
157 profiles to χ versus elevation (z) plots (here and thereafter defined as χ - z plot). Consequently, the
158 slope in χ space is defined as the steepness index (k_s) when the reference area (A_0) is equal to 1. To
159 compare k_s values of longitudinal profiles, a reference concavity index (θ_{ref}) of 0.45 is applied to
160 derive the normalized channel steepness indices (k_{sn} ; Kirby and Whipple, 2012 and references
161 therein). The θ_{ref} was calculated by minimising the scatter in χ space of all rivers draining the study
162 area (Perron and Royden, 2013; Goren et al., 2014; Fig. S1). Our θ_{ref} coincides with the value typically
163 used to derive normalised channel steepness (Snyder et al., 2000; Kirby and Whipple, 2012).

164 Knickpoints were identified and discretized based on the shape of the streams in the longitudinal
165 profiles and the χ - z plots and using available geological maps (Service Géologique du Maroc) and
166 satellite imagery (Google Earth). Importantly, non-lithological knickpoints cluster around similar χ
167 values and separate stream segments with different k_{sn} values (Perron and Royden, 2013;
168 Clementucci, 2022). Using this approach, the knickpoints were divided into non-lithological, minor,
169 lithological and man-made dam. Streams with knickpoints identified as transient (non-lithological)
170 were used for reconstructing river projections to estimate the magnitude of long-term incision. Such
171 profiles can be reconstructed using the k_{sn} values ($\theta_{ref} = 0.45$) of the river segments upstream of the
172 non-lithological knickpoints in the χ space, assuming that the paleo-river network remained
173 approximately the same through time (Perron and Royden, 2013; Smith et al., 2022). Full details of
174 equations and stream analysis are provided in the Supplementary material.

175

176 **3.2. ^{10}Be -derived denudation rates**

177 We measured the concentration of in situ-produced cosmogenic ^{10}Be in quartz from seven river-borne
178 samples collected in active channels and from four bedrock outcrops (Fig. 1c; Table 1). The only
179 lithologies exposed in the WMM that contain quartz are Paleozoic basement rocks such as schists,
180 metasedimentary rocks, and granites. Our sampled catchments cover most of this area (Figs 1b and
181 S2), from the elevated relict to the steeper landscape portions. In one case we sampled a catchment
182 with Paleozoic rocks at the periphery of the uplifted area (sample 53) to compare denudation rates
183 from different sectors of the WMM (Fig 1c). The basin-wide denudation rates were determined only
184 from quartz-bearing rocks, which usually represent 50 to 100% of the total area of the sampled
185 catchments (Table S1). The portions of the sampled catchments with Mesozoic limestones were not
186 considered in the scaling modelling (Clementucci et al., 2022 and references therein). Details of
187 sample preparation, from the initial mineral separation to the final conversion of in situ-produced
188 ^{10}Be to basin-wide denudation rates are provided in the Supplementary material S2. Catchments 58,
189 59, 60 and 62 are sub-catchments of two larger basins named 61 and 63, while catchment 53
190 corresponds to the westernmost sampling location (Fig. 1c). The upstream areas range between 30
191 and 100 km² and 320 and 350 km² for the sub-catchments and the larger basins, respectively. In
192 particular, we selected samples 58 and 59 upstream of the low-relief surfaces and samples 61 and 63
193 downstream of the relict landscape to extract the denudation rates of the adjusted and steeper
194 topography. Catchments 53, 58, 59, 62 and 63 are characterized by uniform quartz-bearing
195 lithologies. Conversely, the contribution of quartz, in catchments 60 and 61, is mostly from two
196 granite massifs exposed in the highest topography of the Moroccan Meseta region. To quantify the
197 local denudation rates of the granite massifs forming the landscape of the western Meseta, we sampled
198 four bedrock outcrops: two from the summit of an erosional surface (samples 56A, 56B) and two
199 from sub-horizontal ridge crests located close to the margin of the relict topography (samples 57A,
200 57B, Table 1 and Fig. 1c).

201 In situ-produced ^{10}Be concentrations were converted to basin-wide denudation rates and bedrock
202 erosion rates using the MATLAB scripts of Balco et al. (2008), which are based on the time-

203 dependent scaling functions of Stone (2000)'s scheme that consider three different categories of
204 particles responsible for the in-situ production of ^{10}Be (Braucher et al., 2011). The geomorphic
205 shielding (Dunne et al., 1999), scaling factor of Stone (2000) and the mean elevation of sampled
206 catchments were determined for each pixel of the digital elevation model (DEM), considering only
207 the areas contributing with quartz-bearing lithologies. In the scaling model, λ describes the
208 radioactive decay constant of ^{10}Be . Denudation rates from ^{10}Be concentrations were calculated using
209 a ^{10}Be sea level high latitude production rate of 4.02 ± 0.33 atoms ^{10}Be (g-SiO₂)/yr (Borchers et al.,
210 2016), and half-lives of $(1.387 \pm 0.012) \times 10^6$ years (Chmeleff et al., 2010; Korschinek et al., 2010).
211 At cosmogenic steady-state, the cosmogenic ^{10}Be in river-borne quartz mineral records a time
212 integrated spatially averaged denudation rate, which represents fluviially mixed denudation products
213 of all the processes occurring in each watershed (Granger et al., 1996). This integration occurs over
214 the time needed to remove one attenuation length (*e.g.*, roughly 60 cm; Lal, 1991).

215 We also applied a sediment mixing correction to estimate a value of denudation for the downstream
216 knickpoint portion of the catchment 63 (basin 64), knowing the denudation and drainage area of the
217 whole catchment and the upper sub-catchment, upstream of the non-lithological knickpoints (Fig. 2;
218 according to Granger et al., 1996; Miller et al., 2013; Rosenkranz et al., 2018):

$$219 \quad D_{1-2} = \frac{D_1 A_1 - D_2 A_2}{A_1 - A_2} \quad (1)$$

220 where $A_1 - D_1$ and $A_2 - D_2$ are the drainage area and basin-wide denudation rate of the whole
221 catchment and above the non-lithological knickpoint, respectively. D_{1-2} is the basin-wide denudation
222 of the adjusted topography downstream of non-lithological knickpoints.

223

224 **3.3. Predicted ^{10}Be -derived denudations and incision of volcanic bedrock**

225 The basin-averaged k_{sn} values, precipitation rates (IMERG data; Huffman et al., 2015), slope and
226 local relief for the portions of the sampled catchments that include only quartz-bearing lithologies
227 were extracted for comparing ^{10}Be -derived denudation rates. The comparison between denudation
228 rates and k_{sn} allowed extracting the averaged erodibility parameter K for basement rocks in the
229 Western Meseta and constraining n equal to 1. We also evaluate the power best-fit to better explore
230 the relationship between channel steepness and erosion, and evaluate the power-law coefficients, K
231 and n . Finally, values of k_{sn} values were directly converted to denudation rates (see details in Adams
232 et al., 2020, and Clementucci et al., 2022).

233 To compare our inferred and measured, short-term denudation rates, with longer-term rates we
234 estimated fluvial incision rates by using the 1-Myr-old lava flows as a temporal and geometrical
235 benchmark because they were emplaced over the relict and the rejuvenated landscape of the WMM
236 domain (Tephrite and Basanite data from Rachidi, 1995).

237

238 **3.4. Rock uplift history**

239 To unravel the rock uplift rates over time, we performed a linear inversion of the rivers draining the
240 Paleozoic basement in the Moroccan Meseta domain. We constrained the averaged erodibility value
241 (K) of the Paleozoic basement rocks, using the stream power river incision model ($K = E/k_{sn}^n$) for n
242 = 1. The K was estimated using the forced-origin linear regression in the plot k_{sn} and denudation
243 rates. Although the variation of K over time is difficult to constrain, our averaged estimate can be
244 considered reliable because short-term denudation rates are consistent with the long-term
245 exhumation rates derived from thermochronological data over other quiescent sectors of the Atlas-
246 Meseta system (e.g., Anti-Atlas and Haouz basin; Lanari et al, 2020a; Clementucci et al., 2022;
247 Western Meseta; Barbero et al., 2010). Assuming block uplift conditions in a selected catchment
248 (no spatial uplift variations) and $n = 1$, the travel time of a perturbation along a river profile (τ) is
249 given by:

250

251

$$\tau = \int_{x_b}^x \frac{dx'}{K A_0^m A(x')^m} \quad (2)$$

252

253

254

255

By removing K from Equation (1) we obtain χ . Following the approach of Goren et al. (2014), a non-dimensional uplift can be inferred from the integration of the stream-power model by assuming that trunks with same χ share a common uplift history and organizing the equation in a matrix form for each data point:

256

$$A^*U^* = z \quad (3)$$

257

258

259

where A^* is an $N \times q$ matrix, q is the number of χ intervals, U^* is the non-dimensional uplift and z is elevation. This is an overdetermined inverse problem, as there are more known data points than unknown parameters. As such, a least-squares estimate for U^* is used (Goren et al., 2014):

260

$$U^* = U^*_{pri} + (A^{*T}A^* + \Gamma^2I)^{-1} \circ^{*T} (z - A^*U^*_{pri}) \quad (4)$$

261

262

263

264

Where Γ is a dampening coefficient that determines the smoothness imposed on the solution, I is the $q \times q$ identity matrix and U^*_{pri} represents the prior guess for U^* , estimated by taking the average slope of the χ -plot (Goren et al., 2014). In this way, from the integration of the stream-power model, χ and U^* can be scaled for K to infer rock uplift rates (U) and time (τ):

265

$$U = K A_0^m U^* \quad (5)$$

266

$$\tau = \frac{\chi}{K A_0^m} \quad (6)$$

267

268

269

270

271

272

273

274

To reduce the noise, we removed streams with a length lower than 10 km. The linear inversion approach is built on the assumption of spatially uniform uplift, which to a first approximation, can be considered reasonable given that the analysed catchments do not cross any fault that may have been active during the late Cenozoic. Moreover, the analysed catchments are entirely in the WWM where they occupy relatively narrow areas, and hence do not record major along-strike variations in uplift rates. Thus, adopting this approach, by combining the inversions from different catchments, it is also possible to investigate spatial variations in rock-uplift histories (Pavano et al., 2021; Racano et al., 2021; Zhong et al., 2022).

275

276

277 4. Results

278 4.1. Topographic and river analysis

279 The WMM domain consists of a NE-SW striking, ~ 130 x 260 km topographic swell, with a maximum
280 elevation of ~1500 m, and a low relief topography that locally has been deeply dissected by fluvial
281 incision (Figs. 3 and 4). Within the WMM, a series of smaller scale geomorphic features such as
282 summit erosional surfaces, ridge crests and fluvial terraces developed onto and inset into the lava
283 deposits are visible (Figs. 4 and 5). The TMA presents a tabular, low-relief topography while the
284 FMA is characterized by an alternation of steep ridges and wide, low-slope valleys (Figs. 3 and S3).
285 Based on the geometry of the river network, the study area was divided into the Central-Western
286 Meseta (CWM), the North-Western Meseta (NWM), and the South-Western Meseta (SWM). The
287 CWM is characterized by rivers draining directly into the Atlantic Ocean, while the rivers of the
288 NWM and SWM represent the main respective tributaries of the Atlantic draining Oued Sebou and
289 Oum Er Rbia, (Fig. 3a). A set of regionally widespread non-lithological knickpoints occur at different
290 elevations in the three domains. These features mark a sharp variation in the k_{sn} , local relief and slope
291 between the deep valleys and the upstream landscape (dashed line in Figs. 3a, 3b and S3). Particularly,
292 high values of k_{sn} ($> 80 \text{ m}^{0.9}$; Fig. 3b), slope ($> 25^\circ$; Fig. S3a), and local relief ($> 500 \text{ m}$; Fig. S3b),
293 are found in the FMA domain and in the downstream portions of the non-lithological knickpoints of
294 the WMM and the TMA domains. The lowest values of topographic and channel parameters are
295 observed in the lower portions of the landscape (Oum Er Rbia and Sebou basins) and upstream of the
296 non-lithological knickpoints (Figs. 3 and S3).

297 The NWM sector is characterized by transient longitudinal profiles. Most of the rivers flowing from
298 the TMA and the WMM present lithological knickpoints between 1500 and 2000 m of elevation at χ
299 values of 20 to 25 (Fig. 6a and 6b). A pattern of widespread non-lithological knickpoints is located

300 between 700 and 1500 m and is associated with a gradual increase in elevation from 200 to 450 km
301 of distance from the river mouth (Fig. 6a and Table S2). These non-lithological knickpoints cluster
302 at χ values of 13 – 17 and separate steep downstream segments from low-gradient upstream channels.
303 A cluster of minor knickpoints is located at lower elevations and separates gentle downstream river
304 segments from steeper upstream portions (Figs. 6a and 6b).
305 Rivers of the CWM sector show concave-up longitudinal profiles downstream of the non-lithological
306 knickpoints and straight low-gradient upper segments (Fig. 6c). These non-lithological knickpoints
307 present a progressive increase in elevation from 300 m close to the river mouth to ~1000 m upstream
308 toward the TMA domain. Most of these non-lithological knickpoints cluster at χ values of 10 – 15,
309 and separate downstream steep segments from low-gradient upstream portions (Fig. 6d). Downstream
310 of these non-lithological knickpoints, the river segments are characterized by gentle low-gradient
311 river profiles, which contain several minor knickpoints (Fig. 6c).
312 The SWM sector is characterized by both equilibrated and transient longitudinal profiles. The main
313 slope break is located at high elevation, where two non-lithological knickpoints lie in the FMA
314 domain (~1900 m and ~1500 m; Fig. 6e and 6f). The average elevations of the non-lithological
315 knickpoints are 627 ($\sigma = 116$) m, 1255 ($\sigma = 86$) m and 1724 ($\sigma = 135$) m for the WMM, TMA and
316 FMA domains, respectively (Table S2).

317

318 **4.2. Long-term river incision**

319 The magnitude of long-term river incision can be estimated by the difference between the
320 reconstructed river profiles projected from the non-lithological knickpoints and the modern base
321 level. Rivers draining the WMM domain (mainly from CWM and NWM sectors) show a mean long-
322 term incision of 415 ($2\sigma = 55$) m, based on 33 river projections (Fig. 7a and Table S3). The mean
323 values of long-term incision of rivers draining the TMA domain is 823 ($2\sigma = 66$) m, based on 8 river
324 projections (Fig. 7a and Table S3). The river projections from the FMA domain show an average
325 long-term incision of 1086 ($2\sigma = 64$) m, although based only on 2 rivers (Fig. 7a and Table S3).

326 There, rivers flow along the SWM sector. Overall, both elevations of the non-lithological knickpoints
327 and magnitude of long-term incision increase progressively from the WMM to the TMA and FMA
328 domains (Fig. 7b).

329 A few valleys of the WWM have been filled by Quaternary lava flows downstream and upstream of
330 major non-lithological knickpoints (Fig. 1a). Subsequently, these lava flows were dissected through
331 fluvial incision leaving well developed terraces. Downstream of major non-lithological knickpoints,
332 the current valley floor is ~ 50 and ~ 100 m below the base and the top of the lava flows, respectively
333 (Figs. 4 and 5b). Considering that the lava yielded radiometric ages of 1 ± 0.1 Ma (Rachdi, 1995),
334 fluvial incision rates vary from $\sim 100 \pm 10$ to $\sim 50 \pm 5$ m/Myr, respectively. Importantly, the latter
335 estimate ($\sim 50 \pm 5$ m/Myr) represents the amount of river channel lowering after the lava emplacement
336 using the unconformity as a reference surface (strath of the terrace). Conversely, in the relict surfaces
337 of the WMM domain the lava flows are poorly incised and the fluvial incision rate are <10 m/Myr
338 (Fig. 4).

339

340 **4.3. ^{10}Be -derived denudation rates**

341 ^{10}Be concentrations from the largest catchments of the WMM yield denudation rates of 19.8 ± 1.4
342 and 32.2 ± 2.3 m/Myr (samples 61 and 63; Fig. 1c and Table 1). Denudation rates from the sub-
343 catchments located upstream of the relict landscape are 17.1 ± 1.3 and 19.5 ± 1.4 m/Myr (samples 58
344 and 59), whereas the sub-catchments downstream of non-lithological knickpoints present rates of
345 17.4 ± 1.2 and 40.0 ± 5.6 m/Myr (samples 60 and 62; Fig. 1c and Table 1). In the western side of the
346 Moroccan Meseta, sample 53 yields a denudation rate of 14.8 ± 1.0 m/Myr. The unmixed rate from
347 catchment 64 (*i.e.*, the denudation rate of the incised portion of the landscape of catchment 63, which
348 was obtained by subtracting the contribution of the relict landscape of sub-catchment 59 and the area
349 of the steep sub-catchment 62 downstream of non-lithological knickpoints; Fig. 2), yields a
350 denudation rate of 35.2 ± 2.6 m/Myr. This shows a good match with the estimates of denudation from

351 the same sub-catchment 62 (Fig. 2 and Table S1). The sampled catchments are characterized by a
352 uniform distribution of quartz, which is mostly sourced from the lower Paleozoic metamorphic
353 basement, except for samples 60 and 61. These two basins are influenced by the occurrence of two
354 large granite outcrops located in the high-standing low relief surface of the WMM (Fig. 4), which
355 most likely provided more quartz than the Paleozoic metamorphic rocks. Local denudation rates of
356 bare bedrock surfaces on the summit plateau (relict surface) and on the ridge crests of the granite
357 massifs vary from 3.5 ± 0.2 to 4.1 ± 0.3 m/Myr (samples 56A, 56B), and from 13.1 ± 0.9 to 14.4 ± 1.0
358 m/Myr (samples 57A, 57B), respectively (Table 1). It is worth noting that the denudation rates of the
359 granitic ridge crests: 1) are significantly higher than the rates of the flat summit surfaces, and 2) are
360 close to the values determined for the watersheds of samples 60 and 61, where the two granite massifs
361 are exposed and may provide most of the quartz that yielded the measured ^{10}Be concentration.

362

363 **4.4. Basin-wide denudation rates versus topographic metrics**

364 The basin-averaged k_{sn} values range from 16.7 to 53.5 $\text{m}^{0.9}$ (Fig. 8a and Table S1), while basin-
365 averaged slope values vary from 6 to 13° (Fig. 8b and Table S1) and the basin-averaged local relief
366 range from 222 to 440 m (Fig. 8c and Table S1). The topographic and channel metrics exhibit a
367 medium to low correlation with the basin-wide denudation rates with R^2 of 0.13, 0.61 and 0.71 for
368 k_{sn} , slope and local relief, respectively (dashed lines in Fig. 8a, 8b, 8c). However, a stronger
369 correlation between the topographic metrics and basin-wide denudation rates is obtained considering
370 only the catchments characterized by a homogeneous distribution of bearing-quartz lithologies of the
371 Paleozoic basement (*i.e.*, excluding samples 60 and 61, which quartz is mostly sourced from the
372 granites above the elevated flat surfaces). In this case, the coefficient of correlation increases to 0.76,
373 0.87 and 0.97 for the basin-averaged k_{sn} values, slope, and local relief, respectively (Fig. 8a, 8b, 8c).
374 The lowest topographic and channel metrics are observed for samples 53, 58 and 59, which represent
375 the westernmost sample (53) and the samples upstream of the non-lithological knickpoints.

376 Conversely, samples 62 and 64 (downstream of the non-lithological knickpoints) show the highest
377 values of basin-averaged k_{sn} values, slope and local relief, whereas samples 60, 61 and 63, show
378 slightly lower values, because they represent the mixed estimates, giving that both the upstream and
379 downstream portions of the catchments are contributing (Fig. 8a, 8b, 8c and Table S1). Finally, basin-
380 wide denudation rates are poorly correlated with the basin-averaged precipitation ($R^2=0.38$; Fig. 8d).

381

382 **4.5. Timing: base-level fall history**

383 To constrain the base level history, the linear inversion of six river profiles has been performed using
384 $n = 1$, $m/n = 0.45$ and K of $7.34 \pm 0.58 \times 10^{-7} \text{ m}^{0.1}/\text{yr}$. The parameter $n = 1$ describes the linear
385 relationship between the erosion rates and k_{sn} (Fig. 8a), as expected for inactive to slow tectonic
386 settings (Godard et al., 2019; Clementucci et al., 2022). The estimation of n has been made testing
387 the stream-power law fits, confirming that the assumption of $n = 1$ is reasonable (Fig. 8a). The six
388 catchments were selected as they flow over the same rocks from which the K value was estimated
389 (Figs. 9a and S5, S6). Results show similar rock uplift histories (Figs. S5 and S6), which are
390 summarized in Figure 9b with the mean and the standard error and deviation of the six inversions.
391 Overall, we observe almost constant rock uplift rates of 15-20 m/Myr from 30-33 until ~ 25 Ma,
392 followed by an increase up to ~ 40 m/Myr between 20 and 15 Ma and by a general decrease to ~ 25
393 m/Myr during the last 10 Ma (Fig. 9b). Importantly, within this decreasing trend, a peak in rock-uplift
394 occurs around 5-6 Ma with values of up to 35 m/Myr despite the large standard deviation.

395

396 **5. Discussion**

397 **5.1. Transient topography in Western Moroccan Meseta and Middle Atlas**

398 The study area is characterized by high-standing relict surfaces bounded by non-lithological
399 knickpoints and deeply incised valleys downstream of these knickpoints. This elevated landscape is
400 characterized by low k_{sn} and include low-topographic slopes and local relief surfaces carved onto
401 Paleozoic metamorphic rocks and Mesozoic limestones that gently dip toward the western sectors
402 (Figs. 3, 4, 5; Barbero et al., 2010; Barcos et al., 2014). A few lithological knickpoints lie at about
403 1500 m of elevation at the boundary between the WMM and TMA domains (see bold yellow points
404 in Fig. 6a, 6b). These knickpoints are marked by a local variation in k_{sn} values and other topographic
405 metrics (slope and local relief) and coincide with a ~500-m-high morphological escarpment
406 representing the stratigraphic contact between the Paleozoic basement and the undeformed Jurassic
407 limestones, reflecting a sharp change in rock erodibility. Conversely, downstream of non-lithological
408 knickpoints the landscape is characterized by steep slopes and k_{sn} values at regional scale (Figs. 3, 4,
409 5 and S2). This topographic dichotomy suggests that the landscape and hence the river network is not
410 in equilibrium (Crosby and Whipple, 2006; Kirby and Whipple, 2012).

411 The disequilibrium condition can be explained through several processes, such as drainage
412 reorganization, climate change, eustatic sea-level fall and tectonic uplift (Snyder et al., 2002; Crosby
413 and Whipple, 2006; Kirby and Whipple, 2012; Miller et al., 2013; Boulton et al., 2014; Ballato et al.,
414 2015; Rixhon et al., 2017; Stokes et al., 2017). In the WMM, a base-level fall due to eustatic forcing
415 is unlikely to be the source of the observed non-lithological knickpoints, because it cannot explain
416 the pattern of k_{sn} values of the river network. Specifically, a wave of knickpoints due to eustatic causes
417 cannot exhibit any major slope break in the chi-space because it is not associated with variations in
418 uplift rate or rock erodibility (Wobus et al., 2006; Kirby and Whipple, 2012; Perron and Royden,
419 2013). Moreover, there is no geological data documenting a major and protracted sea level drop
420 during the late Cenozoic (Miller et al., 2020) and the glacial-interglacial cycles occurred over the last
421 1 Ma would result in multiple phases of incision and filling of the coastal area (*e.g.*, Marra et al.,
422 2016), rather than a wave of incision reaching the upper reaches of the Meseta. Drainage
423 reorganization could explain the increase in long-term fluvial incision downstream of the knickpoints

424 due to a gain in drainage area and hence in water discharge (*e.g.*, Yang et al., 2015). However, the
425 non-lithological knickpoints are observed in the watersheds of all domains (WMM, TMA and FMA;
426 Fig. 3), even in small catchments where there is no evidence of drainage reorganization (*e.g.*, wind
427 gaps, knickpoints due to capture processes among others), as instead observed in the Moulouya basin
428 to the east of FMA (Bartz et al., 2018). Furthermore, increased erosion due to enhanced of average
429 precipitation can also be discarded because it should produce a decrease in the channel slopes
430 downstream of non-lithological knickpoints (Miller et al., 2013; Marder and Gallen, 2023; Figs. 3
431 and S2). Importantly, the non-lithological knickpoints are characterized by a major upstream decrease
432 in k_{sn} values even across the same lithology as documented in the χ space plots (Fig. 6) and by changes
433 in the topographic slope and local relief at the regional scale (Figs. 3 and S2). This indicates that the
434 knickpoints separate portions of the landscape eroding at different rates (*e.g.*, Crosby and Whipple,
435 2006; Kirby and Whipple, 2012; Boulton et al., 2014; Olivetti et al., 2016). Consequently, the portion
436 with steep channels downstream of the knickpoints records the response of the fluvial system to an
437 increase in rock uplift rates, while the upstream low relief topography (relict landscape; Fig. 4)
438 records the pre-uplift conditions, as also observed in the others domains of the Atlas-Meseta system
439 (*e.g.*, High Atlas, Siroua and Anti-Atlas range; Boulton et al., 2014; Zondervan et al., 2020;
440 Clementucci et al., 2023). Overall, these observations indicate that the study area experienced a recent
441 topographic rejuvenation (regional, large-scale surface uplift) as documented in other regions
442 worldwide exhibiting similar topographic patterns (Calvet et al., 2015; Miller et al., 2013).

443 Conversely, minor knickpoints lying at lower elevations can be attributed to Quaternary climate
444 change or a second increase in uplift rates (Hancock and Kirwan, 2007; Molnar and England, 1990).
445 A similar pattern of minor knickpoints has been described also for the Anti-Atlas domain, where their
446 origin is still enigmatic (Clementucci et al., 2023). In the Western Meseta, these less elevated
447 knickpoints do not present a pronounced break point along the river profiles, but they are evenly
448 distributed along the rivers. Moreover, an increase in rock-uplift is shown around 5-6 Ma in the river-

449 profile inversions. These observations suggest a possible upward knickpoint migration following a
450 second topographic rejuvenation event (*i.e.*, a second increase in uplift rates at 5-6 Ma), which most
451 likely occurred at a larger wavelength from the WMM to the TMA and the FMA (and possibly the
452 HP), as testified by uplifted Messinian marine sediments of the Skoura and Saiss basins (Babault et
453 al., 2008).

454

455 **5.2. Erosional dynamics in the rejuvenated Western Moroccan Meseta**

456 The characterization of a transient landscape is critical for understanding the dynamics of a landscape
457 that includes the coexistence of portions that are in equilibrium with past and present uplift rates. In
458 the Western Meseta the lowest bedrock denudation rates are from granite exposed on the tops of the
459 summit surfaces (3.5 ± 0.2 and 4.1 ± 0.3 m/Myr) while the highest rates are from a similar granite
460 exposed along small ridge crests (13.1 ± 0.9 to 14.4 ± 1.0 m/Myr in Table 1). This four-fold difference
461 is presumably caused by different hillslope processes and higher physical and chemical weathering
462 assuming the same exposure time (Riebe et al., 2004). This demonstrates the key role of hillslope
463 topographic gradients and weathering processes in controlling denudation rates from bedrocks (*e.g.*,
464 Meyer et al., 2010; Godard et al., 2019). In our case, the highest denudation rates occur on the ridge
465 morphologies, at the margin of relict landscape, suggesting an abrupt increase in denudation and
466 hillslope erosion at the transition between the rejuvenated and relict topography, indicating a strong
467 control of the geomorphic domain on the ^{10}Be -derived denudation (Figs. 4 and 5).

468 Interestingly, the rates of denudation from the ridge crests (samples 57A and 57B) are very similar to
469 those from the watersheds of the same areas (samples 60 and 61). This may be due to the widespread
470 exposure within the catchments of the relict surface sculptured onto the granite rocks that contains a
471 higher amount of quartz than the adjacent metamorphic rocks. These quartz-rich granite rocks erode
472 at relatively low rates (~ 4 m/Myr) with respect to the basin-wide denudation rates from the

473 catchments composed of metamorphic rocks ($> \sim 17$ m/Myr) and hence have a higher ^{10}Be
474 concentration than the metamorphic basement. The greater amount of quartz sourced from the relict
475 landscape appears to increase the final ^{10}Be concentration for catchments 60 and 61, producing a
476 decrease in denudation rate estimates. This process is further corroborated by the poor sensitivity of
477 these two watersheds to the linearity between basin-wide denudation rates and topographic metrics
478 that are systematically lower than expected (Fig. 8). Low denudation rates are also observed in
479 watersheds 58 and 59, located upstream of non-lithological knickpoints within the relict portion of
480 the WMM landscape where values of basin-averaged k_{sn} , slope and relief are low. These low
481 denudation rates are consistent with those inferred from the thermochronological data (Barbero et al.,
482 2010) averaged over the last 70 Ma and reflect the erosional decay of topography during a long period
483 of relative base level stability (Calvet et al., 2015). Conversely, watersheds 62 and 63 present higher
484 denudation rates because they receive most of the quartz from the incised landscape downstream of
485 non-lithological knickpoints since the distribution of quartz-bearing rocks is uniform. In this case
486 higher rates reflect the increase in uplift rates as documented in other regions that experienced a
487 topographic rejuvenation (e.g., Olivetti et al., 2016; Miller et al., 2013; Clementucci et al., 2022).

488 In slow tectonic settings, there is usually a poor correlation between basin-averaged k_{sn} values and
489 denudation rates, and this is generally attributed either to the lithological control on the landscape
490 dynamics or to a decoupling between hillslope and channel incision (e.g., Olivetti et al., 2016; Peifer
491 et al., 2021; Clementucci et al., 2022). Our data show a good correlation between denudation rates
492 and topographic metrics, despite the high degree of transience in the landscape, suggesting that the
493 landscape is not affected by any lithological control and/or decoupling process, except for catchments
494 60 and 61. In particular, the correlation between basin-wide denudation and k_{sn} suggests that the n
495 parameter of the stream power model can be approximated to 1, as typically observed in slow tectonic
496 settings, where basin-averaged slope and k_{sn} are usually lower than 25° and $200 \text{ m}^{0.9}$, respectively
497 (Fig. S4). Finally, the basin-wide denudation rates are insensitive to averaged annual precipitation
498 rates (Fig. 8d), testifying the lack of a clear dependency of denudation processes to precipitation, as

499 also observed in other slow tectonic ranges characterized by different climatic conditions (Fig. S4;
500 Burbank and Anderson, 2003; Godard et al., 2019; Clementucci et al., 2022).

501

502 **5.3. Topographic evolution and uplift history**

503 The axial zone of the WMM was likely above sea level before the Lutetian (early Cenozoic / middle
504 Eocene) as suggested by the lack of Eocene sediments (Phosphate Plateau), which can be found only
505 along its periphery (Er Rbia Basin) where the phosphatic series shows a deepening and thickening
506 south-westward trend (Fig. 9a; El Kiram et al., 2019). This sedimentation pattern indicates that, while
507 western Morocco was locally subjected to Eocene subsidence and marine sedimentation, the WMM
508 was affected by different processes (Charton et al., 2020). In addition, ~ 200- to 270-Myr-old apatite
509 fission track cooling ages and associated thermal histories indicate limited vertical movements since
510 the Triassic and suggest that the relict landscape might have started to form even earlier than the
511 Lutetian, possibly from the latest Cretaceous (Barbero et al., 2010; Charton et al., 2020). Our inverted
512 rock uplift rates for the oldest resolved time interval (35-22 Ma) of 10-20 m/Myr are in the same
513 order of magnitude as the ¹⁰Be-derived denudation rates averaged over 50 ka (Table 1) and the lava
514 incision rates averaged over the last 1 Ma (see U₁ in Fig. 10a), both for the relict landscape, and the
515 exhumation rates averaged over the last 70 Ma. The occurrence of similar modelled rock uplift and
516 denudation/incision rates across different time scales, could indicate the persistence of a long-term
517 balance between rock uplift and denudation (*i.e.*, steady state denudation; *e.g.*, Whipple, 2001; Willett
518 and Brandon, 2002). However, this observation is based on the overall patterns of denudation at
519 different time scales and does not consider short-term climate fluctuations, which usually precludes
520 the attainment of steady-state conditions over time, possibly triggering rapid changes in denudation
521 rates (Whipple, 2001; Champagnac et al., 2014).

522 Our fluvial inversion results indicate that in the WMM domain, the onset of uplift and associated
523 headward migration of the knickpoints likely started at about 22 Ma and continued until ~10 Ma, with

524 rates ranging from 30 to 50 m/Myr (see U_2 in Fig. 10b). This range of rock uplift rates falls also within
525 the range of ^{10}Be -derived denudation rates and lava flow incision rates downstream of major non-
526 lithological knickpoints ($E_2 = 35\text{-}55$ m/Myr, Figs. 9a and 10b). Barcos et al. (2014) described a
527 maturity stage of the Sebou Basin through a stream length-gradient analysis, which can be a sign of
528 the partial re-equilibrium of these basins since this early Miocene perturbation (Fig. 10b). These
529 portions of the landscape are still eroding at rates that could be representative of the uplift rate U_2
530 (Fig. 10b). Thus, our inverted rates suggest that the entire WMM started to be uplifted from 22-20
531 Ma, in agreement with the occurrence of uplifted Eocene marine deposits in the WMM and in the
532 axial zone of the FMA and coeval with the onset of Miocene alkaline volcanism (Fig. 10b; Salvan,
533 1954; El Kiram et al., 2019; El Attmani et al., 2021). Early Miocene uplift initiation could be also
534 coeval with the onset of widespread contractional deformation in the FMA (Gomez et al., 1996;
535 1998), in the High Atlas Mountains (El Harfi et al., 2001; Lanari et al., 2020a; 2020b) and in the Rif
536 mountains (Fig. 10b; Abbassi et al., 2020). At the same time, the southward migration of the Rifian
537 frontal thrust produced foreland subsidence with the deposition of unconformable marine deposits in
538 the South Rifian Corridor (Saiss and Skoura Basins; Fig. 10b; Capella et al., 2018). Particularly, the
539 Skoura Basin contains a Messinian shallow-water marine sequence possibly deposited in a gulf
540 bounded by the TMA and FMA (Babault et al., 2008; Capella et al., 2018). In the northern Middle
541 Atlas and the Rif Belt, uplift started sometime after 5-7 Ma (Charriere and Saint-Martin, 1989;
542 Babault et al., 2008) at rates of 170-220 m/Myr as testified by the modern elevation of the Messinian
543 deposits of the Skoura Basin and the closure of the marine South Rifian Corridor (Krijgsman et al.,
544 1999; Babault et al., 2008). This episode of more recent regional uplift was accompanied by a Plio-
545 Quaternary peak in volcanism (Fig. 1) and is also registered by the Western Meseta fluvial network
546 (Fig. 10c). There, our inferred rock uplift rates for the last 5-6 Ma increase up to 35 m/Myr (see U_3
547 in Fig. 10c). Thus, our modelling rates appear to document a two-stage uplift history of the WMM
548 that includes uplift initiation at 22-20 Ma (with maximum rates reached at about 15 Ma) followed by
549 a second pulse at 5-6 Ma.

550 Alternatively, the WMM, could have shared an uplift history like that one recorded by the uplifted
551 marine sediment of the Skoura and Saiss Basins (Babault et al., 2008). This implies that the highest
552 knickpoints bounding the elevated relict landscape were originated during the regional 5- to 7-My-
553 old uplift phase (Babault et al., 2008). Although our linear inversion model is calibrated for ^{10}Be data,
554 which integrates erosion rates for the last tens 50.000 years, there are strong indications that short-
555 and long-term, thermochronologically derived, exhumation rates from the relict landscape for the
556 Cenozoic are similar (Barbero et al., 2010). This is also valid for the denudation rates downstream of
557 the most elevated knickpoints, which are consistent with the incision rates inferred from the lava
558 flows over the last 1 Ma. Thus, we can conclude that our inversion is likely not biased by the time
559 scale of the ^{10}Be rates. Moreover, if we apply the approach of profile reconstruction for estimating
560 the timing of the increase in uplift rates, for a surface uplift of 415 \pm 55 m (Fig. 7), an erosion rate
561 for the relict landscape up to 20 m/Ma and an uplift (equal to erosion) rate downstream of the
562 knickpoint of 40-50 m/Ma (Fig. 8) we would expect an minimum onset of uplift between 13 and 20
563 Ma (see equation in figure 5 of Kirby and Whipple 2012). These estimates indicate that the uplift
564 rates downstream of the relict landscape are too low to be consistent with a Messinian onset of surface
565 uplift. Thus, the uplift in WMM should have started earlier than 7 Ma. Taken together, these
566 observations suggest that the WMM and the northern sectors of the FMA, where the Skoura Basin is
567 located, did not share the same uplift history pre 7 Ma. Consequently, a two-stage uplift history for
568 the WMM appears to be the most likely scenario.

569

570 **5.4. Implications for uplift mechanisms**

571 The northern sector of Atlas-Meseta system displays a shorter (10-50 km) and longer (>100 km)
572 wavelength topography associated with processes operating at crustal and deeper scale, respectively
573 (Fig. 11; Gomez et al., 1998; Boulton et al., 2014; Miller et al., 2015; Pastor et al., 2015). In particular,
574 the WMM consists of a NE-SW oriented, \sim 130 x 260 km, \sim 1500-m-high, topographic swell devoid

575 of Cenozoic deformation (Frizon de Lamotte et al., 2009; Barbero et al., 2010). Barbero et al. (2010)
576 interpreted the WMM as an open, lithospheric-scale fold that formed in response of shortening
577 processes (*i.e.*, lithospheric buckling). Although this mechanism has been documented for specific
578 rheological settings (*e.g.*, Cloetingh and Burov 2011), there are no geophysical data proving
579 lithospheric folding in the WMM. Alternative mechanisms which could explain the absence of crustal
580 deformation in the WMM and the occurrence of uplift with a topographic wavelength of a several
581 tens of km (up to more than 100 km) is the emplacement of igneous intrusion or magma underplating
582 (*e.g.*, Thybo and Artemieva, 2013; Townsend, 2022) and the removal of mantle lithosphere
583 (delamination; *e.g.*, Garzzone et al., 2006; Göğüş, and Pysklywec, 2008).

584 Following the scenario constrained by our river-profile inversions, the uplift in the WMM started at
585 ~22-20 Ma, shortly before the onset of alkaline volcanism (~16 Ma; El Azzouzi et al., 2010; Fig. 9)
586 and up to 15 Myr earlier than the rise of the northern sectors of the FMA and the South Rifian Corridor
587 (5-7 Ma; Krijgsman et al., 1999; Babault et al., 2008; Fig 10). These observations suggest that uplift
588 in the WMM would predate the more recent regional uplift and the increase in uplift rates observed
589 at 22-20 Ma must be ascribed to localized processes. The cumulative surface uplift in the WMM is
590 ~415 m with uplift rates varying from 35 to 55 m/Myr during the middle Miocene, and ~30 m/Myr
591 over the last 5-6 Ma (Figs. 9 and 10). These rates indicate that the amount of rock uplift in the last 5-
592 6 Ma would correspond to ~150-180 m. If we consider steady constant denudation of the relict
593 landscape at ~15 m/Myr, the resulting surface uplift during the last 6 Ma will be ~90 m (180 – 90 m,
594 where 90 m is the surface lowering over the last 6 Ma at 15 m/Myr). Thus, accounting for all the
595 above considerations the cumulative surface uplift pre-6 Ma would be 325 m (415 – 90 m). Assuming
596 that Airy isostasy is maintained and considering 2.8×10^3 and 3.23×10^3 kg/m³ as mean crustal (ρ_c)
597 and asthenospheric mantle (ρ_a) density respectively, a ~2000-m-thick intrusion will generate ~300 m
598 of surface uplift (note that $(\rho_a - \rho_c)/\rho_c$ is the ratio between elevation gain and crustal thickening;
599 Turcotte and Schubert, 2002). Similarly, if we assume Airy isostasy and a density of the lithospheric
600 mantle (ρ_m) of 3.3×10^3 kg/m³ the same magnitude of surface uplift will be generated by the removal

601 of a ~14-km-thick blob of mantle (note that $(\rho_m - \rho_a)/\rho_a$ is the ratio between elevation gain and
602 thickness of the removed lithospheric mantle; Molnar and Stock, 2009). A similar magnitude of
603 surface uplift (~500 m) has been also documented for the HP, although timing, rates and mechanism
604 of uplift are unknown (Fig. 11; Pastor et al., 2015).

605 In the FMA the topographic growth reflects the combination of deep-seated and crustal shortening
606 and thickening processes, that led to a cumulative surface uplift of about 1200 m as documented by
607 the river projections from the eastern FMA (Fig 11). Although we do not know the timing of this
608 surface uplift, the same magnitude of uplift occurred in the northern FMA during the last ~5-7 Ma as
609 documented by the uplifted, Messinian marine deposits of the Skoura Basin that are in unconformity
610 over folded Mesozoic limestones (Babault et al., 2008). Thus, it is reasonable to assume that most of
611 the 1200 m of surface uplift in the entire FMA, were gained between 5-7 Ma and the present day.
612 Recent surface uplift is also visible in the southern Rifian Corridor (Saiss and Taza-Guercif basins)
613 where marine sediments experienced ~850 m of surface uplift during the last 6-8 Ma (Krijgsman et
614 al., 1999; Babault et al., 2008). Overall, this configuration suggests that the northern FMA is
615 characterized by a complex long-term history of deformation, uplift, and subsidence predating the
616 last pulse of uplift, which produced ~1200 m of elevation gain at rates of ~170-220 m/Myr (Fig. 10).
617 This renewed uplift is almost coeval with the major, second phase of volcanism centred in the Middle
618 Atlas Mountains (Missenard et al., 2006; El Azzouzi et al., 2010) and could reflect large-scale
619 processes coupled with the reactivation of older tectonic structures (Frizon de Lamotte et al., 2009;
620 Babault et al., 2008) and possibly with the injection of large magmatic bodies within the crust.

621 Specifically, if we assume that shortening produced crustal thickening and surface uplift without
622 surface erosion and/or sediments deposition while Airy isostasy was approximately maintained,
623 thrusting in the FMA would have produced a surface elevation gain of 150 to 730 m over the last 7
624 Ma at rates of 20 to 100 m/Myr (Garzzone et al., 2006; see equations and assumptions in Table 2).
625 The assumption of no erosion (or very limited and hence neglectable erosion) over the last 7 Ma is

626 justified by the occurrence of the well-preserved, uplifted, marine deposits. This increase in elevation
627 would correspond to a crustal thickening of 0.8 to 4.0 km, and if we assume an initial crustal thickness
628 of 28-32 km, we will expect a ~29- to 36.7-km-thick crust (Table 2). These predictions agree with
629 crustal thickness estimates obtained by Gomez et al. (1999) and Miller and Backer (2014; Fig. 11).
630 Consequently, the remaining 470-1050 m of surface uplift should reflect deep-seated processes (*i.e.*,
631 mantle driven) at uplift rates of 65-150 m/Myr.

632 These simple calculations indicate two possible end-member scenarios for the topographic growth of
633 the FMA during the last 5-7 Ma: 1) shortening was the main cause of surface uplift (730 m), which
634 occurred at ~100 m/Myr, while dynamic uplift (470 m) took place at ~65 m/Myr; 2) shortening
635 accounted for a smaller fraction of elevation gain (150 m) and led to uplift at ~20 m/Myr, while
636 dynamic uplift (1050 m) took place at ~150 m/Myr (Table 2). In both cases magma addition may
637 have also, at least locally, influenced to the topographic growth of the FMA, although it is not possible
638 to isolate such a contribution. In any case, if we consider that the volcanic lavas younger than 4 Ma
639 crossing major faults are undeformed, and that the marine deposits of the Skoura Basin are only
640 slightly tilted ($< 10^\circ$), the amount of shortening and thickening during the Plio-Quaternary should be
641 marginal (El Azzouzi et al., 2010; Babault et al., 2008). Thus, the dynamic-supported scenario (option
642 2) seems to be the most reliable possibility (Fig. 10c profile 2) for the topographic growth of the FMA
643 (Babault et al., 2008).

644 Finally, the spatial pattern of surface and uplift rates over the last 5-7 Ma, document a large-scale
645 topographic bulge (wavelength > 300 km) centred in the FMA (Fig. 11). In this configuration, the
646 WMM appears to represent the western shoulder of a bulge with uplift at rates that are 2.5 to 6 times
647 lower than those inferred from the central bulge (Figs. 10c and 11b). This large wavelength signal
648 can be explained as results of lithospheric delamination/dripping or upwelling of asthenospheric
649 mantle associated with even deeper processes like the rise of plumes possibly sourced from the
650 Canary hotspot, or a combination of both (Duggen et al., 2009; Miller and Becker, 2014; Miller et

651 al., 2015). Although a large number of geophysical studies robustly detected a shallow LAB and
652 Moho beneath the Middle Atlas, the nature of the regional, deep dynamic processes is still under
653 debate (*e.g.*, Missenard et al., 2006; Clementucci et al., 2023). Given the wavelength and the
654 amplitude of the most recent uplift event and the characteristics of coeval volcanism, we tend to
655 favour the plume hypothesis. Dynamic rising of asthenospheric mantle can reconcile the occurrence
656 of large-scale surface uplift, non-isostatically compensated topography, anomalous thinning of the
657 crust and lithospheric mantle and intermittent Mio-pliocene alkaline volcanism.

658 In essence, we conclude that the regional topographic evolution of the northern Atlas-Meseta system
659 can result from the contribution of deep components (plume activity and mantle lithospheric
660 delamination) coupled with smaller-scale processes such as possible injection of magmatic bodies
661 within the crust and crustal shortening and thickening, at least for the structural domains characterized
662 by contractional deformation (*e.g.*, Middle and High Atlas).

663

664 **6. Conclusion**

665 Our study allows the characterisation of the magnitude, timing, and rates of the long-term topographic
666 evolution of the northern domains of the Atlas-Meseta system. The preserved relict landscape
667 standing at a high elevation and the occurrence of deeply incised valleys downstream of major non-
668 lithological knickpoints document a transience in topography and hence different stages of
669 topographic evolution. Specifically, the ancient relict topography displays low relief, slope, and k_{sn}
670 values in association with low rates of denudation on both bedrock and watersheds (from 4 to 20
671 m/Myr). Conversely, the downstream portion of the landscape exhibits higher topographic metrics
672 and denudation rates (up to 40 m/Myr) in response to an increase in rock uplift rate that led to a
673 topographic rejuvenation. Within this general trend a few exceptions emerge. In certain catchments
674 of the Western Meseta, the granite massifs exposed on top of the relict landscape provide a higher

675 amount of quartz than the downstream metamorphics, resulting in a mixed signal of ^{10}Be
676 concentration, giving lower denudation rates than expected. These findings highlight that the uneven
677 distribution of quartz-bearing rocks in the sampled catchments and their relative location within the
678 topographic sectors (*i.e.*, relict vs rejuvenated landscape) represent a critical point for deriving basin-
679 wide denudation rates.

680 Combined with geological data, our river-profile inversion results suggest that the elevated relict
681 surfaces of the WMM have been forming since the Eocene. Moreover, our inverted uplift rates
682 indicate that surface uplift and associated knickpoints migrations started at ~22-20 Ma and reached
683 peak values of ~55 m/Myr at ~15 Ma, while the relict portion of the WMM has been eroding steadily
684 at an average of 15 m/Myr. These values agree with our basin-wide denudation and fluvial incision
685 rates derived from 1-Myr-old lava flows for the relict landscape and the catchment area downstream
686 of non-lithological knickpoints. This Miocene uplift (first uplift phase) of the WMM led to a
687 cumulative surface uplift of ~400 m over an extent of ~ 130 x 260 km, and is interpreted to reflect
688 crustal-scale processes such as igneous intrusion or magma underplating.

689 Subsequently, from 5-7 Ma, regional uplift occurred as documented by uplifted, Messinian marine
690 deposits of the Skoura and Saiss basins and minor knickpoints over the WMM domain. Surface uplift
691 inferred from the modern elevation of these marine deposits in the northern FMA (Skoura Basin) is
692 ~1200 m, that agree also with estimates from the western FMA obtained from the river projections.
693 The larger scale (wavelength > 200 km) and the higher magnitude of uplift suggest that mantle
694 processes like upwelling of deep asthenospheric mantle (*i.e.*, plume) may have driven the topographic
695 rejuvenation of the entire Atlas-Meseta system including the WMM (second uplift phase). Uplift
696 occurred at rates ranging from of ~30 to ~170 m/Myr from the WMM to the FMA over a pattern
697 describing a bulge centred in the FMA, where the Plio-Quaternary volcanism is also centred. The
698 WMM represents the western shoulder of this bulge. Finally, minor crustal thickening induced by
699 shortening may have also contributed to the post 7-Ma topographic growth of the FMA. Overall,

700 these results suggest that uplift in the WMM may have occurred in two major steps at 22-20 and 5-6
701 Ma, while in the FMA, at least in its northern and western sectors, the major uplift pulse occurred
702 during the last 5-7 Ma.

703

704 **Acknowledgments**

705 This study is part of the PhD thesis of RC at the University of Roma Tre (PhD Cycle XXXIV). It was
706 supported by the PhD School of Roma Tre and grant “Vinci 2020” awarded to RC (Number: C2-
707 1403). PB was supported by the MIUR (Ministry of Education University and Research), Excellence
708 Department Initiative, Art. 1, com. 314-337, Law 232/2016. The ASTER (CEREGE, Aix-en-
709 Provence) AMS national facility, is supported by the INSU/CNRS, the ANR (Projets thématiques
710 d’excellence programme for the Equipements d’Excellence) and the IRD. We thank G. Aumaître, K.
711 Keddadouche and F. Zaidi for their expertise in AMS measurements, A. Yaaqoub, A. Essaifi, N.
712 Youbi, A. Abbassi and R. Braucher for the stimulating discussions. We are indebted to P-H Blard, S.
713 Willett and T. Schildgen for revising this manuscript as a chapter of the PhD Thesis of the first author,
714 to J. Babault, G. Rixhon and anonymous Reviewers for the excellent and detailed revisions, to the
715 Editor M. Stokes for managing the submission of the manuscript. Finally to L. Benedetti, M. Della
716 Seta, P. Molin and O. Bellier for providing valuable comments during the PhD defense of RC.

717

718 **References**

- 719 Abbassi, A., Cipollari, P., Zaghoul, M. N., Cosentino, D., 2020. The Rif Chain (northern Morocco)
720 in the late Tortonian-early Messinian tectonics of the Western Mediterranean orogenic belt:
721 Evidence from the Tanger-Al Manzla wedge-top basin. *Tectonics*. 39(9).
722 <https://doi.org/10.1029/2020TC006164>.
- 723 Abbassi, A., Cipollari, P., Fellin, M. G., Zaghoul, M. N., Guillong, M., El Mourabet, M., &
724 Cosentino, D. (2022). The Numidian sand event in the Burdigalian foreland basin system of
725 the Rif, Morocco, in a source-to-sink perspective. *GSA Bulletin*, 134(9-10), 2280-2304.
726 <https://doi.org/10.1130/B36136.1>.

727 Adams, B. A., Whipple, K. X., Forte, A. M., Heimsath, A. M., & Hodges, K. V. 2020. Climate
728 controls on erosion in tectonically active landscapes. *Sci. adv.* 6(42), eaaz3166.
729 <https://www.science.org/doi/10.1126/sciadv.aaz3166>.

730 Agharroud, K., Siame, L. L., Ben Moussa, A., Bellier, O., Guillou, V., Fleury, J., and El Kharim, Y.
731 (2021). Seismo-tectonic model for the southern Pre-Rif border (Northern Morocco): Insights
732 from morphochronology. *Tectonics*. 40(4), e2020TC006633.
733 <https://doi.org/10.1029/2020TC006633>.

734 Arboleya, M. L., Teixell, A., Charroud, M., Julivert, M., 2004. A structural transect through the High
735 and Middle Atlas of Morocco. *J. Afr. Earth. Sci.* 39(3-5), 319-327.
736 <https://doi.org/10.1016/j.jafrearsci.2004.07.036>.

737 Babault, J., Teixell, A., Arboleya, M. L., Charroud, M., 2008. A Late Cenozoic age for long-
738 wavelength surface uplift of the Atlas Mountains of Morocco. *Terra nova*. 20(2), 102-107.
739 <https://doi.org/10.1111/j.1365-3121.2008.00794.x>.

740 Babault, J., Van Den Driessche, J., and Charco, M., 2022, 2.10 - Plateau Uplift, Regional Warping,
741 and Subsidence, in Shroder, J. F., ed., *Treatise on Geomorphology (Second Edition)*: Oxford,
742 Academic Press, p. 223-269. <https://doi.org/10.1016/B978-0-12-818234-5.00119-X>.

743 Balco, G., Stone, J. O., Lifton, N. A., Dunai, T. J., 2008. A complete and easily accessible means of
744 calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements. *Quat.*
745 *Geochronol.* 3(3), 174-195. <https://doi.org/10.1016/j.quageo.2007.12.001>. Ballato, P.,
746 Landgraf, A., Schildgen, T. F., Stockli, D. F., Fox, M., Ghassemi, M. R., Kirby, E., Strecker,
747 M. R., 2015. The growth of a mountain belt forced by base-level fall: Tectonics and surface
748 processes during the evolution of the Alborz Mountains, N Iran. *Earth and Planet. Sci. Lett.*
749 425, 204-218. <https://doi.org/10.1016/j.epsl.2015.05.051>.

750 Barbero, L., Jabaloy, A., Gómez-Ortiz, D., Pérez-Peña, J. V., Rodríguez-Peces, M. J., Tejero, R.,
751 Estupinan, J., Azdimousa, A., Vazquez, M., Asebriy, L., 2011. Evidence for surface uplift of
752 the Atlas Mountains and the surrounding peripheral plateaux: Combining apatite fission-track
753 results and geomorphic indicators in the Western Moroccan Meseta (coastal Variscan
754 Paleozoic basement). *Tectonophysics*. 502(1-2), 90-104.
755 <https://doi.org/10.1016/j.tecto.2010.01.005>.

756 Borchers, B., Marrero, S., Balco, G., Caffee, M., Goehring, B., Lifton, N., Nishiizumi, K., Phillips,
757 F., Schaefer, J., Stone, J., 2016. Geological calibration of spallation production rates in the
758 CRONUS-Earth project. *Quat. Geochronol.* 31, 188–198.
759 <https://doi.org/10.1016/j.quageo.2015.01.009>.

760 Boulton, S. J., Stokes, M., & Mather, A. E. (2014). Transient fluvial incision as an indicator of active
761 faulting and Plio-Quaternary uplift of the Moroccan High Atlas. *Tectonophysics*, 633, 16-
762 33. <https://doi.org/10.1016/j.tecto.2014.06.032>.

763 Braucher, R., Merchel, S., Borgomano, J., Bourlès, D. L., 2011. Production of cosmogenic
764 radionuclides at great depth: A multi element approach. *Earth Planet. Sci. Lett.* 309(1-2), 1-9.
765 <https://doi.org/10.1016/j.epsl.2011.06.036>.

766 Burbank, D. W., and Anderson, R. S., 2009. *Tectonic geomorphology*. John Wiley & Sons.
767 <https://doi.org/10.1002/9781444345063>.

768 Calvet, M., Gunnell, Y., Farines, B., 2015. Flat-topped mountain ranges: Their global distribution
769 and value for understanding the evolution of mountain topography. *Geomorphology*. 241, 255-
770 291. <https://doi.org/10.1016/j.geomorph.2015.04.015>.

771 Capella, W., Barhoun, N., Flecker, R., Hilgen, F. J., Kouwenhoven, T., Matenco, L. C., Sierro L.C.,
772 Tulbure, M.A., Yousfi, M.Z., and Krijgsman, W., 2018. Palaeogeographic evolution of the
773 late Miocene Rifian Corridor (Morocco): reconstructions from surface and subsurface data.
774 *Earth-Science Reviews*, 180, 37-59.

775 Carretier, S., Regard, V., Vassallo, R., Martinod, J., Christophoul, F., Gayer, E., Audin L., Lagane,
776 C., 2015. A note on ¹⁰Be-derived mean erosion rates in catchments with heterogeneous
777 lithology: Examples from the western Central Andes. *Earth Surf. Processes Landforms*.
778 40(13), 1719-1729. <https://doi.org/10.1002/esp.3748>.

779 Champagnac, J. D., Valla, P. G., & Herman, F., 2014. Late-Cenozoic relief evolution under evolving
780 climate: A review. *Tectonophysics*. 614, 44-65. <https://doi.org/10.1016/j.tecto.2013.11.037>

781 Charriere, A. and Saint-Martin, J.P., 1989. Relations entre les formations re'cifales du Miocene
782 superieur et la dynamique d'ouverture et de fermeture des communications marines a la
783 bordure me'ridionale du sillon sud-rifain (Maroc). *C. R. Acad. Sci. II*, 309, 611–614.

784 Charriere, A., 1984. Évolution néogène de bassins continentaux et marins dans le Moyen Atlas central
785 (Maroc). *Bull. Soc. Géol. Fr.* 7(6), 1127-1136.

786 Charton, R., Bertotti, G., Arnould, A. D., Storms, J. E., and Redfern, J., 2021. Low-temperature
787 thermochronology as a control on vertical movements for semi-quantitative source-to-sink
788 analysis: A case study for the Permian to Neogene of Morocco and surroundings. *Basin*
789 *Research*, 33(2), 1337-1383. <https://doi.org/10.1111/bre.12517>.

790 Chmeleff, J., von Blanckenburg, F., Kossert, K., Jakob, D., 2010. Determination of the ¹⁰Be half-life
791 by multicollector ICP-MS and liquid scintillation counting. *Nucl. Instrum. Methods Phys.*
792 *Res., Sect. B, Beam Interact. Mater. Atoms.* 268(2), 192–199.
793 <https://doi.org/10.1016/j.nimb.2009.09.012>.

794 Clementucci, R., 2022. Deciphering mantle contribution on surface uplift in the Atlas-Meseta system
795 (Morocco) (Doctoral dissertation). *Earth Sciences*. Università degli studi Roma Tre; Aix
796 Marseille Univ. <https://tel.archives-ouvertes.fr/tel-03630297/document>.

797 Clementucci R., Ballato P., Siame L., Faccenna C., Yaaqoub A., Essaifi A., Leanni L., Guillou V.,
798 2022. Lithological control on topographic relief evolution in a slow tectonic setting (Anti-
799 Atlas, Morocco). *Earth and Planet. Sci. Lett.* <https://doi.org/10.1016/j.epsl.2022.117788>.

800 Clementucci, R., Ballato, P., Siame, L., Fox, M., Lanari, R., Sembroni, A., Faccenna, C., Yaaqoub,
801 A., Essaifi, A., (2023). Surface uplift and topographic rejuvenation of a tectonically inactive
802 range: Insights from the Anti-Atlas and the Siroua Massif (Morocco). *Tectonics*. 42,
803 e2022TC007383. <https://doi.org/10.1029/2022TC007383>

804 Cloetingh, S., and Burov, E., 2011.
805 Lithospheric folding and sedimentary basin evolution: a review and analysis of formation
806 mechanisms. *Basin Research*, 23(3), 257-290. <https://doi.org/10.1111/j.1365-2117.2010.00490.x>.

807 Crosby, B. T., and Whipple, K. X., 2006. Knickpoint initiation and distribution within fluvial
808 networks: 236 waterfalls in the Waipaoa River, North Island, New Zealand. *Geomorphology*.
809 82(1-2), 16-38. <https://doi.org/10.1016/j.geomorph.2005.08.023>.

810 Duggen, S., Hoernle, K. A., Hauff, F., Kluegel, A., Bouabdellah, M., Thirlwall, M. F., 2009. Flow of
811 Canary mantle plume material through a subcontinental lithospheric corridor beneath Africa
812 to the Mediterranean. *Geology*. 37(3), 283-286. <https://doi.org/10.1130/G25426A.1>.

813 El Attmani, M., Bouwafoud, A., Elouariti, S., Mhamdi, H. S., Bouziane, A. B., Mouflih, M., 2021.
814 The Eocene Bryozoans of Bekrit-Timahdite Formation, Middle-Atlas, Morocco: Taxonomy,
815 Paleoenvironments and Paleoclimates Implications. *J. Geosci. Env. Prot.* 9(3), 136-153.
816 <https://doi.org/10.4236/gep.2021.93008>.

817 El Azzouzi, M. H., Maury, R. C., Bellon, H., Youbi, N., Cotten, J., Kharbouch, F., 2010. Petrology
818 and K-Ar chronology of the Neogene-quaternary Middle Atlas basaltic province, Morocco.
819 *Bull. Soc. Géol. Fr.* 181(3), 243-257. <https://doi.org/10.2113/gssgfbull.181.3.243>.

820 El Kiram, N., Jaffal, M., Kchikach, A., El Azzab, D., El Ghorfi, M., Khadiri, O., Jourani, E., Manar,
821 A., and Nahim, M., 2019. Phosphatic series under Plio-Quaternary cover of Tadla Plain,
822 Morocco: Gravity and seismic data. *C. R. Geosci.* 351(6), 420-429.

823 England, P., Molnar, P., 1990. Surface uplift, uplift of rocks, and exhumation of rocks. *Geology*.
824 18(12), 1173-1177. [https://doi.org/10.1130/0091-7613\(1990\)018<1173:SUUORA>2.3.CO;2](https://doi.org/10.1130/0091-7613(1990)018<1173:SUUORA>2.3.CO;2).

825 Faccenna, C., Becker, T. W., 2020. Topographic expressions of mantle dynamics in the
826 Mediterranean. *Earth-Science Reviews*. 209, 103327.
827 <https://doi.org/10.1016/j.earscirev.2020.103327>.

828 Fernandes, V. M., Roberts, G. G., White, N., & Whittaker, A. C., 2019. Continental-scale landscape
829 evolution: A history of North American topography. *J. Geophys. Res., Earth Surface*, 124(11),
830 2689-2722. <https://doi.org/10.1029/2018JF004979>.

831 Forte, A.M., Whipple, K.X., 2019. Short communication: the Topographic Analysis Kit (TAK) for
832 TopoToolbox. *Earth Surf. Dyn.* 7, 87–95. <https://doi.org/10.5194/esurf-7-87-2019>.

833 Frizon de Lamotte, D., Leturmy, P., Missenard, Y., Khomsi, S., Ruiz, G., Saddiqi, O., Guillocheau
834 F., Michard A., 2009. Mesozoic and Cenozoic vertical movements in the Atlas system
835 (Algeria, Morocco, Tunisia). An overview: *Tectonophysics*. 475, 9–28.
836 <https://doi.org/10.1016/j.tecto.2008.10.024>.

837 Godard, V., Dosseto, A., Fleury, J., Bellier, O., Siame, L., ASTER Team., 2019. Transient landscape
838 dynamics across the Southeastern Australian Escarpment. *Earth and Planet. Sci. Lett.* 506,
839 397-406. <https://doi.org/10.1016/j.epsl.2018.11.017>.

840 Gallen, S. F., Wegmann, K. W., 2017. River profile response to normal fault growth and linkage: An
841 example from the Hellenic forearc of south-central Crete, Greece. *Earth Surf. Dyn.* 5(1), 161-
842 186. <https://doi.org/10.5194/esurf-5-161-2017>.

843 Gallen, S. F., 2018. Lithologic controls on landscape dynamics and aquatic species evolution in post-
844 orogenic mountains. *Earth and Planet. Sci. Lett.* 493, 150-160.
845 <https://doi.org/10.1016/j.epsl.2018.04.029>.

846 Garzione, C. N., Molnar, P., Libarkin, J. C., MacFadden, B. J., 2006. Rapid late Miocene rise of the
847 Bolivian Altiplano: Evidence for removal of mantle lithosphere. *Earth and Planet. Sci. Lett.*
848 241(3-4), 543-556. <https://doi.org/10.1016/j.epsl.2005.11.026>.

849 Gomez, F., Barazangi, M., Bensaid, M., 1996. Active tectonism in the intracontinental Middle Atlas
850 Mountains of Morocco: synchronous crustal shortening and extension. *J. Geol. Soc.* 153(3),
851 389-402. <https://doi.org/10.1144/gsjgs.153.3.0389>.

852 Gomez, F., Allmendinger, R., Barazangi, M., Er-Raji, A., Dahmani, M., 1998. Crustal shortening
853 and vertical strain partitioning in the Middle Atlas Mountains of Morocco. *Tectonics*. 17(4),
854 520-533. <https://doi.org/10.1029/98TC01439>.

855 Göğüş, O. H., Pysklywec, R. N., 2008. Near-surface diagnostics of dripping or delaminating
856 lithosphere. *J. Geophys. Res., Solid Earth*. 113(B11). <https://doi.org/10.1029/2007JB005123>.

857 Goren, L., Fox, M., Willett, S. D., 2014. Tectonics from fluvial topography using formal linear
858 inversion: Theory and applications to the Inyo Mountains, California. *J. Geophys. Res. F:*
859 *Earth Surf.* 119(8), 1651-1681. <https://doi.org/10.1002/2014JF003079>.

860 Hoepffner, C., Soulaimani, A., Piqué, A., 2005. The moroccan hercynides. *J. Afr. Earth. Sci.* 43(1-
861 3), 144-165. <https://doi.org/10.1016/j.jafrearsci.2005.09.002>.

862 Huffman, G. J., Bolvin, D. T., Braithwaite, D., Hsu, K., Joyce, R., Xie, P., & Yoo, S. H., 2015. NASA
863 global precipitation measurement (GPM) integrated multi-satellite retrievals for GPM

864 (IMERG). Algorithm Theoretical Basis Document (ATBD) Version, 4, 26.
865 http://pmm.nasa.gov/sites/default/files/document_files/IMERG_ATBD_V4.5.pdf.

866 Jolivet, L., & Faccenna, C., 2000. Mediterranean extension and the Africa-Eurasia collision.
867 *Tectonics*. 19(6), 1095-1106. <https://doi.org/10.1029/2000TC900018>.

868 Kirby, E., Whipple, K. X., 2012. Expression of active tectonics in erosional landscapes. *J. Struct.*
869 *Geol.* 44, 54-75. <https://doi.org/10.1016/j.jsg.2012.07.009>.

870 Korschinek, G., Bergmaierb, A., Faestermann, T., Gerstmann, U.C., Kniew, K., Rugela, G.,
871 Wallner, A., Dillmann, I., Dollinger, G., Lierse von Gostomski, Ch., Kossert, K., Maitia,
872 M., Poutivtseva, M., Remmer, A., 2010. A new value for the half-life of ¹⁰Be by heavy ion
873 elastic recoil detection and liquid scintillation counting. *Nucl. Instr. Meth Res. B.*
874 <https://doi.org/10.1016/j.nimb.2009.09.020>. Krijgsman, W., Langereis, C. G., Zachariasse, W. J.,
875 Boccaletti, M., Moratti, G., Gelati, R., Iaccarino, F., Papani, G., Villa, G., 1999. Late Neogene
876 evolution of the Taza–Guercif Basin (Rifian Corridor, Morocco) and implications for the
877 Messinian salinity crisis. *Mar. Geol.* 153(1-4), 147-160. [https://doi.org/10.1016/S0025-](https://doi.org/10.1016/S0025-3227(98)00084-X)
878 [3227\(98\)00084-X](https://doi.org/10.1016/S0025-3227(98)00084-X).

879 Lal, D., 1991. Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion
880 models. *Earth Planet. Sci. Lett.* 104, 424–439. [http://dx.doi.org/10.1016/0012-](http://dx.doi.org/10.1016/0012-821X(91)90220-C)
881 [821X\(91\)90220-C](http://dx.doi.org/10.1016/0012-821X(91)90220-C)

882 Lanari, R., Fellin, M. V., Faccenna, C., Balestrieri, M. L., Pazzaglia, F. J., Youbi, N., Maden, C.,
883 2020a. Exhumation and surface evolution of the western high atlas and surrounding regions
884 as constrained by low-temperature thermochronology. *Tectonics*. 39(3), e2019TC005562.
885 <https://doi.org/10.1029/2019TC005562>.

886 Lanari, R., Faccenna, C., Fellin, M. G., Essaifi, A., Nahid, A., Medina, F. and Youbi, N., 2020b.
887 Tectonic evolution of the western high Atlas of Morocco: oblique convergence, reactivation,
888 and transpression. *Tectonics*, 39(3), e2019TC005563.
889 <https://doi.org/10.1029/2019TC005563>.

890 Lanari, R., Reitano, R., Giachetta, E., Pazzaglia, F.J., Clementucci, R., Faccenna, C., Fellin, M.G.,
891 2022. Is the Anti-Atlas of Morocco still uplifting? *J. Afr. Earth Sci.* 188, 104481. [https://doi](https://doi.org/10.1016/j.jafrearsci.2022.104481)
892 [.org/10.1016/j.jafrearsci.2022.104481](https://doi.org/10.1016/j.jafrearsci.2022.104481).

893 Lanari, R., Boutoux, A., Faccenna, C., Herman, F., Willett, S. D., & Ballato, P., 2023. Cenozoic
894 exhumation in the Mediterranean and the Middle East. *Earth-Science Reviews*, 104328

895 Marder, E., & Gallen, S. F., 2023. Climate control on the relationship between erosion rate and fluvial
896 topography. *Geology*. <https://doi.org/10.1130/G50832.1>.

897 Marra, F., Rohling, E. J., Florindo, F., Jicha, B., Nomade, S., Pereira, A., & Renne, P. R., 2016.
898 Independent $^{40}\text{Ar}/^{39}\text{Ar}$ and ^{14}C age constraints on the last five glacial terminations from the
899 aggradational successions of the Tiber River, Rome (Italy). *Earth and Planetary Science*
900 *Letters*, 449, 105-117. <https://doi.org/10.1016/j.epsl.2016.05.037>.

901 Meyer, H., Hetzel, R., Fügenschuh, B., Strauss, H., 2010. Determining the growth rate of topographic
902 relief using in situ-produced ^{10}Be : a case study in the Black Forest, Germany. *Earth and*
903 *Planet. Sci. Lett.* 290(3-4), 391-402. <https://doi.org/10.1016/j.epsl.2009.12.034>

904 Miller, S. R., Sak, P. B., Kirby, E., Bierman, P. R., 2013. Neogene rejuvenation of central
905 Appalachian topography: Evidence for differential rock uplift from stream profiles and erosion
906 rates. *Earth and Planet. Sci. Lett.* 369, 1-12. <https://doi.org/10.1016/j.epsl.2013.04.007>.

907 Miller, M. S., Becker, T. W., 2014. Reactivated lithospheric-scale discontinuities localize dynamic
908 uplift of the Moroccan Atlas Mountains. *Geology*. 42(1), 35-38.
909 <https://doi.org/10.1130/G34959.1>.

910 Miller, M. S., O'Driscoll, L. J., Butcher, A. J., Thomas, C., 2015. Imaging Canary Island hotspot
911 material beneath the lithosphere of Morocco and southern Spain. *Earth and Planet. Sci. Lett.*
912 431, 186-194. <https://doi.org/10.1016/j.epsl.2015.09.026>.

913 Miller, K. G., Browning, J. V., Schmelz, W. J., Kopp, R. E., Mountain, G. S., & Wright, J. D., 2020.
914 Cenozoic sea-level and cryospheric evolution from deep-sea geochemical and continental
915 margin records. *Science advances*, 6(20), eaaz1346. <https://doi.org/10.1126/sciadv.aaz1346>.

916 Missenard, Y., Zeyen, H., Frizon de Lamotte, D., Leturmy, P., Petit, C., Sébrier, M., Saddiqi, O.,
917 2006. Crustal versus asthenospheric origin of relief of the Atlas Mountains of Morocco. *J.*
918 *Geophys. Res., Solid Earth*. 111(B3). <https://doi.org/10.1029/2005JB003708>.

919 Olivetti, V., Godard, V., Bellier, O., and ASTER team., 2016. Cenozoic rejuvenation events of Massif
920 Central topography (France): Insights from cosmogenic denudation rates and river profiles.
921 *Earth and Planet. Sci. Lett.* 444, 179-191. <https://doi.org/10.1016/j.epsl.2016.03.049>.

922 Pastor, A., Babault, J., Owen, L. A., Teixell, A., Arboleya, M. L., 2015. Extracting dynamic
923 topography from river profiles and cosmogenic nuclide geochronology in the Middle Atlas
924 and the High Plateaus of Morocco. *Tectonophysics*. 663, 95-109.
925 <https://doi.org/10.1016/j.tecto.2015.06.007>.

926 Pavano, F., & Gallen, S. F., 2021. A geomorphic examination of the Calabrian forearc translation.
927 *Tectonics*, 40(7), e2020TC006692. <https://doi.org/10.1029/2020TC006692>.

928 Peifer, D., Persano, C., Hurst, M. D., Bishop, P., & Fabel, D., 2021. Growing topography due to
929 contrasting rock types in a tectonically dead landscape. *Earth Surface Dynamics*. 9(2), 167-
930 181. <https://doi.org/10.5194/esurf-9-167-2021>.

- 931 Perron, J. T., Royden, L., 2013. An integral approach to bedrock river profile analysis. *Earth Surf.*
932 *Processes Landforms.* 38(6), 570-576. <https://doi.org/10.1002/esp.3302>.
- 933 Racano, S., Schildgen, T. F., Cosentino, D., Miller, S. R., 2021. Temporal and spatial variations in
934 rock uplift from river-profile inversions at the Central Anatolian Plateau southern margin. *J.*
935 *Geophys. Res. F: Earth Surf.* 126(8). <https://doi.org/10.1029/2020JF006027>.
- 936 Rachdi, H., 1995. Etude du volcanisme plio-quaternaire du Maroc central: pétrographie, géochimie
937 et minéralogie. *Notes Mém. Serv. Géol. Maroc.* 381, 157.
- 938 Riebe, C. S., Kirchner, J. W., Finkel, R. C., 2004. Erosional and climatic effects on long-term
939 chemical weathering rates in granitic landscapes spanning diverse climate regimes. *Earth and*
940 *Planet. Sci. Lett.* 224(3-4), 547-562. <https://doi.org/10.1016/j.epsl.2004.05.019>.
- 941 Roberts, G. G., & White, N., 2010. Estimating uplift rate histories from river profiles using African
942 examples. *J. Geophys. Res., Solid Earth.* 115(B2). <https://doi.org/10.1029/2009JB006692>.
- 943 Rosenkranz, R., Schildgen, T., Wittmann, H., Spiegel, C., 2018. Coupling erosion and topographic
944 development in the rainiest place on Earth: Reconstructing the Shillong Plateau uplift history
945 with in-situ cosmogenic ¹⁰Be. *Earth and Planet. Sci. Lett.* 483, 39-51.
946 <https://doi.org/10.1016/j.epsl.2017.11.047>.
- 947 Salvan, H., Farkhany, M., 1982. Les phosphates de chaux sédimentaires du Maroc. Leurs
948 caractéristiques et leurs problèmes (essai de synthèse). *Notes Mem. Serv. geol. Maroc* 14,
949 7e20.
- 950 Schwanghart, W., Scherler, D., 2014. Short communication: TopoToolbox 2 – MATLAB-based
951 software for topographic analysis and modeling in Earth surface sciences. *Earth Surf. Dyn.* 2,
952 1–7. <https://doi.org/10.5194/esurf-2-1-2014>.
- 953 Sembroni, A., Kiraly, A., Faccenna, C., Funicicello, F., Becker, T. W., Globig, J., & Fernandez, M.,
954 2017. Impact of the lithosphere on dynamic topography: Insights from analogue modeling.
955 *Geophysical Research Letters*, 44(6), 2693-2702. <https://doi.org/10.1002/2017GL072668>.
- 956 Sembroni, A., Molin, P., Soligo, M., Tuccimei, P., Anzalone, E., Billi, A., Franchini, S., Rinaldi, M.,
957 and Tarchini, L., 2020. The uplift of the Adriatic flank of the Apennines since the Middle
958 Pleistocene: New insights from the Tronto River basin and the Acquasanta Terme Travertine
959 (central Italy). *Geomorphology*, 352, 106990.
960 <https://doi.org/10.1016/j.geomorph.2019.106990>.
- 961 Smith, A. G., Fox, M., Schwanghart, W., Carter, A., 2022. Comparing methods for calculating
962 channel steepness index. *Earth-Science Reviews*, 227, 103970.
963 <https://doi.org/10.1016/j.earscirev.2022.103970>.

964 Snyder, N. P., Whipple, K. X., Tucker, G. E., & Merritts, D. J., 2000. Landscape response to tectonic
965 forcing: Digital elevation model analysis of stream profiles in the Mendocino triple junction
966 region, northern California. *Geological Society of America Bulletin*, 112(8), 1250-1263.
967 [https://doi.org/10.1130/0016-7606\(2000\)112<1250:LRTTFD>2.0.CO;2](https://doi.org/10.1130/0016-7606(2000)112<1250:LRTTFD>2.0.CO;2).

968 Snyder, N. P., Whipple, K. X., Tucker, G. E., Merritts, D. J., 2002. Interactions between onshore
969 bedrock-channel incision and nearshore wave-base erosion forced by eustasy and tectonics.
970 *Basin Research*. 14(2), 105-127. <https://doi.org/10.1046/j.1365-2117.2002.00169.x>.

971 Stokes, M., Mather, A. E., Belfoul, M., Faik, F., Bouzid, S., Geach, M. R., Cunha, P.P., Boulton, S.J.,
972 & Thiel, C. (2017). Controls on dryland mountain landscape development along the NW
973 Saharan desert margin: Insights from Quaternary river terrace sequences (Dadès River, south-
974 central High Atlas, Morocco). *Quaternary Science Reviews*, 166, 363-
975 379. <https://doi.org/10.1016/j.quascirev.2017.04.017>.

976 Townsend, M., 2022. Linking surface deformation to thermal and mechanical magma chamber
977 processes. *Earth and Planet. Sci. Lett.* 577, 117272.
978 <https://doi.org/10.1016/j.epsl.2021.117272>.

979 Turcotte, D. L., & Schubert, G., 2002. *Geodynamics*. Cambridge university press.

980 Vernant, P., Fadil, A., Mourabit, T., Ouazar, D., Koulali, A., Davila, J. M., Garate, J., McClusky, A.,
981 Reilinger, R., 2010. Geodetic constraints on active tectonics of the Western Mediterranean:
982 Implications for the kinematics and dynamics of the Nubia-Eurasia plate boundary zone.
983 *Journal of Geodynamics*. 49(3-4), 123-129. <https://doi.org/10.1016/j.jog.2009.10.007>.

984 Whipple, K. X., 2001. Fluvial landscape response time: How plausible is steady-state denudation?.
985 *American Journal of Science*, 301(4-5), 313-325. <https://doi.org/10.2475/ajs.301.4-5.313>.

986 Willett, S. D., & Brandon, M. T., 2002. On steady states in mountain belts. *Geology*. 30(2), 175-178.
987 [https://doi.org/10.1130/0091-7613\(2002\)030<0175:OSSIMB>2.0.CO;2](https://doi.org/10.1130/0091-7613(2002)030<0175:OSSIMB>2.0.CO;2).

988 Wobus, C., Whipple, K. X., Kirby, E., Snyder, N., Johnson, J., Spyropolou, K., Crosby B., &
989 Sheehan, D., 2006. Tectonics from topography: Procedures, promise, and pitfalls. *Special*
990 *Papers - Geological Society of America*, 398, 55. [https://doi.org/10.1130/2006.2398\(04\)](https://doi.org/10.1130/2006.2398(04)).

991 Yang, R., Willett, S. D., Goren, L., 2015. In situ low-relief landscape formation as a result of river
992 network disruption. *Nature*. 520(7548), 526-529. <https://doi.org/10.1038/nature14354>.

993 Zhong, Y., Willett, S. D., Picotti, V., Xiong, J., & Zhang, H., 2022. Spatial and temporal variations
994 of incision rate of the middle Yellow River and its tributaries. *Journal of Geophysical*
995 *Research: Earth Surface*, 127(1), e2021JF006327. <https://doi.org/10.1029/2021JF006327>.

996 Zondervan, J. R., Stokes, M., Boulton, S. J., Telfer, M. W., & Mather, A. E., 2020. Rock strength and
997 structural controls on fluvial erodibility: Implications for drainage divide mobility in a

998 collisional mountain belt. *Earth and Planetary Science Letters*, 538,
999 116221. <https://doi.org/10.1016/j.epsl.2020.116221>.




Figure 1. (a) Topographic hillshade of the Rif, Atlas and Meseta domains. **(b)** Morpho-structural domains in the western central Morocco and selected. FMA: Folded Middle Atlas. TMA: Tabular Middle Atlas. NMAF: North Middle Atlas fault. SMAF: South Middle Atlas fault. **(c)** DEM of the Western Moroccan Meseta. The white transparent areas show the sampled catchments while the dots show the location of the sampling points. Note the occurrence of two major catchment and several sub-catchments. The white hexagons show the location of bedrock samples. The thick dashed line indicates the boundary between the TMA, the FMA and the Western Meseta.




Figure 2. (a) Topographic and geological map of the sampled catchment 63. Note, sampling point 59 is located upstream the non-lithological knickpoint, 62 is downstream the non-lithological knickpoint and 63 represents the mixed rate of basin-wide denudation. **(b)** Schematic illustration of the catchment 63, showing the correction method to estimate the denudation rate of the downstream portion (64) of the catchment 63, by using drainage area and denudation rate of the basin 63 and 59, respectively. Note, basin-wide denudation derived from basin 62 is consistent with estimate value of the basin 64, both located in the downstream of non-lithological knickpoints.




Figure 3. (a) DEM (b) k_{sn} map with main knickpoints. The bold knickpoints demark the escarpment between the TMA and Western Moroccan Meseta (WMM), while the thick dashed line indicates the boundary between the main morpho-structural domains. The solid lines demark the tributaries of the Oued El Rbia (South Western Meseta, SWM), of the Oued Sebou (North Western Meseta, NWM) and rivers draining directly from the

Western Meseta to the Atlantic ocean (CWM: Central Western Meseta). TMA and FMA indicate the Tabular Middle Atlas and the Folded Middle Atlas, respectively.




Figure 4. Prospective Google Earth view of the Western Moroccan Meseta. The granite rocks are red polygons and are located at the maximum topography over the summit erosional surface within the sampled catchments 60 and 61 (*i.e.*, relict landscape). The Pleistocene lava flows are deeply incised and well preserved in the deep incisions on the relict topography, respectively. The location of the picture is shown in figure 3.




Figure 5. (a) View of the summit erosional surface of the Western Moroccan Meseta near the locality of Oulmes. The typical morphologies of the granite rocks lying at the maximum elevation are shown in the foreground (black lines). (b) View of the Aguenour River valley. Note the incision across the lava flow deposits (~100 and ~50 m from the top and the base of the lava flow, respectively). (c1-c2) View of the granite outcrops close to the plateau margin. Note the rugged morphologies related to the granite incisions (green lines). The location of the pictures is shown in figure 4.




Figure 6. Longitudinal river profiles and χ -z plots with knickpoints for the NWM (a-b), CWM (c-d), and SWM (e-f). The knickpoints legend is in figure 2a. The non-lithological knickpoints are located at the slope-break zone separating segments with different k_{sn} value. (g) Longitudinal river profiles with the major non-lithological knickpoints for the three morpho-structural domains. Red rivers: CWM sector; black rivers: NWM sector; green rivers: SWM sector. The non-lithological knickpoints mark the boundary between the downstream steep river profiles and the upstream low-gradient river segments, describing a paleo-surface margin. Note, the rivers of NWM and SWM sectors come mainly from the most elevated TMA and FMA domains, respectively.




Figure 7. (a) Modern longitudinal profiles and projections of the relict landscape (black dashed segments) upstream of the highest knickpoints using $\theta_{ref} = 0.45$ and the k_{sn} of the relict portion in the χ plot. The main peaks of long-term fluvial incision are at ~ 400 m and ~ 800 m in the Western Moroccan Meseta and Tabular Middle Atlas domains, respectively (see details in Table S2). The density distribution legend is relative to the data point of the projected segments. Inset shows the frequency diagrams of the magnitude of long-term incision for the entire dataset (NWM, WMM, SWM). Note, the rivers of NWM and SWM sectors come mostly from the TMA and FMA domains, respectively. **(b)** Along-strike topographic swath profile, knickpoints elevation and magnitude of long-term incision obtained from the reconstructed river projections. SWM : South Western Meseta; CWM : Central Western Meseta; NWM : North Western Meseta; TMA : Tabular Middle Atlas. The location of the swath profile is shown in figure 3a.




Figure 8. Basin-wide denudation rate vs basin-averaged a) k_{sn} , b) slope, c) local relief (2.5 km radius), and d) precipitation rate for the sampled basins. The black and grey solid lines show the bootstrap linear best-fit, and forced linear best-fit without considering basins with exposed granite rocks. The blue trend is the power best-fit. The dashed black lines denote the best-fit considering all sampled basins. The red circles indicate catchments characterized by the contribution of quartz from granite massifs (details in fig. 4).




Figure 9. a) Predicted denudation rates based on the distribution of normalized channel steepness in the basement rocks. b) Rock-uplift history for selected rivers (bold blue rivers in figure 9a). Mean and standard deviation and error results from the river-profile inversion of six basins of the Western Meseta (details in Figs. S5 and S6 in supplementary material).




Figure 10. Schematic evolution of the Western Meseta and the Middle Atlas domains since the Lutetian (45-40 Ma). **a)** Erosional steady-state topography of the Western Meseta with marine conditions in the Middle Atlas and surrounding regions. **b)** Miocene topographic rejuvenation associated with volcanism and upstream migration of an erosional wave in response to the increase in rock-uplift rates at 22-20 Ma. Note also the southward propagation of the frontal thrusts of the Rif orogenic system with subsidence and marine deposition in the Saiss and the Skoura basins. **c)** Second slight increase in rock-uplift rates in the Western Meseta. Abrupt increase in uplift rates in the Middle Atlas as testified by uplifted marine Messinian deposits at rates of $\sim 170 \text{ m/Myr}$. Inset in figures **b** and **c** show the river profiles and the inverse modelling results of the rivers flowing through the Western Meseta.




Figure 11. a) Tridimensional view of the northern Atlas-Meseta system. **b)** Swath profile (details and location are shown in Fig. 1b) with filtered topography at 10 – 50 – 100 km wavelengths and data of long-term surface uplift of WMM, TMA, FMA and HP (this work and Pastor et al., 2015). WMM: Western Moroccan Meseta; TMA and FMA: Tabular and Folded Middle Atlas; HP: High Plateaus; SMAF: South Middle Atlas Fault; NMAF: North Middle Atlas Fault (after Gomez et al., 1996). The bottom panel indicates the projected position of Moho and lithosphere-asthenosphere boundary (LAB) beneath the Atlas-Meseta system from S-received functions (data from Miller and Becker, 2014).

Table 1. Denudation rates from cosmogenic ^{10}Be in river borne sand and bedrock samples.

| ID | Geomorphic position | Long. | Lat. | Elev. ^a | Quarz mass ^b | Carrier mass (⁹ Be) | 10Be/9Be ^c | | [¹⁰ Be] ^d 10 ³ (at./g) | | Production rate ^e | | | Denudation (m/Myr) | | Integration time ^f (yr) | |
|-------|---------------------|--------|---------|--------------------|-------------------------|---------------------------------|----------------------------|-----------|---|-------|------------------------------|------------|------------|--------------------|-------|------------------------------------|---------|
| | | | | | | | Value (10 ⁻¹⁵) | Error (%) | Value | Error | Neutrons | Slow muons | Fast muons | Value | Error | Value | Error |
| 53 | Watershed | 7.1500 | 33.5364 | 455 | 20.70 | 0.151 | 195.1 | 3.4 | 284.7 | 9.8 | 5.08 | 0.015 | 0.04 | 14.76 | 1.02 | 48081 | 3330.1 |
| 58 | Watershed | 5.7512 | 33.2640 | 1173 | 21.82 | 0.150 | 292.6 | 4.3 | 406.0 | 17.6 | 8.69 | 0.02 | 0.05 | 17.11 | 1.27 | 41624 | 3080.5 |
| 59 | Watershed | 5.5714 | 33.3807 | 1083 | 21.68 | 0.151 | 241.6 | 3.4 | 338.5 | 11.7 | 8.16 | 0.02 | 0.05 | 19.49 | 1.35 | 36639 | 2535.6 |
| 60 | Watershed | 6.2790 | 33.4496 | 914 | 21.20 | 0.150 | 238.8 | 3.4 | 339.4 | 11.8 | 7.28 | 0.02 | 0.05 | 17.36 | 1.20 | 41023 | 2845.5 |
| 62 | Watershed | 5.9743 | 33.7520 | 721 | 21.12 | 0.151 | 95.5 | 12.3 | 135.0 | 17.0 | 6.32 | 0.02 | 0.05 | 39.94 | 5.57 | 18044 | 2514.7 |
| Blank | | | | | | | 0.45 | 51.3 | | | | | | | | | |
| 61 | Watershed | 6.365 | 33.698 | 928 | 21.47 | 0.152 | 211.5 | 3.3 | 302.7 | 10.3 | 7.35 | 0.02 | 0.05 | 19.80 | 1.37 | 36076 | 2488.0 |
| 63 | Watershed | 5.931 | 33.829 | 1049 | 21.74 | 0.151 | 146.6 | 4.1 | 204.8 | 8.5 | 7.95 | 0.02 | 0.05 | 32.15 | 2.34 | 22369 | 1630.0 |
| 56A | Summit | 6.0535 | 33.4123 | 1124 | 21.84 | 0.150 | 1055.6 | 3.1 | 1467.4 | 46.0 | 8.43 | 0.02 | 0.05 | 3.46 | 0.23 | 163446 | 11064.1 |
| 56B | Summit | 6.0535 | 33.4123 | 1124 | 21.59 | 0.150 | 900.1 | 2.5 | 1259.7 | 31.2 | 8.43 | 0.02 | 0.05 | 4.11 | 0.27 | 139446 | 9053.0 |
| 57A | Ridge crest | 5.8906 | 33.1922 | 1098 | 19.45 | 0.150 | 251.2 | 3.3 | 389.0 | 12.9 | 8.23 | 0.02 | 0.05 | 14.36 | 0.98 | 41958 | 2874.7 |
| 57B | Ridge crest | 5.8913 | 33.1919 | 1099 | 17.24 | 0.151 | 239.1 | 3.5 | 423.6 | 14.9 | 8.24 | 0.02 | 0.05 | 13.13 | 0.91 | 45808 | 3185.9 |
| Blank | | | | | | | 0.788 | 23.05 | | | | | | | | | |

^a Basin mean elevation.

^b Mass of pure quartz dissolved.

^c Isotopic ratio uncertainties include statistical error on counted ^{10}Be events (N) and analytical error correction.

^d Uncertainties on ^{10}Be concentrations are calculated using the standard error propagation method using the quadratic sum of the relative errors associated to the counting statistics, AMS internal error (0.5% for ASTER), and error associated to the process blank.

^e Production rate is the spatially averaged production rate over the river basin, scaled for elevation, latitude (Stone, 2000) and geomorphic shielding (Dunne et al., 1999). The values, relative to watersheds, are calculated for a non-uniform distribution of quartz-lithologies in the sample basins (e.g., Carretier et al., 2015; Clementucci et al., 2022).

^f Integration times are calculated after Lal (1991) while their uncertainties are calculated propagating the error of the uncertainties associated with denudation rates.

Table 2. Structural topographic relief and lithospheric thickening of the Folded Middle Atlas based on the equation of Garzione et al., (2006): $dh/dt = (\rho_m - \rho_c) / \rho_c * (UH/W)$, assuming that Airy isostasy is maintained during shortening and for crustal (ρ_c) and asthenospheric mantle (ρ_m) densities of 2.8 and 3.23 kg/m³, respectively. Shortening produces thickening and surface uplift at a rate of dh/dt (assuming no surface erosion and subsidence with deposition). The parameters H (crustal thickness) and W (orogen width) at 7 Ma and the inferred thickness after 7 Ma of shortening at rates of 0.5 to 1.5 mm/yr are consistent with modern crustal thickness data reported by Miller and Becker, (2014).

| W^a min (km) | W^a max (km) | H^b min (km) | H^b max (km) | U^c min (mm/yr) | U^c max (mm/yr) | dh/dt^d min (mm/yr) | dh/dt^d max (mm/yr) |
|-------------------|-------------------|----------------|-------------------|----------------------|----------------------|--------------------------|--------------------------|
| 70 | 100 | 28 | 32 | 0.5 | 1.5 | 0.021 | 0.105 |

| Since 7 Ma | | | | | |
|-----------------------------|-----------------------------|--|--|---|---|
| Elevation gain, min (km) | Elevation gain, max (km) | Crustal root ^e min (km) | Crustal root ^e max (km) | Mean thickness of crust ^f min (km) | Mean thickness of crust ^f max (km) |
| 0.15 | 0.73 | 0.81 | 3.96 | 28.96 | 36.7 |

^a Width of the deforming Middle Atlas range.

^b Initial crustal thickness

^c Shortening rate (data from Vernant et al., 2010 and Gomez et al., 1999).

^d Surface uplift rate assuming lithospheric (ρ_c) and mantle (ρ_m) density of 2.8 and 3.23 kg/m³ (after Garzione et al., 2006).

^e Expected crustal root of the thickened crust according to Airy isostatic model (Mean gain elevation* ρ_c)/(ρ_c * ρ_m).

^f Expected crustal thickness of the thickened crust (H + Elevation gain + Crustal root)

Supplementary materials

S1. Topographic, stream analysis

A Shuttle Radar Topography Mission Digital Elevation Model (SRTM DEM, pixel size of 30 m) was used to extract the drainage network. Hillslope metrics, such as slope and local relief with a circular sampling window of 2.5 km radius, as well as basin and channel morphometric characteristics were extracted using ArcGIS and the MATLAB software packages TopoToolbox (Schwanghart and Scherler, 2014) and Topographic Analysis Kit (TAK, Forte and Whipple, 2019), together with additional MATLAB functions (*e.g.*, Gallen, 2017).

Regions characterized by recent changes in base level retain a topographic signal in the upstream river network that provides the opportunity to evaluate the magnitude and the spatial variation of such perturbations.

The relationships between channel steepness and erosion/uplift rates have been demonstrated in various geological settings (Kirby and Whipple, 2012; Miller et al., 2013; Adams et al., 2020). In detachment-limited conditions, these relationships are modulated by the stream-power model (Howard, 1994; Whipple and Tucker, 1999). Under steady-state conditions, erosion and rock uplift rates are balanced and dz/dt is equal to zero, therefore the power-law can be solved as (Flint et al., 1974):

$$S = \left(\frac{E}{K}\right)^{\frac{1}{n}} A^{-\left(\frac{m}{n}\right)} \quad (1)$$

where E is the erosion rate (*i.e.*, change in elevation (z) of a stream with respect to time (t)), A is the upstream contributing drainage area, S is the local channel slope, K is the erosional efficiency (which is function of lithology and climate), and m and n are constant values (which depend on basin hydrology, channel geometry and erosional processes). Equation (1) predicts a power relation between slope and drainage area and can be integrated, assuming E and K as spatially invariant (Perron and Royden, 2013):

$$z(x) = z(x_b) + \left(\frac{U}{KA_0^m}\right)^{\frac{1}{n}} \int_{x_b}^x \left(\frac{A_0}{A(x')}\right)^{\frac{m}{n}} dx = z(x_b) + k_s A_0^{-\left(\frac{m}{n}\right)} \chi \quad (2)$$

where:

$$\chi = \int_{x_b}^x \left(\frac{A_0}{A(x')}\right)^{\frac{m}{n}} dx' \quad (3)$$

where x_b is the base level and χ is an integral quantity, A_0 is the reference drainage area, and m/n describes the concavity of the stream. Equation (2) allows transforming river longitudinal profiles to χ versus elevation plots (here and thereafter defined as χ - z plot, where z and χ are the dependent and independent variable, respectively). Consequently, $(U/K)^{1/n}$ is the slope in the χ space, where the slope is defined as steepness index (k_s) by imposing A_0 equal to 1. The steepness index (k_s) and the concavity (m/n) can be determined through the integral approach proposed by Perron and Royden, (2013). To compare k_s values of longitudinal profiles, a typical reference concavity index (θ_{ref}) of 0.45, is applied to derive normalized channel steepness indices (k_{sn} ; Snyder et al., 2000; Wobus et al., 2006). This value represents the best-fit concavity for all rivers draining the study area that minimizes the scatter in the χ space (Perron and Royden, 2013; Goren et al., 2014; Fig. 3h).

Knickpoints were identified and discretized along the stream network based on the shape of the streams in the longitudinal profiles, the χ - z plots and using available geological maps (Service Géologique du Maroc) and satellite imagery (Google Earth). Importantly, non-lithological, transient knickpoints cluster around similar χ values (Perron and Royden, 2013) and separate stream segments with different k_{sn} values (Kirby and Whipple, 2012; Olivetti et al., 2016; Clementucci et al., 2022). Conversely, knickpoints associated with heterogeneities along the profile and lithological contrasts are usually anchored in space and do not show a significant break in χ space (Fig. S1; Kirby and Whipple, 2012). Using this approach, the knickpoints were divided into four categories: non-lithological knickpoints, minor, lithological and dam related.

S2. ^{10}Be concentration measurements and denudation rates calculations

Samples were analyzed following a protocol including: (1) crushing (only for bedrock samples) and sieving of sand to extract the 250 – 1000 μm fraction. (2) Magnetic separation with the Frantz magnetic separator

(three steps: 0.5 - 1 - 1.5 Ampere). (3) Decarbonation of carbonate-rich samples through the reaction with HCl. (4) Leaching with HCl and H₂SiF₆ solution to remove all remaining minerals except quartz. (5) Decontamination from atmospheric ¹⁰Be in the quartz grains by means of a three sequential leaching in HF. (6) Total dissolution of quartz grains, after addition of 150 µl spike of an in-house carrier of ⁹Be carrier solution (3025 ±9 µg/g), from deep-mined phenakite crystal (Merchel et al., 2008). (7) Extraction of the final amount of Be using hydrofluoric and perchloric fuming to remove fluorides; cation and anion exchange chromatography was used to eliminate iron, aluminum, manganese, and other elements. (8) Mixing of beryllium oxide with a 325-mesh niobium powder. Measurements of in situ-produced ¹⁰Be concentrations were performed at the French Accelerator mass spectrometry (AMS) national facility ASTER (Aix-en-Provence). The obtained ¹⁰Be/⁹Be ratios were corrected for procedural blanks and calibrated directly against the in-house STD-11 standard (¹⁰Be/⁹Be = (1.191±0.013) × 10⁻¹¹; Braucher et al., 2015). Analytical uncertainties (1σ) are related to external uncertainties, variation of isotopic ratios of standards during the runs, and AMS counting statistics, and were propagated into the uncertainty of the calculated denudation rates. Finally, the basin-wide denudation rates were calculated considering the three different categories of particles responsible for the in situ production of ¹⁰Be (Braucher et al., 2011), solving analytically:

$$C(x, t) = \frac{P_n}{\lambda + \frac{\rho \cdot \varepsilon}{\Lambda_n}} \cdot e^{\frac{\rho \cdot x}{\Lambda_n}} \cdot \left(1 - e^{-\left(\lambda + \frac{\rho \cdot \varepsilon}{\Lambda_n}\right)t}\right) + \frac{P_{\mu s}}{\lambda + \frac{\rho \cdot \varepsilon}{\Lambda_{\mu s}}} \cdot e^{\frac{\rho \cdot x}{\Lambda_{\mu s}}} \cdot \left(1 - e^{-\left(\lambda + \frac{\rho \cdot \varepsilon}{\Lambda_{\mu s}}\right)t}\right) + \frac{P_{\mu f}}{\lambda + \frac{\rho \cdot \varepsilon}{\Lambda_{\mu f}}} \cdot e^{\frac{\rho \cdot x}{\Lambda_{\mu f}}} \cdot \left(1 - e^{-\left(\lambda + \frac{\rho \cdot \varepsilon}{\Lambda_{\mu f}}\right)t}\right) \quad (4)$$

where $C(x, t)$ corresponds to the nuclide concentration as a function of depth x (g/cm²), ε (g/cm²/yr) represents the denudation rate and t (yr) the exposure time. P_n , $P_{\mu s}$, $P_{\mu f}$ and Λ_n , $\Lambda_{\mu s}$, $\Lambda_{\mu f}$ are the production rates and effective apparent attenuation lengths of neutrons, slow muons and fast muons, respectively. Λ_n , $\Lambda_{\mu s}$, $\Lambda_{\mu f}$ values used in this paper are 160, 1500 and 4320 g/cm², respectively (Braucher et al., 2011), λ is the radioactive decay constant of ¹⁰Be and ρ is the rock density.

Table S1. Averaged denudation rate, channel steepness, topographic slope, local relief, precipitation rate of the quartz-bearing areas exposed in the sampled basins

| Sample | Basin area | Quartz-bearing area | Quartz-bearing area | Denudation rate (m/ Myr) | | k_{sn} (m ^{0.9}) ^a | | Slope ^b | Local Relief ^c | Precipitation rate ^d |
|--------|--------------------|---------------------|---------------------|--------------------------|-------|---|-------|--------------------|---------------------------|---------------------------------|
| ID | (km ²) | (km ²) | (%) | Value | Error | Value | Error | (°) | (m) | (m/ yr) |
| 53 | 535.5 | 228.0 | 42.6 | 14.76 | 1.02 | 22.7 | 0.5 | 8.01 | 222.2 | 411.5 |
| 58 | 339.3 | 339.3 | 100.0 | 16.66 | 1.23 | 16.7 | 0.6 | 5.9 | 228.4 | 517.9 |
| 59 | 970.1 | 496.2 | 51.1 | 19.49 | 1.35 | 27.6 | 0.7 | 6.9 | 288.4 | 599.8 |
| 60 | 1007.2 | 1007.2 | 100.0 | 17.60 | 1.22 | 46.4 | 0.8 | 10.0 | 337.8 | 472.1 |
| 61 | 3195.8 | 2775.8 | 86.9 | 19.80 | 1.37 | 43.6 | 1.0 | 10.1 | 381.9 | 495.2 |
| 62 | 354.5 | 238.9 | 67.4 | 39.94 | 5.57 | 42.4 | 0.9 | 12.9 | 439.9 | 549.6 |
| 63 | 3456.3 | 2591.5 | 75.0 | 32.15 | 2.34 | 44.4 | 1.0 | 9.8 | 377.4 | 586.3 |
| 64 | - | 1646.2 | - | 35.15 | 2.58 | 53.5 | 1.5 | 11.5 | 426.7 | 579.3 |

^a Basin-averaged k_{sn} of quartz-bearing lithologies in the sampled basins.

^b Basin-averaged slope of quartz-bearing lithologies in the sampled basins.

^c Basin-averaged local relief (2.5 km radius) of quartz-bearing lithologies in the sampled basins.

^d Averaged precipitation rate for the quartz-bearing lithologies areas exposed in the sampled basins.




Figure S1. Plot showing the best-fit concavity index for the rivers of the study area, which is ~ 0.45 .




Figure S2. Morpho-structural domains in the northwestern central Morocco and selected . FMA: Folded Middle Atlas. TMA: Tabular Middle Atlas. NMAF: North Middle Atlas fault. SMAF: South Middle Atlas fault. The white polygons are the sampling catchments. The bold blue rivers are the selected streams to perform the linear inversion.




Figure S3. The slope (**a**) and local relief (2.5 km radius; **b**) maps with main knickpoints. The bold knickpoints describe the escarpment between the TMA and Moroccan Meseta, while the thick dashed line indicates the boundary between the main morpho-structural domains. The white solid lines demark the tributaries of the Oued El Rbia (South Western Meseta, SWM), of the Oued Sebou (North Western Meseta, NWM) and rivers draining directly from the Western Meseta to the Atlantic ocean (CWM: Central Western Meseta). TMA and FMA indicate the Tabular Middle Atlas and the Folded Middle Atlas, respectively.

Table S2. Non-lithological knickpoints and river profile data

| Stream ^a | Long. | Lat. | Elevation | DFM ^b | DFD ^c | Drainage area | Location ^d |
|---------------------|----------|-----------|------------|------------------|------------------|--------------------|-----------------------|
| | °W | °N | (m) | (km) | (km) | (km ²) | |
| cwm1 | 798790.6 | 3686682.0 | 952 | 201.10 | 53.50 | 402.6 | Meseta |
| cwm2 | 789008.6 | 3686828.0 | 1080 | 183.89 | 13.15 | 23.1 | Meseta |
| cwm3 | 782292.6 | 3707195.0 | 870 | 158.74 | 34.06 | 130.0 | Meseta |
| cwm4 | 771342.6 | 3723036.0 | 579 | 133.13 | 9.46 | 10.1 | Meseta |
| cwm5 | 788862.6 | 3677776.0 | 1036 | 218.14 | 23.20 | 91.6 | Meseta |
| cwm6 | 769298.6 | 3688215.0 | 829 | 175.38 | 13.76 | 22.8 | Meseta |
| cwm7 | 764407.6 | 3683616.0 | 846 | 171.84 | 9.94 | 19.5 | Meseta |
| cwm8 | 756596.6 | 3677849.0 | 780 | 179.56 | 12.64 | 34.1 | Meseta |
| cwm9 | 788643.6 | 3651058.0 | 821 | 245.55 | 48.31 | 420.5 | Meseta |
| cwm 10 | 737178.6 | 3697121.0 | 558 | 180.84 | 113.02 | 1659.8 | Meseta |
| cwm 11 | 720242.6 | 3689894.0 | 624 | 117.30 | 30.97 | 151.4 | Meseta |
| cwm 12 | 725571.6 | 3703691.0 | 587 | 99.21 | 13.76 | 24.6 | Meseta |
| cwm 13 | 730316.6 | 3704348.0 | 570 | 104.67 | 10.10 | 31.1 | Meseta |
| cwm 14 | 727615.6 | 3716685.0 | 372 | 86.58 | 16.31 | 64.2 | Meseta |
| cwm 15 | 714694.6 | 3695588.0 | 575 | 109.39 | 16.58 | 32.6 | Meseta |
| cwm 16 | 709949.6 | 3690259.0 | 550 | 108.05 | 19.47 | 68.3 | Meseta |
| cwm 17 | 711993.6 | 3704932.0 | 417 | 87.22 | 10.62 | 16.5 | Meseta |
| cwm 18 | 711190.6 | 3718875.0 | 380 | 68.00 | 10.91 | 16.9 | Meseta |
| cwm 19 | 706737.6 | 3678141.0 | 573 | 128.93 | 28.52 | 197.5 | Meseta |
| cwm 20 | 699656.6 | 3723255.0 | 309 | 64.17 | 14.03 | 28.9 | Meseta |
| cwm 21 | 703379.6 | 3652810.0 | 715 | 133.85 | 27.29 | 177.1 | Meseta |
| cwm 22 | 691918.6 | 3652883.0 | 683 | 125.31 | 18.69 | 110.4 | Meseta |
| cwm 23 | 682063.6 | 3662008.0 | 594 | 100.79 | 15.43 | 48.6 | Meseta |
| cwm 24 | 676369.6 | 3660548.0 | 625 | 98.17 | 20.55 | 84.8 | Meseta |
| cwm 25 | 664032.6 | 3670330.0 | 450 | 98.74 | 37.46 | 187.6 | Meseta |
| cwm 26 | 657389.6 | 3665585.0 | 486 | 97.54 | 8.08 | 9.7 | Meseta |
| cwm 27 | 652936.6 | 3657920.0 | 510 | 92.40 | 23.13 | 69.1 | Meseta |
| cwm 28 | 662791.6 | 3643247.0 | 641 | 124.46 | 30.87 | 215.0 | Meseta |
| cwm 29 | 638628.6 | 3649817.0 | 450 | 97.60 | 14.43 | 42.9 | Meseta |
| cwm 30 | 663813.6 | 3658869.0 | 586 | 109.10 | 17.52 | 84.7 | Meseta |
| cwm 31 | 698415.6 | 3665439.0 | 702 | 122.12 | 20.19 | 65.5 | Meseta |
| cwm 32 | 715059.6 | 3728876.0 | 208 | 60.65 | 87.61 | 766.1 | Meseta |
| cwm 33 | 748347.6 | 3692522.0 | 742 | 142.19 | 9.29 | 18.5 | Meseta |
| cwm 34 | 737178.6 | 3697121.0 | 558 | 133.22 | 28.75 | 124.6 | Meseta |
| cwm 35 | 747836.6 | 3695880.0 | 695 | 145.04 | 9.45 | 16.7 | Meseta |
| cwm 36 | 771561.6 | 3703691.0 | 1012 | 155.34 | 15.27 | 33.6 | Meseta |
| cwm 37 | 753092.6 | 3732745.0 | 290 | 105.40 | 33.15 | 175.7 | Meseta |
| cwm 38 | 742945.6 | 3747564.0 | 331 | 72.90 | 14.93 | 42.9 | Meseta |
| Mean wmm | | | 621 ± 121 | | | | |
| nwm1 | 800323.6 | 3705808.0 | 618 | 243.97 | 118.61 | 2097.4 | Meseta |
| nwm2 | 892887.6 | 3707268.0 | 1475 | 454.62 | 84.77 | 1351.2 | TMA |
| nwm3 | 895734.6 | 3741505.0 | 1202 | 383.38 | 48.15 | 290.2 | TMA |
| nwm4 | 869965.6 | 3738074.0 | 1369 | 378.39 | 33.77 | 198.9 | TMA |
| nwm5 | 857336.6 | 3743038.0 | 1013 | 304.82 | 20.06 | 76.4 | TMA |
| nwm6 | 856752.6 | 3724350.0 | 1408 | 316.88 | 31.22 | 148.9 | TMA |
| nwm7 | 820544.6 | 3717415.0 | 1175 | 255.61 | 10.79 | 35.3 | TMA |
| nwm8 | 836385.6 | 3733110.0 | 1045 | 237.99 | 19.01 | 59.4 | TMA |
| nwm9 | 783898.6 | 3718510.0 | 632 | 217.02 | 15.62 | 45.1 | Meseta |
| nwm10 | 928949.6 | 3707560.0 | 1348 | 441.63 | 49.22 | 410.3 | TMA |
| nwm11 | 812514.6 | 3724131.0 | 830 | 232.77 | 19.43 | 105.9 | Meseta |
| nwm12 | 801272.6 | 3737344.0 | 545 | 201.53 | 31.10 | 103.6 | Meseta |
| nwm13 | 790687.6 | 3710261.0 | 761 | 225.87 | 21.69 | 64.6 | Meseta |
| Mean nwm | | | 1032 ± 188 | | | | |

| | | | | | | | |
|-------------|----------|-----------|------------|--------|-------|------|-----|
| swm1 | 859161.6 | 3653467.0 | 1915 | 595.72 | 13.29 | 57.7 | FMA |
| swm2 | 847262.6 | 3670914.0 | 1532 | 595.71 | 19.90 | 84.9 | FMA |
| Mean swm | 798790.6 | 3686682.0 | 1724 ± 135 | | | | |
| Mean Meseta | | | 627 ± 116 | | | | |
| Mean TMA | | | 1255 ± 86 | | | | |
| Mean FMA | | | 1724 ± 135 | | | | |

^a cwm : river draining directly to the ocean; nwm : river draining into the Oued Sebou; swm : river draining into the Oued Er Rbia.

^b Distance from divide.

^c Distance from river mouth

^d Meseta : rivers draining the Western Moroccan Meseta sectors (WMM); TMA: rivers draining the Tabular Middle Atlas (NWM sector). FMA: rivers draining the Folded Middle Atlas (SWM sector).

Table S3. Reconstructed relict stream profile and 95% confidence interval

| Stream ^a | m/n | Elevation knickpoint | ΔZ^b | Error (2σ) | Location ^b |
|---------------------|------|----------------------|--------------|------------------------|-----------------------|
| | | (m) | (m) | (m) | |
| cwm1 | 0.45 | 952 | 608.2 | 10.1 | Meseta |
| cwm2 | 0.45 | 1080 | 845.3 | 2.7 | Meseta |
| cwm3 | 0.45 | 870 | 404.8 | 6.3 | Meseta |
| cwm4 | 0.45 | 579 | 252.3 | 11.8 | Meseta |
| cwm5 | 0.45 | 1036 | 570.8 | 14.2 | Meseta |
| cwm6 | 0.45 | 829 | 447.0 | 15.7 | Meseta |
| cwm7 | 0.45 | 846 | 506.5 | 6.9 | Meseta |
| cwm8 | 0.45 | 780 | 356.8 | 9.5 | Meseta |
| cwm9 | 0.45 | 821 | 396.7 | 3.8 | Meseta |
| cwm 10 | 0.45 | 558 | 332.5 | 6.5 | Meseta |
| cwm 11 | 0.45 | 624 | 396.6 | 4.1 | Meseta |
| cwm 12 | 0.45 | 587 | 423.5 | 4.7 | Meseta |
| cwm 13 | 0.45 | 570 | 341.5 | 13.1 | Meseta |
| cwm 14 | 0.45 | 372 | 310.7 | 14.1 | Meseta |
| cwm 15 | 0.45 | 575 | 457.8 | 4.1 | Meseta |
| cwm 16 | 0.45 | 550 | 497.4 | 7.0 | Meseta |
| cwm 17 | 0.45 | 417 | 309.5 | 13.9 | Meseta |
| cwm 18 | 0.45 | 380 | 341.4 | 4.1 | Meseta |
| cwm 19 | 0.45 | 573 | 399.0 | 22.0 | Meseta |
| cwm 20 | 0.45 | 309 | 193.7 | 14.8 | Meseta |
| cwm 21 | 0.45 | 715 | 602.0 | 1.1 | Meseta |
| cwm 22 | 0.45 | 683 | 521.9 | 1.3 | Meseta |
| cwm 23 | 0.45 | 594 | 430.3 | 6.8 | Meseta |
| cwm 24 | 0.45 | 625 | 475.9 | 2.5 | Meseta |
| cwm 25 | 0.45 | 450 | 203.9 | 1.4 | Meseta |
| cwm 26 | 0.45 | 486 | 319.2 | 9.0 | Meseta |
| cwm 27 | 0.45 | 510 | 413.5 | 8.7 | Meseta |
| cwm 28 | 0.45 | 641 | 428.1 | 4.6 | Meseta |
| cwm 29 | 0.45 | 450 | 313.6 | 4.9 | Meseta |
| Mean wmm | | | 417.3 | 54.9 | |
| nwm1 | 0.45 | 618 | 328.4 | 17.8 | Meseta |
| nwm2 | 0.45 | 1475 | 961.3 | 26.9 | TMA |
| nwm3 | 0.45 | 1202 | 722.4 | 22.6 | TMA |
| nwm4 | 0.45 | 1369 | 725.1 | 23.8 | TMA |
| nwm5 | 0.45 | 1013 | 574.3 | 41.3 | TMA |
| nwm6 | 0.45 | 1408 | 917.7 | 23.3 | TMA |
| nwm7 | 0.45 | 1175 | 944.3 | 12.6 | TMA |
| nwm8 | 0.45 | 1045 | 544.8 | 28.1 | TMA |
| nwm9 | 0.45 | 632 | 363.1 | 24.6 | Meseta |
| nwm10 | 0.45 | 1348 | 1196.1 | 17.6 | TMA |
| nwm11 | 0.45 | 830 | 635.6 | 24.1 | Meseta |
| nwm12 | 0.45 | 545 | 265.1 | 27.1 | Meseta |
| Mean nwm | | | 681.5 | 86.4 | |
| swm1 | 0.45 | 1915 | 1067.0 | 37.8 | FMA |
| swm2 | 0.45 | 1532 | 1105.4 | 89.6 | FMA |
| Mean swm | | | 1086.2 | 63.7 | |
| Mean Meseta | | | 414.9 | 54.9 | |
| Mean TMA | | | 823.2 | 66.5 | |
| Mean FMA | | | 1086.2 | 63.7 | |

^a cwm : river draining directly to the ocean; nwm : river draining into the Oued Sebou; swm : river draining into the Oued Er Rbia.

^b Estimated long-term fluvial incision from reconstructed relict stream profiles.

^c Meseta domains : rivers draining the Western Moroccan Meseta domain (mostly from cwm and nwm sector); TMA: rivers draining the Tabular Middle Atlas (mostly nwm sectors); FMA: rivers draining the Folded Middle Atlas (from swm sector).




Figure S4. Global compilation of basin-averaged slope versus basin-wide denudation rates for slow tectonic settings (modified from Olivetti et al., 2016; Clementucci et al., 2022).




Figure S5. Predicted denudation rates based on the distribution of normalized channel steepness in the basement rocks. The colored rivers represent the catchments selected for the linear inversion (details in fig. S6).




Figure S6. River-profile inversion of six basins of the Western Meseta. The river inversions are constrained using the mean erodibility value of basement rocks, $n = 1$ and $m/n = 0.45$. In the bottom τ -plots, the thick transparent trends represent the river inversion results from $K \pm 1\sigma$ ($7.34 \pm 0.58 * 10^{-7} \text{ m}^{0.1}/\text{yr}$). Location of selected rivers is shown in figure S5 through the colour scale.

References

- Adams, B. A., Whipple, K. X., Forte, A. M., Heimsath, A. M., Hodges, K. V., 2020. Climate controls on erosion in tectonically active landscapes. *Sci. adv.* 6(42), eaaz3166. <https://www.science.org/doi/10.1126/sciadv.aaz3166>.
- Braucher, R., Merchel, S., Borgomano, J., Bourlès, D. L., 2011. Production of cosmogenic radionuclides at great depth: A multi element approach. *Earth Planet. Sci. Lett.* 309(1-2), 1-9. <https://doi.org/10.1016/j.epsl.2011.06.036>.
- Braucher, R., Guillou, V., Bourlès, D. L., Arnold, M., Aumaître, G., Keddadouche, K., Nottoli, E., 2015. Preparation of ASTER in-house $^{10}\text{Be}/^{9}\text{Be}$ standard solutions. *Nucl. Instrum. Methods Phys. Res., Sect. Beam Interactions with Materials and Atoms.* 361, 335-340. <https://doi.org/10.1016/j.nimb.2015.06.012>.
- Carretier, S., Regard, V., Vassallo, R., Martinod, J., Christophoul, F., Gayer, E., Audin L., Lagane, C., 2015. A note on ^{10}Be -derived mean erosion rates in catchments with heterogeneous lithology: Examples from the western Central Andes. *Earth Surf. Process. Landf.* 40(13), 1719-1729. <https://doi.org/10.1002/esp.3748>.
- Clementucci R., Ballato P., Siame L., Faccenna C., Yaaqoub A., Essaifi A., Leanni L., Guillou V., 2022. Lithological control on topographic relief evolution in a slow tectonic setting (Anti-Atlas, Morocco). *Earth and Planet. Sci. Lett.* <https://doi.org/10.1016/j.epsl.2022.117788>.
- Dunne, J., Elmore, D., Muzikar, P., 1999. Scaling factors for the rates of production of cosmogenic nuclides for geometric shielding and attenuation at depth on sloped surfaces. *Geomorphology.* 27(1-2), 3-11. [https://doi.org/10.1016/S0169-555X\(98\)00086-5](https://doi.org/10.1016/S0169-555X(98)00086-5).
- Duxbury, J., Bierman, P. R., Portenga, E. W., Pavich, M. J., Southworth, S., Freeman, S. P., 2015. Erosion rates in and around Shenandoah National Park, Virginia, determined using analysis of cosmogenic ^{10}Be . *American Journal of Science*, 315(1), 46-76. <https://doi.org/10.2475/01.2015.02>.
- Flint, J. J., 1974. Stream gradient as a function of order, magnitude, and discharge. *Water Resources Research*, 10(5), 969-973. <https://doi.org/10.1029/WR010i005p00969>.
- Forte, A.M., Whipple, K.X., 2019. Short communication: the Topographic Analysis Kit (TAK) for TopoToolbox. *Earth Surf. Dyn.* 7, 87–95. <https://doi.org/10.5194/esurf-7-87-2019>.
- Gallen, S. F., 2018. Lithologic controls on landscape dynamics and aquatic species evolution in post-orogenic mountains. *Earth and Planet. Sci. Lett.* 493, 150-160. <https://doi.org/10.1016/j.epsl.2018.04.029>.
- Godard, V., Dosseto, A., Fleury, J., Bellier, O., Siame, L., ASTER Team., 2019. Transient landscape dynamics across the Southeastern Australian Escarpment. *Earth and Planet. Sci. Lett.* 506, 397-406. <https://doi.org/10.1016/j.epsl.2018.11.017>.
- Goren, L., Fox, M., Willett, S. D., 2014. Tectonics from fluvial topography using formal linear inversion: Theory and applications to the Inyo Mountains, California. *J. Geophys. Res., Earth Surface*, 119(8), 1651-1681. <https://doi.org/10.1002/2014JF003079>.
- Granger, D. E., Kirchner, J. W., Finkel, R., 1996. Spatially averaged long-term erosion rates measured from in situ-produced cosmogenic nuclides in alluvial sediment. *The J. Geol.* 104(3), 249-257. <http://www.jstor.org/stable/30068190>.

- Howard, A.D., Kerby, G., 1983. Channel changes in badlands. *Geol. Soc. Am. Bull.* 94, 739–752. [http://dx.doi.org/10.1130/0016-7606\(1983\)94<739:CCIB>2.0.CO;2](http://dx.doi.org/10.1130/0016-7606(1983)94<739:CCIB>2.0.CO;2).
- Kirby, E., Whipple, K. X., 2012. Expression of active tectonics in erosional landscapes. *J. Struct. Geol.* 44, 54–75. <https://doi.org/10.1016/j.jsg.2012.07.009>.
- Legrain, N., Dixon, J., Stüwe, K., von Blanckenburg, F., Kubik, P., 2015. Post-Miocene landscape rejuvenation at the eastern end of the Alps. *Lithosphere*, 7(1), 3–13. <https://doi.org/10.1130/L391.1>.
- Mandal, S. K., Lupker, M., Burg, J. P., Valla, P. G., Haghypour, N., Christl, M., 2015. Spatial variability of ¹⁰Be-derived erosion rates across the southern Peninsular Indian escarpment: A key to landscape evolution across passive margins. *Earth and Planet. Sci. Lett.*, 425, 154–167. <https://doi.org/10.1016/j.epsl.2015.05.050>.
- Matmon, A., Bierman, P. R., Larsen, J., Southworth, S., Pavich, M., Caffee, M., 2003. Temporally and spatially uniform rates of erosion in the southern Appalachian Great Smoky Mountains. *Geology*, 31(2), 155–158. [https://doi.org/10.1130/0091-7613\(2003\)031<0155:TASURO>2.0.CO;2](https://doi.org/10.1130/0091-7613(2003)031<0155:TASURO>2.0.CO;2).
- Merchel, S., Arnold, M., Aumaître, G., Benedetti, L., Bourlès, D. L., Braucher, R., Alfimov V., Freeman S.P.H.T., Steier P., Wallner, A., 2008. Towards more precise ¹⁰Be and ³⁶Cl data from measurements at the 10– 14 level: Influence of sample preparation. *Nucl. Instrum. Methods Phys. Res., Sect. Beam Interactions with Materials and Atoms.* 266(22), 4921–4926. <https://doi.org/10.1016/j.nimb.2008.07.031>.
- Miller, S. R., Sak, P. B., Kirby, E., Bierman, P. R., 2013. Neogene rejuvenation of central Appalachian topography: Evidence for differential rock uplift from stream profiles and erosion rates. *Earth and Planet. Sci. Lett.*, 369, 1–12.
- Olivetti, V., Godard, V., Bellier, O., ASTER team., 2016. Cenozoic rejuvenation events of Massif Central topography (France): Insights from cosmogenic denudation rates and river profiles. *Earth and Planet. Sci. Lett.*, 444, 179–191. <https://doi.org/10.1016/j.epsl.2016.03.049>.
- Perron, J.T., Royden, L., 2013. An integral approach to bedrock river profile analysis. *Earth Surf. Process. Landf.* 38, 570–576. <https://doi.org/10.1002/esp.3302>.
- Salgado, A. A. R., de Andrade Rezende, E., Bourles, D., Braucher, R., da Silva, J. R., Garcia, R. A., 2016. Relief evolution of the Continental Rift of Southeast Brazil revealed by in situ-produced ¹⁰Be concentrations in river-borne sediments. *Journal of South American Earth Sciences*, 67, 89–99. <https://doi.org/10.1016/j.jsames.2016.02.002>
- Scharf, T. E., Codilean, A. T., De Wit, M., Jansen, J. D., Kubik, P. W., 2013. Strong rocks sustain ancient postorogenic topography in southern Africa. *Geology*, 41(3), 331–334. <https://doi.org/10.1130/G33806.1>.
- Schwanghart, W., Scherler, D., 2014. Short communication: TopoToolbox 2 – MATLAB-based software for topographic analysis and modeling in Earth surface sciences. *Earth Surf. Dyn.* 2, 1–7. <https://doi.org/10.5194/esurf-2-1-2014>.
- Stone, J. O., 2000. Air pressure and cosmogenic isotope production. *J. Geophys. Res., Solid Earth.* 105(B10), 23753–23759. <https://doi.org/10.1029/2000JB900181>.
- Whipple, K.X., Tucker, G.E., 1999. Dynamics of the stream-power river incision model: implications for height limits of mountain ranges, landscape response timescales, and research needs. *J. Geophys. Res., Solid Earth.* 104, 17661–17674. <https://doi.org/10.1029/1999JB900120>.