

Taphonomy of early life (2.1 Ga) in the francevillian basin (Gabon): Role of organic mineral interactions

Julie Ngwal'Ghoubou Ikouanga, Claude Fontaine, Franck Bourdelle, Ahmed

Abd Elmola, Jérémie Aubineau, Olabode Bankole, Laurie Reisberg, Anne-Catherine Pierson-Wickmann, Armelle Riboulleau, Alain Trentesaux, et

al.

► To cite this version:

Julie Ngwal'Ghoubou Ikouanga, Claude Fontaine, Franck Bourdelle, Ahmed Abd Elmola, Jérémie Aubineau, et al.. Taphonomy of early life (2.1 Ga) in the francevillian basin (Gabon): Role of organic mineral interactions. Precambrian Research, 2023, 395, pp.107155. 10.1016/j.precamres.2023.107155. insu-04182352

HAL Id: insu-04182352 https://insu.hal.science/insu-04182352

Submitted on 24 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2	
3	Taphonomy of early life (2.1 Ga) in the Francevillian Basin (Gabon): Role
4	of organic mineral interactions
5 6	Julie Ngwal'ghoubou Ikouanga ^a , Claude Fontaine ^a , Franck Bourdelle ^b , Ahmed Abd Elmola ^a , Jérémie
7 8	Aubineau ^c , Olabode M. Bankole ^a , Laurie Reisberg ^d , Anne-Catherine Pierson-Wickmann ^e , Armelle Riboulleau ^f , Alain Trentesaux ^f , Claude Laforest ^a , Alain Meunier ^a , Abderrazak El Albani ^a *
9	
10	*University of Poitiers, UMR 7285 CNRS, IC2MP, F-86073, Poitiers Cedex 9, France.
11 12	sur-Oise, France
13 14	^c Geosciences Montpellier, UMR 5243, CC 60 – University of Montpellier, F-34095 Montpellier, France
15	^d Centre de Recherches Pétrographiques et Géochimiques, UMR 7358 CNRS - Université de Lorraine,
16	F-54500, Vandoeuvre-lès-Nancy, France.
17	^e University of Rennes, CNRS, Géosciences Rennes - UMR 6118, F-35000 Rennes, France.
18 19	^f UMR 8187 LOG CNRS, University of Lille, ULCO, 59655 Villeneuve d'Ascq, France.
20	*Corresponding author.
21	E-mail address: <u>abder.albani@univ-poitiers.fr</u> (A. El Albani)
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	

35 Abstract

36 The taphonomy of early soft-bodied organisms in Palaeoproterozoic sediments is not yet clearly 37 understood, even though some locations where these fossils are found present all the conditions for 38 exceptional fossil preservation. Indeed, the degree of fossil preservation has received attention, but better knowledge of the environmental conditions and associated taphonomic processes is also essential. 39 40 In the Gabonese Francevillian Basin, the discovery of macrofossils (2.1 Ga) of multicellular organisms 41 in black shales is an outstanding example of this degree of preservation. Indeed, the biological diversity, as a wide variety of fossil morphologies are observed, is associated with two major taphonomic 42 processes - moulding (lenticular-shaped forms) or early pyritization, while the fossil host rocks were 43 not deeply buried and were affected only by weak to moderate diagenesis. However, usually, the 44 mechanisms of this preservation remain difficult to assess, as the original taphonomic processes are 45 impacted by diagenesis and still misunderstood. In this way, by closely observing fossil mineralization 46 47 in four morphotypes of macrofossils and associated host rocks from mineralogical and textural points 48 of view, this work aims to provide some keys to a taphonomic comprehension of soft-bodied organism 49 preservation. After the deposition of dead organisms on the clayey sediment, an illitization process, which depends on the availability of dissolved K driven by bacterial activity, started from the first stages 50 51 of preservation by moulding the lenticular-shaped forms and proceeded in the pores of the other 52 macrofossils after their pyritization. At the fossil level, the intensity of illitization is controlled by the mode of preservation and the evolution of the associated permeability. In the nonpyritized lenticular-53 54 shaped specimens, illitization was not achieved, preserving I-S mixed-layer minerals, while in the 55 pyritized forms, the illitization degree was more extensive. In comparison, I-S mixed-layer minerals are almost absent in pyritized abiotic concretions. A second process, which occurred later, consists of 56 general chloritization from fluid circulation. Our detailed results show that each specimen behaved like 57 microsystem with its own physico-chemical and mineralogical evolution during 58 a 59 preservation/diagenesis. This finding allows us to propose a conceptual model of the taphonomic 60 history, describing the fossilization stages for each type of specimen.

61

63 1. Introduction

Knowledge of early Precambrian biospheres is primarily based on the study of exceptionally wellpreserved fossilized remains of soft-bodied organisms (Knoll, 1985). Shales appear to be very favourable host rocks for the preservation of the earliest forms of life (Ngombi-Pemba et al., 2014; Wilson and Butterfield, 2014), but the mechanisms of this preservation remain difficult to assess, as the original taphonomic processes are still largely misunderstood and can be impacted by various chemical and mineralogical transformations.

70 In fact, the preservation of soft bodies involves sufficiently slow biological degradation for the more 71 resistant organic molecules to be fossilized by imprinting or mineralization (Briggs, 2003). The early stage of this preservation is mainly controlled by the nature of the sediments and their rate of 72 73 accumulation (Briggs and Kear, 1993) in the presence or absence of biological agents (bacteria) together 74 with the water column chemistry (major elements, dissolved O2, etc.) of the depositional environment 75 (Gaines et al., 2012; Gaines and Droser, 2010). Then, diagenetic processes (e.g., compaction, 76 cementation, fluid-rock interaction, and potential hypothermal and magmatic activities) can also impact 77 the quality of fossilization (Muscente et al., 2015). The quality of the conserved fossils, i.e., the degree of resemblance to the original organisms, is controlled by the intensity of all these predominant factors, 78 79 with a direct impact on reconstructing the initial form of soft-bodied organisms and deriving information 80 on their lifestyles based on the geometric (morphologies, tracks, burrows, etc.) and chemical (organic 81 molecules, isotopes, etc.) signatures that have been preserved.

82 The Palaeoproterozoic fossil-rich black shales of the Francevillian Basin (Gabon) constitute a 83 unique case study to fill these gaps. They hold exceptionally fossilized fauna, the oldest ecosystem is 84 2.1 billion years, they consist of decimetric macrofossils (El Albani et al., 2023, 2014, 2010; Ossa et al., 2023) of soft-bodied organisms capable of movement (El Albani et al., 2019a), and they are in close 85 association with photosynthetic microbial mats (Aubineau et al., 2019, 2018; Reynaud et al., 2018). 86 87 However, even if this fossiliferous system was not affected by metamorphic conditions - thereby saving a large amount of taphonomic information (El Albani et al., 2014, 2019a; Gauthier-Lafaye and Weber, 88 1989; Ngombi-Pemba et al., 2014; Ossa Ossa et al., 2013; Weber, 1968a) - the fossilization/preservation 89 conditions and mechanisms are still unclear. An adapted mineralogical and petrographic study of these 90 91 organic matter-/clay-rich rocks should remove these uncertainties.

92 In this way, by closely observing fossil mineralization and host rocks from mineralogical and
93 textural points of view and by describing organo-mineral interactions, this work aims to provide some
94 keys to a taphonomic comprehension of soft-bodied organism preservation in Francevillian black shales.
95 The study strategy involves the following:

96 (i) Remains of both biomineralized and nonmineralized taxa in fossil deposits were discriminated97 (Conway Morris, 1986).

98 (ii) Pyritic concretions that have no apparent links with the fossil specimens were observed (El
99 Albani et al., 2023, 2019b, 2014, 2010; Ossa et al., 2023). This can help us to better understand the
100 conditions that prevailed during fossilization.

101 (iii) Organo-mineral interactions were characterized. This can serve as indirect evidence for 102 understanding the taphonomy and biogenicity of the earliest organisms (Lepot, 2020), especially clay mineral transformation, which can record palaeobiological and palaeoecological variations 103 104 (Behrensmeyer et al., 2000). In this sense, resolving the role of minerals in the fossilization of soft bodies is also a key element in understanding any environmental bias in fossil assemblages. Illustrating the 105 106 importance of organo-mineral interactions, Burges Shales fossils are certainly preserved primarily as 107 carbonaceous compressions (Butterfield, 1995), but diagenetic mineralization may have stabilized organic matter or replicated the morphology (Anderson et al., 2011; Briggs, 2003). 108

109 Although the mineralogical study of Palaeoproterozoic Francevillian rocks has already been 110 carried out in part (e.g., Aubineau et al., 2021), a systematic study such as the one presented here is 111 unprecedented. Clay minerals, specifically illite, mixed-layer minerals (MLMs) and chlorite, have been documented within or near the fossils but not on a large scale with as many samples as in this study 112 (more than 40 fossils + host rock samples). In this respect, our approach is resolutely innovative: it 113 114 systematically characterizes the mineralogy and textural arrangement of clay minerals and pyrite in the 115 host rock as well as in the fossils but independently assesses nonpyritized lenticular-shaped specimens, pyritized lobate forms, string-shaped specimens, and segmented morphologies. The results should allow 116 117 us to propose a conceptual model depicting the taphonomy and the mechanisms of mineralization during 118 fossilization and diagenesis according to the submillimetre-scale physico-chemical conditions.

119

120 2. Geological settings

121 The Franceville Basin is filled by a series of unmetamorphosed sedimentary deposits of 122 Palaeoproterozoic age resting unconformably on an Archean basement (Figs. 1 and 2) (Aubineau et al., 123 2021; Bankole et al., 2015; Bros et al., 1992; Gauthier-Lafave, 1986; Ossa Ossa et al., 2013; 124 Thiéblemont et al., 2014; Weber, 1968a). The Francevillian siliciclastic fluvial and marine sediments 125 are subdivided into four major lithostratigraphic formations, FA to FD, from oldest to youngest. The FA 126 Formation dominantly consists of fluvio-deltaic conglomerates and sandstones (Bankole et al., 2015; Gauthier-Lafaye and Weber, 1989), while the overlying FB Formation is dominated by heterolithic 127 128 marine rocks deposited during a period of tectonic subsidence and basin deepening (Reynaud et al., 129 2018). The significant lithological differences and sedimentary structures within this formation resulted 130 in several lithostratigraphic subdivisions into the FB₁ (including a, b, and c units) and FB₂ (a and b units) 131 members (Azziley Azzibrouck, 1986; Weber, 1968b). The FC Formation hosts shallow marine dolostones and cyanobacteria-hosting stromatolitic cherts interbedded with black shale horizons (Lekele 132 Baghekema et al., 2017; Préat et al., 2011), while the FD Formation is made of marine black shales with 133

interbedded volcanic tuffs, which were deposited in a transgressive phase (Thiéblemont et al., 2014).
Although large-scale synclinals and anticlinals have been formed in the Francevillian subbasin,
sediments were not deformed, which is marked by the exceptional conditions of preservation of the
entire Francevillian sedimentary stratum.

138 The Francevillian fossil deposits, which belong to the FB2 Member, are in the Moulendé quarry close to Franceville town on the northern side of the Mvengué syncline. The latter stretches along an NNW-139 140 SSE axis parallel to the edge of the basin represented by the eastern boundary of the Chaillu Massif (Fig. 141 1). The tectonic activity with the generation of normal faults predominantly in the centre of the 142 Francevillian subbasin led to the formation of well-exposed outcrops (Fig. 2); (Gauthier-Lafaye and 143 Weber, 1989; Mayaga-Mikolo, 1996; Thiéblemont et al., 2009; Weber, 1968b). Sea-level fall that began 144 during the deposition of the FB_{1c} unit and extended during the deposition of the overlying FB₂ Formation 145 caused contemporary-level to increasingly oxic conditions in the water column with transient oxygen-146 deficient bottom waters (Aubineau et al., 2021; Canfield et al., 2013; Ossa Ossa et al., 2018). Specifically, the FB₂ deposits aged 2092 ± 48 Ma (Ngwal'ghoubou Ikouanga, 2022) are composed of 147 148 Poubara sandstones interbedded with thin black shale levels (FB2a unit) and 5-m thick black shales with 149 cm-thick siltstone layers (FB_{2b} unit) (Fig. 3). Lithofacies analyses indicated that the black shale deposits of FB_{2b} were emplaced in a calm, low-energy marine environment after a large discharge of sands 150 151 (Reynaud et al., 2018). The relatively oxygenated water column, together with the cyanobacterial mats, 152 may have favoured the development and preservation of complex macroorganisms (Aubineau et al., 153 2021, 2018; Canfield et al., 2013; El Albani et al., 2019a; Reynaud et al., 2018).

154 3. Materials and Methods

165

168

The studied specimens were collected from the FB_{2b} black shales. Four morphotypes of macrofossils, including different lobate-shaped, string-shaped, segmented, and lenticular-shaped forms, as well as microbial mat communities, described as mat-related structures (MRSs), are preserved throughout this entire rock unit (Aubineau et al., 2018; El Albani et al., 2019b, 2014, 2010). Ten specimens of each morphology (total of 40) were analysed.

160 The vertical distribution of the fossiliferous content is irregular (Fig. 3). While, the interbedded siltstones

are completely devoid of macrofossils, MRSs have been described inside (Aubineau et al., 2018).

162 Finally, pyrite concretions (Reynaud et al., 2018) were also found throughout the black shales. They are

aligned in the lower part, are sparse and smaller in the middle part and form almost continuous pyritic

164 beds in the upper part of the FB_{2b} unit. Polished slab sections were made for petrographic

166 with a diamond tip to precisely determine the mineralogical and chemical nature of the specimens.

characterization at the macroscale. Microsamples were separated using a Dremel microdrill equipped

167 Specimens of all macrofossil morphological types were analysed: one unmineralized morphological

type (round shape) and three mineralized types (lobed, tubular, and segmented shapes). For each of these

Code de champ modifié

types, the specimen (SP) and the host black shale (BS) have been characterized in detail. Pyriticconcretions (clusters and nodules) present in the host black shales have also been investigated.

171 3.1. Petrographic observations

172 Petrographic observations of polished thin sections were performed to study the textural relationships,

173 morphology, mineral assemblages, and chemical composition. Polished thin sections were first

174 examined by a Nikon ECLIPSE E600 POL optical microscope coupled with a Nikon Digital Sight DS-

175 U1 camera installed with NIS-Element D software. Selected polished thin sections were carbon-coated

176 and examined using a JEOL JSM IT500 Scanning Electron Microscope at the University of Poitiers,

177 France. The scanning electron microscopy (SEM) instrument was equipped with secondary electron

178 (SE) and backscatter electron (BSE) detectors.

179 3.2. Bulk and clay mineralogy

The bulk and clay mineralogical compositions were obtained by X-ray diffraction (XRD) from randomly 180 181 oriented bulk powders and oriented clay preparations, respectively, using conventional procedures 182 (Brindley and Brown, 1980; Moore and Reynolds, 1997). The X-ray diffractograms were obtained using 183 a Bruker D8 Advance diffractometer equipped with a copper source ($\lambda CuK\alpha = 1.5418$ Å). The analytical 184 conditions were 40 kV and 40 mA; the acquisition parameters were 1 s with a step of 0.025 °20 for angular ranges of 2-65 °20 and 2-30 °20 for randomly oriented powder and oriented clay samples, 185 respectively. XRD patterns were deconvoluted in the 7-10° 2 θ range using Fitik software to 186 semiquantify mixed layer clay minerals (using peak areas). 187

188 4. Results

189 4.1. Macroscopic description of studied biotic and abiotic structures.

Here, we report a multiscale description of specimens previously described by El Albani et al. (2023,
2019b, 2014 and 2010).

192 *4.1.1 Lenticular-shaped forms*

The lenticular-shaped fossils look like subcircular capsules with a more or less pronounced central bulge (Fig. 4A-B). They can be found associated with other macrofossils (Fig. 4A), form aggregates overlapping each other, or are preserved alone (Fig. 4B). The diameters range from 0.3 to 5 cm. The inner surface comprises a core and a flat collar containing fine radial striations. The length of the collar represents 25% of the total radius. The observed shape of the lenticular fossils results from epirelief moulding, i.e., light silty and dark clayey laminae of the host sediments mould the specimens (convexity of laminae shown in Fig. 4C).

200 *4.1.2. Lobate form*

The lobate macrofossils are characterized by a rounded (Fig. 4D) or ovoid (Fig. 4E) inner part and an 201 202 outer part composed of a succession of smaller lobes delimited by deep radial striae. The lengths of the 203 specimens vary from 2 to 4 cm, and their widths vary from 1 to 2 cm. They are almost entirely 204 mineralized by phenocrystal of euhedral pyrite (2 to 3 mm). The central lobes lack striae and may be more or less smooth. The lobes are mineralized, consisting of aggregates of euhedral pyrite grains (Fig. 205 206 4F). The mineralized specimens locally display a submillimetre-sized border of more "diffuse" pyrite, 207 particularly at the lobed end - where smaller euhedral pyrite crystals are packed in a clay matrix - which 208 rapidly fades at the outer edge of the specimen. At the contact zone between the specimens and host 209 sediments, textural changes in the sediments are visible (Fig. 4F): the host sediment presents alternating 210 light silty laminae and dark clay laminae that follow the border of the contact zone, suggesting that the 211 host sediment was not consolidated at the time of fossil mineralization.

212 4.1.3. String-shaped forms

The string-shaped specimens have a millimetre-scale ovoid diameter and centimetre-scale elongated body fossils (Fig. 4G, H). The edges are clear (Fig. 4G) but may be bifid (Fig. 4H). They have been interpreted as multicellular organisms capable of moving vertically or horizontally with respect to the sediment to reach nutrients (El Albani et al., 2019a). The string-shaped structures are largely pyritized, but the remaining porosity is filled by a clayey material that is more abundant in the cores of the fossiliferous structures (Fig. 4I). The host black shale sediments are slightly deformed in the contact zone with the fossils.

220 4.1.4. Segmented forms

The segmented specimens, which are pyritized and often associated with pyritized MRSs, appear as segmented, sinuous cylinders (Fig. 5A). The length of such fossilized structures reaches several centimetres, while the maximum width is up to 2 cm. The segmented forms may have a homogeneous shape over the entire length, and a narrowing of a few millimetres has been observed. Pyritization is likely continuous between segmented fragments. Perpendicular to the elongation, cylindric fossils appear completely pyritized, although the core host septaria-type fractures are filled with a dark clayey matrix (Fig. 5B). Intergranular porosity is strongly reduced at the edges of segmented specimens.

228 4.1.5. Mat-related structures

Mat-related structures are ubiquitous in both black shale laminae and intercalated siltstone levels.
Numerous mat morphotypes, up to 10, were formed by either mat growth or mat-protected patterns
(Aubineau et al., 2018). The latter resulted from the mat preservation of abiotic structures. One of the

most common MRSs in the FB2b unit is the fairy-ring structure (Fig. 5C). In addition, the FB2b MRS
may also be associated with macrofossils.

234 4.1.6. Pyritized mineral concretions

235 Mineral concretions consist of aggregates of euhedral pyrite phenocrystals that display specific 236 morphologies. Concretions may form isolated clusters or discontinuous beds (Fig. 5D-E). The polished cross-section shows that the pyritization of concretions was intense where the crystal size increases from 237 the core to the edge (Fig. 5F). Early small euhedral crystals likely served as templates for the later 238 crystals, which are now observable. A reduction in the intergranular porosity is strongly apparent at the 239 edges. While the light silty and dark clayey laminae of the host black shales are slightly deformed near 240 241 the concretions, a general flexure following the overall structure is lacking. This suggests that the 242 sediment was partially consolidated when the pyrite crystallized.

243 4.2. Mineralogy

244 4.2.1. Host sediments

245 Based on XRD data and SEM observations, the bulk mineralogy of the host sediments close to each fossil mainly consists of quartz and clay minerals (Fig. 6A-7A-8A). Carbonates, Do (peak at 2.89 Å -246 247 dolomite, and the peak is sometimes slightly shifted to 2.90 Å - dolomite-Fe-/ankerite), are minor, and 248 pyrite, Py (peak at 2.71 Å), is in extremely low amounts. The clay fraction (Fig. 6B-7B-8B) is mainly composed of chlorite, C (characteristic peaks at 14.2 Å and 7.09 Å); illite/mica, I/M (at 10.00 Å); mixed 249 250 layers of illite/smectite, MLs and MLi (~ 10.8 Å); and smectite, Sm (16.9 Å), to a lesser extent, which was identified after ethylene glycol saturation (Fig. 6C-7C-8C). This treatment also causes a slight shift 251 towards higher 00l spacing and a reduction in the intensity of the ML peak coupled with an appearance 252 253 of a shoulder at approximately 9.8 Å.

The deconvolution of the XRD patterns for mixed layer ML phases (Fig. 9A, Tab. 1-2) shows a clear contribution of an illite-rich term in mixed layers (MLi) at 10.3 Å, which may be slightly influenced by a term richer in illite (illite-like) at 10.1 Å. Assuming that the smectite interlayers are entirely occupied by divalent cations (Ca, Mg), one can approximate the I/S ratios in both types of ML: MLi (I=94%; S=6%) and MLs (I=80%; S=20%). In both cases, illite is always the dominant component of these mixed-layer minerals (Tab. 1).

260 4.2.2. Lenticular-shaped forms (nonpyritized macrofossils)

261	The lenticular-shaped forms a	re mainly filled by quartz (3.34 Å	A) associated with clay minerals (4.49 Å)
-----	-------------------------------	------------------------------------	---

- dolomite (2.89 Å), and pyrite to a lesser extent (2.71 Å) (Fig. 6A). The clay fraction consists of chlorite

264 6B-6C). The mixed-layer MLs are characterized by a d₀₀₁ reflection centred at 10.9 Å after glycolation,

which corresponds to ordered MLs with the persistence of some smectite-rich layers (I=82%; S=18%).

266 Overall, the mineralogical assemblage is identical to the host sediment (Fig. 9B); the illite/smectite ratio

seems to be lower, indicating a lower illite content than in the sediment.

268 4.2.3. Pyritized fossils

The lobate, string-shaped and segmented specimens are dominantly filled by pyrite (Fig. 7A). Quartz and clay minerals are identified only after separation of the <2 μm clay fraction. The clay mineral assemblage, consisting of chlorite, mixed layer MLi and MLs, and illite/mica, is similar to that of the host black shale sediments (Fig. 7B). Nonetheless, smectite is present in the string-shaped structures (Fig. 7C). The illite/(MLi) ratio is globally lower than that of the host sediment (Fig. 9B, Tab. 1). Importantly, the main mineralogical differences between pyritized fossils and their host sediment are the higher abundance of mixed layers compared to illite/mica combined with the absence of carbonates.

276 4.2.4. Pyritized concretions

277 Unsurprisingly, these concretions are made of pyrite with few associated phases (Fig. 8A). Nonetheless,

278 chlorite, illite, mixed layers of MLi and quartz are observed in the $\leq 2 \mu m$ clay fraction (Fig. 8B-8C).

279 Unlike the sediment and all fossils, low amounts of MLi (10.3-10.4 Å), together with the lack of MLs,

characterize these concretions, as indicated by the high illite/MLi ratio (Fig. 9C, Tabs. 1-2).

281 4.3. Petrography

282 4.3.1. Host sediments

Along the stratigraphic sequence of the Moulendé quarry, the host sediments always consist of a clay 283 matrix, quartz, chlorite, pyrite, and dolomite, as illustrated by XRD data and SEM observations (Fig. 284 10). The clay phase proportion varies according to the granulometry of the laminae. The clayey facies 285 (Fig. 10A) correspond to deposits of a densely packed clayey matrix with chlorites, disseminated pyrite 286 287 and rounded to subangular quartz grains densely packed. The transitional facies - composed of silty clay (Fig. 10B) or clayey silt (Fig. 10C and 10D) - is marked by an increase in the grain size of quartz and 288 chlorite. Compared to the clayey facies, the quartz morphology here is closer to polycrystalline 289 aggregates. Chlorites display three 2-D morphologies, including lamellar structures, rectangular 290 platelets, and ovoidal shapes, which can exceed 50 µm in length and diameter (Fig. 10A-10B-10). In all 291 292 cases, the particles are composed of poly-lamellar stacks. In addition, the edges of the ovoids show thick grey halos, probably due to iron depletion from postdepositional alteration and/or recrystallization 293 processes. The porosity can be filled by intertwined or stacked secondary chlorite lamellae, which 294 295 developed at grain contacts with primary phases (quartz, chlorite in platelets) or as coatings (Fig. 10D). 296 In these facies, pyrite is either disseminated, corresponding to euhedral to subeuhedral crystals, or intergrown with chlorites (Fig. 10C). Although pyrite is a minor phase, its proportion is higher than inthe clayey facies.

299 4.3.2. Nonpyritized macrofossils: Lenticular-shaped forms

The clay material of the lenticular-shaped forms is mainly filled by illite particles, while quartz grains are scarce (Fig. 10E). On the other hand, chlorites in flexuous lamellae, oriented parallel to the stratification, seem to be much more abundant than in the sediments, although chlorite with platelet and ovoid shapes are present in low amounts. The size (10 to $20 \,\mu\text{m}$) of the latter is generally smaller than that of the host sediment. Finally, tiny anhedral, disseminated pyrite grains are observed.

305 *4.3.3. Pyritized fossils*

Clayey assemblages identified by XRD fill the intergranular pores between pyrite crystals that preserved
the mineralized macrofossils, while detrital minerals (e.g., quartz, primary chlorite, and dolomite) are
almost absent (Fig. 10F-10G).

The lobate, string-shaped, and segmented specimens are essentially composed of illite I/M (Fig. 10F). Micrometric illite (I/M) is mainly in the form of stacked sheets (Fig. 10) and shows signs of free growth within the pore spaces. Chlorite has developed as short platelets in the intragranular porosity, probably at the expense of illite as encompassing relict patches of illitic matrix (Fig. 10F). Inside the porosity, chlorite lamellae are interbedded with illite as "accordion" structures. Secondary quartz patches are also observed (Fig. 10G).

315 4.3.3. Pyritized mineral concretions

Pyrite concretions has allowed the development of strongly interlocking pyrite phenocrystals, reducing the porosity (Fig. 10H-10I). The clay assemblage that fills the residual porosity of plated concretions seems to be diagenetic considering their size and shape, and does not contain an illite-rich matrix. In the nodules, part of the porosity is occupied by the illite. However, chlorite is more abundant than illite (Fig. 10I).

321 5. Discussion

322 Processes impacting soft-bodied organism preservation

323 Studying the degree of conservation of biological bodies is essential for understanding taphonomy.

- 324 Moreover, it is of great interest to elucidate the environmental conditions and taphonomic mechanisms
- that prevailed during the different stages of preservation (Behrensmeyer et al., 2000).
- 326 The preservation of soft bodies involves sufficiently slow biological degradation for the more resistant
- 327 organic molecules to be fossilized by imprinting or mineralization (Briggs, 2003). The early stage of

328 preservation is mainly controlled by the nature of the sediments and their rate of accumulation (Briggs 329 and Kear, 1993) in the presence or absence of biological agents (bacteria) together with the water column 330 chemistry (major elements, dissolved O₂, etc.) of the depositional environment (Gaines et al., 2012; Gaines and Droser, 2010). Compaction, cementation, fluid-rock interaction, and potential hypothermal 331 332 and magmatic activities can also impact the quality of fossilization (Muscente et al., 2015).. Such parameters have a direct impact on reconstructing the initial form of soft-bodied organisms and deriving 333 334 information on their lifestyles based on the geometric (morphologies, tracks, burrows, etc.) and chemical (organic molecules, isotopes, etc.) signatures that have been preserved. 335

336

337 Taphonomic information from pyritization

338 As previously presented, in the FB_{2b} formation (El Albani et al., 2010, 2014, 2019, Aubineau et al. 2019), pyritization is not specific to the body fossils but is also associated with abiotic concretions in 339 the host black shales throughout the stratigraphic sequence. This is expected because the black shale 340 341 sediments contained sufficient organic matter and iron to promote pyritization independently of the carbon origin (Canfield et al., 2013). Petrographic observations of lenticular nonpyritized specimens 342 343 indicate that moulding materials are inherited from the host sediments. The similarity of the detrital and 344 diagenetic clay materials in both the host sediments and lenticular infilling is not observed when 345 pyritized macrofossils and their associated host sediments are compared.

346 Pyritization of soft bodies related to bacterial sulfate reduction (Gabbott et al., 2004) can occur when 347 organic biodegradation is relatively rapid (Briggs et al., 1996) and takes place in clay-rich sediments (Farrell, 2014) under the oxic-anoxic interface (Berner, 1984). Under these conditions, the formation of 348 349 hydrogen sulfide and the occurrence of reactive iron concentration gradients in or around the labile 350 organic matter of dead organisms (Guan et al., 2017) favour the nucleation of iron sulfide (Allen, 2002) or iron-rich clays (Petrovich, 2001). Rapid organic matter remineralization produces carbon dioxide (or 351 352 methane) and water. The degradation rate is a function of several factors, such as (i) the physico-353 chemical properties of the sediments, (ii) the characteristics of the decomposing organic matter, and (iii) 354 the microbial populations involved (Muscente et al., 2017). In the studied case, the coexistence of 355 different taphonomic styles in the same sedimentary laminae cannot be explained by variations in criteria 356 (i) and/or (ii). On the other hand, the speed of degradation can play an important role in favouring 357 biomineralization. This mechanism depends on the biodegradability of the main molecular compounds (carbohydrates, lipids, proteins) of the organic matter (e.g., (Stankiewicz et al., 2000), i.e., their ability 358 359 to be photodegraded, hydrolysed or incorporated into the biomass by serving as nutrients for 360 microorganisms (Briggs, 1999). Rapidly decomposing organic matter, known as labile material, is easily 361 attacked by enzymatic catabolism, and its chemical constituents are easily reused by microbial metabolism. In contrast, the organic materials that are part of the composition of structural bodies 362

(envelopes, partitions, etc.) can be recalcitrant to decomposition (*e.g.*, Butterfield, 2003; Tegelaar et al.,
1989).

365

366 Taphonomic information from clay minerals and their evolution during diagenesis

367 The Francevillian biota lived in a shallow marine environment where periodically resuspended fine 368 particles eventually sedimented into fine-grained laminae (Reynaud et al., 2018) in an oxic to suboxic water column (Aubineau et al., 2021; Canfield et al., 2013). The finely laminated horizons are composed 369 370 of quartz and clay minerals associated with low amounts of dolomite and pyrite. The primary clay 371 mineralogical assemblage of Francevillian sediments evolved differently depending on whether it was 372 located within the preserved specimens by (i) simple compression casting or by (ii) mineralization 373 (biogenic or abiotic). The former mechanism can be compared to the Burgess Shale-type preservation 374 (Butterfield, 1995; Gaines et al., 2005) but without the overprint of metamorphism. Regardless of the preservation mechanism, all macrofossils are in laminae where clayey microfacies prevail in overall 375 376 low-permeability rocks. This suggests a direct relationship between the nature of the original organism, 377 its taphonomy, and the rock morphological elements that have been preserved.

378 The degree, condition, and mechanism of fossil preservation are strongly controlled by mineral 379 transformations and/or neo-formations, particularly clay minerals (e.g., (Anderson et al., 2011; Gabbott 380 et al., 2001; Martin et al., 2004) and associated phases (sulfides, carbonates, etc.). During early 381 diagenesis, anoxic microenvironments develop in association with organic matter degradation; Fe² ions 382 adsorbed on the preserved biopolymers may favour nucleation of crystals of iron(II)-rich clay minerals 383 (berthierine or ferroan saponite), which form a coating on the organic remains and/or replace parts of the organism (Petrovich, 2001). Nevertheless, these minerals associated with both pyritized and 384 385 nonpyritized specimens evolve during burial diagenesis, leading to the formation of new phases 386 inherited from their initial chemical composition. Therefore, the specific study of these minerals is 387 justified to reveal the conditions and mechanisms of fossil preservation in ancient sediments.

388 At the millimetre to centimetre scale of the fossil microsites and their host sediments, the dominant 389 mechanism of clay mineral transformation associated with the early stages of preservation of Francevillian soft-bodied organisms is illitization. This authigenesis is represented first by smectite 390 391 sensu stricto and then smectite-rich mixed-layer illite/smectite, namely, MLs. Since only a few MLs are still preserved, extensive illitization occurred. The intensity of this mechanism is primarily controlled 392 by the potassium availability in the system (Cuadros and Linares, 1996). Considering the absence of K-393 394 feldspars and evidence of K-rich fluids in the study area, the K supply is of paramount importance. Near 395 the surface, dissolved potassium in the overlying water column was theoretically available (Santiago 396 Ramos et al., 2018; Sun et al., 2015). However, seawater's sodium concentration and ionic strength are 397 not very favourable for direct solid-solution exchange and absorption of potassium in the interlayer spaces of swelling minerals (smectites and illite/smectite mixed layers). Alternatively, potassium may 398

399 have been supplied by the degradation of MRSs that naturally fixed K for cellular machinery (Aubineau 400 et al., 2019). The latter mechanism, in the marine environment, has long been known to regulate 401 microorganism growth and cellular osmotic pressure (Rhoads and Epstein, 2003) and to collaborate in 402 many metabolic reactions, such as protein synthesis and enzyme activation (Stautz et al., 2021). Thus, 403 a similar mechanism of K concentration can be argued for bacteria that degrade the organic matter of dead organisms, which is the most plausible explanation. In this case, heterotrophic microbial respiration 404 405 degrading the organic matter would have promoted K release and concentration in the interstitial pore waters of the unconsolidated sediments. This K increase would have favoured the conversion of swelling 406 407 smectite into MLs in the early stages of burial and, ultimately, illite, which would correspond to induced 408 biogenic mineralization. Despite the presence of K in the marine system, in both cases, the end of bacterial activity stopped K accumulation in the preserving microsystem and thus reduced the illitization 409 410 mechanism.

411 Illitization is a general mineral transformation that affects both the host sediments and the studied specimens. This mechanism involves the transformation of MLs into MLi as transitional phases and 412 413 then into illite. Nevertheless, the degree of this transformation is variable regarding the proportion of 414 illite-MLi-MLs phases. When the K content was sufficient, the MLi could be completely transformed into illite-like and/or illite phases. However, here, all samples have remaining MLs and/or MLi. In the 415 416 case of the nonpyritized fossils, illitization was not achieved, as testified by the preservation of 417 illite/smectite mixed layers (MLs) and the lower illite/MLi ratio. The same occurred in the pyritized 418 specimens, but illitization was more intense, i.e., a higher illite/MLi ratio, except for segmented forms. 419 For other forms, the host sediments present higher illite/MLi ratios than fossils, while the pyrite 420 concretions show MLi that is very enriched in K and a higher illite/MLi ratio. These heterogeneous 421 results highlight that each fossil represents a microsystem whose location, nature and rate of degradation influence the availability of potassium, controlling the intensity of illitization. 422

The illitization mechanism was followed by chloritization, as shown by the development of secondary chlorites from the microcrystalline illitic matrices and/or recrystallized MLi in "accordion-like" structures. The presence of many chlorite shapes is due to the mixing of primary and secondary phases (Fig. 10). In the fossils, ovoids and laths are observed in low amounts on the outer rims associated with reduced porosity and are absent in the central part. Conversely, "accordion-like" structures and radial shapes are much more abundant in the central parts of the pyritized specimens but absent in the nonpyritized circular forms.

In host sediments and fossils, chlorite shapes indicate different origins. Ovoid forms are clearly detrital. The other shapes are probably secondary products from direct precipitation (fibroradial) or from transformation of a precursor (accordion-like shapes). In the latter case and in the illitic clay matrix, chloritization seems to be developed at the expense of the smectitic part of the clay phases and encapsulates residual S–MLs–MLi. Under these conditions, both transformation and dissolution/precipitation mechanisms are probably involved. In contrast, in the concretions,
chloritization results from direct precipitation in a free pore space, leading to the development of
chlorite-dolomite paragenesis in the intergranular porosity. Regardless of the study area, chloritization
occurred after the early phase of illitization and during burial diagenesis, benefiting from fluid
circulation that supplied the necessary Fe and Mg.

440

441 Towards a taphonomic conceptual model

442 Our results allow us to summarize the taphonomic history by a conceptual model (Fig. 11), describing443 the fossilization stages of each type of specimen within a microsystem.

After their death, the specimens are deposited on clayey sediment that is mainly composed of quartz, clay minerals, small amounts of dolomites and pyrite. The specimens are quickly buried within specific microenvironments, where all of the fossilization mechanisms take place. In this way, lenticular specimens undergo nonpyritized fossilization, while the lobed, segmented and tubular forms are pyritized.

After decomposition of the organism's body, lenticular forms are trapped in a sediment that is still loose
but coherent enough to maintain the mould. The voids left by decomposition are progressively filled by
particles via size sorting, favouring a high concentration of fine material within the specimen.

452 The mode of fossilization of the pyritized forms is approximately identical. After their deposition, the 453 specimens are degraded progressively by sulfate-reducing bacteria during burial activity. The 454 conservation of the specimens is conditioned by the speed of degradation of the internal structures and 455 favoured by the bacterial activity. Early pyritization permits 3D preservation of each specimen.

After this early stage of fossilization, a secondary diagenetic overprint participates in the stabilization 456 of the specimens. This is represented by the illitization of the smectitic phases. This illitization is 457 458 controlled by the availability of K and the permeability of each microsystem. Due to the high 459 concentration of finer materials, nonpyrite specimens present microsystems in which exchanges with 460 the outside are more reduced compared to pyrite specimens and concretions. The intensity of the illitization reaction increases progressively from the nonpyritic forms to the abiotic concretions through 461 the pyritic specimens. These microsystems are later subjected to generalized chloritization, reflecting 462 463 the circulation of fluids rich in Fe and Mg. Abiotic pyrite concretions are formed during late diagenesis.

464 As summarized here, our data provide robust evidence that clay minerals, such as smectite, illite, 465 illite/smectite mixed layers, and chlorite, can be used as indicators for the physico-chemical changes 466 that occur at the local scale during the preservation of soft-bodied organisms.

467 6. Conclusion

468 The 2.1 Ga Francevillian fossils preserved in black shales show two early preservation modes corresponding to moulding/compression and pyritization. In the pore spaces of the sediments and fossils, 469 470 the mechanisms of clay transformation associated with the evolution of the system are successively illitization and chloritization. The intensity of these processes and the habit of the authigenic clay 471 472 minerals depend on the degree of porosity and key element availability, such as K. Thus, the intensity 473 of illitization at the early stage differs between the sediment, pyritized and nonpyritized fossils and pyrite 474 concretions of the same age. The potassium availability may have been controlled by the microbial 475 activity of bacterial mats everywhere in the black shale levels, leading to the biodegradation of organic 476 matter from soft bodies. In this scheme, each fossil represents a microsystem. At a late stage, 477 chloritization begins, and its intensity is determined by the contents of soluble iron and magnesium from 478 fluid circulation.

In the same depositional setting, we observe heterogeneous clay mineralogical assemblages, depending 479 on the presence or absence of organic matter remains. We also observe contrasting clay mineralogy by 480 481 comparing biotic and abiotic pyritized specimens. In fact, preservation (moulding and pyritization) 482 occurred within microsites, whose properties significantly impacted the clay mineral reactions, which 483 cemented the pore spaces and thus contributed to the final fossilization. Fossil microsites must be 484 considered microsystems that have undergone physical and chemical changes that are different from 485 those of their immediate surrounding environment, which is similar to additional indirect evidence of 486 the existence of biological remains encapsulated in the sediment. These data could represent a diagnostic 487 tool for detecting traces of life in rocks from the early Earth.

488

489

Author Contributions: A.E.A. designed the research. A.E.A., J.N.I., C.F and F.B. wrote the
manuscript. A.E.A, J.N.I, O.B., A.T and C.L. did the field work. J.N.I., C.F., A.M., A.A.E., A.R. and
A.E.A. prepared the samples and performed XRD analyses and models. L.R., A.C.W. Performed Re/Os
datations.

494

495 Acknowledgment

Funding was provided by La Region Nouvelle Aquitaine, French Embassy at Libreville (Gabon). They
are acknowledged for their support and collaboration. The authors are also grateful to the National
Agency of National Parcs, le CENAREST, Socoba BTP for their assistance and access to the localities.

Agency of National Parcs, le CENAREST, Socoba BTP for their assistance and access to the localities.
We appreciate C. Lebailly, R. Oslisly, J.L. Albert, C. Boissard, L.Tromas S. Ventalon, J.C. Baloche for

500 their assistance and support.

510 Table 1 – Ratios of areas of d_{001} reflections of illite/mica and illite/smectite mixed-layers

511 rich in illite, measured from diffractogram patterns of air-dried oriented preparations.

Host sediment and specimen				Major peak area	
associated		Illite (10.0 Å)	MLi (≈10.5 Å)	Ratio	
		(cts.°2q)	(cts.°2q)	Aillite/AMLi	
Circular (P)	max.	469	3774	0.12	
	min.	215	3654	0.06	
Lobated (L)	max.	658	2719	0.24	
	min.	118	1100	0.11	
Segmented (S)	max.	1540	3664	0.42	
Segmented (S)	min.	41	2303	0.02	
Tubular (T)	max.	1475	1432	1.03	
	min.	578	2706	0.21	
Congrations (Co)	max.	119	42	2.85	
Concretions (Co)	min.	427	1603	0.27	

515 and deconvolution of diffractograms. Empty circles: traces.

		Smectite s.s.	MLs	MLi and Illite like	Illite s.s. Mica
Host rock		0	0	••	•••
Lenticular fossil n	on pyritized	••	•••	••••	0
Lobate fossil p	oyritized	•	••	•••	•
Segmented fossil p	oyritized	•	••	••••	0

Table 2 – Semi-quantitative overview of the mineralogy of the clay phases based on XRD analysis

	Tubular fossil Concretions	pyritized pyrite	•	••	••	•
516						
517						

- 519 Figure 1 Geological map of the Francevillian Basin (Gabon).
- 520 Figure 2 Geological section showing the structure and the lithostratigraphic sequence of the
- 521 Francevillian deposits on the northern edge of Mvengué syncline.
- 522 Figure 3 Lithostratigraphy of the studied section (Moulendé quarry).
- 523 Figure 4 Photographs showing the spatial relations between: A) pyritized lobate (pl) and non-
- 524 pyritized lenticular shaped specimens. Arrows represent at top left a lenticular-shaped specimen and
- 525 pyritised specimen at the right side. B) top view and C) cross section of unpyritized lenticular-shaped
- 526 specimen. Arrows represent a sharp limit between a lenticular-shaped specimen and host-sediment.
- 527 **D**) pyritized round lobated shapes, **E**) pyritized oval lobated shape, **F**) cross-section of a pyritized
- 528 lobated shape. Arrows represent a sharp limit between the pyritized specimen and host-sediment with
- the evidence of pre-compaction. G) simple tubular shape, H) bifid tubular shape, I) cross-section of
 tubular shape. Abbreviations: (SP) specimen, (BS) black shale. Scale 1 cm.
- tubular shape. Abbreviations. (51) specificit, (55) black shale. Scale I chi.

531 Figure 5 – Photographs showing: A) pyritized segmented shape in contact with the bacterial mat

532 (arrow), **B**) cross-section of segmented shape. Arrow represent the sharp and pre-compactional limit

- 533 between specimen and host-sediment. C) Fairy ring structure type of bacterial mat (Aubineau et al
- 534 2018), **D**) plated coarse pyrite concretion, **E**) top view and **F**) cross-section of nodular pyrite
- 535 concretion. Arrow represent the contact between coarse pyrite crystals cross-cutting the host-sediment.
- 536 Abbreviations: (SP) specimen, (BS) black shale host. Scale 1 cm.
- 537 Figure 6 A) Powder diffractograms showing bulk mineral composition of the host sediments and
- 538 non pyritized macro-fossils (ciculars), **B**) Diffractograms of oriented air-dry preparations **C**)
- 539 Diffractograms of oriented preparations after ethylene glycol solvation. Minerals: (C) Chlorite, (Do)
- 540 Dolomite, (I/M) illite and/or Mica, (MLi) Mixed-layer illite-rich, (MLs) Mixed-layers smectite-rich,
- 541 (Py) Pyrite, (Q) Quartz.
- 542 Figure 7 A) Powder diffractograms showing the bulk mineral composition of the host sediments and
- 543 pyritized macro-fossils (lobates, tubulars and segmenteds), **B**) Diffractograms of oriented air-dry
- 544 preparations C) Diffractograms of oriented preparations after ethylene glycol solvation. Minerals: (C)
- Chlorite, (Do) Dolomite, (I/M) illite and/or Mica, (MLi) Mixed-layer illite-rich, (MLs) Mixed-layers
 smectite-rich, (Py) Pyrite, (Q) Quartz, (Sm) Smectite.
- 547 Figure 8-A) Powder diffractograms showing the bulk mineral composition of the host sediments and
- 548 concretion, B) Diffractograms of oriented air-dry preparations and C) Diffractograms of oriented
- 549 preparations after ethylene glycol solvation. Minerals: (C) Chlorite, (Do) Dolomite, (I/M) illite and/or
- 550 Mica, (MLi) Mixed-layer illite-rich, (MLs) Mixed-layers smectite-rich, (Py) Pyrite, (Q) Quartz, (Sm)
 551 Smectite.
- SSI Shieetite.
- 553 (A) host sediment, (B) macrofossil samples, and (C) concretion. Distribution of clay minerals

- 554 proportions in quaternary plot with illite (I), Chlorite (C), Mixed-layers smectite-richer (MLs) and
- 555 illite-rich (Mli) end-members. Minerals: (C) Chlorite, (I) Illite and (MLi) Mixed-layers illite-rich
- 556 and/or illite-like, (MLs) Mixed-layers smectite-rich.
- 557 Figure 10 Micrographs in backscattered electron (BSE) of the host sediments (A, B, C, D), non-
- 558 pyritized macrofossils (E), infilling of intergranular porosity of pyritized macrofossils (F, G), and
- 559 infillings of intergranular porosity of plated (H) and nodular (I) pyrite concretions. Abbreviations: (C)
- 560 Chlorite, (Do) Dolomite, (IM) Illite/Mica, (MA) Undifferentiated Clay Matrix, (Py) pyrite, (Q) Quartz.
- Figure 11 Conceptual model illustrating the taphonomy and mechanisms of mineralization during
 burial diagenesis.
- 563
- 564







613 Figure 4















664 Figure 9:



669 Figure 10:



672 Figure 11 (Suite)





698 Figure 12



702 References

703	Allen, R.E., 2002. Role of diffusion-precipitation reactions in authigenic pyritization. Chemical
704	Geology 182, 461-472. https://doi.org/10.1016/S0009-2541(01)00334-5
705	Anderson, E.P., Schiffbauer, J.D., Xiao, S., 2011. Taphonomic study of Ediacaran organic-walled
706	fossils confirms the importance of clay minerals and pyrite in Burgess Shale-type
707	preservation. Geology 39, 643-646. https://doi.org/10.1130/G31969.1
708	Aubineau, J., Bankole, O.M., Baron, F., Grégoire, B., El Albani, A., 2021. Authigenic kaolinite and
709	sudoite in sandstones from the Paleoproterozoic Franceville sub-basin (Gabon). Comptes
710	Rendus. Géoscience 353, 209-226. https://doi.org/10.5802/crgeos.62
711	Aubineau, J., El Abani, A., Bekker, A., Somogyi, A., Bankole, O., Macchiarelli, R., Meunier, A.,
712	Riboulleau, A., Reynaud, JY., Konhauser, K.O., El Albani, A., 2019. Microbially induced
713	potassium enrichment in Paleoproterozoic shales and implications for reverse weathering on
714	early Earth. Nature Communications 10, 2370. https://doi.org/10.1038/s41467-019-10620-3
715	Aubineau, J., El Albani, A., Chi Fru, E., Gingras, M.K., Batonneau, Y., Buatois, L., Geffroy, C.,
716	Labanowski, J., Laforest, C., Lemée, L., Mángano, M., Meunier, A., Pierson-Wickmann, A
717	C., Recourt, P., Riboulleau, A., Trentesaux, A., Konhauser, K.O., 2018. Unusual microbial
718	mat-related structural diversity 2.1 billion years ago and implications for the Francevillian
719	biota. Geobiology 16, 476-497. https://doi.org/10.1111/gbi.12296
720	Azziley Azzibrouck, G., 1986. Sedimentologie et geochimie du francevillien b (proterozoique
721	inferieur). Metallogenie des gisements de manganese de moanda, gabon (These de doctorat).
722	Strasbourg 1.
723	Bankole, O.M., El Albani, A., Meunier, A., Gauthier-Lafaye, F., 2015. Textural and paleo-fluid flow
724	control on diagenesis in the Paleoproterozoic Franceville Basin, South Eastern, Gabon.
725	Precambrian Research 268, 115–134. https://doi.org/10.1016/j.precamres.2015.07.008
726	Behrensmeyer, A.K., Kidwell, S.M., Gastaldo, R.A., 2000. Taphonomy and Paleobiology.
727	Paleobiology 26, 103–147.
728	Berner, R.A., 1984. Sedimentary pyrite formation: An update. Geochimica et Cosmochimica Acta 48,
729	605-615. https://doi.org/10.1016/0016-7037(84)90089-9
730	Birck, J.L., Barman, M.R., Capmas, F., 1997. Re-Os Isotopic Measurements at the Femtomole Level
731	in Natural Samples - Birck - 1997 - Geostandards Newsletter - Wiley Online Library [WWW
732	Document]. URL https://onlinelibrary.wiley.com/doi/10.1111/j.1751-908X.1997.tb00528.x
733	(accessed 11.4.22).
734	Briggs, D., 2003. The Role of Decay and Mineralization in the Preservation of Soft-Bodied Fossils.
735	Annual Review of Earth and Planetary Sciences 31, 275–301.
736	https://doi.org/10.1146/annurev.earth.31.100901.144746

737	Briggs, D., Raiswell, R., Bottrell, S., Hatfield, D., Bartels, C., 1996. Controls on the pyritization of
738	exceptionally preserved fossils: an analysis of the Lower Devonian Hunsrück Slate of
739	Germany. American Journal of Science 296, 633-663.
740	Briggs, D.E.G., 1999. Molecular taphonomy of animal and plant cuticles: selective preservation and
741	diagenesis. Philos Trans R Soc Lond B Biol Sci 354, 7–17.
742	https://doi.org/10.1098/rstb.1999.0356
743	Briggs, D.E.G., Kear, A.J., 1993. Decay and Preservation of Polychaetes: Taphonomic Thresholds in
744	Soft-Bodied Organisms. Paleobiology 19, 107-135.
745	Brindley, G.W., Brown, G., 1980. Crystal Structures of Clay Minerals and their X-Ray Identification.
746	https://doi.org/10.1180/mono-5
747	Bros, R., Stille, P., Gauthier-Lafaye, F., Weber, F., Clauer, N., 1992. Sm-Nd isotopic dating of
748	Proterozoic clay material: An example from the Francevillian sedimentary series, Gabon.
749	Earth and Planetary Science Letters 113, 207-218. https://doi.org/10.1016/0012-
750	821X(92)90220-P
751	Butterfield, N.J., 2003. Exceptional Fossil Preservation and the Cambrian Explosion1. Integrative and
752	Comparative Biology 43, 166–177. https://doi.org/10.1093/icb/43.1.166
753	Butterfield, N.J., 1995. Secular distribution of Burgess-Shale-type preservation. Lethaia 28, 1–13.
754	https://doi.org/10.1111/j.1502-3931.1995.tb01587.x
755	Canfield, D.E., Ngombi-Pemba, L., Hammarlund, E.U., Bengtson, S., Chaussidon, M., Gauthier-
756	Lafaye, F., Meunier, A., Riboulleau, A., Rollion-Bard, C., Rouxel, O., Asael, D., Pierson-
757	Wickmann, AC., El Albani, A., 2013. Oxygen dynamics in the aftermath of the Great
758	Oxidation of Earth's atmosphere. Proceedings of the National Academy of Sciences 110,
759	16736–16741. https://doi.org/10.1073/pnas.1315570110
760	Conway Morris, S., 1986. The community structure of the Middle Cambrian Phyllopod Bed (Burgess
761	Shale) The Palaeontological Association [WWW Document]. URL
762	https://www.palass.org/publications/palaeontology-journal/archive/29/3/article_pp423-467
763	(accessed 11.4.22).
764	Creaser, R.A., Papanastassiou, D.A., Wasserburg, G.J., 1991. Negative thermal ion mass spectrometry
765	of osmium, rhenium and iridium. Geochimica et Cosmochimica Acta 55, 397-401.
766	https://doi.org/10.1016/0016-7037(91)90427-7
767	Cuadros, J., Linares, J., 1996. Experimental kinetic study of the smectite-to-illite transformation.
768	Geochimica et Cosmochimica Acta 60, 439-453. https://doi.org/10.1016/0016-
769	7037(95)00407-6
770	El Albani, A., Bengtson, S., Canfield, D.E., Bekker, A., Macchiarelli, R., Mazurier, A., Hammarlund,
771	E.U., Boulvais, P., Dupuy, JJ., Fontaine, C., Fürsich, F.T., Gauthier-Lafaye, F., Janvier, P.,
772	Javaux, E., Ossa, F.O., Pierson-Wickmann, AC., Riboulleau, A., Sardini, P., Vachard, D.,
773	Whitehouse, M., Meunier, A., 2010. Large colonial organisms with coordinated growth in

774	oxygenated environments 2.1 Gyr ago. Nature 466, 100–104.
775	https://doi.org/10.1038/nature09166
776	El Albani, A., Bengtson, S., Canfield, D.E., Riboulleau, A., Rollion Bard, C., Macchiarelli, R.,
777	Ngombi Pemba, L., Hammarlund, E., Meunier, A., Moubiya Mouele, I., Benzerara, K.,
778	Bernard, S., Boulvais, P., Chaussidon, M., Cesari, C., Fontaine, C., Chi-Fru, E., Garcia Ruiz,
779	J.M., Gauthier-Lafaye, F., Mazurier, A., Pierson-Wickmann, A.C., Rouxel, O., Trentesaux,
780	A., Vecoli, M., Versteegh, G.J.M., White, L., Whitehouse, M., Bekker, A., 2014. The 2.1 Ga
781	Old Francevillian Biota: Biogenicity, Taphonomy and Biodiversity. PLoS ONE 9, e99438.
782	https://doi.org/10.1371/journal.pone.0099438
783	El Albani, A., Mangano, M.G., Buatois, L.A., Bengtson, S., Riboulleau, A., Bekker, A., Konhauser,
784	K., Lyons, T., Rollion-Bard, C., Bankole, O., Lekele Baghekema, S.G., Meunier, A.,
785	Trentesaux, A., Mazurier, A., Aubineau, J., Laforest, C., Fontaine, C., Recourt, P., Chi Fru, E.,
786	Macchiarelli, R., Reynaud, J.Y., Gauthier-Lafaye, F., Canfield, D.E., 2019a. Organism
787	motility in an oxygenated shallow-marine environment 2.1 billion years ago. Proc Natl Acad
788	Sci U S A 116, 3431–3436. https://doi.org/10.1073/pnas.1815721116
789	El Albani, A., Mangano, M.G., Buatois, L.A., Bengtson, S., Riboulleau, A., Bekker, A., Konhauser,
790	K., Lyons, T., Rollion-Bard, C., Bankole, O., Lekele Baghekema, S.G., Meunier, A.,
791	Trentesaux, A., Mazurier, A., Aubineau, J., Laforest, C., Fontaine, C., Recourt, P., Chi Fru, E.,
792	Macchiarelli, R., Reynaud, J.Y., Gauthier-Lafaye, F., Canfield, D.E., 2019b. Organism
793	motility in an oxygenated shallow-marine environment 2.1 billion years ago. Proc Natl Acad
794	Sci USA 116, 3431-3436. https://doi.org/10.1073/pnas.1815721116
795	A. El Albani, K.O. Konhauser, A. Somogyi, J. Ngwalghoubou Ikouanga, A. Lamboux, J. Blichert-
796	Toft, E. Chi-Fru, C. Fontaine, A. Mazurier, A. Riboulleau, AC. Pierson-Wickmann, and F.
797	Albarède, 2023. A search for life in Palaeoproterozoic marine sediments using Zn isotopes and
798	geochemistry. Earth Planetary Sciences Letter (In Press).
799	Farrell, Ú.C., 2014. Pyritization of Soft Tissues in the Fossil Record: An Overview. The
800	Paleontological Society Papers 20, 35-58. https://doi.org/10.1017/S1089332600002795
801	Gabbott, S.E., Norry, M.J., Aldridge, R.J., Theron, J.N., 2001. Preservation of fossils in clay minerals;
802	a unique example from the Upper Ordovician Soom Shale, South Africa. Proceedings of the
803	Yorkshire Geological Society 53, 237-244. https://doi.org/10.1144/pygs.53.3.237
804	Gabbott, S.E., Xian-guang, H., Norry, M.J., Siveter, D.J., 2004. Preservation of Early Cambrian
805	animals of the Chengjiang biota 4.
806	Gaines, R.R., Droser, M.L., 2010. The paleoredox setting of Burgess Shale-type deposits.
807	Palaeogeography, Palaeoclimatology, Palaeoecology 297, 649-661.
808	https://doi.org/10.1016/j.palaeo.2010.09.014
809	Gaines, R.R., Hammarlund, E.U., Hou, X., Qi, C., Gabbott, S.E., Zhao, Y., Peng, J., Canfield, D.E.,
810	2012. Mechanism for Burgess Shale-type preservation. Proceedings of the National Academy
811	of Sciences 109, 5180-5184. https://doi.org/10.1073/pnas.1111784109

812	Gaines, R.R., Kennedy, M.J., Droser, M.L., 2005. A new hypothesis for organic preservation of
813	Burgess Shale taxa in the middle Cambrian Wheeler Formation, House Range, Utah.
814	Palaeogeography, Palaeoclimatology, Palaeoecology, Interpretation of Biological and
815	Environmental Changes across the Neoproterozoic-Cambrian Boundary 220, 193-205.
816	https://doi.org/10.1016/j.palaeo.2004.07.034
817	Gauthier-Lafaye, F., 1986. Les gisements d'uranium du Gabon et les réacteurs d'Oklo. Modèle
818	métallogénique de gîtes à fortes teneurs du Protérozoïque inférieur. Sciences Géologiques,
819	bulletins et mémoires 78.
820	Gauthier-Lafaye, F., Weber, F., 1989. The Francevillian (Lower Proterozoic) uranium ore deposits of
821	Gabon. Economic Geology 84, 2267–2285. https://doi.org/10.2113/gsecongeo.84.8.2267
822	Guan, C., Wang, W., Zhou, C., Muscente, A.D., Wan, B., Chen, X., Yuan, X., Chen, Z., Ouyang, Q.,
823	2017. Controls on fossil pyritization: Redox conditions, sedimentary organic matter content,
824	and Chuaria preservation in the Ediacaran Lantian Biota. Palaeogeography,
825	Palaeoclimatology, Palaeoecology, Biosedimentary records of China from the Precambrian to
826	present. 474, 26-35. https://doi.org/10.1016/j.palaeo.2016.05.013
827	Karhu, J., Holland, H., 1996. Carbon isotopes and the rise of atmospheric oxygen. Geology 24.
828	https://doi.org/10.1130/0091-7613(1996)024<0867:CIATRO>2.3.CO;2
829	Knoll, A.H., Whittington, H.B., Morris, S.C., 1985. Exceptional preservation of photosynthetic
830	organisms in silicified carbonates and silicified peats. Philosophical Transactions of the Royal
831	Society of London. B, Biological Sciences 311, 111-122.
832	https://doi.org/10.1098/rstb.1985.0143
833	Lekele Baghekema, S.G., Lepot, K., Riboulleau, A., Fadel, A., Trentesaux, A., El Albani, A., 2017.
834	Nanoscale analysis of preservation of ca. 2.1 Ga old Francevillian microfossils, Gabon.
835	Precambrian Research 301, 1-18. https://doi.org/10.1016/j.precamres.2017.08.024
836	Martin, D., Briggs, D., Parkes, R., 2004. Experimental attachment of sediment particles to invertebrate
837	eggs and the preservation of soft-bodied fossils. Journal of The Geological Society - J GEOL
838	SOC 161, 735-738. https://doi.org/10.1144/0016-764903-164
839	Mayaga-Mikolo, F., 1996. Chronologie des evenements sedimentaires, magmatiques et tectono-
840	metamorphiques du precambrien d'afrique centrale occidentale (gabon) : tectogenese ogooue
841	et heritage archeen (These de doctorat). Clermont-Ferrand 2.
842	Moore, D.M., Reynolds, R.C., 1997. X-ray diffraction and the identification and analysis of clay
843	minerals, 2nd ed. ed. Oxford University Press, Oxford.
844	Muscente, A.D., Hawkins, A.D., Xiao, S., 2015. Fossil preservation through phosphatization and
845	silicification in the Ediacaran Doushantuo Formation (South China): a comparative synthesis.
846	Palaeogeography, Palaeoclimatology, Palaeoecology, Ediacaran Environments and
847	Ecosystems 434, 46-62. https://doi.org/10.1016/j.palaeo.2014.10.013
848	Muscente, A.D., Schiffbauer, J.D., Broce, J., Laflamme, M., O'Donnell, K., Boag, T.H., Meyer, M.,
040	Hauking A.D. Hundley, I.W. McNamore, M. MacKamie, I.A. Stanlay, C.D. Hinmon

849 Hawkins, A.D., Huntley, J.W., McNamara, M., MacKenzie, L.A., Stanley, G.D., Hinman,

850	N.W., Hofmann, M.H., Xiao, S., 2017. Exceptionally preserved fossil assemblages through
851	geologic time and space. Gondwana Research 48, 164–188.
852	https://doi.org/10.1016/j.gr.2017.04.020
853	Ngombi-Pemba, L., Albani, A.E., Meunier, A., Grauby, O., Gauthier-Lafaye, F., 2014. From detrital
854	heritage to diagenetic transformations, the message of clay minerals contained within shales of
855	the Palaeoproterozoic Francevillian basin (Gabon). Precambrian Research 255, 63-76.
856	https://doi.org/10.1016/j.precamres.2014.09.016
857	Ngwal'ghoubou Ikouanga J. A. Les interactions argiles-biota paléoprotérozoïque (Gabon) et leurs
858	implications dans la préservation taphonomique. Thèse de doctorat. Université de Poitiers.
859	UFR des sciences fondamentales et appliquées (faculté). 235p
860	Ossa, F., Pons, ML., Bekker, A., Hofmann, A., Poulton, S., Andersen, M., Agangi, A., Gregory, D.,
861	Reinke, C., Steinhilber, B., Marin-Carbonne, J., Schoenberg, R., 2023. Zinc enrichment and
862	isotopic fractionation in a marine habitat of the c. 2.1 Ga Francevillian Group: A signature of
863	zinc utilization by eukaryotes? https://doi.org/10.1016/j.epsl.2023.118147
864	Ossa Ossa, F., El Albani, A., Hofmann, A., Bekker, A., Gauthier-Lafaye, F., Pambo, F., Meunier, A.,
865	Fontaine, C., Boulvais, P., Pierson-Wickmann, AC., Cavalazzi, B., Macchiarelli, R., 2013.
866	Exceptional preservation of expandable clay minerals in the ca. 2.1Ga black shales of the
867	Francevillian basin, Gabon and its implication for atmospheric oxygen accumulation.
868	Chemical Geology 362, 181–192. https://doi.org/10.1016/j.chemgeo.2013.08.011
869	Ossa Ossa, F., Hofmann, A., Wille, M., Spangenberg, J.E., Bekker, A., Poulton, S.W., Eickmann, B.,
870	Schoenberg, R., 2018. Aerobic iron and manganese cycling in a redox-stratified Mesoarchean
871	epicontinental sea. Earth and Planetary Science Letters 500, 28-40.
872	https://doi.org/10.1016/j.epsl.2018.07.044
873	Petrovich, R., 2001. Mechanisms of Fossilization of the Soft-Bodied and Lightly Armored Faunas of
874	the Burgess Shale and of Some Other Classical Localities. American Journal of Science 301,
875	683-726. https://doi.org/10.2475/ajs.301.8.683
876	Préat, A., Bouton, P., Thiéblemont, D., Prian, JP., Ndounze, S.S., Delpomdor, F., 2011.
877	Paleoproterozoic high δ 13C dolomites from the Lastoursville and Franceville basins (SE
878	Gabon): Stratigraphic and synsedimentary subsidence implications. Precambrian Research
879	189, 212-228. https://doi.org/10.1016/j.precamres.2011.05.013
880	Reynaud, JY., Trentesaux, A., El Albani, A., Aubineau, J., Ngombi-Pemba, L., Guiyeligou, G.,
881	Bouton, P., Gauthier-Lafaye, F., Weber, F., 2018. Depositional setting of the 2·1 Ga
882	Francevillian macrobiota (Gabon): Rapid mud settling in a shallow basin swept by high-
883	density sand flows. Sedimentology 65, 670-701. https://doi.org/10.1111/sed.12398
884	Rhoads, D., Epstein, W., 2003. Osmotic control ofkdp operon expression in Escherichia coli (
885	potassium transport / lac fusion / turgor pressure Semantic Scholar [WWW Document]. URL
886	https://www.semanticscholar.org/paper/Osmotic-control-ofkdp-operon-expression-in-coli-

887	%28-%2F-Rhoads-Epstein/abbf75eb6c4081904997d6a057b597a991a72604 (accessed
888	11.4.22).
889	Santiago Ramos, D.P., Morgan, L.E., Lloyd, N.S., Higgins, J.A., 2018. Reverse weathering in marine
890	sediments and the geochemical cycle of potassium in seawater: Insights from the K isotopic
891	composition (41K/39K) of deep-sea pore-fluids. Geochimica et Cosmochimica Acta,
892	Chemistry of oceans past and present: A Special Issue in tribute to Harry Elderfield 236, 99-
893	120. https://doi.org/10.1016/j.gca.2018.02.035
894	Selby, D., Creaser, R., 2004. Macroscale NTIMS and microscale LA-MC-ICP-MS Re-Os isotopic
895	analysis of molybdenite: Testing spatial restrictions for reliable Re-Os age determinations, and
896	implications for the decoupling of Re and Os within molybdenite. Geochimica et
897	Cosmochimica Acta 68, 3897–3908. https://doi.org/10.1016/j.gca.2004.03.022
898	Selby, D., Creaser, R., 2003. Re-Os geochronology of organic rich sediments: An evaluation of
899	organic matter analysis methods. Chemical Geology 200, 225-240.
900	https://doi.org/10.1016/S0009-2541(03)00199-2
901	Stankiewicz, B.A., Briggs, D.E.G., Michels, R., Collinson, M.E., Flannery, M.B., Evershed, R.P.,
902	2000. Alternative origin of aliphatic polymer in kerogen. Geology 28, 559-562.
903	https://doi.org/10.1130/0091-7613(2000)28<559:AOOAPI>2.0.CO;2
904	Stautz, J., Hellmich, Y., Fuss, M.F., Silberberg, J.M., Devlin, J.R., Stockbridge, R.B., Hänelt, I., 2021.
905	Molecular Mechanisms for Bacterial Potassium Homeostasis. J Mol Biol 433, 166968.
906	https://doi.org/10.1016/j.jmb.2021.166968
907	Sun, H., Chen, M., Zou, L., Shu, R., Ruan, R., 2015. Study of the kinetics of pyrite oxidation under
908	controlled redox potential. Hydrometallurgy 155, 13-19.
909	https://doi.org/10.1016/j.hydromet.2015.04.003
910	Tegelaar, E.W., de Leeuw, J.W., Derenne, S., Largeau, C., 1989. A reappraisal of kerogen formation.
911	Geochimica et Cosmochimica Acta 53, 3103-3106. https://doi.org/10.1016/0016-
912	7037(89)90191-9
913	Thiéblemont, D., Bouton, P., Préat, A., Goujou, JC., Tegyey, M., Weber, F., Ebang Obiang, M.,
914	Joron, J.L., Treuil, M., 2014. Transition from alkaline to calc-alkaline volcanism during
915	evolution of the Paleoproterozoic Francevillian basin of eastern Gabon (Western Central
916	Africa). Journal of African Earth Sciences 99, 215–227.
917	https://doi.org/10.1016/j.jafrearsci.2013.12.007
918	Thiéblemont, D., Castaing, C., Billa, M., Bouton, P., Préat, A., 2009. NOTICE EXPLICATIVE DE
919	LA CARTE GÉOLOGIQUE ET DES RESSOURCES MINÉRALES DE LA RÉPUBLIQUE
920	GABONAISE À 1/1 000 000 384.
921	Völkening, J., Walczyk, T., G. Heumann, K., 1991. Osmium isotope ratio determinations by negative
922	thermal ionization mass spectrometry. International Journal of Mass Spectrometry and Ion
923	Processes 105, 147-159. https://doi.org/10.1016/0168-1176(91)80077-Z

924	Weber, F., 1968a. Une série précambrienne du Gabon : le Francevillien. Sédimentologie, géochimie,
925	relations avec les gites minéraux associés. Sciences Géologiques, bulletins et mémoires 28.
926	Weber, F., 1968b. SÉDIMENTOLOGIE, GÉOCHIMIE, RELATIONS AVEC LES GITES
927	MINÉRAUX ASSOCIÉS 331.
928	Wilson, L.A., Butterfield, N.J., 2014. Sediment Effects on the Preservation of Burgess Shale-Type
929	Compression Fossils. PALAIOS 29, 145-153.
930	