



HAL
open science

Sedimentary evidence of tsunamis in New Caledonia, southwest Pacific

Raphael Paris, Bernard Pelletier, Jean Roger, Patrick Wassmer, Pierre
Sabatier

► **To cite this version:**

Raphael Paris, Bernard Pelletier, Jean Roger, Patrick Wassmer, Pierre Sabatier. Sedimentary evidence of tsunamis in New Caledonia, southwest Pacific. *Marine Geology*, 2023, 463, 10.1016/j.margeo.2023.107116 . insu-04197033

HAL Id: insu-04197033

<https://insu.hal.science/insu-04197033v1>

Submitted on 16 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Sedimentary evidence of tsunamis in New Caledonia, southwest 2 Pacific

3

4 Raphaël Paris¹, Bernard Pelletier², Jean Roger³, Patrick Wassmer⁴, Pierre Sabatier⁵

5 ¹ Université Clermont Auvergne, CNRS, IRD, OPGC, Laboratoire Magmas et Volcans, F-63000 Clermont-
6 Ferrand, France.

7 ² IRD, UMR Géoazur (UCA, CNRS, IRD, OCA), antenne de Nouméa , BP A5, 98848 Nouméa, Nouvelle-
8 Calédonie

9 ³ GNS Science, 1 Fairway Drive, Avalon 5010, PO Box 30368, Lower Hutt 5040, New Zealand

10 ⁴ Retired from Université de Strasbourg

11 ⁵ EDYTEM, Université de Savoie Mont-Blanc, CNRS, F-73376 Le Bourget du Lac, France.

12

13 Abstract

14 This study aims to identify tsunamis in the coastal sedimentary archives of New Caledonia (Grande Terre and the
15 Loyalty Islands) in the southwest Pacific, to extend a relatively short historical tsunami record (150 years), and
16 to bring more constraints on the frequency of the main tsunamigenic earthquakes in the region. Using a
17 combination of sedimentary and geochemical approaches on sediment core samples, we document three types of
18 deposits related to marine inundation: (1) bioclastic sand layers; (2) crypto-deposits with a marine chemical
19 signature; and (3) accumulations of rounded pumice lapilli. Most of the sampling sites being located outside the
20 morpho-dynamic impact zone of the cyclones, a tsunami origin is preferred. The spatial distribution of the
21 deposits suggests tsunami runups exceeding 2.5 m a.s.l. on the northeastern coast of Grande Terre, 6 m on the
22 eastern coast of Maré Island, and 10 m on the northeastern coast of Lifou Island. Finally, at least six tsunamis
23 were preserved in the sedimentary record: one event soon after 1000 BCE, two events during the first
24 millennium CE, and at least three events during the last millenium (including a 15th century tsunami, the 28
25 March 1875 tsunami, and a tsunami possibly related to the 1729 earthquake in Vanuatu).

26 **Keywords:** tsunami deposits, paleotsunami, New Caledonia, Loyalty Islands, Melanesia, Pacific Islands

27

28 1. Introduction

29 Oceania is exposed to a high diversity and high frequency of natural hazards, both of climatic
30 (hurricanes, floods, landslides, droughts, fires, sea-level rise) and tectonic origin (earthquakes,
31 volcanic eruptions, landslides, tsunamis) (Pararas-Carayannis, 2007). The Melanesian region
32 (including Papua New Guinea, the Solomon Islands, Vanuatu, Fiji, and New Caledonia) is

33 particularly exposed to volcanic and tectonic hazards because it is located at the junction
34 between the Australian and Pacific plates, resulting in two high-convergence subduction
35 zones (Fig. 1): the Vanuatu Arc (former New Hebrides) and the Tonga-Kermadec Arc (Bevis
36 et al., 1995; Pelletier et al., 1998; Calmant et al., 2003). As a consequence, this region is
37 among the most seismic areas on Earth, with ~20 Mw 8.0+ earthquakes having occurred over
38 the last three centuries with related tsunamis (NCEI/WDS Global Historical Tsunami
39 Database: https://www.ngdc.noaa.gov/hazard/tsu_db.shtml), and it is also exposed to trans-
40 Pacific tsunamis generated by remote earthquakes (Goff et al., 2022).

41 In this paper, we focus on tsunami hazard in New Caledonia through the study of the
42 sedimentary record of paleo-tsunamis. Three catalogues dedicated to tsunamis that have
43 affected New Caledonia were published over the last thirty years (Louat and Baldassari, 1989;
44 Sahal et al., 2010; Roger et al., 2019). Sahal et al. (2010) enlarged the original catalogue of
45 Louat and Baldassari (1989) to 18 events based on archives and witness reports,
46 distinguishing local, regional, and trans-oceanic tsunamis. Roger et al. (2019) later updated
47 the catalogue using a decision table based on seismic and sea-level data, thus reaching a list of
48 37 tsunamis reported with certainty or recorded in New Caledonia from March 1875 to
49 January 2019. Since then, 10 tsunamis with amplitudes lower than 1 m were detected on the
50 coastal gauge data provided by REFMAR (<https://data.shom.fr>). Among all tsunamis listed in
51 the catalogues, only 8 generated local wave runups around or higher than 2 m on the coasts of
52 New Caledonia. Some tsunamis are reported with certainty and clearly related to local or
53 regional earthquakes (1875, 2007, 2018), while others are reported by testimonials and their
54 origin and dates are uncertain (1936, 1942-1943, 1950-1951). The 28 March 1875 tsunami
55 caused by a large earthquake with an estimated magnitude Mw 8.1 in southern Vanuatu was
56 the oldest and the largest, and also the only deadly event reported, with 25 deaths at Lifou,
57 Loyalty Islands (Louat and Baldassari, 1989; Sahal et al., 2010; Le Duff et al., 2016;
58 Ioualalen et al., 2017). The two most recent tsunamis triggered by local earthquakes on the
59 southern part of the Vanuatu subduction zone (the 5 December 2018 Mw 7.5 Tatine
60 earthquake, and the 11 February 2021 Mw 7.7 Matthew earthquake) were particularly well
61 recorded by the tide gauge network, and thus used to validate and/or improve local and
62 regional tsunami hazard models and warning systems (Gusman et al., 2022; Roger et al.,
63 2021, 2023).

64 However, the New Caledonia tsunami catalogue covers a time period of ~150 years, which is
65 relatively short compared to the timescales of geophysical processes and the return period of

66 large earthquakes. The collection of written information about natural hazards began with the
67 arrival of European settlers in 1854. The first settlers (Lapita) arrived approximately 3500
68 years ago, then followed by the Kanaks. Nothing from the Lapita knowledge was transmitted
69 and most of the Kanak oral tradition was lost due to wars and diseases brought by Europeans.
70 Hence, this study aims to identify tsunamis in the sedimentary archives to extend the tsunami
71 record of New Caledonia beyond historical data, and thus bringing more constraints on the
72 tsunamigenic earthquakes frequency in the region.

73

74 **2. Methods**

75 We surveyed coastal lowland sites on Grande Terre (northeastern coast), Ile des Pins, and the
76 Loyalty Islands (Maré and Lifou Islands) (Fig. 1). More than 40 stratigraphic logs were
77 obtained based on hand-held trenching and coring (using an Edelman auger, a gouge auger,
78 push cores, and U-channels). The cores retrieved were 63 mm large and up to 1 m long.
79 Topographic transects were locally realized using a clinometer and a GPS. Sites were selected
80 based on their morphology, preservation from anthropic activities, and space available for
81 sediment accumulation (e.g., coastal marshes behind sand ridges). Because the coasts of New
82 Caledonia are regularly exposed to tropical cyclones (Bruyère et al., 2022), we tried to focus
83 our investigations on sites that are spared by the morpho-dynamic impact of wave surges
84 during cyclones, although a non-turbulent water level rise cannot systematically be ruled out.

85 Non-destructive chemical analyses of 7 core samples were performed at down core resolution
86 of 2 mm using the Avaatech X-ray Fluorescence (XRF) core scanner of the EDYTEM
87 laboratory (Université Savoie Mont Blanc). A first run at 10 kV and 0.6 mA for 20 s was
88 performed to detect lightweight elements, such as Al, Si, Mg, K, Ca, Ti, Mn, S and Fe, and a
89 second run was performed at 30 kV and 0.1 mA for 30 s to detect Br, Rb, Sr, Ni, Cu, and Zr
90 (Richter et al., 2006). Each spectrum was converted by a deconvolution process into relative
91 components (intensities) expressed in counts per second (cps). Filtered data were mean-
92 centred and normalized to the standard deviation (Supplementary data 1). Principal
93 Component Analysis (PCA) was then realized using Past software Version 4.01 (Hammer et
94 al., 2001) to determine correlations between the different measured elements and to identify
95 sediment endmembers. Element ratios are expressed as logarithms (ln) of XRF count ratios to
96 avoid potential matrix effects (Weltje and Tjallingii, 2008). Previous studies on storm and
97 tsunami deposits have demonstrated that XRF core-scanner data is particularly relevant for

98 identifying marine flooding events, characterising the vertical variations of the chemical
99 composition of the sediment, and the different sediment sources (Sabatier et al., 2010; Cuven
100 et al., 2013; Chagué-Goff et al., 2017; Biguenet et al., 2020; Paris et al., 2021).

101 Grain size analysis was performed on bulk samples (100 g for each unit and/or subunit)
102 collected from trenches or cores. Organic matter was removed using hydrogen peroxide.
103 Sediments were then washed with KCl or HCl to remove flocculating ions and agitated for 4-
104 6 h in distilled water and sodium hexametaphosphate. A T34 sieve size was used to remove
105 particles exceeding 2 mm in diameter. Grain size measurements from 0.04 to 2000 µm were
106 performed using a Beckman Coulter LS-230 (EOST Laboratory, University of Strasbourg).

107 Pumice samples were collected on the eastern coast of Maré Island (Fig. 1: Patho site) to
108 determine their origin. Their whole-rock major element composition was determined using
109 Inductively-Coupled Plasma Atomic Emission Spectroscopy (ICP-AES, Jobin-Yvon Ultima
110 C) at LMV. Whole-rock trace element composition was obtained by solution Inductively-
111 Coupled Plasma Mass-Spectrometry (ICP-MS, Agilent-7500, Agilent Technologies) at LMV.

112 A total of 27 samples of wood fragments, organic sediment, and marine shells were ¹⁴C dated
113 at LMC14 (Laboratoire de Mesure du Carbone 14, Gif-sur-Yvette) using accelerator mass
114 spectrometry (Dumoulin et al., 2017; Moreau et al., 2013). Calibration at 2σ was performed
115 using the SHCal2020 calibration curve (Hogg et al., 2020) for wood and sediment samples
116 and the Marine20 calibration curve (Heaton et al., 2020) for marine shell samples (Table 1).

117

118 **3. Results**

119 3.1. Taaré Bay, Grande Terre

120 Taaré Bay is located on the southeastern coast of Grande Terre between Yaté and Goro (Fig.
121 2). Coring was performed in the northern part of the bay, where an old isthmus is filled by
122 lagoon sediments. The main sedimentary facies is a dark brown peat that turns reddish at the
123 base of the sequence. No evidence of high-energy marine events could be found. The coral
124 substrate (gravel), corresponding to the inner facies of an old back-reef platform, was reached
125 at a depth of 0.6 m. This old coral platform was found at many other sites along the north-
126 eastern coast of Grande Terre (sites that are reported herein), thus limiting the preservation
127 potential of tsunami deposits in the sedimentary record.

128

129 3.2. Bwa, Lebris Bay, Grande Terre

130 Lebris Bay or Bwa Bay is located on the eastern coast of Grande Terre (Fig. 2). The site is a
131 good candidate for preserving tsunami deposits because (1) the coral barrier reef is
132 discontinuous on this coast, thus facilitating tsunami propagation inside the lagoon and to the
133 coast, and (2) the elongated shape of the bay (1.3 km large and 7 km long, opened to the
134 Southeast) makes it a good trap for long waves. The NW-SE orientation of the bay (i.e.
135 perpendicular to the coastline) reduces the possibility for having strong wave surges during
136 cyclones, even though its altitude is very low (< 1 m a.s.l.). A 1 m long core (BAW-01) was
137 retrieved from a small swamp at the very end of the bay (Fig. 4). From 0 to 50 cm depth, the
138 sediment is a coarse sand (medium sand to very fine pebbles) in a dark brown silty matrix,
139 turning to gravel in a brown silty matrix from 50 to 100 cm depth. Varied clast lithology
140 reflects the location of the bay at the transition between peridotites to the East and volcano-
141 sedimentary rocks to the West (Maurizot et al., 2020). Wood fragments are observed all along
142 the core. The abundant silty matrix makes it difficult to read the structure of the core, but
143 there are subtle variations in the grain size. Rounded pumice lapilli are well preserved in
144 coarser horizons (up to medium pebble size) at depths of 10 cm, 15 cm, and 22.5 cm (Fig. 3).

145 XRF chemical analysis of core BAW-01 was realized from 0 to 50 cm depth. The results
146 obtained from PCA with the two first dimensions (representing 53% of the total variance)
147 allow to distinguish two different chemical endmembers: (a) Si, Al, K, Ca, Ti (with a positive
148 loading on PCA first dimension – PCA 1), and (b) Fe, Mn, Zr, Zn, Cu, Cl (with a positive
149 loading on PCA second dimension – PCA 2). Other elements, such as Sr and Br, have their
150 own signatures. Br has a strong negative loading on PCA 2 (29% of the total variance),
151 whereas Sr has a positive loading on PCA 3 (14% of the total variance). Coarse-grained
152 horizons and pumice clasts are associated with a decrease in Ti and Ca (PCA 1), a slight
153 decrease in Fe (PCA 2) and an increase in Sr (PCA 3 and Sr/Ti ratio), as seen, for instance, at
154 15 cm depth in Fig. 3.

155 Two wood fragments were ¹⁴C dated, giving inverted ages of 773-903 cal. CE at 21 cm depth
156 and 1501-1598 cal. CE at 47 cm depth (Table 1).

157

158 3.3. Koulnoue – Womi, Grande Terre

159 The Koulnoue (KOU) and Womi (WOM) sampling sites are located 600-700 m from the
160 shoreline in a mangrove South-East of the Léwé Bay on the northeastern coast of Grande
161 Terre (Fig. 4). The mangrove is protected from the sea by a 300 m large, 2.5 m high sand
162 ridge that is eroded but not overtopped during cyclones. Nowadays the bay has a narrow
163 estuary cut in the Palaeocene limestones (Maurizot et al., 2020). This small pass was opened
164 in the 1930s for economical use. The locals told us that previously there was another pass
165 north of the lagoon at least until the 1950s. The water level rises in the bay during cyclones,
166 but our sampling sites are not affected by the wave surges due to their distal location. Cores
167 from Koulnoue and Womi are typically 50-120 cm long, and a coarse coral gravel at their
168 base often prevented any deeper coring. Here, we focus on the KOUL-03 (Fig. 5) and WOM-
169 02 (Fig. 6) sections, which are the most representative of the dataset.

170 At Koulnoue, the basal coral gravel is covered by a 45-110 cm thick sequence of brown silty
171 sediment, with poorly defined coarser-grained layers or horizons enriched in fine-to-medium
172 sand and plant fragments (e.g., silt-sand layer at 39 cm depth on core KOUL-03, Fig. 5). The
173 sand size fraction is mainly composed of angular grains of calcite (~80%) and quartz (~20%).
174 A rounded pumice lapilli is found buried in a silt-sand layer at 26-27 cm depth. No marine
175 bioclasts were detected. Intense bioturbation and roots are serious limitations for ¹⁴C dating in
176 mangrove environments, but we could obtain an age of 1809-1947 cal. CE on well-preserved
177 silty sediment at a depth of 35 cm in core KOUL-03 (Table 1). PCA of XRF data shows that
178 (1) Ca and Sr and (2) Cl and Br are clearly distinct from the other elements (Si, Ti, K, Fe, Zr,
179 Cu, etc.). Sr and Ca are strongly correlated, with a negative loading on PCA 2, which
180 represents 14% of the variance (Fig. 7). Br and Cl are the only elements that have a negative
181 loading on PCA 1 (53% of the variance), and they are clearly opposed to Ca and Sr on PCA 2.
182 There is little overlap between the different sedimentary facies on the PCA biplot (Fig. 7). The
183 vertical profile of chemical composition shows little variation, except for small peaks of Ca/Ti
184 and Sr/Ti that are associated with negative PCA 1 at depths of 15 cm, 27-29 cm and 39-40 cm
185 in the silts (Fig. 5). The basal coral gravel is characterized by high ratios of Ca/Ti and Sr/Ti.

186 At Womi, the 120 cm-long sequence is made of brown-to-grey poorly sorted silty sediment
187 with intercalated layers of poorly sorted bioclastic sand or coarse silt (Figs. 6). The grain size
188 distributions of both the background silt sediment and the bioclastic layers are finely skewed,
189 and they often show a slope break at the silt-sand transition. The number of coarse-grained
190 bioclastic layers varies from one section to another (up to 5 layers identified in core WOM-
191 02), but there is a good stratigraphic correlation between them. Among the many marine

192 bioclasts visible under the microscope, we distinguish gastropod and bivalve shells, fragments
193 of coral and calcareous algae, and foraminifera (with different levels of abrasion). The great
194 majority of the shells are broken. The lithogenic sand size fraction is composed of calcite and
195 quartz. For Womi as well as for Kouloune cores, Br and Cl have a negative loading on PCA 1
196 (53% of the variance), while Sr and Ca have a negative loading on PCA 3 (15%), together
197 with Zr (Fig. 7). Coarse-grained bioclastic layers and isolated shells are associated with peaks
198 of Sr (e.g., Sr/Ti ratio), and they are clearly distinct from the silty facies (Fig. 6).

199 The sequence is time-controlled by four ¹⁴C ages. Dark-brown silty sediment sampled using
200 the Edelman auger is dated to 3384-3321 cal. BCE at 200 cm depth on section WOM-01
201 (Table 1). The 1213-978 cal. BCE age obtained at 110 cm depth in the same section provides
202 an intermediate age that coincides stratigraphically with the base of WOM-02 (Fig. 6). In core
203 WOM-02, wood sampled from a bioclastic sand layer at 85 cm depth is dated 1218-983 cal.
204 BCE, and a charcoal found in light-brown silt 1451-1515 cal. CE at 61 cm depth.

205

206 3.4. Pouébo

207 The Pouébo site is a “*tanne*” (or “*tan*”). This Senegalese word (Wolof language) refers to a
208 specific type of tropical coastal marsh and is commonly used in the French literature (Lebigre,
209 2004). The *tanne* corresponds to a bare or sparsely vegetated salt marsh on the inner edge of a
210 mangrove, inundated during high water springs, cyclones, or tsunamis. During the dry season,
211 the *tanne* is covered by saline deposits formed by efflorescence. The Pouébo *tanne* is bounded
212 to the Southeast by the Pouébo River estuary, to the Northeast by a mangrove and a fringing
213 reef and to the West by the coastal plain. The 8 sampling sites are distributed along a WNW-
214 ESE transect at distances of 400-900 m from the estuary and 1300-1600 m from the reef (Fig.
215 8).

216 The sedimentary sequences show an accumulation of 60-100 cm of laminated sandy deposits
217 overlying yellowish bioclastic gravel with abundant coral fragments and marine shells (Fig.
218 9). Traces of bioturbation are rare, and the sequences are generally well preserved. Grain size
219 distributions range from fine to coarse sand. The dominant facies is a moderately- sorted fine
220 sand, but some layers or laminae are slightly coarser (medium sand) and less sorted. The
221 colour of the sediment is light-brown to grey in the upper part of the sequences and grey to
222 yellowish or reddish in the lower part. Quartz from the gneiss substratum (Maurizot et al.,

223 2020) is the dominant mineral. Marine bioclasts (fragments of coral and calcareous algae,
224 bivalve and gastropod shells, foraminifera, etc.) are present at all levels of the stratigraphy in
225 varying abundance (Fig. 10). Pumices are found at depths of 21 cm and 28 cm in core POU-
226 02 (Fig. 11) and at depths of 15 cm and 37 cm in core POU-08 (Fig. 12).

227 XRF core-scanner analysis was performed on two cores: POU-02 and POU-08. On both
228 cores, Sr and Ca contents are correlated, and they have a negative loading on PCA 1 (Fig. 7:
229 PCA 1 represents 39% of the total variance on POU-02 and 42% on POU-08). Br and Cl
230 again form another group with a positive loading on PCA 2, in association with Zr and Si for
231 POU-02. A third group is represented by Ti, Al, and K, sometimes with Fe (POU-02) or Si
232 (POU-08). On vertical variations in chemical composition, negative peaks of PCA 1 and
233 positive peaks of the Ca/Ti and Sr/Ti ratios correspond to layers or laminae enriched in marine
234 bioclasts, as seen, for instance, at 20-22 cm depth in core POU-02 (Fig. 11) and at 12-15 cm
235 depth in core POU-08 (Fig. 12).

236 The ^{14}C ages obtained both on sediment and wood are often incoherent (Table 1 and Fig. 9),
237 although there is little bioturbation and an overall continuity of the stratigraphy from one
238 sampling site to another. As an example, wood collected at a depth of 88 cm in core POU-08
239 is dated to 1669-1784 cal. CE, whereas another wood collected at 40 cm is dated 770-459 cal.
240 BCE. In core POU-02, sediment is dated to 120-251 cal. CE at 65 cm depth, 1130-1224 cal.
241 CE at 20 cm depth, and 409-541 cal. CE at 18 cm depth.

242

243 3.5. Ile-des-Pins

244 On the northern coast of the Ile-des-Pins (Fig. 13), we selected a site on the western side of
245 the Baie des Crabes, close to Gadji, 1150 m away from the crest of the fringing reef. The bay
246 is very shallow and almost transverse to the reef crest, thus reducing the energy of the waves
247 during cyclones. The sampling site is located 50 m inland. A 10 cm thick bioclastic sand layer
248 is buried at depths of 48-58 cm in the dark-brown silty soil. White bioclastic sand turning to
249 coral gravel at 90-100 cm depth prevented any deeper penetration of the auger. A ^{14}C age of
250 1720-1811 cal. CE is obtained on bulk sediment (soil) at a depth of 43 cm, just above the
251 bioclastic sand layer (Table 1).

252 On the southwestern coast at Baie des Rouleaux (Fig. 1), we surveyed a small marsh protected
253 by a 4 m high dune. The sequence shows no evidence of high-energy marine inundation. The

254 dark-brown mangrove peat (0-84 cm depth) overlies a paleodune. A ^{14}C age of 2059-1889 cal.
255 BCE is obtained at 90 cm depth on bulk sediment in a thin organic horizon of the sand dune
256 (Table 1).

257

258 3.6. Maré Island

259 On Maré Island, there are very few sites that meet our criteria. The island is mostly bounded
260 by rocky coasts, and thus, there is little accommodation space for preserving tsunami deposits
261 (i.e., no coastal marshes) between the coastline and the paleocliffs of Plio-Quaternary uplifted
262 coral reef terraces (Maurizot et al., 2020). We focused on the eastern coast of the island
263 around Patho (Fig. 14). Following the recommendations of locals, we selected sites that were
264 not inundated during cyclones. In Nawa Redje, 2.4 km south of Patho at an altitude of 8 m
265 a.s.l., there is a small cave, 4 m deep, carved in the reefal limestone. The bottom of the cave is
266 filled by a 1 m thick fan-type accumulation of coarse bioclastic sediment. The upper part of
267 the sequence (0-65 cm depth) consists of a massive poorly sorted gravel with abundant coral
268 fragments (up to 15 cm large), gastropod and bivalve shells (including oysters up to 12 cm
269 large). The deposit also includes rounded pumice lapilli, which are more abundant at depths of
270 15 and 50 cm. A charcoal found at a depth of 50 cm, similar to the lower pumices, is dated
271 1796-1896 cal. CE (Table 1). The lower part of the sequence (65-100 cm depth) is a
272 moderately sorted greyish coarse sand with some fragments of coral and shells (< 1 cm large).

273 North of Patho village, well-preserved sections of a paleodune system are exposed along the
274 walls of a sand quarry north of Patho, 80 m from the shoreline and 380 m from the fringing
275 reef (Fig. 14). The site faces a 320 m-long break in the reef crest. The top of the dune is here
276 at an altitude of 9 m a.s.l., which greatly reduces possible interference from cyclones. The
277 dominant facies of sand is coarse, and the great majority of the clasts are derived from marine
278 species (fragments of shells and coral). Light-brown pedogenetic horizons enriched in organic
279 matter emerge in the upper part of the sections (e.g., at 60 cm depth on sections PAT-04 and
280 PAT-05: Fig. 15). A thick dark-brown paleosol cuts the paleodune on section PAT-04. This soil
281 covers a period of four centuries, according to ^{14}C ages of 1128-1220 cal. CE at 110 cm depth
282 and 1497-1650 cal. CE at 75 cm depth on bulk sediment. A thin pedogenetic horizon at 55 cm
283 depth is dated 1621-1673 cal. CE on charcoal (Table 1).

284 The most striking feature of the paleodune is the abundance of pumices in the form of layers
285 and lines of rounded pumice lapilli. The geometry of these pumice deposits is complex:
286 discontinuous, undulating, horizontal or gently sloping seaward, one layer turning to several
287 layers with sand in between, etc. The lower contact with the sand is sharp and erosive,
288 whereas the upper contact is diffuse. We distinguish three main pumice units up to 30 cm
289 thick over a section of 2.8 m: pumice P1 at ~140-170 cm, pumice P2 at ~100-130 cm, and
290 pumice P3 at ~20-50 cm depth (Fig. 15). The P2 unit is clast-supported and exclusively
291 composed of rounded pumice clasts, whereas units P1 and P3 display both clast-supported
292 and matrix-supported facies. The matrix is made of a sand that is similar to the bioclastic sand
293 of the paleodune. Whole-rock major element chemistry of pumice clasts indicates a dacitic
294 composition (65-70 wt. % SiO₂ and 3.5-5.2 wt. % of alkali Na₂O+K₂O), but it is possible to
295 distinguish different types of pumice (Fig. 16). In the P1 unit, there is only one type of pumice
296 - greyish, crystal-poor, with evidence of magma mingling (enclaves of a darker lower-SiO₂
297 component). The P2 unit is a mixture of three types of pumice: (1) pumices similar to P1, (2)
298 pumices close to P1 in terms of texture and composition but with crystals of pyroxene, and (3)
299 rare white pumice with a higher SiO₂ content (69 wt. %). Different types of pumice are also
300 observed in unit P3, but the P1-type is ultradominant. There are two possible sources of
301 pumice in the region: the Tonga and the Vanuatu volcanic arcs. Both major and trace elements
302 suggest that the pumices deposited in the Patho paleodune come from the Tonga Islands (Fig.
303 16), which is consistent with the direction of the trade winds, and the westwards direction of
304 the South Equatorial oceanic current.

305 In the upper part of the PAT-05 section, pumice unit P3 is eroded by dark brown 30 cm thick
306 gravel. This detritic deposit has a sandy matrix (similar to the underlying bioclastic sand of
307 the dune) and includes both terrestrial (snails) and marine bioclasts (broken shells and corals)
308 and rounded pumice lapilli. The abundance of pumice decreases from the base to the top of
309 the gravel. A charcoal near the base of the gravel (25 cm depth) is dated 1728-1809 cal. CE
310 (Table 1).

311

312 3.7. Lifou Island

313 As for Maré Island, Lifou is not the ideal place for preserving tsunami deposits. We focused
314 our investigations on two sites: (1) Mu on the southeastern coast (Fig. 1), which was severely
315 impacted by the 1875 tsunami (Le Duff et al., 2016; Louat and Baldassari, 1989), and (2) Wé,

316 Châteaubriand Bay, on the eastern coast (Fig. 17). A survey around the two small crescent-
317 shaped bays of Mu was not very successful. At only one site located 105 m from the
318 shoreline, we found from top to base (1) a brown sandy soil up to 40 cm thick; (2) fine
319 bioclastic sand with pumice lapilli and coral cobbles relying on (3) the coral substratum at a
320 depth of 80 cm. The contact between the substratum and the sand is irregular. Although the
321 deposits could be likely related to the 1875 tsunami, we could not find suitable material for
322 ^{14}C dating.

323 Sampling sites at Wé are situated on the western shore of the Châteaubriand Bay on the lee
324 side of a vegetated dune (Fig. 17). The wide bay is oriented towards the open ocean. The most
325 complete sequence reported here (WE-03) is located 250 m from the shoreline at an altitude
326 of 2 m a.s.l., where the dune is 10 m high. The substratum (coral gravel) was reached at a
327 depth of 2.65 m (Fig. 18). The lower part of the sequence (120-265 cm depth) consists of
328 orange-to-yellowish clay silt without pumice or marine bioclasts. Between 120 and 90 cm
329 depth, the stratigraphy is unclear due to the presence of a water table in a layer of non
330 bioclastic. From 90 to 50 cm depth, dark-brown silt-clay sediment turns to light brown and
331 then to brown silt at 66 cm depth. An age of 52 cal. BCE – 79 cal. CE was obtained on silt
332 sediment at a depth of 66 cm (Table 1). Altered pumice lapilli are found at various depths, but
333 they never form well-defined layers or lenses. Between 50 and 40 cm, the sequence of silt is
334 discontinued by a yellowish bioclastic sand with abundant rounded pumice lapilli on top.
335 Pumice clasts are still frequent in the overlying brown silt-sand, sometimes associated with
336 marine shells (e.g., at 29-30 cm depth on Fig. 18). In the upper part of the sequence,
337 background silt-sand sediment is again interrupted by a lens-shaped layer of rounded pumice
338 lapilli with few marine shells (at 15-19 cm depth on Fig. 18). Dark brown silt sand is dated
339 781-882 cal. CE at 18 cm depth (Table 1).

340 XRF core-scanner analysis was realized on a U-channel between 19 and 72 cm depth on
341 section WE-03. The PCA biplot clearly separates the different sedimentary facies identified
342 along the core (Fig. 7). In terms of chemical composition, silt-clay and silt sediments of the
343 lower part of the sequence (50-72 cm depth) are different from the yellowish bioclastic sand
344 (45-50 cm depth) and the upper silt-sand sediment (19-40 cm depth). All chemical elements
345 have a positive loading on PCA 1 (50%), except Ca and Sr, which are both well correlated
346 with a strong positive loading on PCA 2 (18%). Vertical profiles of PCA dimensions and
347 elemental ratios (Fig. 18) show that the lower silts (50-72 cm depth) are characterized by
348 negative PCA 2, and low Ca/Ti. The bioclastic sand coincides with positive PCA 2,

349 decreasing PCA 1, and high Ca/Ti. Upper silt-sand sediment (0-45 cm depth) is characterized
350 by vertical variations in composition that are essentially due to the presence of pumice and
351 fine marine shells dispersed on the sediment.

352

353 **4. Discussion**

354

355 4.1. Types of tsunami deposits

356 The identification of paleotsunamis in the coastal sedimentary sequences of New Caledonia is
357 essentially based on the presence of sediments with a clear marine signature, which is linked
358 to the presence of marine bioclasts in most cases (gastropod and bivalve shells, coral
359 fragments, foraminifera, etc.). In addition to the textural data, the vertical variations in the
360 geochemical composition of the sediments make it possible to detect the main phases of
361 marine flooding. Ca and Sr are dominant elements in marine biogenic carbonates, shells and
362 coral skeletons, as well as fine carbonated mud from more protected area (deeper or internal
363 carbonated platform) (Chagué-Goff, 2010; Sabatier et al., 2010). Therefore, Ca and Sr are
364 often used as proxies for tsunami inundation due to their higher concentrations in marine
365 sediments (Chagué-Goff et al., 2011; Cuven et al., 2013; Font et al., 2013; Chagué-Goff et al.,
366 2017; Biguenet et al., 2020; Paris et al., 2021). The Ca/Ti ratio also represents a relevant
367 indicator of marine sediment input in a terrestrial environment, as Ti is mainly present in
368 continental clay and oxides (e.g., Moreira et al., 2017). In the sedimentary sequences studied
369 here, Ca and Sr are strongly correlated, and opposed to other more siliciclastic element such
370 as Si, Al, K, Fe, and Ti on at least one of the PCA dimensions (Fig. 7).

371 All sites investigated are located beyond the limit of debris (drift wood, corals, pumice, algae,
372 plastic, etc.) that were left by recent large cyclones such as Cook (April 2017), Donna (May
373 2017) and Oma (February 2019), as inferred from our field observations and eyewitness
374 accounts. Although our study sites are located outside the morpho-sedimentary impact zone of
375 the cyclones, we cannot definitely rule out a cyclone origin for some of the marine event
376 identified (e.g. crypto-deposits with a marine chemical signature at Bwa).

377 While the different tsunami deposits identified here have similarities in terms of geochemical
378 signature, their textural characteristics differ from one site to another. Three types of deposits
379 can be distinguished:

- 380 1. Bioclastic sands that are coarser grained and less sorted than the background
381 sediments, as observed at Womi (Fig. 6, Grande Terre), Baie des Crabes (Fig. 13, Ile-
382 des-Pins), Nawa Redje and Patho (Fig. 15, Maré Island), and Wé (Fig. 18, Lifou
383 Island). This is the most common type of tsunami deposit, as described in many other
384 places (Engel et al., 2020). The preservation of a distinct tsunami sand unit in the
385 stratigraphy requires (a) a source of sand in the nearshore and/or the inner shelf, (b) a
386 tsunami wave high enough to overtop the coastal ridge, and (c) a space to
387 accommodate the sand brought by the tsunami (e.g., a marsh). As an example, these
388 conditions are particularly well met in Womi (Fig. 5) and Wé (Fig. 17).
- 389 2. Crypto-deposits or horizons that (a) are slightly coarser grained than the clayey
390 sediment of the marsh and (b) have a marine signature (peaks of Sr/Ti and Ca/Ti due
391 to the presence of fine marine bioclasts) but (c) do not form a distinct layer.
392 Interestingly, these deposits are often associated with rounded pumice lapilli, as seen
393 in Bwa (Fig. 3) or Koulnoue (Fig. 5). Tsunami crypto-deposits reflect either a lack of
394 preservation of a distinct (macroscopic) tsunami deposit in the stratigraphy (e.g., due
395 to bioturbation in a mangrove), a lack of accommodation space, or the tsunami's
396 inability to transport a volume of sediment large enough to form a significant deposit
397 (e.g., in the distal zone of the marsh, close to the limit of inundation).
- 398 3. Accumulations of rounded pumice lapilli in the form of layers, lines or lenses, whose
399 characteristics differ completely from a juvenile pumice fallout deposit, both at the
400 deposit scale (erosive contact with basal sediment, cross-laminae, bioclastic sand
401 matrix) and clast scale (high degree of rounding of the pumice clasts, different types of
402 pumice are sometimes mixed in a single unit, indicating reworking from one unit to
403 another). Based on their chemical composition, all pumices found in the coastal
404 sedimentary sequence of New Caledonia have a Tonga origin (Fig. 16). These pumices
405 drifted to the coasts of New Caledonia in the form of rafts, as observed following
406 submarine eruptions in Tonga in 2002 (Bryan et al., 2004), 2006 (Bryan et al., 2012),
407 and 2019 (Jutzeler et al., 2020). The pumices were then stranded on the beach at high
408 tide, reworked back to the ocean, and occasionally pushed further inland by storm
409 surges and tsunamis. These successive transports very quickly give a rounded shape to

410 the pumice clasts. Rounded pumice lapilli were found to be associated with tsunami
411 deposits of the 1883 eruption of Krakatau in the Sunda Strait, Indonesia (Carey et al.,
412 2001; Paris et al., 2014), and the 1650 eruption of Kolumbo in the Aegean Sea, Greece
413 (Ulvrova et al., 2016).

414

415 4.2. Paleotsunami inundation distance and runup

416 The spatial distribution of the paleotsunami deposits gives a minimum estimate of the
417 inundation limit and runup height, assuming a coastal morphology close to the present-day
418 one. In a low topography context such as in the Koulnoue mangrove or Pouébo tanne, the
419 tsunami reached a minimum distance of 700 and 1600 m inland, respectively. Unfortunately,
420 there is no information available on the inundation extent at these two sites during historical
421 and recent tsunamis (e.g., 1875, Mw ~8 earthquake in southern Vanuatu; 2007, Solomon
422 Islands Mw 8.0 earthquake).

423 However, at other sites, it is clear that the runup heights inferred from the altitude of the
424 tsunami deposits (which correspond to minimum values) are often higher than the runups
425 mentioned in the tsunami catalogue, which do not exceed 3.5 m a.s.l. (Sahal et al., 2010). This
426 catalogue covers the last 150 years, with the oldest event being the 1875 tsunami. At Womi
427 (Fig. 4, northeastern coast of Grande Terre), a tsunami must first propagate through 300 m of
428 dune ridge culminating at an altitude of 2.5 m a.s.l., and then 300 m of mangrove before
429 reaching our sampling site at a distance of 600 m from the shoreline. On the eastern coast of
430 Maré Island, traces of tsunamis were found in a paleodune and a karstic cave at present-day
431 altitudes of 5-7 m a.s.l. (Figs. 14 and 15). At Wé in the Châteaubriand Bay (Fig. 17, eastern
432 coast of Lifou Island), two tsunamis were able to overtop a 10 m-high dune before
433 accumulating sand and pumice deposits on its lee side (Fig. 18). Numerical simulations of
434 tsunamis suggest that a $M_w \geq 8.3$ earthquake in southern Vanuatu is required to generate
435 runups >10 m in the Châteaubriand Bay (Ioualalen et al., 2017). The tsunami generated by the
436 March 28, 1875 earthquake in southern Vanuatu severely damaged the southeastern coasts of
437 Lifou, particularly at Mu and Thoth where the damages and loss of lives were the highest. At
438 Thoth the waves penetrated up to 400 m inland and nothing remained of the village (Rousset,
439 1975 in (Le Duff et al., 2016)). At Mu, a minimum runup of 3.6 m was reported (Ella, 1890 in
440 (Louat and Baldassari, 1989)). Important waves were also noticed northward along the eastern
441 coast at Luengoni and We in the Chateaubriand Bay. However, its impact there is not well

442 documented. At We, the sea flooded the catholic church, which is located approximately 50 m
443 away from the coastline at high tide (Gaide, 1875 in (Louat and Baldassari, 1989)).

444 These estimates of paleotsunami runup and inundation distance are given for information
445 only, because the coastal landscapes may have undergone significant transformations (e.g.,
446 dune erosion, coastal retreat, mangrove degradation) due to climatic, tectonic, and
447 anthropogenic factors.

448

449 4.3. A chronology of paleotsunamis in New Caledonia

450 Figure 19 provides a chronology of paleotsunamis identified in the coastal stratigraphy of
451 Grande Terre and the Loyalty Islands based on the probability distribution of ^{14}C ages. Ages
452 obtained on samples taken from tsunami deposits were considered possibly reworked and thus
453 represent minimum ages. As an example, the inconsistency of the ages obtained at Pouébo
454 suggests that some samples were reworked and transported, in a context where the
455 stratigraphy nevertheless seems well preserved (Fig. 9). Little is known about the sedimentary
456 dynamics of tannes, but it seems that high-energy events represent the major input of
457 sediments, which complicates the construction of reliable chronological models.

458 The Womi sequence is the most complete one, with five distinct tsunami deposits during the
459 last two millennia (Fig. 6: bioclastic sand-silt units): a thick tsunami deposit dated ~1000
460 BCE, three other tsunamis between ~1000 BCE and ~1500 CE (the last one being very close
461 to 1450 CE), and a post-1500 CE tsunami deposit that is not precisely dated. The two tsunami
462 deposits identified at Wé on Lifou Island (Fig. 18), both correspond to major events since the
463 waves were potentially able to overtop a 10-m high dune (assuming that the dune was as high
464 as the present-day one). An archeological site at 4 m below the top of the dune was dated 260-
465 425 cal. CE (Sand, 1998), which confirms that the dune existed long before. The age of the
466 oldest tsunami deposit in Wé is poorly constrained (between ~100 CE and ~800 CE), whereas
467 the youngest deposit occurred at ~900 CE. Thus, there were at least two major tsunamis
468 impacting both Grande Terre (Womi) and the Loyalty Islands (Wé) during the first
469 millennium CE (Fig. 19). Our knowledge of the paleotsunami chronology in the Melanesian
470 region is very sporadic (Goff et al., 2022, 2011a, 2011b), and it is therefore difficult to
471 correlate the paleotsunami deposits described here with other events on other islands.

472 The 15th century tsunami described in Tonga (Lavigne et al., 2021), Wallis and Futuna (Goff
473 et al., 2011b), and many other places in Melanesia and Polynesia (Goff et al., 2022) is
474 possibly recorded at two sites, on the northeastern coast of Grande Terre (Womi), and on the
475 eastern coast of Maré Island (Patho). However, the origin of this 15th century regional
476 tsunami is still debated (Goff et al., 2022; Lavigne et al., 2021).

477 Another major tsunami possibly occurred at approximately 1700 CE, as recorded at Maré
478 (Patho), Grande Terre (Koulnoue, Womi, and possibly Bwa), and Ile-des-Pins (Baie des
479 Crabes). The oldest earthquake in the Vanuatu - New Caledonia catalogue dates back to 1729
480 \pm 3 CE (Louat and Baldassari, 1989). In fact, this event was inferred from ²³⁰Th ages on
481 emerged coral heads that died during coseismic uplift in Malekula, Vanuatu (Edwards et al.,
482 1988). It is a good candidate for the ~1700 CE tsunami identified here in New Caledonia.

483 Then, a tsunami in the second half of the XIXth century, which most likely corresponds to the
484 March 1875 tsunami, is recorded both in the Loyalty Islands (Maré) and in Grande Terre
485 (Pouébo, and perhaps Womi and Koulnoue). A more recent event also appears in the Pouébo
486 sequences (Fig. 9), but it is not temporally constrained because ¹⁴C ages obtained for this site
487 are not reliable for the reasons mentioned above.

488

489 **5. Conclusion**

490 In this study, we document for the first time tsunami deposits on the coasts of New Caledonia
491 (Grande Terre and the Loyalty Islands). Three types of deposits are distinguished: (1)
492 Bioclastic sand layers, similar to many other tsunami deposits described elsewhere in the
493 world; (2) Crypto-deposits that do not form a distinct layer, but have a marine chemical
494 signature; (3) Accumulations of rounded pumice lapilli originating from the Tonga Islands,
495 transported as pumice rafts and later deposited inland by tsunamis. The spatial distribution of
496 the tsunami deposits suggests tsunami runups exceeding 2.5 m a.s.l. on the northeastern coast
497 of Grande Terre (Womi), 6 m on the eastern coast of Maré Island (Patho), and 10 m on the
498 northeastern coast of Lifou Island (Wé). At least five tsunamis occurred during the last two
499 millennia, plus an additional event (recorded only at one site in Grande Terre) during the first
500 millennium BCE. We thus extend the tsunami catalogue of New Caledonia, which was
501 previously limited to the last 150 years since European colonization (Sahal et al., 2010). This
502 work also lays the foundations for an integrated approach to the tsunami hazard in New

503 Caledonia by allowing a comparison of the different timescales of tsunami records. At the
504 timescale of the sedimentary archives, we found at least five tsunamis since ~100 CE, which
505 means an annual probability of ~0.0025. This gives an idea of the recurrence of major
506 tsunamis such as the 28 March 1875 tsunami, which makes the link between the geological
507 timescale, i.e. tsunamis that left a deposit in the sedimentary record, and the historical
508 timescale, i.e. tsunamis observed and/or recorded on instruments (Roger et al., 2019).

509

510 **Acknowledgements**

511 This work was funded by the CNRS-IRD AAP 2019 “Risques” of the MITI (Mission pour les
512 Initiatives Transverses et Interdisciplinaire) and by the Government of New Caledonia under
513 the TSUCAL project. ¹⁴C analyses were acquired thanks to the CNRS-INSU ARTEMIS
514 (Accelerator for Research in Earth Sciences, Environment, Museology) National Radiocarbon
515 AMS Measurement Program at LMC14 (Laboratoire de Mesure ¹⁴C, CEA, Saclay). The
516 authors acknowledge the support of Mhammed Benbakkar and Krzysztof Suchorski (major
517 and trace element chemistry, LMV), Martine Trautman (grain size analysis, LAS-EOST), and
518 Anne-Lise Develle (XRF core scanner, EDYTEM), as well as all people who have given us
519 their support during the field campaign, namely: Jean-Maurice, Joseph et Daniel Bwarat
520 (Hienghène), Mr Bouillot, Ms Hiamparemane (Hienghène city hall), Benoît Maudet (Head of
521 the Gendarmerie at Hienghène), Father Vincent (Hienghène Church), Fideli and his brother
522 Maurice (Ile-des-Pins), Gérard Bashman (Technical Director at the Hôtel Méridien, Ile-des-
523 Pins), Gaby (Baie des Rouleaux, Ile-des-Pins), Raynald Washetine and Wademio Richard
524 (Patho, Maré Island), Boula Eanes (Chief of the Mu tribe, Lifou Island), and Lola
525 Whashetyne (Wabao, Maré Island). The Nataiwatch Guesthouse (Ile-des-Pins) and the
526 Koulnoue Village Resort (Grande Terre) provided respectively free and reduced-cost
527 accommodation to support this work. The authors also thank Pedro Costa and an anonymous
528 colleague for reviewing this work. This is ClerVolc contribution n° 611.

529

530 **Data availability**

531 All data generated or analysed during this study are included in this article (and its
532 supplementary information files) or available from the corresponding author on reasonable
533 request.

535 **References**

- 536 Bevis, M., Taylor, F.W., Schutz, B.E., Recy, J., Isacks, B.L., Helu, S., Singh, R., Kendrick, E.,
 537 Stowell, J., Taylor, B., Calmantli, S., 1995. Geodetic observations of very rapid
 538 convergence and back-arc extension at the Tonga arc. *Nature* 374, 249–251.
 539 <https://doi.org/10.1038/374249a0>
- 540 Biguenet, M., Sabatier, P., Chaumillon, E., Chagué, C., Arnaud, F., Jorissen, F., Coulombier,
 541 T., Geba, E., Cordrie, L., Vacher, P., Develle, A.L., Chalmin, E., Soufi, F., Feuillet, N.,
 542 2020. A 1600 year-long sedimentary record of tsunamis and hurricanes in the Lesser
 543 Antilles (Scrub Island, Anguilla). *Sedimentary Geology* 105806.
 544 <https://doi.org/10.1016/j.sedgeo.2020.105806>
- 545 Bruyère, O., Soulard, B., Lemonnier, H., Laugier, T., Hubert, M., Petton, S., Desclaux, T., Van
 546 Wynsberge, S., Le Tesson, E., Lefèvre, J., Dumas, F., Kayara, J.-F., Bourassin, E.,
 547 Lalau, N., Antypas, F., Le Gendre, R., 2022. Hydrodynamic and hydrological
 548 processes within a variety of coral reef lagoons: field observations during six cyclonic
 549 seasons in New Caledonia. *Earth Syst. Sci. Data* 14, 5439–5462.
 550 <https://doi.org/10.5194/essd-14-5439-2022>
- 551 Bryan, S.E., Cook, A., Evans, J.P., Colls, P.W., Wells, M.G., Lawrence, M.G., Jell, J.S., Greig,
 552 A., Leslie, R., 2004. Pumice rafting and faunal dispersion during 2001–2002 in the
 553 Southwest Pacific: record of a dacitic submarine explosive eruption from Tonga. *Earth
 554 and Planetary Science Letters* 227, 135–154.
 555 <https://doi.org/10.1016/j.epsl.2004.08.009>
- 556 Bryan, S.E., Cook, A.G., Evans, J.P., Hebden, K., Hurrey, L., Colls, P., Jell, J.S., Weatherley,
 557 D., Firm, J., 2012. Rapid, Long-Distance Dispersal by Pumice Rafting. *PLoS ONE* 7,
 558 e40583. <https://doi.org/10.1371/journal.pone.0040583>
- 559 Calmant, S., Pelletier, B., Lebellegard, P., Bevis, M., Taylor, F.W., Phillips, D.A., 2003. New
 560 insights on the tectonics along the New Hebrides subduction zone based on GPS
 561 results: GPS-CONSTRAINED TECTONICS OF THE NEW HEBRIDES ARC. *J.
 562 Geophys. Res.* 108. <https://doi.org/10.1029/2001JB000644>
- 563 Carey, S., Morelli, D., Sigurdsson, H., Bronto, S., 2001. Tsunami deposits from major
 564 explosive eruptions: An example from the 1883 eruption of Krakatau 5.
- 565 Chagué-Goff, C., 2010. Chemical signatures of palaeotsunamis: A forgotten proxy? *Marine
 566 Geology* 271, 67–71. <https://doi.org/10.1016/j.margeo.2010.01.010>
- 567 Chagué-Goff, C., Schneider, J.-L., Goff, J.R., Dominey-Howes, D., Strotz, L., 2011.
 568 Expanding the proxy toolkit to help identify past events — Lessons from the 2004
 569 Indian Ocean Tsunami and the 2009 South Pacific Tsunami. *Earth-Science Reviews*
 570 107, 107–122.
- 571 Chagué-Goff, C., Szczuciński, W., Shinozaki, T., 2017. Applications of geochemistry in
 572 tsunami research: A review. *Earth-Science Reviews* 165, 203–244.
 573 <https://doi.org/10.1016/j.earscirev.2016.12.003>
- 574 Cuven, S., Paris, R., Falvard, S., Miot-Noirault, E., Benbakkar, M., Schneider, J.-L., Billy, I.,
 575 2013. High-resolution analysis of a tsunami deposit: Case-study from the 1755 Lisbon
 576 tsunami in southwestern Spain. *Marine Geology* 337, 98–111.
 577 <https://doi.org/10.1016/j.margeo.2013.02.002>
- 578 Dumoulin, J.-P., Comby-Zerbino, C., Delqué-Količ, E., Moreau, C., Caffy, I., Hain, S.,
 579 Perron, M., Thellier, B., Setti, V., Berthier, B., al, et, 2017. Status Report on Sample
 580 Preparation Protocols Developed at the LMC14 Laboratory, Saclay, France: From

581 Sample Collection to 14C AMS Measurement. *Radiocarbon* 59, 713–726.
582 <https://doi.org/10.1017/RDC.2016.116>

583 Edwards, R.L., Taylor, F.W., Wasserburg, G.J., 1988. Dating earthquakes with high-precision
584 thorium-230 ages of very young corals. *Earth and Planetary Science Letters* 90, 371–
585 381. [https://doi.org/10.1016/0012-821X\(88\)90136-7](https://doi.org/10.1016/0012-821X(88)90136-7)

586 Engel, M., Pilarczyk, J., May, S.M., Brill, D., Garrett, E., 2020. Geological records of
587 tsunamis and other extreme waves.

588 Font, E., Veiga-Pires, C., Pozo, M., Nave, S., Costas, S., Ruiz Muñoz, F., Abad, M., Simões,
589 N., Duarte, S., Rodríguez-Vidal, J., 2013. Benchmarks and sediment source(s) of the
590 1755 Lisbon tsunami deposit at Boca do Rio Estuary. *Marine Geology* 343, 1–14.
591 <https://doi.org/10.1016/j.margeo.2013.06.008>

592 Goff, J., Borrero, J., Easton, G., 2022. In search of Holocene trans-Pacific palaeotsunamis.
593 *Earth-Science Reviews* 233, 104194. <https://doi.org/10.1016/j.earscirev.2022.104194>

594 Goff, J., Chagué-Goff, C., Dominey-Howes, D., McAdoo, B., Cronin, S., Bonté-Grapetin, M.,
595 Nichol, S., Horrocks, M., Cisternas, M., Lamarche, G., Pelletier, B., Jaffe, B., Dudley,
596 W., 2011a. Palaeotsunamis in the Pacific Islands. *Earth-Science Reviews* 107, 141–
597 146. <https://doi.org/10.1016/j.earscirev.2010.10.005>

598 Goff, J., Lamarche, G., Pelletier, B., Chagué-Goff, C., Strotz, L., 2011b. Predecessors to the
599 2009 South Pacific tsunami in the Wallis and Futuna archipelago. *Earth-Science*
600 *Reviews* 107, 91–106. <https://doi.org/10.1016/j.earscirev.2010.11.003>

601 Gusman, A.R., Roger, J., Power, W., Fry, B., Kaneko, Y., 2022. The 2021 Loyalty Islands
602 Earthquake (Mw 7.7): Tsunami Waveform Inversion and Implications for Tsunami
603 Forecasting for New Zealand. *Earth and Space Science* 9.
604 <https://doi.org/10.1029/2022EA002346>

605 Hammer, O., Harper, D.A.T., Ryan, P.D., 2001. PAST: Paleontological Statistics Software
606 Package for Education and Data Analysis 9.

607 Heaton, T.J., Köhler, P., Butzin, M., Bard, E., Reimer, R.W., Austin, W.E.N., Bronk Ramsey,
608 C., Grootes, P.M., Hughen, K.A., Kromer, B., Reimer, P.J., Adkins, J., Burke, A.,
609 Cook, M.S., Olsen, J., Skinner, L.C., 2020. Marine20—The Marine Radiocarbon Age
610 Calibration Curve (0–55,000 cal BP). *Radiocarbon* 62, 779–820.
611 <https://doi.org/10.1017/RDC.2020.68>

612 Hogg, A.G., Heaton, T.J., Hua, Q., Palmer, J.G., Turney, C.S., Southon, J., Bayliss, A.,
613 Blackwell, P.G., Boswijk, G., Bronk Ramsey, C., Pearson, C., Petchey, F., Reimer, P.,
614 Reimer, R., Wacker, L., 2020. SHCal20 Southern Hemisphere Calibration, 0–55,000
615 Years cal BP. *Radiocarbon* 62, 759–778. <https://doi.org/10.1017/RDC.2020.59>

616 Ioualalen, M., Pelletier, B., Solis Gordillo, G., 2017. Investigating the March 28th 1875 and
617 the September 20th 1920 earthquakes/tsunamis of the Southern Vanuatu arc, offshore
618 Loyalty Islands, New Caledonia. *Tectonophysics* 709, 20–38.
619 <https://doi.org/10.1016/j.tecto.2017.05.006>

620 Jutzeler, M., Marsh, R., Sebille, E., Mittal, T., Carey, R.J., Fauria, K.E., Manga, M., McPhie,
621 J., 2020. Ongoing Dispersal of the 7 August 2019 Pumice Raft From the Tonga Arc in
622 the Southwestern Pacific Ocean. *Geophys. Res. Lett.* 47.
623 <https://doi.org/10.1029/2019GL086768>

624 Lavigne, F., Morin, J., Wassmer, P., Weller, O., Kula, T., Maea, A.V., Kelfoun, K., Mokadem,
625 F., Paris, R., Malawani, M.N., Faral, A., Benbakkar, M., Saulnier-Copard, S., Vidal,
626 C.M., Tu’I’afitu, T., Kitekei’aho, F., Trautmann, M., Gomez, C., 2021. Bridging
627 Legends and Science: Field Evidence of a Large Tsunami that Affected the Kingdom
628 of Tonga in the 15th Century. *Front. Earth Sci.* 9, 748755.
629 <https://doi.org/10.3389/feart.2021.748755>

- 630 Le Duff, M., Dumas, P., Sabinot, C., Allenbach, M., 2016. Le risque tsunami en Nouvelle-
631 Calédonie : Évolutions des facteurs de vulnérabilités et de résiliences à Lifou en
632 territoire coutumier kanak. *vertigo*. <https://doi.org/10.4000/vertigo.17951>
- 633 Lebigre, J.M., 2004. Les marais à mangrove de Nouvelle-Calédonie, un exemple de milieu
634 « naturel » lagunaire, Nouméa, Centre de Documentation Pédagogique, Scérén.
- 635 Louat, R., Baldassari, C., 1989. Chronologie des séismes et des tsunamis ressentis dans la
636 région Vanuatu Nouvelle Calédonie (1729-1989) = Chronology of felt earthquakes and
637 tsunamis in the region Vanuatu New Caledonia (1729-1989) 49.
- 638 Maurizot, P., Robineau, B., Vendé-Leclerc, M., Cluzel, D., 2020. Chapter 1 Introduction to
639 New Caledonia: geology, geodynamic evolution and mineral resources. *Memoirs* 51,
640 1–12. <https://doi.org/10.1144/M51-2019-33>
- 641 Moreau, C., Caffy, I., Comby, C., Delqué-Količ, E., Dumoulin, J.-P., Hain, S., Quiles, A.,
642 Setti, V., Souprayen, C., Thellier, B., al, et, 2013. Research and Development of the
643 Artemis 14C AMS Facility: Status Report. *Radiocarbon* 55, 331–337.
644 <https://doi.org/10.1017/S0033822200057441>
- 645 Pararas-Carayannis, G., 2007. Natural Disasters in Oceania, in: Lidstone, J., Dechano, L.M.,
646 Stoltman, J.P. (Eds.), *International Perspectives on Natural Disasters: Occurrence,*
647 *Mitigation, and Consequences*. Springer Netherlands, Dordrecht, pp. 193–210.
648 https://doi.org/10.1007/978-1-4020-2851-9_10
- 649 Paris, R., Sabatier, P., Biguenet, M., Bougouin, A., André, G., Roger, J., 2021. A tsunami
650 deposit at Anse Meunier, Martinique Island: Evidence of the 1755 CE Lisbon tsunami
651 and implication for hazard assessment. *Marine Geology* 439, 106561.
652 <https://doi.org/10.1016/j.margeo.2021.106561>
- 653 Paris, R., Wassmer, P., Lavigne, F., Belousov, A., Belousova, M., Iskandarsyah, Y.,
654 Benbakkar, M., Ontowirjo, B., Mazzoni, N., 2014. Coupling eruption and tsunami
655 records: the Krakatau 1883 case study, Indonesia. *Bull Volcanol* 76, 814.
656 <https://doi.org/10.1007/s00445-014-0814-x>
- 657 Pelletier, B., Calmant, S., Pillet, R., 1998. Current tectonics of the Tonga–New Hebrides
658 region. *Earth and Planetary Science Letters* 164, 263–276.
659 [https://doi.org/10.1016/S0012-821X\(98\)00212-X](https://doi.org/10.1016/S0012-821X(98)00212-X)
- 660 Richter, T.O., Gaast, S.V.D., Koster, B., Vaars, A., Gieles, R., Stigter, H.C. de, Haas, H. de,
661 Weering, T.C.E. van, 2006. The Avaatech XRF Core Scanner: technical description
662 and applications to NE Atlantic sediments, in: Rothwell, R.G. (Ed.), *New Techniques*
663 *in Sediment Core Analysis*. Geological Society of London, p. 0.
664 <https://doi.org/10.1144/GSL.SP.2006.267.01.03>
- 665 Roger, J., 2020. Données bathymétriques et topographiques de Nouvelle- Calédonie :
666 Réalisation d'un MNT terre- mer pour l'étude de l'aléa tsunami. TSUCAL project,
667 IRD - New Caledonia Government. 42 pp.
- 668 Roger, J., Pelletier, B., Aucan, J., 2019. Update of the tsunami catalogue of New Caledonia
669 using a decision table based on seismic data and marigraphic records. *Nat. Hazards*
670 *Earth Syst. Sci.* 19, 1471–1483. <https://doi.org/10.5194/nhess-19-1471-2019>
- 671 Roger, J., Pelletier, B., Duphil, M., Lefèvre, J., Aucan, J., Lebellegard, P., Thomas, B.,
672 Bachelier, C., Varillon, D., 2021. The Mw 7.5 Tadine (Maré, Loyalty Islands)
673 earthquake and related tsunami of 5 December 2018: seismotectonic context and
674 numerical modeling. *Nat. Hazards Earth Syst. Sci.* 21, 3489–3508.
675 <https://doi.org/10.5194/nhess-21-3489-2021>
- 676 Roger, J., Pelletier, B., Gusman, A., Power, W., Wang, X., Burbidge, D., Duphil, M., 2023.
677 Potential tsunami hazard of the southern Vanuatu subduction zone: tectonics, case
678 study of the Matthew Island tsunami of 10 February 2021 and implication in regional

679 hazard assessment. *Nat. Hazards Earth Syst. Sci.* 23, 393–414.
680 <https://doi.org/10.5194/nhess-23-393-2023>
681 Sabatier, P., Dezileau, L., Briquieu, L., Colin, C., Siani, G., 2010. Clay minerals and
682 geochemistry record from northwest Mediterranean coastal lagoon sequence:
683 Implications for paleostorm reconstruction. *Sedimentary Geology* 228, 205–217.
684 <https://doi.org/10.1016/j.sedgeo.2010.04.012>
685 Sahal, A., Pelletier, B., Chatelier, J., Lavigne, F., Schindel , F., 2010. A catalog of tsunamis in
686 New Caledonia from 28 March 1875 to 30 September 2009. *Comptes Rendus*
687 *Geoscience* 342, 434–447. <https://doi.org/10.1016/j.crte.2010.01.013>
688 Sand, C., 1998. Recent Archaeological Research in the Loyalty Islands of New Caledonia 37,
689 194–223.
690 Ulvrova, M., Paris, R., Nomikou, P., Kelfoun, K., Leibrandt, S., Tappin, D.R., McCoy, F.W.,
691 2016. Source of the tsunami generated by the 1650 AD eruption of Kolumbo
692 submarine volcano (Aegean Sea, Greece). *Journal of Volcanology and Geothermal*
693 *Research* 321, 125–139. <https://doi.org/10.1016/j.jvolgeores.2016.04.034>
694 Weltje, G.J., Tjallingii, R., 2008. Calibration of XRF core scanners for quantitative
695 geochemical logging of sediment cores: Theory and application. *Earth and Planetary*
696 *Science Letters* 274, 423–438. <https://doi.org/10.1016/j.epsl.2008.07.054>
697

698

699 **Figure captions**

700

701 Fig. 1 – A: Location map of New Caledonia and the Loyalty Islands. Squares indicate
702 historical earthquakes in the Solomon, Vanuatu, Fiji, and Tonga Islands. Epicenters of
703 historical earthquakes that generated tsunamis in New Caledonia, including far-field sources
704 such as Japan, Kamchatka, and Chile, are indicated in yellow. B: Location map of the
705 sampling sites in New Caledonia and the Loyalty Islands. Background relief image from
706 (Roger, 2020).

707

708 Fig. 2 – Location map of a sampling site in Bwa, Lebris Bay, northeastern coast of Grande-
709 Terre. Background satellite image provided by Google Earth.

710

711 Fig. 3 – ^{14}C ages and XRF core-scanner data of core BWA-01 (PCA 1, PCA 2 and PCA 3
712 correspond to the first three dimensions of the Principal Component Analysis). See Table 1 for
713 details on ^{14}C ages.

714

715 Fig. 4 - Location map of sampling sites in Koulnoue and Womi, northeastern coast of Grande-
716 Terre. White dotted lines indicate limestone hills with steep slopes. Background satellite
717 image provided by Google Earth.

718

719 Fig. 5 - Sedimentary log, ^{14}C ages, and XRF core scanner data of core KOUL-03. See PCA
720 biplots in Fig. 8.

721

722 Fig. 6 - Sedimentary log, grain size distribution of selected bulk samples, ^{14}C ages, and XRF
723 core-scanner data of core WOM-02 and sedimentary log of core WOM-01.

724

725 Fig. 7 – PCA biplots of XRF core-scanner data. Sedimentary facies are defined from the top
726 to the base of the cores. The circle indicates the 95% confidence interval.

727

728

729 Fig. 8 - Location map of sampling sites in Pouébo, northeastern coast of Grande-Terre.
730 Background satellite image provided by Google Earth.

731

732 Fig. 9 – Sedimentary logs, stratigraphic correlations, and ^{14}C ages of sampling sites in
733 Pouébo.

734

735 Fig. 10 – Microscope views of different marine bioclasts found in core POU-08, including
736 coral fragments, bivalve and gastropod shells, foraminifera (both fresh or abraded), and
737 rounded pumice clasts.

738

739 Fig. 11 - Sedimentary log and XRF core scanner data of core POU-02.

740

741 Fig. 12 - Sedimentary log and XRF core scanner data of core POU-08.

742

743 Fig. 13 – Location map, sedimentary log and ^{14}C age of a trench section at Baie des Crabes,
744 northern coast of Ile-des-Pins. Background satellite image provided by Google Earth.

745

746 Fig. 14 – Location maps of sampling sites in Patho, eastern coast of Maré Island (Loyalty
747 Islands), and photograph of the PAT-05 trench section showing units of rounded pumice lapilli
748 in a paleodune. Background satellite image provided by Google Earth.

749

750 Fig. 15 – Sedimentary logs and ^{14}C ages of the Patho sampling sites.

751

752 Fig. 16 – Major and trace element compositions of pumice sampled in Patho (site PA-03)
753 compared to other samples of volcanic rocks from Tonga and Vanuatu (GEOROC database).

754

755 Fig. 17 – Location map of sampling sites in Wé, Châteaubriand Bay, Lifou Island (Loyalty
756 Islands). Background satellite image provided by Google Earth.

757

758 Fig. 18 – Sedimentary log, ^{14}C ages and XRF core scanner data of core WE-03.

759

760 Fig. 19 – Probability distribution of ^{14}C ages calibrated using the SHCal2020 calibration
761 curve (Hogg et al., 2020) for wood and bulk sediment samples and the Marine20 calibration
762 curve (Heaton et al., 2020) for marine shell samples. Blue areas correspond to tsunami
763 deposits as recorded in the stratigraphic record relative to the ^{14}C ages. Question marks
764 indicate tsunami events that are not well-constrained by the ages.

765

766

767

768

769 Suppl. 1 – Vertical profiles of Ca, Ti, and Sr determined using an XRF core scanner, with
770 energies of 10 kV for Ca and Ti, and 30 kV for Sr. Data mean-centred and normalized to the
771 standard deviation.

772