



HAL
open science

**A new Mesozoic record of the pseudoscorpion family
Garypinidae from Upper Cretaceous (Santonian)
Ajkaite amber, Ajka area, Hungary**

János Novák, Mark S Harvey, Márton Szabó, Jörg U Hammel, Danilo Harms,
Ulrich Kotthoff, Christoph Hörweg, Manuel Brazidec, Attila Ósi

► **To cite this version:**

János Novák, Mark S Harvey, Márton Szabó, Jörg U Hammel, Danilo Harms, et al.. A new Mesozoic record of the pseudoscorpion family Garypinidae from Upper Cretaceous (Santonian) Ajkaite amber, Ajka area, Hungary. *Cretaceous Research*, 2024, 153, pp.105709. 10.1016/j.cretres.2023.105709 . insu-04205956

HAL Id: insu-04205956

<https://insu.hal.science/insu-04205956>

Submitted on 13 Sep 2023


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proof

A new Mesozoic record of the pseudoscorpion family Garypinidae from Upper Cretaceous (Santonian) Ajkaite amber, Ajka area, Hungary

János Novák, Mark S. Harvey, Márton Szabó, Jörg U. Hammel, Danilo Harms, Ulrich Kotthoff, Christoph Hörweg, Manuel Brazidec, Attila Ósi



PII: S0195-6671(23)00237-9

DOI: <https://doi.org/10.1016/j.cretres.2023.105709>

Reference: YCRES 105709

To appear in: *Cretaceous Research*

Received Date: 11 June 2023

Revised Date: 7 September 2023

Accepted Date: 8 September 2023

Please cite this article as: Novák, J., Harvey, M.S., Szabó, M., Hammel, J.U., Harms, D., Kotthoff, U., Hörweg, C., Brazidec, M., Ósi, A., A new Mesozoic record of the pseudoscorpion family Garypinidae from Upper Cretaceous (Santonian) Ajkaite amber, Ajka area, Hungary, *Cretaceous Research*, <https://doi.org/10.1016/j.cretres.2023.105709>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Ltd.

1 **A new Mesozoic record of the pseudoscorpion family Garypinidae from Upper**
2 **Cretaceous (Santonian) Ajkaite amber, Ajka area, Hungary**

3 János Novák ^{a*}, Mark S. Harvey ^b, Márton Szabó ^{c,d}, Jörg U. Hammel ^e, Danilo Harms ^f,

4 Ulrich Kotthoff ^g, Christoph Hörweg^h, Manuel Brazidecⁱ, Attila Ósi ^d

5
6 ^a Eötvös Loránd University, Department of Systematic Zoology and Ecology, Pázámány Péter
7 Sétány 1/C, Budapest 1117, Hungary.

8 ^b Collections & Research, Western Australian Museum, 49 Kew Street, Welshpool, Western
9 Australia 6986, Australia; School of Biological Sciences, University of Western Australia,
10 Crawley, Western Australia 6009, Australia.

11 ^c Hungarian Natural History Museum, Department of Paleontology and Geology, Ludovika tér
12 2-6., Budapest 1083, Hungary

13 ^d ELTE Eötvös Loránd University, Institute of Geography and Earth Sciences, Department of
14 Palaeontology, Pázmány Péter sétány 1/C, Budapest 1117, Hungary.

15 ^e Helmholtz-Zentrum Hereon, Institute of Materials Physics, Max-Planck-Str. 1 Geesthacht,
16 21502, Germany.

17 ^f Leibniz-Institute for the Analysis of Biodiversity Change (LIB), Martin-Luther-King-Platz 3
18 Hamburg, 20146, Germany; Honorary Research Fellow: Harry Butler Centre, Murdoch
19 University, Murdoch, Western Australia; Australian Museum Research Institute, Australian
20 Museum, Sydney, Australia; SARChI Chair on Biodiversity Value and Change, Centre for
21 Invasion Biology, University of Venda, Thohoyandou, South Africa.

22 ^g Leibniz-Institute for the Analysis of Biodiversity Change (LIB), Martin-Luther-King-Platz 3
23 Hamburg, 20146, Germany; Institute for Geology, Universität Hamburg, Bundesstraße 55,
24 20146 Hamburg, Germany.

25 ^h Natural History Museum Vienna, 3. Zoology (Invertebrates), Burgring 7, 1010 Vienna,
26 Austria.

27 ⁱUniversity of Rennes, CNRS, Géosciences Rennes, UMR 6118, 35000 Rennes, France.

28

29 * novakjanos01@gmail.com

30

31 **Abstract.** Pseudoscorpions have a sparse fossil record although they are among the oldest
32 terrestrial lineages with origins extending to the Devonian (ca. 390 Ma). Amongst the 25 extant
33 families of pseudoscorpions, only 14 are known from fossils, most of which are preserved in
34 European ambers from the Eocene. Fossil pseudoscorpions from the Cretaceous of Europe have
35 only been reported from three localities, Archingeay amber (France), from the Rhenish Massif
36 (Germany), and from Álava and Teruel amber (Spain), but only one of these has ever been
37 formally described. Here we add a new fossil pseudoscorpion genus and species, *Ajkagarypinus*
38 *stephani* gen. et sp. nov., from an Upper Cretaceous (Santonian, ca. 86.3–83.6 Ma) amber
39 deposit in the Ajka Coal Formation (Ajka area, Hungary), the so-called Ajkaite. This fossil
40 extends the spatial range of the pseudoscorpion family Garypinidae and suggests that they
41 might have been widespread in the whole Eurasian landmasses under humid, subtropical
42 climatic conditions.

43

44 **Keywords:** Santonian, *Ajkagarypinus*, Pseudoscorpiones, amber inclusion, Late Cretaceous.

45

46 1. Introduction

47 Pseudoscorpions are small predatory arachnids, with more than 3900 species in 25 recent and
48 one fossil families (World Pseudoscorpiones Catalog, 2023 – as WPC, 2023 in the followings).

49 The earliest known species, *Dracochela deprehendor* Schawaller et al. 1991, was found in the

50 Gilboa mudstones, New York, United States and is of Middle Devonian (ca. 390 Ma) age
51 (Schawaller et al., 1991; Judson, 2012). The next oldest fossil species, *Archaeofeaella*
52 *henderickxi* Kolesnikov et al. 2022, was recently described as a claystone fossil from the
53 Ukraine, originating from the Late Triassic (Carnian – Norian) (Kolesnikov et al., 2022). These
54 older records aside, fossil pseudoscorpions are comparatively rare, and due to their small size
55 and relatively soft bodies, almost all of them are preserved as amber inclusions rather than
56 compression fossils. Moreover, most of the fossil specimens are restricted to Cretaceous and
57 Cenozoic ambers (Harms and Dunlop, 2017).

58 Even though each pseudoscorpion family most likely originated in the Palaeozoic or
59 Mesozoic (Benavides et al., 2019), only eight pseudoscorpion families have been recorded from
60 the Mesozoic, including Chthoniidae, Feaellidae, Pseudogarypidae, Ideoroncidae, Cheiridiidae,
61 Garypinidae, Cheliferidae, and Chernetidae (Fig. 1; Harms and Dunlop, 2017; Geißler et al.,
62 2022). The majority of Mesozoic pseudoscorpions have been reported from Burmese amber
63 (Cenomanian, Cretaceous, ca. 99 Ma) (e.g., Harms and Dunlop, 2017; Harvey et al., 2018; Porta
64 et al. 2020; Wriedt et al., 2021; Geißler et al., 2022). In contrast, only a handful of Mesozoic
65 pseudoscorpions have been reported in Europe. *Heurtaulia rossiorum* Judson, 2009 was
66 described from upper Albian Archingeay amber (ca. 100 Ma) from south-western France
67 (Perrichot, 2004; Judson, 2009). Furthermore, a carbonised fossil chela of a pseudogarypid
68 pseudoscorpion was found in the mid-Cretaceous infills of the Rhenish Massif (Judson, 2017),
69 which basically consists of upper Devonian limestone and middle Cretaceous karst fillings
70 (upper Albian – lower Cenomanian) (Viehofen et al., 2008). Two unidentified pseudoscorpions
71 have also been reported from the Álava amber deposit from Spain (Albian, Cretaceous, ca. 100–
72 113 Ma) by Delclòs et al. (2007) but not formally described. Sánchez-García et al. (2023)
73 reported a third Spanish specimen consisting of an isolated pedipalp from Albian Teruel amber,
74 identified as a Pseudogarypidae.

75 The pseudoscorpion family Garypinidae Daday, 1889 belongs to the superfamily
76 Garypinoidea Daday, 1889, with Larcidae Chamberlin, 1930 as its sister group (Benavides et
77 al., 2019), although two independent molecular studies failed to find evidence for a
78 monophyletic Garypinidae (Murienne et al., 2008; Benavides et al., 2019). The family was
79 originally created by Daday (1889) and subsequently regarded as a subfamily of Olpiidae
80 Banks, 1895 (Chamberlin 1930), but returned to family level by Judson (1993, 2005).
81 Garypinidae currently consists of 21 genera and 81 species (WPC, 2023). Modern
82 representatives of the family are present in all continents with the exception of Antarctica. They
83 reach their highest diversity in tropical and subtropical areas of the world occurring
84 predominantly in leaf litter, under bark, and under stones (Beier, 1963). Phoresy, a passive
85 method of dispersal, has been suggested for the fossil species *Garypinus electri* Beier, 1937
86 (Judson 2005), found in Eocene Baltic amber (Beier 1937). The oldest formal fossil member of
87 the family is *Amblyolpium burmiticum* (Cockerell, 1920) which was reported from Burmese
88 amber (Cockerell, 1920). Lately, Sánchez-García et al. (2023) restudied the Álava material
89 (Albian, Cretaceous, 100–113 Ma) and ascribed the two specimens to the family, pending
90 description. A further specimen of the family was also reported from Oise (Le Quesnoy) amber
91 (Ypresian, Eocene, 53 Ma), but lacks a species description (Nel et al. 1999; Judson 2009).

92 The Ajka amber (so-called *ajkaite*) is a recently re-discovered Upper Cretaceous
93 (Santonian) amber from Hungary, found to be very rich in arthropod inclusions. In this
94 assemblage, spiders, cockroaches, dipterans, hymenopterans and coleopterans have been
95 described so far (Borkent, 1997; Szabó et al., 2022a,b,c,d), and further inclusions are still
96 awaiting detailed study. The present paper describes the oldest currently known European
97 representative of Garypinidae, discovered in ajkaite, together with its taxonomic and faunistic
98 implications.

99

100

101 **2. Geological background**

102 Ajkaite, a type of often heavily oxidized amber containing organic inclusions, occurs in
103 the layers of the Upper Cretaceous Ajka Coal Formation (Fig. 2A). The first outcrops of the
104 formation were discovered in the Csinger Valley (Figs 2B; C), four kilometres southeast of the
105 city of Ajka in the 1860's, where the earliest researchers noticed the fossil resin content of the
106 coal-bearing layers (Hantken, 1878; Szabó, 1871) (Fig. 2A). The Ajka Coal Formation occurs
107 in the Bakony Mountains (Transdanubian Range), deposited in three different sub-basins (Ajka,
108 Magyarpolány-Devecser, and Gyepükaján) (see Császár and Góczán, 1988; Siegl-Farkas,
109 1988).

110 Upper Triassic (or, more rarely, Jurassic or Lower Cretaceous) limestones form the
111 geological base of the Ajka Coal Formation. Ajka coal is overlain by either the marine Jákó
112 Marl Formation (Haas et al., 1992), or the Eocene shallow marine deposits of the Szőc
113 Limestone Formation. The Ajka Coal Formation is built up by an alternation of coal or clayey
114 coal beds, marls, sands and sandstone beds, as well as light grey to brownish carbonaceous to
115 argillaceous pelitic rocks with interbedded molluscan lumachelles (Haas, 1983). The thickness
116 of the formation sometimes reaches 100 metres. The depositional process of the formation
117 resulted in at least seven main coal beds (from the oldest coal bed nr. 6 to the youngest coal bed
118 nr. 0). The thickness of the coal beds ranges from 80 to 360 cm (Kozma, 1991). Based on
119 palynological and nannoplankton investigations, the formation has been dated as Santonian,
120 86–83 Ma (Bodor and Baranyi, 2012; Bodrogi et al., 1998; Ósi et al., 2016; Siegl-Farkas and
121 Wagreich, 1996). Ajkaite is known to occur (in varying amounts) across all coal-bearing layers
122 of the formation.

123

124

125 3. Material and methods

126 Coal mining in the Ajka area commenced in 1865 (Kozma, 1991) and ceased in 2004.
127 Amber-bearing coal seams are no longer available, and ajkaites are no longer collectible. In the
128 last years of the operation of the Ajka coal mines, coal wastes (e.g. Jókai coal refuse) were a
129 popular destination for amateur fossil collectors. The pseudoscorpion described below is
130 located in a piece of reddish ajkaite, provided by the amateur collector István Szabó, who found
131 it most probably on the Jókai coal refuse on 27 March 1997. The specimen is now housed in
132 the invertebrate paleontological collection of the Department of Paleontology and Geology of
133 the Hungarian Natural History Museum (NHMUS; Budapest).

134 The amber stone was originally fused to a piece of the original embedding coal matrix
135 (Fig. 3A), which was removed during the preparation of the inclusion-bearing part of the amber.
136 The piece was then carefully cut into a near-cubic shape, the coal was removed, and polished
137 using thin silicon carbid sanding papers on a grinder polisher (Buehler MetaServ 3000) in order
138 to facilitate better observations of the specimen using light microscopes (Fig. 3B).

139 Line drawings were made with the use of Photoshop CS5 software. Photographs and
140 measurements were taken partly by a Nikon SMZ25 stereomicroscope and a Nikon DS-Ri2
141 camera unit, using Nikon NIS-Elements BR v.5.02.00 camera analysis software (all Nikon
142 Instruments Inc., New York, NY, USA) at the Natural History Museum of Vienna, and partly
143 with a Leica DMC4500 camera attached to a Leica M205C stereomicroscope (University of
144 Rennes) (the origin of the images are marked in Figure 3A–F). Additionally, a Zeiss Axioskop
145 2 was used for detailed examination (Eötvös Loránd University).

146 To achieve 3D details, the amber piece containing the pseudoscorpion was examined
147 further at the Deutsches Elektronen Synchrotron (Hamburg, Germany) where it was studied
148 using synchrotron radiation based micro-computed tomography (SR μ CT) (Figs 4A–F).
149 Imaging was performed at the Imaging Beamline P05 (IBL) (Greving et al., 2014; Haibel et al.,

2010; Wilde et al., 2016) operated by the Helmholtz-Zentrum Hereon at the storage ring
PETRA III (Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany). Imaging was
done at a photon energy of 32.6 keV and a sample to detector distance of 80 mm. Projections
were recorded using a custom developed 20 MP CMOS camera system (Lytaev et al., 2014)
with an effective pixel size of 1.28 μm . For each tomographic scan 2401 projections at equal
intervals between 0 and π have been recorded. Tomographic reconstruction has been done by
applying a transport of intensity phase retrieval approach and using the filtered back projection
algorithm (FBP) implemented in a custom reconstruction pipeline (Moosmann et al., 2014)
using Matlab (Math-Works) and the Astra Toolbox (van Aarle et al., 2015, 2016; Palenstijn et
al., 2011). For the processing, raw projections were binned for further processing two times,
resulting in an effective pixel size of the reconstructed volume of 2.56 μm . Subfamily level
classification follows Harvey's work (Harvey, in press).

162

163

164 **4. Systematic paleontology**

165 Order: Pseudoscorpiones de Geer, 1778

166 Family: Garypinidae Daday, 1889

167 Subfamily: Garypininae Daday, 1889

168 Remarks: the new taxon can be clearly attributed to the family Garypinidae by the combination
169 of the following characteristics: venom apparatus present in both chelal fingers; movable
170 cheliceral finger with just 1–2 subapical teeth; *gs* subdistal on movable cheliceral finger;
171 carapace sub-rectangular; pseudosternum absent; surface of pedipalps and carapace smooth;
172 tergites divided; rallum with four blades; basal concentration of the trichobothria of the internal
173 series on fixed chelal finger; legs I and II diplotarsate; arolia divided.

174

175 Genus *Ajkagarypinus* gen. nov.

176 urn:lsid:zoobank.org:act:031AF2D1-ADDC-4721-B62A-17AAAE948126

177 *Etymology.* The genus is named after the settlement of Ajka (Hungary), the locality of Ajkaite.

178 *Type species.* *Ajkagarypinus stephani* sp. nov.

179 *Diagnosis.* *Ajkagarypinus* gen. nov. differs from all other genera of the family Garypinidae by
180 the setation of the cheliceral hand, in which *es* is duplicated, and the combination of the
181 following characteristics: legs I and II with patella about same size as femur; cheliceral setae *ls*
182 and *is* situated apart from each other; fixed chelal finger with 8, movable with 4 trichobothria;
183 trichobothrium *est* situated approximately midway between *et* and *esb*; trichobothrium *st* dorsal
184 to *b*; pedipalpal femur without trichobothria.

185 *Remarks.* There is one character that requires further discussion, which is the position of the
186 trichobothria on the movable chelal finger. In most garypinids, they are situated in a straight
187 line with *st* situated between *sb* and *t*, whereas in species of *Serianus*, *Protogarypinus* and
188 *Teratolpium*, trichobothrium *st* is situated directly dorsal to *sb* (e.g. Beier, 1954, 1959; Hoff,
189 1956, 1964; Mahnert 1988; Muchmore, 1968; Piedra-Jiménez, et al., 2019). It is difficult to
190 ascertain the homology of the trichobothria in *Ajkagarypinus*, but it is not inconceivable that
191 the dorsalmost trichobothrium is *st*, and its position dorsal to *b* represents a very unique
192 morphological modification.

193

194 *Ajkagarypinus stephani* sp. nov.

195 urn:lsid:zoobank.org:act:C7FEB2F3-F3DB-4A6D-B8E4-74DF49CCDAC2

196 Figs 3–5.

197

198 *Etymology.* This species is named in honour of the amateur fossil collector István Szabó
199 (Stephan is the Latin translation of the Hungarian name István), who provided the amber piece
200 for study and donated it to the collection of the Hungarian Natural History Museum.

201 *Holotype.* NHMUS PAL 2022.91.1., adult, sex unknown.

202 *Locality and horizon.* Ajka Coal Formation (Santonian, Upper Cretaceous), unknown shaft of
203 the Ajka coal mine; Ajka (Hungary).

204 *Diagnosis.* See the diagnosis of the genus.

205 *Description.* Colour: body and pedipalps reddish brown; carapace slightly darker than the
206 pedipalps; legs lighter yellowish brown; opisthosoma partly missing (Figs 3C, D, E, F).

207 Carapace (Figs 3F, 4A, 5A). Conical and 1.55× longer than wide, reaching its maximal width
208 at the base. Two pairs of corneate eyes present near the anterior margin. Vestitural setae
209 acuminate. Lyrifissures of carapace not visible. Surface smooth.

210 Chelicera (Fig. 5B). Hand with 6 acuminate setae, two of which situated at the externolateral
211 side of the hand: *es* duplicated; *ls* and *is* situated apart from each other; movable finger with 1
212 acuminate subdistal seta. Rallum with four blades. Serrula exterior: ca. 15 blades; lamina
213 exterior not visible. Galea long and simple, probably broken on left chelicera.

214 Coxal region and genitalia not visible due to a crack on the surface of the amber piece.

215 Pedipalps (Fig. 5C). Robust. Vestitural setae acuminate, surface smooth. Femur broadening
216 gradually from the pedicel, widest at its middle part. Femur 2.95, patella 1.53, chela (with
217 pedicel) 3.09, hand 1.58 × longer than broad, movable finger 0.79 times longer than hand
218 (without pedicel). Chelal fingers slightly shorter than the hand (without pedicel) (Fig. 5D).

219 Fixed chelal finger with 8, movable with 4 trichobothria. Trichobothria *eb* and *esb* grouped
220 basally; *est* medially, halfway between *et* and *esb*; *et* subterminally; *ib*, *isb*, *ist*, and *it* grouped
221 near the base of finger; trichobothria *b*, *sb*, and *st* forming a group near to base; *t* distad to them,
222 approximately halfway between *sb* and the finger's tip. Venom glands and venom ducts present

223 in both fingers, exact length of venom ducts not determinable due to the preservation of the
224 fossil. Chelal teeth small, triangular, slightly retrose, and closely spaced; equal in form and size
225 although the exact number cannot be determined due to bubbles partly covering them.
226 Accessory teeth absent.

227 Legs. Junction between femora and patellae I and II oblique. Leg I and II diplotarsate; arolia
228 long and divided; claws not visible; vestitural setae acuminate. Left legs III, IV, and right IV
229 missing, but right leg III visible only on synchrotron model.

230 Abdomen: tergites divided. First three not clearly visible, remaining ones missing.

231 Measurements (in mm). Body length (without the missing abdominal segments) 0.682.
232 Carapace 0.39×0.31 . Chelicerae 0.13×0.07 ; movable finger length 0.08. Pedipalp: femur 0.39
233 $\times 0.13$; patella: 0.27×0.18 ; chela 0.57×0.19 ; hand length (without pedicel) 0.29; movable
234 finger length 0.23.

235 *Remarks.* The specimen preserved muscle filaments as imprints in the pedipalps that are clearly
236 visible in both trochanters, femora, patellae and chelae (Figs 3E-F). Muscle imprints are not
237 rare in amber (e.g., Henwood, 1992; van de Kamp et al., 2014; Li et al., 2021) but this is the
238 first time that such an exceptional preservation is recorded in Ajkaite.

239

240

241 **5. Discussion**

242 Pseudoscorpion fossils are very rare in the Carpathian Basin. Sziráki and Dulai (2002)
243 reported a poorly preserved compression fossil pseudoscorpion specimen from the upper
244 Miocene (Sarmatian) of the Tokaj Mts. (Hungary). *Ajkagarypinus stephani* gen. et sp. nov.
245 represents the second fossil pseudoscorpion from the Carpathian Basin, further highlighting the
246 importance of the present discovery (not counting that Mesozoic pseudoscorpions are overall
247 rare in the European fossil record, see above).

248 The new garypinid fossil from the Santonian of Hungary is the second described species
249 of this family from the Mesozoic following *Amblyolpium burmiticum* (Cockerell, 1920) and the
250 third described Mesozoic pseudoscorpion species from Europe following *Heurtaulia*
251 *rossiorum* Judson, 2009 and *Archaeofearella henderickxi* Kolesnikov, Turbanov, Eskov,
252 Propistsova et Bashkuev, 2022. It is also one of the oldest fossil pseudoscorpions in amber.

253

254 *Palaeoecology*

255 The discovery of *Ajkagarypinus stephani* gen. et sp. nov. is an interesting addition to
256 our knowledge on the Ajkaite paleofauna, and also the Ajka coal paleobiota. The depositional
257 area of the Ajka Coal likely reflects an arboreous-swampy and lacustrine environment. The
258 climate has been estimated to be a tropical monsoon climate with presumably seasonal, high
259 precipitation (2000–2500 mm per year) (L. Rákosi, pers. comm.; Siegl-Farkas, 1988). This
260 conforms with the ecological needs of the closest living relatives of the fossil plant forms
261 documented from the Ajka Coal, so the palaeoclimate must have been tropical to subtropical
262 (Bodor et al., 2012). Based on palynological and palaeocarpological studies, Normapolles-
263 related forests are suggested with a fern-dominated understorey (Bodor and Baranyi, 2012).

264 The arthropod fauna studied in detail so far conforms with this environment. Among the
265 spider fossils, a hersiliid spider, *Hungarosilia verdesi* Szabó et al., 2022a, has been described
266 (Szabó et al., 2022a). Modern members of this family generally live in warm, tropical,
267 subtropical climates where many of these are associated with tree bark habitats. The
268 ceratopogonid dipterans *Adelohelea magyarica* Borkent, 1997 and *Leptoconops clava* Borkent,
269 1997, the hymenopterans *Ajkanesia harmincipsziloni* Szabó et al., 2022b, *Amissodigitus belai*
270 Szabó et al., 2022b and *Spathiopteryx soosi* Szabó et al., 2022b, the click-beetle *Ajkaelater*
271 *merkli* Szabó et al. 2022c and the cockroach *Alienopterix santonicus* Szabó et al. 2022d also fit
272 well into the climatic conditions mentioned above (Borkent, 1997; Szabó et al., 2022b, c, d).

273 The occurrence of a garypinid pseudoscorpion further enriches this diverse arthropod world and
274 help to reconstruct the former environment more accurately. Extant representatives of the
275 family Garypinidae have been found in diverse habitats, such as in rainforests (Harvey, 1988;
276 Dashdamirov, 2008), temperate forests (Hoff, 1961; Benedict and Malcolm, 1978; Harvey and
277 Štáhlavský, 2010; Piedra-Jiménez et al. 2019), woodlands (Gardini and Gavalas, 2021; Hoff,
278 1959), semi-arid deserts (Chamberlin, 1923; Heurtault, 1970; Nassirkhani and Doustaresharaf,
279 2019) and sea-shore habitats (Lee, 1979; Mahnert 2014). Habitat types include moist leaf litter,
280 tree bark and rocks. In the light of occurrences in rainforests and other mesic habitats, the
281 occurrence of Garypinidae fits well into the ecosystem of the Ajka Coal paleohabitat.

282 Phoresy as a means of passive dispersal might occur in the Garypinidae and was
283 suggested for *Garypinus electri*, an Eocene fossil member of the family Garypinidae (Judson,
284 2005) from Baltic amber. Phoresy would be an ideal form of dispersal in a system of islands,
285 just like the western Tethyan archipelago had been during the Late Cretaceous (Csiki-Sava et
286 al. 2015). *Ajkagarypinus stephani* might have been living under or near tree bark, using phoresy
287 on insects as a means of passive dispersal similar to many modern pseudoscorpions. That said,
288 the Garypinidae are ecologically diverse and also occur in leaf litter habitats whereas their
289 morphology seems quite conserved both over time and between recent genera. The suggestion
290 of phoresy is therefore hypothetical and can only be tested if additional specimens become
291 available.

292

293 **6. Conclusions**

294 A new fossil pseudoscorpion genus and species, *Ajkagarypinus stephani* gen. et sp. nov., is
295 described from an Upper Cretaceous (Santonian, ca. 86.3–83.6 Ma) amber deposit in the Ajka
296 Coal Formation (Ajka area, Hungary), the so-called Ajkaite. *Ajkagarypinus stephani* gen. et sp.
297 nov., is the third described Mesozoic pseudoscorpion species from Europe following

298 *Heurtaulia rossiorum* Judson, 2009 and *Archaeofeaella henderickxi* Kolesnikov, Turbanov,
299 Eskov, Propistsova et Bashkuev, 2022. This fossil extends the spatial range of the family
300 Garypinidae and suggests that they might have been widespread in the whole Eurasian
301 landmasses under humid, subtropical climatic conditions.

302

303 **7. Acknowledgements**

304 Dr. Jason A. Dunlop and an anonymous reviewer are thanked for their constructive suggestions
305 and comments, by which they highly improved an earlier version of the manuscript. Special
306 thanks belong to Constantin Mey for his help in refining the 3D Synchrotron model. We thank
307 the Department of Palaeontology and the Department of Systematic Zoology and Ecology of
308 the Eötvös Loránd University, to Dr. Klára Dózsa-Farkas, the Plant Protection Institute of
309 Centre for Agricultural Research (with the technical assistance of Dr. Jenő Kontschán) and the
310 Natural History Museum Vienna for the support during light microscope investigations and
311 photography of the amber stone specimen. This research was supported by the MTA-ELTE
312 Lendület Dinosaur Research Group (Grant No. 95102), National Research, Development and
313 Innovation Office (NKFIH K 116665, K 131597, PD 130190, FK 130627). The authors
314 acknowledge the DESY (Hamburg, Germany), a member of the Helmholtz Association HGF,
315 for the provision of experimental facilities. Beamtime was allocated for proposal I-20190010.
316 Danilo Harms and Ulrich Kotthoff thank the German Research Foundation (DFG) for financial
317 support (HA 8785/5-1 and KO 3944/10-1).

318

319 **8. References**

320 Aarle, W. van, Palenstijn, W.J., De Beenhouwer, J., Altantzis, T., Bals, S., Batenburg, K.J.,
321 Sijbers, J. 2015. The ASTRA Toolbox: A platform for advanced algorithm development in

- 322 electron tomography. *Ultramicroscopy* 157, 35–47,
323 <http://dx.doi.org/10.1016/j.ultramic.2015.05.002>
- 324 Aarle, W. van, Palenstijn, W.J., Cant, J., Janssens, E., Bleichrodt, F., Dabravolski, A., De
325 Beenhouwer, J., Batenburg, K.J., Sijbers, J. 2016. Fast and flexible X-ray tomography using
326 the ASTRA toolbox. *Optics Express* 24(22), 25129–25147,
327 <http://dx.doi.org/10.1364/OE.24.025129>
- 328 Beier, M. 1937. Pseudoscorpione aus dem baltischen Bernstein. Festschrift zum 60.
329 Geburtstage von Professor Dr. Embrik Strand, Riga, 2, 302–316.
- 330 Beier, M. 1954. Report from Prof. T. Gislén's expedition to Australia in 1951–1952. 7.
331 Pseudoscorpionidea. *Acta Universitatis Lundensis, nova series* (2) 50, 1–26.
- 332 Beier, M. 1959. Zur Kenntnis der Pseudoscorpioniden-Fauna des Andengebietes. *Beiträge zur*
333 *Neotropischen Fauna* 1, 185–228.
- 334 Beier, M. 1963. Ordnung Pseudoscorpionidea (Afterscorpione). *Bestimmungsbücher zur*
335 *Bodenfauna Europas*, 1. Akademie-Verlag, Berlin, 313.
- 336 Benavides, L. R., Cosgrove, J. G., Harvey, M. S., Giribet, G. 2019. Phylogenetic interrogation
337 resolves the backbone of the Pseudoscorpiones tree of life. *Molecular Phylogenetics and*
338 *Evolution* 139: 106509. <http://dx.doi.org/10.1016/j.ympev.2019.05.023>
- 339 Benedict, E.M., Malcolm, D.R. 1978. Some garypoid false scorpions from western North
340 America (Pseudoscorpionida: Garypidae and Olpiidae). *Journal of Arachnology* 5, 113–132.
- 341 Bodor, E.R., Baranyi, V. 2012. Palynomorphs of the Normapolles group and related plant
342 mesofossils from the Iharkút vertebrate site, Bakony Mountains (Hungary). *Central*
343 *European Geology* 55 (3), 259–292. <https://doi.org/10.1556/CEuGeol.55.2012.3.3>
- 344 Bodor, E.R., Baranyi, V., Hermanová, Z. 2012. The earliest Sabiaceae fruit remains of Hungary.
345 *Hantkeniana* 7, 11–18.

- 346 Bodrogi, I., Fogarasi, A., Yazikova, E.A., Sztanó, O., Báldi-Beke, M. 1998. Upper Cretaceous
347 of the Bakony Mts. (Hungary): sedimentology, biostratigraphy, correlation. *Zentralblatt für*
348 *Geologie und Paläontologie*, Teil I 11/12, 1179-1194.
- 349 Borkent, A. 1997. Upper and Lower Cretaceous biting midges (Ceratopogonidae: Diptera) from
350 Hungarian and Austrian amber and the Koonwarra Fossil Bed of Australia. *Stuttgarter*
351 *Beiträge zur Naturkunde Serie B* 249, 10.
- 352 Chamberlin, J.C. 1923. New and little known pseudoscorpions, principally from the islands and
353 adjacent shores of the Gulf of California. *Proceedings of the California Academy of Sciences*
354 (4) 12, 353–387.
- 355 Chamberlin, J.C. 1930. A synoptic classification of the false scorpions or chela-spinners, with
356 a report on a cosmopolitan collection of the same. Part II. The Diplosphyronida (Arachnida-
357 Chelonethida). *Annals and Magazine of Natural History* (10) 5, 1–48, 585–620.
- 358 Cockerell, T.D.A. 1920. Fossil arthropods in the British Museum-I. *Annals and Magazine of*
359 *Natural History*, ser. 9, 5, 273–279.
- 360 Császár, G.H., Góczán, F. 1988. A Bakony felső-kréta kőszénkutató és kőszén lág vizsgálata.
361 (in Hungarian. English translation: Upper Cretaceous coal prospecting and peat bog studies
362 in the Bakony Mts). In: *Magyar Állami Földtani Intézet Évi Jelentése 1986-ról*, pp. 155–
363 178.
- 364 Csiki-Sava, Z., Buffetaut, E., Ősi, A., Pereda-Suberbiola, X., Brusatte, S.L. 2015. Island life in
365 the Cretaceous - faunal composition, biogeography, evolution, and extinction of land-living
366 vertebrates on the Late Cretaceous European archipelago. *ZooKeys* 469, 1–161.
367 <https://doi.org/10.3897/zookeys.469.8439>
- 368 Daday, E. 1889. A Magyar Nemzeti Múzeum álskorpióinak áttekintése. *Természetrázi Füzetek*
369 11, 111–136, 165–192.

- 370 Dashdamirov, S. 2008. A new species of false scorpion from Baltic amber with a redescription
371 of *Electrochelifer balticus* Beier, 1955 and remarks on some fossil Cheliferidae (Arachnida:
372 Chelonethida). *Acta Biologica Benrodis* 14, 1–13.
- 373 Delclòs, X., Arillo, A., Peñalver, E., Barrón, E., Soriano, C., López Del Valle, R., Bernárdez,
374 E., Corral, C., Ortuño, V.M. 2007. Fossiliferous amber deposits from the Cretaceous
375 (Albian) of Spain. *Compte Rendus - Palevol* 6, 135–149.
376 <https://doi.org/10.1016/j.crpv.2006.09.003>
- 377 Gardini, G., Gavalas, I. 2021. The pseudoscorpions of the island of Iraklia (Cyclades, Greece).
378 *Parnassiana Archives* 9, 43–78.
- 379 Geißler, C., Kotthoff, U., Hammel, J., Harvey, M.S., Harms, D. 2022. The first fossil of the
380 pseudoscorpion family Ideoroncidae (Arachnida: Pseudoscorpiones): a new taxon from the
381 mid-Cretaceous of northern Myanmar. *Cretaceous Research* 130 (105030). 1–12.
- 382 Greving, I., Wilde, F., Ogurreck, M., Herzen, J., Hammel, J. U., Hipp, A., Friedrich, F.,
383 Lottermoser, L., Dose, T., Burmester, H., Müller, M., Beckmann, F. 2014. P05 imaging
384 beamline at PETRA III: first results. In: Stuart, R. S., (Ed.), *Proceedings of SPIE -*
385 *Developments in X-Ray Tomography IX*, Vol. 9212, San Diego, California, United States,
386 pp. 92120O-8.
- 387 Haas, J. 1983. Senonian in the Transdanubian Central Range. *Acta Geologica Hungarica* 26,
388 21–40.
- 389 Haas, J., Jocha-Edelényi, E., Császár, G. 1992. Upper Cretaceous coal deposits in Hungary. In:
390 McCabe, P., Parris, J.T. (Eds.), *Controls on the distribution and quality of Cretaceous coals*,
391 vol. 267. Geological Society of America, Special Paper, pp. 245–262.
- 392 Haibel, A., Ogurreck, M., Beckmann, F., Dose, T., Wilde, F., Herzen, J., Müller, M., Schreyer,
393 A., Nazmov, V., Simon, M., Last, A., Mohr, J. 2010. Micro- and nano-tomography at the

- 394 GKSS Imaging Beamline at PETRA III. SPIE Optical Engineering + Applications, Vol.
395 7804. SPIE, pp. 8.
- 396 Hantken, M. 1878. A magyar korona országainak széntelepei és szénbányászata. Légrády,
397 Budapest, p. 331.
- 398 Harms, D., Dunlop, J.A. 2017. The fossil history of pseudoscorpions (Arachnida:
399 Pseudoscorpiones). Fossil Record 20, 215–238.
- 400 Harvey, M.S. 1988. Pseudoscorpions from the Krakatau Islands and adjacent regions, Indonesia
401 (Chelicerata: Pseudoscorpionida). Memoirs of the Museum of Victoria 49, 309–353.
- 402 Harvey, M.S. A preliminary phylogeny for the pseudoscorpion family Garypinidae
403 (Pseudoscorpiones: Garypinoidea), with new taxa and remarks on the Australasian fauna.
404 Invertebrate Systematics, in press.
- 405 Harvey, M.S., Cosgrove, J.G., Harms, D., Selden, P.A., Shih, C., Wang, C.-C. 2018. The oldest
406 chthonioid pseudoscorpion (Arachnida: Pseudoscorpiones: Chthonioidea: Chthoniidae. a
407 new genus and species from mid-Cretaceous Burmese amber. Zoologischer Anzeiger 273,
408 102–111.
- 409 Harvey, M.S., Štáhlavský, F. 2010. A review of the pseudoscorpion genus *Oreolpium*
410 (Pseudoscorpiones: Garypinidae), with remarks on the composition of the Garypinidae and
411 on pseudoscorpions with bipolar distributions. Journal of Arachnology 38, 294–308.
- 412 Henwood, A. 1992. Exceptional preservation of Dipteran flight muscle and the taphonomy of
413 insects in amber. Palaios 7, 203–212. <https://doi.org/10.2307/3514931>
- 414 Heurtault, J. 1970. Pseudoscorpions du Tibesti (Tchad). I. Olpiidae. Bulletin du Muséum
415 National d'Histoire Naturelle, Paris, (2) 41, 1164–1174.
- 416 Hoff, C.C. 1956. Diplosphyronid pseudoscorpions from New Mexico. American Museum
417 Novitates 1780, 1–49.
- 418 Hoff, C.C. 1959. The ecology and distribution of the pseudoscorpions of north-central New

- 419 Mexico. University of New Mexico Publications in Biology 8, 1–68.
- 420 Hoff, C.C. 1961. Pseudoscorpions from Colorado. Bulletin of the American Museum of Natural
421 History 122, 409–464.
- 422 Hoff, C.C. 1964. The pseudoscorpions of Jamaica. Part 3. The suborder Diplosphyronida.
423 Bulletin of the Institute of Jamaica, Science Series 10(3), 1–47.
- 424 Judson, M.L.I. 1993. African Chelonethi: studies on the systematics, biogeography and natural
425 history of African pseudoscorpions (Arachnida) [Abstract]. Index to theses, with abstracts,
426 accepted for higher degrees by the universities of Great Britain and Ireland 42(2), 697.
- 427 Judson, M.L.I. 2005. Baltic amber fossil of *Garypinus electri* Beier provides first evidence of
428 phoresy in the pseudoscorpion family Garypinidae (Arachnida: Chelonethi). In 'European
429 Arachnology 2003 (Proceedings of the 21st European Colloquium of Arachnology, St.-
430 Petersburg, 4-9 August 2003).' (Eds. DV Logunov and D Penney) pp. 127–131. (Moscow:
431 KMK Scientific Press Ltd)
- 432 Judson, M. L. I. 2009. Cheliferoid pseudoscorpions (Arachnida, Chelonethi) from the Lower
433 Cretaceous of France. Geodiversitas 31, 61–71.
- 434 Judson, M.L.I. 2012. Reinterpretation of *Dracochela deprehendor* (Arachnida:
435 Pseudoscorpiones) as a stem-group pseudoscorpion. Palaeontology 55, 261–283.
- 436 Judson, M. L. I. 2017. A new subfamily of Feallidae (Arachnida, Chelonethi, Fealloidea)
437 from Southeast Asia. Zootaxa 4258, 1–33.
- 438 Kamp, T. van de, Dos Santos, R., Baumach, T., Krogmann, L. 2014. Scanning the past -
439 Synchrotron microtomography of fossil wasps in amber. Entomology Heute 26, 151–160.
- 440 Kolesnikov, V.B., Turbanov, I.S., Eskov, K.Y., Propistsova, E.A., Bashkuev, A.S. 2022. First
441 non-amber Mesozoic pseudoscorpion from Upper Triassic deposits of Eastern Europe, with
442 a description of two new fossil subfamilies (Arachnida, Pseudoscorpiones, Feallidae).
443 Papers in Palaeontology, e1466, 1–14. <http://dx.doi.org/10.1002/spp2.1466>

- 444 Kozma, K. 1991. Az ajkai szénbányászat története (History of the Coal Mining at Ajka).
445 Veszprémi Szénbányák Kiadó, Veszprém, p. 531 (in Hungarian).
- 446 Lee, V.F. 1979. The maritime pseudoscorpions of Baja California, México (Arachnida:
447 Pseudoscorpionida). Occasional Papers of the California Academy of Sciences 131: i–iv, 1–
448 38.
- 449 Li, Y., Yamamoto, S., Huang, D., Cai, C. 2021. New species of *Paradontomma* from mid-
450 Cretaceous Burmese amber with muscle tissue preservation (Coleoptera: Archostemata:
451 Ommatidae). Papéis Avulsos de Zoologia 61, e20216153. [https://doi.org/10.11606/1807-](https://doi.org/10.11606/1807-0205/2021.61.53)
452 0205/2021.61.53
- 453 Lytaev, P., Hipp, A., Lottermoser, L., Herzen, J., Greving, I., Khokhriakov, I., Meyer-Loges,
454 S., Plewka, J., Burmester, J., Caselle, M., Vogelgesang, M., Chilingaryan, S., Kopmann, A.,
455 Balzer, M., Schreyer, A., Beckmann, F. 2014. Characterization of the CCD and CMOS
456 cameras for grating-based phase-contrast tomography. SPIE Optical Engineering +
457 Applications, Vol. 9212. SPIE, pp. 10.
- 458 Mahnert, V. 1988. Zwei neue Garypininae-Arten (Pseudoscorpiones: Olpiidae) aus Afrika mit
459 Bemerkungen zu den Gattungen *Serianus* Chamberlin und *Paraserianus* Beier. Stuttgarter
460 Beiträge zur Naturkunde (A) 420, 1–11.
- 461 Mahnert, V. (2014). Pseudoscorpions (Arachnida: Pseudoscorpiones) from the Galapagos
462 Islands (Ecuador). Revue Suisse de Zoologie 121: 135–210.
- 463 Moosmann, J., Ershov, A., Weinhardt, V., Baumbach, T., Prasad, M. S., LaBonne, C., Xiao,
464 X., Kashef, J., Hoffmann, R. 2014. Time-lapse X-ray phase-contrast microtomography for
465 in vivo imaging and analysis of morphogenesis. Nature Protocols, 9, 294–304.
- 466 Muchmore, W.B. 1968. A new species of the pseudoscorpion genus *Serianus* (Arachnida,
467 Chelonethida, Olpiidae) from North Carolina. Entomological News 79, 145–150.

- 468 Murienne, J., Harvey, M.S., Giribet, G. 2008. First molecular phylogeny of the major clades of
469 Pseudoscorpiones. *Molecular Phylogenetics and Evolution* 49, 170–184.
470 <https://doi.org/10.1016/j.ympev.2008.06.002>
- 471 Nassirkhani, M., Doustaresharaf, M.M. 2019. Description of a new pseudoscorpion species of
472 the genus *Amblyolpium* Simon (Pseudoscorpiones: Garypinidae) from north-western Iran.
473 *Zoology in the Middle East* 65, 268–273. <https://doi.org/10.1080/09397140.2019.1609174>
- 474 Nel, A., Plöeg, G. de., Dejax, J., Dutheil, D., de Franceschi, D., Gheerbrant, E., Godinot, M.,
475 Hervet, S., Menier, J.-J., Ague, M., Bignot, G., Cavagnetto, C., Duffard, S., Gaudant, J.,
476 Hua, S., Jpsang, A., de Lapparent de Broin, F., Pozzi, J.-P., Paicheler, J.- C., Beuchet, F.,
477 Rage, J.-C. 2009. Un gisement sparnacien exceptionnel à plantes, arthropodes et vertébrés
478 (Éocène basal, MP7. Le Quesnoy (Oise, France), *C. R. Acad. Sci.-Series IIA – Earth and*
479 *Planetary Science* 329, 65–72.
- 480 Ósi, A., Bodor, E.R., Makádi, L., Rabi, M. 2016. Vertebrate remains from the Upper Cretaceous
481 (Santonian) Ajka Coal Formation, western Hungary. *Cretaceous Research* 57, 228-238.
482 <https://doi.org/10.1016/j.cretres.2015.04.014>
- 483 Palenstijn, W. J., Batenburg, K. J., Sijbers, J. 2011. Performance improvements for iterative
484 electron tomography reconstruction using graphics processing units (GPUs). *Journal of*
485 *Structural Biology* 176, 250–253. <https://doi.org/10.1016/j.jsb.2011.07.017>
- 486 Perrichot, V. 2004. Early Cretaceous amber from south-western France: insights into the
487 Mesozoic litter fauna. *Geologica Acta*, 2, 9–22. <https://doi.org/10.1344/105.000001629>
- 488 Piedra-Jiménez, D.F., Alvarez-Padilla, F., and González Santillán, E. 2019. Two new species
489 of pseudoscorpions (Arachnida: Pseudoscorpiones) from a Mexican oak forest near Pico de
490 Orizaba National Park. *Journal of Arachnology* 47: 95–109. [https://doi.org/10.1636/0161-](https://doi.org/10.1636/0161-8202-47.1.95)
491 [8202-47.1.95](https://doi.org/10.1636/0161-8202-47.1.95)

- 492 Porta, A., Michalik, P., Franchi, E., Proud, D. 2020. The first fossil pycnocheiridiine
493 pseudoscorpion (Pseudoscorpiones: Cheiridiidae: *Procheiridium* gen. nov.) from Cretaceous
494 Burmese amber. *Zootaxa* 4801, 142–150. <https://doi.org/10.11646/zootaxa.4801.1.6>
- 495 Sánchez-García, A., Delclòs, X., Peñalver, E. 2023. False scorpions (Arachnida:
496 Pseudoscorpiones) from Cretaceous Spanish amber. 4th Paleontological Virtual Congress
497 [online], p. 34.
- 498 Schawaller, W. 1981. Pseudoskorpione (Cheliferidae) phoretisch auf Käfern (Platypodidae) in
499 Dominikanischen Bernstein (Stuttgarter Bernsteinsammlung: Pseudoscorpionidea und
500 Coleoptera). *Stuttgarter Beiträge zur Naturkunde (B)* 71, 1–17.
- 501 Schawaller, W., Shear, W. A., Bonamo, P. M. 1991. The first Paleozoic pseudoscorpions
502 (Arachnida, Pseudoscorpionida). *American Museum Novitates* 3009, 1–24.
- 503 Schwarze, D., Harms, D., Hammel, J., Kotthoff, U. 2021. The first fossils of the most basal
504 pseudoscorpion family (Arachnida: Pseudoscorpiones: Pseudotyranochthoniidae):
505 evidence for major biogeographical shifts in the European paleofauna. *PalZ* 96: 11–27.
506 <https://doi.org/10.1007/s12542-021-00565-8>
- 507 Siegl-Farkas, Á. 1988. Az Ajkai Kőszén Formáció palynozstratigráfiája és fejlődéstörténete.
508 In: *Magyar Állami Földtani Intézet Évi Jelentése 1986-ról*, 179–209.
- 509 Siegl-Farkas, Á., Wagreich, M. 1996. Correlation of palyno- (spores, pollen, dinoflagellates)
510 and calcareous nannofossil zones in the Late Cretaceous of the Northern Calcareous Alps
511 (Austria) and the Transdanubian Central Range (Hungary). *Advances in Austrian -*
512 *Hungarian Joint Geological Research, Budapest*, 127–135.
- 513 Szabó, J. 1871. Az ajkai kőszénteleg a Bakonyban. (in Hungarian: The coal deposits of Ajka in
514 the Bakony Mountains). *Földtani Közlöny* 1, 124–130.
- 515 Szabó, M., Hammel, J. U., Harms, D., Kotthoff, U., Bodor, E., Novák, J., Kovács, K., Ósi, A.
516 2022a. First record of the spider family Hersiliidae (Araneae) from the Mesozoic of Europe

- 517 (Bakony Mts, Hungary). Cretaceous Research, 131.
518 <https://doi.org/10.1016/j.cretres.2021.105097>
- 519 Szabó, M., Brazidec, M., Perrichot, V., Szenti, I., Kukovecz, Á., Ósi, A. 2022b. A unique record
520 of the Late Cretaceous of East-Central Europe: The first fossil wasps (Hymenoptera:
521 Bethyridae, Spathiopterygidae) from the ajkaite amber (Bakony Mts., western Hungary).
522 Cretaceous Research, 139, 105314. <https://doi.org/10.1016/j.cretres.2022.105314>
- 523 Szabó, M., Kundera, R., Hoffmannová, J., Németh, T., Bodor, E., Szenti, I., Prosvirov, A. S.,
524 Kukovecz, Á., Ósi, A. 2022c. The first mainland European Mesozoic click-beetle
525 (Coleoptera: Elateridae) revealed by X-ray micro-computed tomography scanning of an
526 Upper Cretaceous amber from Hungary. Scientific Reports. [https://doi.org/10.1038/s41598-](https://doi.org/10.1038/s41598-021-03573-512:24)
527 [021-03573-512:24](https://doi.org/10.1038/s41598-021-03573-512:24)
- 528 Szabó, M., Szabó, P., Kóbor, P., Ósi, A. 2022d. *Alienopterix santonicus* sp. n., a metallic
529 cockroach from the Late Cretaceous ajkaite amber (Bakony Mts, western Hungary)
530 documents Alienopteridae within the Mesozoic Laurasia. Biologia.
531 <https://doi.org/10.1007/s11756-022-01265-7>
- 532 Sziráki, G., Dulai, A., 2002. Sarmatian (Late Miocene) arthropods from Tállya and
533 neighbouring localities (Tokaj Mts, Hungary. A preliminary report. Annales Historico-
534 Naturales Musei Nationalis Hungarici 94, 31-44.
- 535 Viehofen, A., Hartkopf-Fröder, C., and Friis, E. M. 2008. Inflorescences and flowers of
536 *Mauldinia angustiloba* sp. nov. (Lauraceae) from Middle Cretaceous karst infillings in the
537 Rhenish Massif, Germany. International Journal of Plant Sciences 169, 871–889.
- 538 Wilde, F., Ogurreck, M., Greving, I., Hammel, J. U., Beckmann, F., Hipp, A., Lottermoser, L.,
539 Khokhriakov, I., Lytaev, P., Dose, T., Burmester, H., Müller, M., Schreyer, A. 2016. Micro-
540 CT at the imaging beamline P05 at PETRA III. AIP Conference Proceedings 1741, 030035.
541 <https://doi.org/10.1063/1.4952858>.

542 World Pseudoscorpiones Catalog 2023. World Pseudoscorpiones Catalog. Natural History
 543 Museum, Bern. <https://wac.nmbe.ch/order/pseudoscorpiones/3>, accessed at: 2023.05.31.

544 Wriedt, A.-L., Harvey, M.S., Hammel, J.U., Kotthoff, U., Harms, D. 2021. The second
 545 chthonioid pseudoscorpion (Pseudoscorpiones: Chthoniidae) from mid-Cretaceous Burmese
 546 amber: a new genus with unique morphological features and potential Gondwanan affinities.
 547 Journal of Arachnology 48, 311–321. <https://doi.org/10.1636/JoA-S-20-017>

548

549

550 Figure 1. The phylogenetic tree of pseudoscorpions according to Benavides et al. (2019) with
 551 known fossils added according to Harms and Dunlop (2017), to Schwarze et al.
 552 (2021), and to Geißler et al. (2022). The record of *Ajkagarypinus* is indicated by a red
 553 star. Vertical colour bands reflect the fossiliferous localities containing fossil Pseudoscorpions, blue
 554 (sedimentary): Devonian Gilboa mudstones, Triassic Ukrainian claystone; yellow (amber):
 555 Cretaceous Myanmar amber, Eocene succinite (Bitterfeld, Baltic amber), Miocene Mexican and
 556 Dominican Republic amber. (COLOURED PICTURE)

557 Figure 2. A: Locality map of Ajka. B: Location of Ajka-Csinger valley and the Ajka Coal area.
 558 C: Simplified geological section of the Ajka Subbasin (modified after Szabó et al., 2022c)
 559 (COLOURED PICTURE)

560 Figure 3. *Ajkagarypinus stephani* gen. et sp. nov. holotype (NHMUS PAL 2022.91.1) images.
 561 A: the Ajkaite piece before polishing (Leica DMC4500; scale bar: 20mm); B: the Ajkaite
 562 piece with the holotype in itself, after polishing, positioned by a plasticine piece (Nikon
 563 SMZ25; scale bar: 1mm); C: the holotype from lateral view (Nikon SMZ25; scale bar:
 564 0.5mm); D: holotype, dorsal view (Leica DMC4500; scale bar: 0.5mm); E: dorsal view
 565 (Leica DMC4500; scale bar: 0.25mm); F: dorsal view (Leica DMC4500; scale bar: 0.1mm).
 566 (COLOURED PICTURE)


567 Figure 4. *Ajkagarypinus stephani* gen. et sp. nov. holotype (NHMUS PAL 2022.91.1),
568 screenshots of the synchrotron model (scalebar: 0.5mm). A: dorsal view; B: ventral view; C:
569 frontal view; D: right lateral view; E: left lateral view; G: ventrolateral view.

570 Figure 5. *Ajkagarypinus stephani* gen. et sp. nov. holotype (NHMUS PAL 2022.91.1). A: body,
571 dorsal view (scale bar: 0.5mm); B: right chelicera, dorsal view (scale bar: 0.1mm); C: left
572 pedipalp, dorsal view (scale bar: 0.25mm); D: right chela, lateral view (scale bar: 0.25mm).


573


574


Journal Pre-proof




Journal Pre-proof









Ajkagarypinus stephani gen. et sp. nov. from Santonian is described

Ajkagarypinus stephani is the second fossil pseudoscorpion from the Carpathian Basin

Ajkagarypinus stephani is the second species of Garypinidae from the Mesozoic

This fossil greatly contributes to Mesozoic pseudoscorpion biogeography

Journal Pre-proof

Author contribution

JN conducted the taxonomical investigation of the specimen and elaborated the first version of the manuscript. MH provided advice regarding the accurate taxonomic placement of the new taxon. AÓ and MS elaborated the first draft of geological and palaeoecological parts of the manuscript. JH, DH and UK organized to the synchrotron model with support of a student contributor. MB conducted the polishing of the amber piece. CH, MB, and JN captured photographs on the specimen. AÓ, MS, UK, and DH provided insight to the paleontological dimensions of the finding. All authors read, finetuned and approved the final version of the manuscript.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof