



HAL
open science

High resolution spectra of CO₂ ice based on SPICAM/MEX observations

Alexander Lomakin, Anna Fedorova, Oleg Korablev, Franck Montmessin,
Gaetan Lacombe, Franck Lefèvre

► **To cite this version:**

Alexander Lomakin, Anna Fedorova, Oleg Korablev, Franck Montmessin, Gaetan Lacombe, et al..
High resolution spectra of CO₂ ice based on SPICAM/MEX observations. European Planetary Science
Congress 2018, Sep 2018, Berlin, Germany. pp.EPSC2018-552-2. insu-04392878

HAL Id: insu-04392878

<https://insu.hal.science/insu-04392878>

Submitted on 14 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High resolution spectra of CO₂ ice based on SPICAM/MEX observations

Alexander Lomakin (1,2), Anna Fedorova (1), Oleg Korabev (1), Franck Montmessin (3), Gaetan Lacombe (3), Franck Lefèvre (3)

(1) Space Research Institute of the Russian Academy of Sciences, Moscow, Russia, (2) Moscow Institute of Physics and Technology (State University), Dolgoprudniy, Russia, (3) CNRS LATMOS, Université Versailles St Quentin en Yvelines, Quartier des Garennes, 11 bd d'Alembert, 78280 Guyancourt, France (aleksander.lomakin96@gmail.com)

1. Introduction

CO₂ cycle on Mars defines fundamental processes both on surface and in atmosphere. On poles condensation of a large part of the atmosphere (up to 30%) results seasonal growth and retreat of polar caps [1], changing reflectance and emissivity of surface, that has dramatic consequences for energy budget [2] and changes local and global climate on the planet [3]. IR spectroscopy is one of the ways to measure abundance of carbon dioxide ice on Mars. CO₂ ice has narrow features in NIR that allows to distinguish it from gaseous absorption in atmosphere. These features allowed to map distribution and seasonal evolution of condensation and sublimation of CO₂ ice on polar caps [4]. Meanwhile, spectrum of carbon dioxide ice in NIR still hasn't been studied in details yet, because low resolution spectrometers cannot resolve weak sharp lines.

2. Observations

Since 2004 SPICAM IR spectrometer on Mars-Express has been carrying out measurements of the Martian atmosphere and surface in near-infrared range from 1 to 1.7 μm with the spectral power about 2000. The observations collected information about ices distribution on the surface for Martian years from MY27 to MY34. The analysis of SPICAM dataset related to surface features had not been made before. To process surface spectra of SPICAM and clean them from atmospheric signatures algorithm was developed based on the latest spectroscopic databases HITRAN2016 and Martian general circulation model v5.3. Here we present accurate high resolution spectra of Martian poles (figure 1) and seasonal maps for multi-annual distribution of carbon dioxide ice on surface of Mars.




Figure 1: Comparison of SPICAM data and laboratory measurements by Hansen et al. (2005)

Acknowledgements

SPICAM operations are funded by CNES and Roscosmos. The data analysis by A.L., A.F. and O.K. has been supported by the Russian Government grant 14.W03.31.0017.

References

- [1] Piqueux S., Byrne S., Kieffer H., Titus T., Hansen C., *Icarus*, Volume 251, 2015, Pages 332-338, <https://doi.org/10.1016/j.icarus.2014.12.014>.
- [2] Wood S.E., David A.Paige, *Icarus* (1992), Volume 99, Issue 1, p. 15-27; doi: 10.1016/0019-1035(92)90167-6.
- [3] Hourdin et al., Workshop on Atmospheric Transport on Mars p 16-17
- [4] Langevin, Y., et al. (2006), *Nature* 442(7104): 790-792.