



HAL
open science

First Ion Insights during Titan's Ionosphere Relevant Gas Mixture EUV Irradiation

Jérémy Bourgalais, Ludovic Vettier, Sarah Tigrine, Guy Cernogora, Nathalie Carrasco

► **To cite this version:**

Jérémy Bourgalais, Ludovic Vettier, Sarah Tigrine, Guy Cernogora, Nathalie Carrasco. First Ion Insights during Titan's Ionosphere Relevant Gas Mixture EUV Irradiation. European Planetary Science Congress 2018, Sep 2018, Berlin, Germany. insu-04398668

HAL Id: insu-04398668

<https://insu.hal.science/insu-04398668>

Submitted on 16 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First Ion Insights during Titan's Ionosphere Relevant Gas Mixture EUV Irradiation

Jérémy Bourgalais, Ludovic Vettier, Sarah Tigrine, Guy Cernogora and Nathalie Carrasco

Université Versailles St-Quentin, Sorbonne Université, UPMC Univ. Paris 06, CNRS/INSU, LATMOS-IPSL, 11 boulevard d'Alembert, 78280 Guyancourt, France (jeremy.bourgalais@latmos.ipsl.fr)

Abstract

In Titan's ionosphere, the chemical growth in the atmosphere is believed to occur through the chemical coupling of radicals and reactive charged species but the complexity of the chemistry is far from being understood. Here we present a series of EUV Titan's atmosphere simulation experiments. We irradiated at 73.6nm different gas mixtures (N_2/CH_4 & He/CH_4) in a photochemical reactor. Our work is focused for the first time on photochemical ionic products.

1. Introduction

The arrival of the Cassini-Huygens spacecraft led to the discovery of an unexpected efficient organic chemistry, including very heavy ionic species [1,2] in the Titan's thermosphere. It showed that large condensable molecules are formed in the upper region of the moon where the flux of photons partly initiate chemical reactions. The photochemistry of CH_4 is one of the dominant chemical processes, but the large amount of N_2 is believed to significantly change the subsequent chemistry. As a matter of fact, the instruments onboard to the Cassini spacecraft showed that nitrogen participates in the chemistry and Titan's aerosols contain significant amounts of nitrogen [3,4]. In addition, Titan's atmosphere simulation experiments [5-7] have also demonstrated that the presence of nitrogen increases the molecular complexity. However, even if the chemical growth in the atmosphere is believed to occur through the chemical coupling of radicals and reactive charged species (ionic species and electrons) the complexity of the chemistry is far from being understood. Indeed, the current photochemical models [8] reasonably explain the formation of small carbon/nitrogen bearing molecules, but fail at giving accurate processes for the formation of heavy C and/or N-based species observed (>4 carbon and/or nitrogen atoms). It is mainly due to the complexity of reaction pathways and the limited set of available

data (branching ratios and coefficient rates) for chemical reactions.

2. Experimental Approach

Numerous laboratory studies investigated the gas phase products in the atmosphere of Titan through irradiation of N_2/CH_4 mixtures by using a variety of energy sources (*e.g.*, laser induced plasma, electrical discharge, UV lamps or synchrotron). However, those sources, despite synchrotron beamline, allow to work only above 110nm. Indeed, it requires EUV wavelengths (<110 nm) in order to address the effect of photoionisation of nitrogen as it might plays a key role in the upper part of the atmosphere of Titan regarding the ionization cross section at these wavelengths and the abundance of the molecule. There is only one EUV synchrotron radiation study on N_2/CH_4 gas mixture [7,9,10]. They have shown that the presence of the dominant nitrogen species in the gas mixture (*ca.* 95%) enhances the formation of unsaturated hydrocarbons (*e.g.*, benzene, toluene) at wavelength below 80nm. The production is believed to initially come from a dissociative charge-transfer reaction between N_2^+ and methane, leading to the formation of unsaturated complex hydrocarbons through production of $C_2H_5^+$ with subsequent dissociative recombination. However, they were not able to look at ions to validate this mechanism assumption.

Our work aims at corroborating the major role, highlighted by Imanaka *et al.*, played by the photoionization of nitrogen in the formation of complex organic molecules by monitoring simultaneously neutral and ion EUV photoproducts.

3. Results

Accordingly, here we present a series of EUV Titan's atmosphere simulation experiments. We irradiated at 73.6nm different gas mixtures (N_2/CH_4 & He/CH_4) in

the APSIS (Atmospheric Photochemistry Simulated by Synchrotron) reactor recently developed in the LATMOS laboratory [11]. The wavelength is obtained by using a surfatron-type discharge with a neon gas flow in the mbar pressure range coupled to the photochemical APSIS reactor. This experimental setup designed to carry research on planetary atmospheres allow *in situ* mass spectrometry analysis of the neutral and ionic species produced in the APSIS reactor.

The first experiment analysis for the photolysis of only methane seems to confirm the assumption of Imanaka *et al.* with CH_5^+ and C_2H_5^+ as the main ion species. However, in the case of a coupled N_2/CH_4 system, CH_3^+ is not the main contributor for the growth of the unsaturated species. The formation seems to be driven by nitrogen species such as N_2^+ , HCNH^+ and NH_4^+ . As shown in Figure 1 illustrating a mass spectrum of the positive ions detected in the photochemical reactor obtained from a N_2/CH_4 gas mixture irradiated 73.6nm. We also notice an efficient ion production up to m/z 150.




Figure 1: Positive ions mass spectrum obtained from a N_2/CH_4 gas mixture irradiated 73.6nm

This work constraints the detailed chemical mechanisms by bringing for the first time information about ionic and neutral species formed in nitrogen-dominated atmospheres by EUV irradiation. This work is important to reduce the sources of uncertainty and/or bias in the model predictions of Titan's ionosphere.

Acknowledgements

This research is supported by the ERC Starting Grant PRIMCHEM, grant agreement n°636829.

References

- [1] Coates, A. J., Crary, F. J., Lewis, G. R., Young, D. T., Waite, J. H., & Sittler, E. C. (2007). *Geophysical Research Letters*, 34(22).
- [2] Crary, F. J., Magee, B. A., Mandt, K., Waite Jr, J. H., Westlake, J., & Young, D. T. (2009). *Planetary and Space Science*, 57(14-15), 1847-1856.
- [3] Israël, G., Szopa, C., Raulin, F., Cabane, M., Niemann, H. B., Atreya, S. K., et al. (2005). *Nature*, 438(7069), 796.
- [4] Vuitton, V., Yelle, R. V., & McEwan, M. J. (2007). *Icarus*, 191(2), 722-742.
- [5] Sebree, J. A., Trainer, M. G., Loeffler, M. J., & Anderson, C. M. (2014). *Icarus*, 236, 146-152.
- [6] Trainer, M. G., Sebree, J. A., Yoon, Y. H., & Tolbert, M. A. (2013). *The Astrophysical Journal Letters*, 766(1), L4.
- [7] Imanaka, H., & Smith, M. A. (2009). *The Journal of Physical Chemistry A*, 113(42), 11187-11194.
- [8] Moses, J. I., Fouchet, T., Bézard, B., Gladstone, G. R., Lellouch, E., & Feuchtgruber, H. (2005). *Journal of Geophysical Research: Planets*, 110(E8).
- [9] Imanaka, H., & Smith, M. A. (2007). *Geophysical research letters*, 34(2).
- [10] Imanaka, H., & Smith, M. A. (2010). *Proceedings of the National Academy of Sciences*, 107(28), 12423-12428.
- [11] Tigrine, S., Carrasco, N., Vettier, L., & Cernogora, G. (2016). *Journal of Physics D: Applied Physics*, 49(39), 395202.