



**HAL**  
open science

## Earth and Venus exospheres as seen by SWAN/SOHO and SPICAV/VEX

Igor I. Baliukin, Jean-Loup Bertaux, Eric Quémerais, Jean-Yves Chaufray,  
Vladislav Izmodenov

► **To cite this version:**

Igor I. Baliukin, Jean-Loup Bertaux, Eric Quémerais, Jean-Yves Chaufray, Vladislav Izmodenov. Earth and Venus exospheres as seen by SWAN/SOHO and SPICAV/VEX. EPSC-DPS Joint Meeting 2019, Sep 2019, Geneva, Switzerland. pp.EPSC-DPS2019-463. insu-04413948

**HAL Id: insu-04413948**

**<https://insu.hal.science/insu-04413948v1>**

Submitted on 24 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Earth and Venus exospheres as seen by SWAN/SOHO and SPICAV/VEX

Igor Baliukin (1,2), Jean-Loup Bertaux (1,3), Eric Quémerais (3), Jean-Yves Chaufray (3) and Vladislav Izmodenov (1,2,4)  
(1) Space Research Institute, Russian Academy of Sciences, Moscow, Russia, (2) Lomonosov Moscow State University, Moscow, Russia, (3) LATMOS-UVSQ, Université Versailles Saint-Quentin, Guyancourt, France, (4) Institute for Problems in Mechanics, Moscow, Russia (igor.baliukin@gmail.com)

### Abstract

The Earth's hydrogen exosphere Lyman- $\alpha$  radiation was mapped with the SWAN/SOHO instrument in January 1996, 1997 and 1998 (low solar activity). The use of a hydrogen absorption cell allowed to disentangle the interplanetary emission from the geocoronal one and to assign the absorbed signal almost entirely to the geocorona. The geocorona was found to extend at least up to 100 Earth Radii ( $R_E$ ) with an intensity of 5 Rayleigh, an unprecedented distance well exceeding the recent results of LAICA imager ( $\sim 50 R_E$ ), and encompassing the orbit of the Moon ( $\sim 60 R_E$ ). We developed a numerical kinetic model of the hydrogen atoms distribution in the exosphere which includes the solar Lyman- $\alpha$  radiation pressure and the ionization. The radiation pressure compresses the H exosphere on the day side, producing a bulge of H density between 3 and 20  $R_E$  which fits observed intensities very well. The SWAN Lyman- $\alpha$  distribution of intensity was compared both to LAICA (2015) and to OGO-5 (1968) measurements. An onion-peeling technique is used to retrieve hydrogen number density in the exosphere for the three SWAN observations. They show an excess of density versus models at large distances, which is likely due to non-thermal atoms (not in the model).

The SPICAV UV spectrometer was used to measure Lyman- $\alpha$  radiation from 2006 to 2014 in orbit around Venus as part of the Venus Express (VEX) mission. We have performed the analysis of the SPICAV data on observations of the Lyman- $\alpha$  radiation scattered on the "night" side of the Venusian exosphere. We have extended the work [1], where only 4 observations of the "night" side of Venusian hydrogen corona have been considered. In the framework of our study, we have found  $\sim 15$  "nightside" observations that have never been analyzed before. A qualitative and quantitative analysis of these data was carried out.

We compare both (Earth and Venus) hydrogen exospheres and discuss the main differences/similarities (extension, number density and temperature at the exobase level, existence of the non-thermal component).

### Acknowledgements

This work has been supported by the Ministry of Education and Science of Russian Federation grant 14.W03.31.001. We thank Rosine Lallement and Olga Katushkina for fruitful discussions and Stéphane Ferron for SWAN data management.

### References

- [1] Chaufray, J.-Y., Bertaux, J.-L., Quémerais, E., Leblanc, F., Sulis, S.: Observations of the nightside venusian hydrogen corona with SPICAV/VEX, *Icarus*, Vol. 262, pp. 1-8, 2015.