



HAL
open science

HCl in the atmosphere of Mars: chlorine isotopic ratio

Alexander V. Trokhimovskiy, Kevin Olsen, Oleg I. Korablev, Franck Montmessin, Franck Lefèvre, Anna A. Fedorova, Juan Alday, Denis A. Belyaev, Andrey Patrakeev, Alexey Shakun

► **To cite this version:**

Alexander V. Trokhimovskiy, Kevin Olsen, Oleg I. Korablev, Franck Montmessin, Franck Lefèvre, et al.. HCl in the atmosphere of Mars: chlorine isotopic ratio. Europlanet Science Congress 2020, Sep 2020, Online, Unknown Region. pp.EPSC2020-673, 10.5194/epsc2020-673 . insu-04421226

HAL Id: insu-04421226

<https://insu.hal.science/insu-04421226v1>

Submitted on 27 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Europlanet Science Congress 2020

Virtual meeting

21 September – 9 October 2020

[Home](#) > EPSC2020-673

EPSC Abstracts

Vol.14, EPSC2020-673, 2020

<https://doi.org/10.5194/epsc2020-673>

Europlanet Science Congress 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



HCl in the atmosphere of Mars: chlorine isotopic ratio

Alexander Trokhimovskiy et al.

The ExoMars Trace Gas Orbiter (TGO) mission had started regular measurements in 2018. Primary goal of the mission is to quantify trace gases that could indicate geologic or biogenic activity on Mars (Vago et al., 2015). Atmospheric Chemistry Suite mid-infrared channel (ACS MIR) is a high resolution cross-dispersion spectrometer operating in solar occultation mode (Korablev et al., 2018). It was designed to make the most sensitive measurements of the atmosphere to date. During each occultation up to 20 diffraction orders are simultaneously recorded at different tangent altitudes. In 2020 ACS MIR reports the discovery of the gaseous hydrogen chloride (HCl). Absorption features are present in several consecutive diffraction orders, with both isotopes H^{37}Cl and H^{35}Cl are clearly observed. HCl was observed by ACS simultaneously in both hemispheres after the main phase of the global dust storm. Though the formation mechanism is not fully clear, we believe that the presence of HCl is associated with the lifted dust and chlorine component in it.

On Earth, in general, the chlorine isotope variations in nature are relatively small, ranging from ~ -2 to $+2$ ‰. However, large variations are observed, e.g. in extraterrestrial materials and volcanic gases, due to kinetic fractionation. On Mars Farley et al. (2016) reported a range from -1 ‰ to -51 ‰ (5% reduction) for the $\delta^{37}\text{Cl}$ in the samples drilled in the Gale Crater. ACS observations demonstrate enrichment of the ^{37}Cl up to $+250$ ‰ on average in the atmospheric gaseous. In principle, most atmospheric elements on Mars have heavy isotope enrichments due to preferential loss of the light isotope to space (e.g. Vandaele et al., 2019). Early hydrodynamic escape during intense extreme ultraviolet radiation followed by prolonged atmospheric 'erosion' explains the heavy isotope enrichment. Chlorine loss, as HCl, would raise the $\delta^{37}\text{Cl}$ value of the residual materials, involved in the dust-atmospheric exchange cycle.

References

Farley K.A., Martina P., Archer P.D., et al.: Light and variable $^{37}\text{Cl}/^{35}\text{Cl}$ ratios in rocks from Gale Crater, Mars: Possible signature of perchlorate, *Earth and Planetary Science Letters* 438:14-24, DOI: 10.1016/j.epsl.2015.12.013, 2016.

Korablev, O., Montmessin, F., Trokhimovskiy, et al.: The Atmospheric Chemistry Suite (ACS) of Three Spectrometers for the ExoMars 2016 Trace Gas Orbiter, *Space. Sci. Rev.*, 214(1), 7, doi:10.1007/s11214-017-0437-6, 2018.

Vago, J., Witasse, O., Svedhem, et al.: ESA ExoMars program: The next step in exploring Mars, *Sol. Syst. Res.*, 49(7), 518–528, doi:10.1134/S0038094615070199, 2015.

Vandaele, A. C., Korablev, O., Daerden, F. et al.: Martian dust storm impact on atmospheric H_2O and D/H observed by ExoMars Trace Gas Orbiter, *Nature*, 568, 521–525, doi:10.1038/s41586-019-1097-3, 2019.

How to cite: Trokhimovskiy, A., Olsen, K., Korablev, O., Montmessin, F., Lefevre, F., Fedorova, A., Alday Parejo, J., Beliaev, D., Patrakeev, A., and Shakun, A.: HCl in the atmosphere of Mars: chlorine isotopic ratio, Europlanet Science Congress 2020, online, 21 September–9 Oct 2020, EPSC2020-673, <https://doi.org/10.5194/epsc2020-673>, 2020



Contact

[Imprint](#)

[Contact](#)



