

## Ma MISS on ExoMars 2020

Maria Cristina de Sanctis, Francesca Altieri, Eleonora Ammannito, Simone de Angelis, Marco Ferrari, Alessandro Frigeri, David Biondi, R. Mugnuolo, S Pirrotta, T. Di Iorio, et al.

## ▶ To cite this version:

Maria Cristina de Sanctis, Francesca Altieri, Eleonora Ammannito, Simone de Angelis, Marco Ferrari, et al.. Ma\_MISS on ExoMars 2020. Ninth International Conference on Mars, Jul 2019, Pasadena, United States. pp.LPI Contribution No. 2089, id.6016. insu-04421457

## HAL Id: insu-04421457 https://insu.hal.science/insu-04421457

Submitted on 27 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ma\_MISS on EXOMARS 2020. M.C. De Sanctis (1), F. Altieri (1), E. Ammannito (2), S. De Angelis (1), M. Ferrari (1), A. Frigeri (1), D. Biondi (1), R. Mugnuolo (2), S. Pirrotta (2), T. Di Iorio (3), F. Capaccioni (1), M.T. Capria (1), V. Ciarletti (4), B. Ehlmann (5), C. Federico (1), G. Magni (1), M. Lavagna (6), M. Formisano (1), S. Fonte (1), M. Giardino (2), G. Piccioni (1), F. Westall (7) and the Ma\_MISS team. (1) Institute for Space Astrophysics and Planetology, IAPS-INAF, Rome, Italy (mariacristina.desanctis@inaf.it);(2) Italian Space Agency, ASI, Italy; (3) ENEA, Italy;(4)LATMOS, France; (5) Calthec, USA; (6) POLI-MI, Italy,(7) CNRS Orleans, France

The ExoMars 2020 launch window opens on 26 July 2020, and ExoMars will touchdown at Oxia Planum, a location with a strong potential for past habitability and for preserving physical and chemical biosignatures. The mission will deliver two science elements to the martian surface: (1) a lander instrumented to conduct environmental and geophysical measurements and (2) a rover with scientific payload to conduct a search for signs of life.

Ma\_MISS is a visible and near infrared (VNIR,  $0.4-2.2~\mu m$ ) micro spectrometer hosted by the drill system of the ExoMars 2020 rover [1]. The Ma\_MISS instrument has been developed to provide hyperspectral images of boreholes excavated by the ExoMars rover drill. Ma\_MISS will characterize the mineralogy and stratigraphy of the shallow subsurface down to two meters [2]. The main objectives of Ma\_MISS are: (1) determine the composition of the subsurface materials; (2) map the distribution of the subsurface H<sub>2</sub>O-bearing and OH-bearing materials and possibly ices; (3) characterize important optical and physical properties of the materials (e.g., grain size); (4) produce a stratigraphic column that will provide information on the subsurface geology.

The drill can reach down to 2 m below the surface, and Ma\_MISS will operate periodically during pauses in drilling activity and will produce hyperspectral images of the drill's borehole. The Ma\_MISS instrument's main driving requirement was its miniaturization because it is embedded within the drill (Fig. 1). The spectrometer is placed in a box on the side wall of the drill box. The spectral range is  $0.4-2.2~\mu m$ , with a spectral sampling of 20 nm a SNR~100 and a spatial resolution of 120  $\mu m$ . The light from a 5W lamp is collected and carried, through an optical fiber bundle, to the miniaturized Optical Head, hosted within the drill tip. A Sapphire Window with high hardness and transparency on the drill tip protects the Ma\_MISS optical head allowing to observe the borehole wall.

Different depths can be reached by the use of three extension rods, 50 cm long, each containing optical fibers and a collimator. The first extension rod is connected to the nonrotating part of the Drill, hosted on the rover, through a Fiber Optical Rotating Joint

(FORJ), that allows the continuity of the signal link between the rotating part of the drill and the spetrometer




Fig. 1: Artistic view of the ExoMars-Pasteur Rover with instruments allocations. Ma\_MISS is integrated within the drill.

The Ma\_MISS instrument has been calibrated and delivered and is now integrated into the drill.

Results obtained in the lab on mineral/rock samples confirm that the Ma\_MISS spectrometer has a spectral range, resolution and imaging capabilities suitable for the Mars subsurface characterization[2,3]. The spectra acquired with the Ma\_MISS fine spatial resolution ( 120 µm) show minerals that are not recognizable at coarser resolution (~6 mm ) (Fig.2). In the case reported in Fig. 2, a lava from *Montiferru/Bonarcado*, Ma\_MISS is able to identify the a great diversity, indicating several different mineralogical phases within the observed area. Phyllosilicate, olivine, opaque phase can be identified with Ma\_MISS but not at coarser resolution.




Fig.2 <u>Left Top</u>: slab of Montiferru/Bonarcado Lava. The letters (A,B,C,D) correspond to the areas analyzed with Spectro-goniometer setup. Each area is 6-mm-size; <u>Left bottom</u>: spectra acquired from the regions A,B,C,D, with the spectro-photometer FieldSpec Pro coupled with a mechanical goniometer, having a spatial resolution of about 6 mm. <u>Right</u>: Spectra acquired with Ma\_Miss BreadBoard setup. In each single 6-mm-sized area, spectra in different positions have been acquired with Ma\_Miss BB setup; Data at 1 µm are not shown in several spectra due to high level of detector noise, as well as data <0.5 µm.

Ma\_MISS is the only instrument in the rover's Pasteur payload able to analyze subsurface material in its natural condition (*in situ*), prior to extracting samples for further analysis. In synergy with other Rover instruments, MA\_MISS findings will help to refine criteria for deciding form where to collect samples.

## References:

[1]Vago J.L. et al. (2017): Astrobiology, 17, 6, 7. [2]De Sanctis et al. (2017): Astrobiology, 17, 6, 7. [3]De Angelis et al. (2017): PSS, 144, DOI: 10.1016/j.pss.2017.06.005