

# Constraining Ganymede's exosphere through numerical simulations of its ionosphere and Galileo observations

Gianluca Carnielli, Marina Galand, Ronan Modolo, François Leblanc, Arnaud Beth, Hans L. F. Huybrighs, Xianzhe Jia

### ► To cite this version:

Gianluca Carnielli, Marina Galand, Ronan Modolo, François Leblanc, Arnaud Beth, et al.. Constraining Ganymede's exosphere through numerical simulations of its ionosphere and Galileo observations. European Planetary Science Congress 2018, Apr 2018, Berlin, Germany. pp.EPSC2018-350. insu-04422373

## HAL Id: insu-04422373 https://insu.hal.science/insu-04422373

Submitted on 28 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. EPSC Abstracts Vol. 12, EPSC2018-350, 2018 European Planetary Science Congress 2018 © Author(s) 2018



# Constraining Ganymede's exosphere through numerical simulations of its ionosphere and Galileo observations

Gianluca Carnielli (1), Marina Galand (1), Ronan Modolo (2), François Leblanc (3), Arnaud Beth (1), Hans L. F. Huybrighs (4,5,6), Xianzhe Jia (7)

(1) Department of Physics, Prince Consort Road, Imperial College London, London SW7 2AZ, UK

(gianluca.carnielli10@imperial.ac.uk), (2) LATMOS/IPSL, UVSQ Université Paris-Saclay, UPMC Univ. Paris 06, Guyancourt, France, (3) LATMOS/IPSL, CNRS, Sorbonne Université, UVSQ, Paris, France (4) Max-Planck-Institut für Sonnensystemforschung, Göttingen, Germany, (5) Swedish Institute of Space Physics, Post Office Box 812, Kiruna, Sweden, (6) Institut für Geophysik und extraterrestrische Physik, Technische Universität, Germany, (7) Department of Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI 48109-2143, USA

#### Abstract

Our current understanding of Ganymede's neutral environment derives from models which are driven by a limited set of observations. These include remote sensing UV measurements from close flybys by the Galileo and Voyager spacecraft and from Earth by the Hubble Space Telescope (HST).

Estimated parameters from observations include the column density of H [1,2] and  $O_2$  [2,3]. For  $O_2$ , we argue that the derivation of the column density was based on strong assumptions on the plasma environment, which is poorly characterized. These estimates have been used as reference values in exospheric models (e.g. [4,5,6]), but cannot be confirmed without additional in situ measurements, which will be made by the JUICE spacecraft during the late phase of the mission.

One way to check the consistency between the estimated column density of the neutral species and Galileo observations, is to develop a model of Ganymede's ionosphere, starting from an exosphere which reproduces the estimated column densities, and check that the plasma properties derived by the model along the flyby trajectories correspond to those measured by Galileo.

We have developed such a model for Ganymede [7] using the exospheric configuration from Leblanc et al. (2018) [4], which reproduces the  $O_2$  column density estimated by Hall et al. (1998) [3]. For the plasma number density along the Galileo flyby trajectories, the model underestimates by more than one order of magnitude the observations. We attribute the cause of this discrepancy to the assumed configuration of the neutral exosphere, which we argue to be underesti-

#### mated.

Here, we present a new possible configuration of the  $O_2$  exosphere which leads to a plasma density in the ionospheric model that is consistent with that measured by Galileo along the close flyby trajectories. To reach agreement, we had to increase the column density of  $O_2$  from the original estimate of Hall et al. (1998) [3].

We conclude that overall Ganymede's O<sub>2</sub> exosphere should be denser than what has been assumed so far.

#### Acknowledgements

We warmly acknowledge the support of the IPSL data centre CICLAD for providing access to their computing resources. Work at Imperial College London was supported by STFC of UK through a postgraduate studentship and under grants ST/K001051/1 and ST/N000692/1.

#### References

- Barth, C.A., Hord, C.W., Stewart, A.I.F., Pryor, W.R., Simmons K.E., McClintock, W.E., Ajello, J.M., Naviaux, K.L. and Aiello, J.J.: Galileo ultraviolet spectrometer observations of atomic hydrogen in the atmosphere at Ganymede, Geophysical Research Letters, Vol. 24, p. 2147, 1997.
- [2] Feldman, P.D., McGrath, M.A., Strobel, D.F., Moos, H.W., Retherford, K.D. and Wolven, B.C.: HST/STIS Ultraviolet Imaging of Polar Aurora on Ganymede, The Astrophysical Journal, Vol. 535, pp. 1085-1090, 2000
- [3] Hall, D.T., Feldman, P.D., McGrath, M.A. and Strobel, D.F.: The Far-Ultraviolet Oxygen Airglow of Europa and

Ganymede, The Astrophysical Journal, Vol. 499, pp. 475-481, 1998

- [4] Leblanc, F., Oza, A.V., Leclercq, L., Schmidt, C., Cassidy, T., Modolo, R., Chaufray, J.Y. and Johnson, R.E.: On the orbital variability of Ganymede's atmosphere, Icarus, Vol. 293, pp. 185-198, 2018
- [5] Marconi, M.L.: A kinetic model of Ganymede's atmosphere, Icarus, Vol. 190, pp. 155-174, 2007
- [6] Turc, L., Leclercq, L., Leblanc, F., Modolo, R. and Chaufray, J.Y.: Modelling Ganymede's neutral environment: A 3D test-particle simulation, Icarus, Vol. 229, pp. 157-169, 2014
- [7] Carnielli, G., Galand, M., Leblanc, F., Leclercq, L., Modolo, R., Beth, A., Huybrighs, H.L.F., Jia, X.: First 3D test particle model of Ganymede's ionosphere, Icarus, under review