



**HAL**  
open science

## High CO<sub>2</sub> in the mantle source of ocean island basanites

Francesco Maria Lo Forte, Federica Schiavi, Estelle F. Rose-Koga, Silvio G Rotolo, Maximilien Verdier-Paoletti, Alessandro Aiuppa, Vittorio Zanon

### ► To cite this version:

Francesco Maria Lo Forte, Federica Schiavi, Estelle F. Rose-Koga, Silvio G Rotolo, Maximilien Verdier-Paoletti, et al.. High CO<sub>2</sub> in the mantle source of ocean island basanites. *Geochimica et Cosmochimica Acta*, 2024, 10.1016/j.gca.2024.01.016 . insu-04431773

**HAL Id: insu-04431773**

**<https://insu.hal.science/insu-04431773v1>**

Submitted on 1 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## Journal Pre-proofs

High CO<sub>2</sub> in the mantle source of ocean island basanites

Francesco Maria Lo Forte, Federica Schiavi, Estelle F. Rose-Koga, Silvio G. Rotolo, Maximilien Verdier-Paoletti, Alessandro Aiuppa, Vittorio Zanon

PII: S0016-7037(24)00039-5  
DOI: <https://doi.org/10.1016/j.gca.2024.01.016>  
Reference: GCA 13298

To appear in: *Geochimica et Cosmochimica Acta*

Received Date: 28 August 2023  
Accepted Date: 17 January 2024

Please cite this article as: Maria Lo Forte, F., Schiavi, F., Rose-Koga, E.F., Rotolo, S.G., Verdier-Paoletti, M., Aiuppa, A., Zanon, V., High CO<sub>2</sub> in the mantle source of ocean island basanites, *Geochimica et Cosmochimica Acta* (2024), doi: <https://doi.org/10.1016/j.gca.2024.01.016>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 The Author(s). Published by Elsevier Ltd.



## High CO<sub>2</sub> in the mantle source of ocean island basanites

Francesco Maria Lo Forte <sup>a</sup>, Federica Schiavi <sup>b</sup>, Estelle F. Rose-Koga <sup>c</sup>, Silvio G. Rotolo <sup>a,d</sup>, Maximilien Verdier-Paoletti <sup>e</sup>, Alessandro Aiuppa <sup>a</sup>, and Vittorio Zanon <sup>f</sup>

<sup>a</sup> Dipartimento di Scienze della Terra e del Mare, Università degli Studi di Palermo, Palermo, 90123, Italy.

<sup>b</sup> Université Clermont Auvergne, CNRS, IRD, OPGC, Laboratoire Magmas et Volcans, F-63000 Clermont-Ferrand, France

<sup>c</sup> Institut des Sciences de la Terre d'Orléans (ISTO) UMR 7327, CNRS-Université d'Orléans-BRGM, 1A rue de la Férollerie, 45071 Orléans, Cedex 2, France.

<sup>d</sup> Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Palermo, Via Ugo La Malfa 153, 90146, Palermo, Italy.

<sup>e</sup> Muséum National d'Histoire Naturelle de Paris (MNHN), Institut de Minéralogie, de Physique des Matériaux, 61 rue Buffon CP52 75005 Paris, France

<sup>f</sup> Instituto de Investigação em Vulcanologia e Avaliação de Riscos, Universidade dos Açores, Rua Mãe de Deus, Ponta Delgada, 9501-801, Portugal.

**corresponding author:** Francesco Maria Lo Forte

**corresponding author mail address:** [francescomaria.loforte@unipa.it](mailto:francescomaria.loforte@unipa.it)

### Abstract

Some of the most CO<sub>2</sub>-rich magmas on Earth are erupted by intraplate ocean island volcanoes. Here, we characterise olivine-hosted melt inclusions from recent (< 10 ky) basanitic tephra erupted by Fogo, the only active volcano of the Cape Verde Archipelago in the eastern Atlantic Ocean. We determine H<sub>2</sub>O, S, Cl, F in glassy melt inclusions and recalculate the total (glass+shrinkage bubbles) CO<sub>2</sub> budget by three independent methodologies. We find that the Fogo parental basanite, entrapped as melt inclusion in forsterite-rich (Fo<sub>80-85</sub>) olivines, contains up to ~2.1 wt% CO<sub>2</sub>, 3-47% of which is partitioned in the shrinkage bubbles. This CO<sub>2</sub> content is among the highest ever measured in melt inclusions in OIBs. In combination with ~2 wt.% H<sub>2</sub>O content, our data constrain an entrapment pressure range for the most CO<sub>2</sub>-rich melt inclusion of 648-1430 MPa, with a most

conservative estimate at 773-1020 MPa. Our results therefore suggest the parental Fogo melt is stored in the lithospheric mantle at minimum depths of ~27 to ~36 km, and then injected into a vertically stacked magma ponding system. Overall, our results corroborate previous indications for a CO<sub>2</sub>-rich nature of alkaline ocean island volcanism. We propose that the Fogo basanitic melt forms by low degrees of melting (F=0.06-0.07) of a carbon-enriched mantle source, containing up to 355-414 ppm C. If global OIB melts are dominantly as carbon-rich as our Fogo results suggest, then OIB volcanism may cumulatively outgas as high as ~16-21 Tg of carbon yearly, hence substantially contributing to the global deep carbon cycle.

**Keywords:** alkaline ocean islands, intraplate volcanism, Fogo volcano, melt inclusions, Cape Verde.

## 1. Introduction

Quantifying the initial amount of volatile elements (H<sub>2</sub>O, CO<sub>2</sub>, S, Cl, and F) transported by OIB magmas is essential for understanding the cycling of volatiles from Earth's deep interior (e.g., Hirschmann, 2006, 2018; Dasgupta and Hirschmann, 2010), and their role for mantle melting, and magma generation and extraction (e.g., Foley, 2011; Dasgupta, 2018). Although sustaining only ~10% of the annual magma output on Earth, OIB magmas can contain more than 5% volatiles, and therefore significantly contribute to the annual magmatic gas output (e.g., Cabral et al., 2014; Longpré et al., 2017; Boudoire et al., 2018; Taracsák et al., 2019; Burton et al., 2023), especially if compared to more voluminous - but less volatile rich - Mid Ocean Ridge Basalts (MORBs) (e.g., Hauri et al., 2019). Oceanic intra-plate magmas have some of the most CO<sub>2</sub>-enriched magmas on Earth (Longpré et al., 2017; Taracsák et al., 2019), likely reflecting their derivation from low melting degree of a metasomatised (C-enriched) mantle source (Foley et al., 2009, 2011; Dasgupta et al., 2013; Foley and Fischer, 2017; Dasgupta, 2018; Aiuppa et al., 2021).

Glassy melt inclusions (MI), small droplets of silicate melt entrapped in crystals during their growth in magmas or during episodes of crystal fracturing, are - at least in principle - physically and chemically isolated systems shielded from the surrounding environment by the host crystal (e.g., Lowenstern 1995; Danyushevsky et al. 2002; Schiano 2003; Wallace 2005; Cannatelli et al. 2016; Wallace et al., 2021; Rose-Koga et al., 2021). As such, the entrapped melt should escape extensive volatile loss upon magma ascent/storage in the crust prior/during eruption. Upon quenching as glass, MI should allow recording the dissolved volatile contents at the P-T-X conditions of entrapment, and eventually the parental melt volatile contents if entrapment has occurred at/near source (and before volatile saturation is reached) (e.g., Métrich and Wallace, 2008; Edmonds and Wallace, 2017). However, the task is complicated by a range of processes that can potentially modify MI composition after it is entrapped in the crystal (e.g., Anderson 1974; Roedder 1979; Danyushevsky et al., 2002; Wallace et al., 2021), including post-entrapment crystallization at the MI-host crystal interface (e.g., Danyushevsky et al., 2000), diffusive exchanges of major, trace and volatile (H<sup>+</sup>) elements (e.g., Gaetani and Watson, 2000, 2002; Danyushevsky, 2001; Cottrell et al., 2002; Gaetani et al. 2012) and - above all - shrinkage bubble formation (Anderson, 1974; Roedder, 1979; MacLennan, 2017). This latter process is especially relevant for volatile species, as it leads to partitioning of poorly soluble CO<sub>2</sub> from the melt inclusion glass into the shrinkage bubble. Shrinkage bubbles can contain up to >90% of the bulk MI CO<sub>2</sub> content (e.g., Mironov and Portnyagin, 2011; Hartley et al., 2014; Mironov et al., 2015; Moore et al., 2015; Esposito et al., 2016; Aster et al., 2016; Rasmussen et al., 2018, 2020), demonstrating the caveats of traditional approaches (Lowenstern, 2015) that focus on measuring volatiles in glass only. Still, accurately quantifying the shrinkage bubble CO<sub>2</sub> content is technically challenging (e.g., Naumov et al., 2003),


and few examples are available of reconstructions of the total (bubble+glass) MI CO<sub>2</sub> budget in OIB (e.g., Taracsák et al., 2019; Tucker et al., 2019, Burton et al., 2023).

Some of the most alkali-rich OIB magmas are erupted in the Cape Verde Archipelago, in the central eastern Atlantic. This archipelago lies on the largest intraplate bathymetric anomaly in the Earth's oceans – the Cape Verde Rise - interpreted as the surface expression of a hotspot sustained by the ascent of a deeply-rooted mantle plume (Holm et al., 2006; French and Romanowicz, 2015). The presence of highly alkaline silica undersaturated rocks (nephelinites, melanephelinites, and melilitites), as well as carbonatites (Hoernle et al., 2002; Bonadiman et al., 2005), are evidence for a heterogeneous, oxidized (Moussallam et al., 2019), potentially C-enriched mantle source, as also indicated by the CO<sub>2</sub>-rich signature of fumaroles (Aiuppa et al., 2020; Melián et al., 2021).

In this study we use primitive MI composition to confirm the carbon-rich signature of the Cape Verde magma source. We report on the major, trace, and volatile composition of olivine-hosted MI from basanitic tephra erupted at Fogo, the youngest active volcano of the Cape Verde Archipelago. Our goal is to determine the initial (parental melt) volatile contents, in the attempt to (i) test the carbon-enriched source hypothesis, (ii) refine our understanding of the architecture of the magma system, and (iii) more broadly, provide additional evidence for the carbon-enriched nature of alkali rich, OIB magmatism.

## 2. Volcanological Background

Fogo is the only active volcano in the Cape Verde Archipelago, in the central eastern Atlantic (Figure 1). Volcanic activity on the islands started ~26 Ma (Torres et al., 2010), and is dominated by silica-undersaturated alkaline melts (e.g., Gerlach et al., 1988), and rarer carbonatites (Hoernle et al., 2002; Mourão et al., 2012). Subaerial volcanic activity has been divided into four eruptive phases: (i) The Ribeira de Almada Group (~4.5 Ma, Foeken et al., 2009); (ii) The Monte Barro Group (of unknown age); (iii) The Monte Amarelo Group, whose activity preceded the lateral collapse of Monte Amarelo at circa 62 to 123 ka (Day et al., 1999; Foeken et al., 2009; Martínez-Moreno et al., 2018); and (iv) The Chã das Caldeira Group (Foeken et al., 2009) related to the recent growth of Pico do Fogo central volcano. The Pico do Fogo volcano erupted 28 times since the 1500s. Eruptions are sourced by a complex, vertically elongated magma plumbing system, with main magma storage/stagnation zones at ~24, ~13-19 and ~9-12 km depth (Hildner et al., 2011, 2012; Klügel et al., 2020; Lo Forte et al., 2023).



**Figure 1.** Modified Satellite map of Fogo Island from Google Earth, showing sampling sites with their relative pictures. Insets show the location of the Cape Verde archipelago (top right), and Fogo Island (bottom right).

### 3. Materials and Methods

#### 3.1. Sample preparation

We report data for 18 MI hosted in 12 olivine phenocrysts from three different tephra samples representative the last eruptive phase of Fogo volcano (see tables S1 and S2 of Supplementary Material S1, for further details). In detail, we analysed (i) 8 MI hosted in 5 olivines from the clinopyroxene- and olivine-rich tephra samples of São Jorge (SJ), of the early Holocene period of activity ( $\sim 10$  ka), and (ii) 10 MI hosted in 7 olivines from the recent basanitic tephra samples of the 1951 and 2014/15 eruptions. Samples were prepared according to the methodology described by Rose-Koga et al., (2021). Tephra samples were initially sieved to separate grain size populations in the 250–4000  $\mu\text{m}$  range. For each sample, about 400 olivine phenocrysts were hand-picked from these grain size fractions. Olivine crystals were mounted on glass slides using Crystal Bond 509®, subsequently they were doubly polished with silicon carbide paper and alumina powders to attain a maximum thickness of 80–100  $\mu\text{m}$ . Twenty phenocrysts, containing 26 MIs with size greater than 25  $\mu\text{m}$  were selected for analysis.

#### 3.2. Analytical techniques

Electron microprobe analysis (EMPA) was used to determine the major element compositions of both host olivines and melt inclusions, as well as Cl, and F concentrations in glasses. Volatile element concentrations ( $\text{H}_2\text{O}$ ,  $\text{CO}_2$ , S, and Cl) in glasses were measured by nano-SIMS. The  $\text{CO}_2$  concentration in shrinkage bubbles in MI was determined by Raman micro-spectroscopy. Finally, Laser Ablation-ICPMS was utilized for trace element analysis. The full compositional dataset is listed in tables S3, S4, S5, S5.1, and S5.2 of Supplementary Material S1.

### 3.2.1. Electron microprobe analysis

Electron microprobe analyses were performed with a Cameca SxFiveTactis with a LaB6 source equipped with five wavelength-dispersive spectrometers at the Laboratoire Magmas et Volcans (LMV), in Clermont-Ferrand, France. Major elements and volatiles were analysed in the same analytical session, with an accelerating voltage of 15 kV, an 8 nA beam current, and a 20  $\mu\text{m}$  defocused beam for major elements. Chlorine, sulphur, and fluorine analyses were conducted using an accelerating voltage of 15 kV, a 40 nA beam current and a 20  $\mu\text{m}$  defocused beam, following the acquisition program outlined in the Cameca Peak Sight software (see Rose-Koga et al., 2020 for more details). The olivines were analysed with an accelerating voltage of 15 kV and 15 nA beam current. The total concentration of volatiles was determined by summing the counts from multiple iterations. Total counting times were 100 s for F and Cl peaks. The detection limits for F and Cl were 300 and 150 ppm, respectively. KE12 LMV, CFA47 (Spilliaert et al., 2006), and ALV-98I R23 (Hekin and Walker, 1987) glasses and the San Carlos olivine were used as standards. Typical errors on the samples ( $1\sigma$  error of mean) are less than 10% for all elements analysed both in MI and in olivines.

### 3.2.2. Nano-SIMS analysis

High spatial resolution measurements of volatiles in minor and trace amounts in MI were conducted using the CAMECA nanoSIMS50 of the Muséum National d'Histoire Naturelle de Paris. All measurements were carried using a focussed  $\text{Cs}^+$  primary ion beam accelerated to 16 keV in high vacuum conditions ( $< 8 \times 10^{-10}$  mbar). Prior to acquisition, a  $5 \times 5 \mu\text{m}^2$  region was pre-sputtered with a 280 pA primary beam for 2 minutes to remove the Au coat and ensure a clean surface. The current was then reduced to 20 pA and the inner  $3 \times 3 \mu\text{m}^2$  were rastered over a  $32 \times 32$  pixels matrix with a 1 ms/px dwell time during 100 cycles ( $\sim 2$  min). A numerical blanking was applied to  $32 \times 32$  px matrix to consider only the innermost  $1 \times 1 \mu\text{m}^2$  region, this in order to limit the contribution of potential contamination. A cloud of electron was generated above the sample's surface to avoid the accumulation of positive charges induced by the implantation of the  $\text{Cs}^+$  beam.  $^{12}\text{C}$ ,  $^{16}\text{O}^1\text{H}$ ,  $^{28}\text{Si}$ ,  $^{32}\text{S}$ , and  $^{35}\text{Cl}$  were acquired simultaneously in multicollection mode on 5 electron multipliers (Ems). Each measurement was corrected for the 44 ns of the detector's deadtime. We used a  $20 \times 140 \mu\text{m}$  entrance slit (#3) with a 200  $\mu\text{m}$  diameter aperture slit (#2) and a 10% energy slit to optimize the Mass-Resolving Power (MRP) without compromising our transmission. This setting enabled a  $\text{MRP} > 8000$  which ensured the separation of  $^{16}\text{O}^1\text{H}$  and its isobaric interference  $^{17}\text{O}$ . The standards used for the calibration of the different elements were: two glasses NIST 610 and 612; a Suprasil crystal; two basaltic glasses MRN-G1 and IND-G2 (Shimizu et al., 2017) and three internal standards STR 10-11-13 (Créon et al., 2018). All of this enables to achieve a statistically robust linear regression spanning a range of concentrations from wt% ( $\text{H}_2\text{O}$ ) to hundreds of ppm ( $\text{CO}_2$ , S, Cl) (Figure S1 in the Supplementary Material S2). The nano-SIMS technique is typically associated with higher analytical precision ( $< 5\%$ ) and lower detection limits ( $< 10$  ppm typically) for Cl (Rose-Koga et al., 2021) relative to electron probe (Rose-Koga et al., 2020). The Suprasil standards is a synthetic crystal known for its purity. It has been extensively used alongside Herasil crystal to assess the blank values for  $\text{H}_2\text{O}$  and  $\text{CO}_2$  (e.g., Shimizu et al., 2021). By setting their concentrations

at 0 in our calibrations we systematically subtract these blank contributions. Errors were estimated as the quadratic sum of the measured  $1\sigma$  Poisson ratio error and the  $1\sigma$  prediction/confidence interval local estimation and were subsequently reported as  $2\sigma$  errors. For each measurement corrected by our determined linear regression we can extract an error associated with the estimation through the 95% confidence interval. This error is then quadratically added to the error on the measurement itself (poissonian error on ratios).


### 3.2.3. Raman Spectroscopy

Raman spectra were collected using an inVia confocal Raman micro-spectrometer manufactured by Renishaw and equipped with a  $532.1 \pm 0.3$  nm diode-pulsed solid state laser (200 mW output power), a Peltier-cooled CCD detector of  $1040 \times 256$  pixels, and a Leica DM 2500 M optical microscope with a motorised XYZ stage, at the LMV. Scattered light was collected via a back-scattered geometry. Laser power was about 10 mW on the sample surface. A 100x microscope objective, a grating of 2400 grooves  $\text{mm}^{-1}$ , and a 20- $\mu\text{m}$  slit aperture (high confocality setting) were used, which resulted in spectral resolution better than  $0.4 \text{ cm}^{-1}$  and in vertical and horizontal spatial resolutions of approximately 2-3 and 1  $\mu\text{m}$ , respectively, near the surface of the sample. Calibration of peak positions was performed at room temperature, based on the  $520.5 \text{ cm}^{-1}$  peak of silicon and the  $1331.5 \text{ cm}^{-1}$  peak of a diamond standard; two Neon emission bands of 568.98163 and 576.44188 nm wavelengths were used to check the linearity of the spectrometer. For  $\text{CO}_2$  measurements in MI bubbles, the spectra were recorded from 725 to  $1880 \text{ cm}^{-1}$  using the WiRE<sup>TM</sup> 4.4 software. Then, spectra were acquired in the 60- $1320 \text{ cm}^{-1}$  wavenumber range to detect presence of mineral phases (e.g., carbonates, sulphates) or fluid  $\text{SO}_2$ . Finally, spectra were collected in the  $2800\text{-}3900 \text{ cm}^{-1}$  region to detect  $\text{H}_2\text{O}$  and/or  $\text{OH}^-$  molecules. Acquisition time for a single  $\text{CO}_2$  analysis was fixed at 60 s. Spectra were initially treated for removal of cosmic rays, and then an appropriate polynomial baseline correction was applied before fitting a combined Gaussian-Lorentzian curve to the  $\text{CO}_2$  Fermi diad (e.g., Moore et al., 2015; Mironov et al., 2020; Schiavi et al., 2020). The analytical uncertainty associated with  $\text{CO}_2$  density measurements is  $< 0.05 \text{ g/cm}^3$  (Boudoire et al., 2023). Examples of Raman spectra acquired on shrinkage bubbles in melt inclusion are shown in Figure 2. In all shrinkage bubbles, only  $\text{CO}_2$  was detected as fluid phase (Figure 2a).

### 3.2.4. Laser Ablation-ICPMS

Trace-element measurements on melt inclusions were carried out using a laser ablation system (193 nm Excimer Resonetics M-50E) associated with an inductively coupled plasma mass spectrometer (Agilent 7500 cs LA-ICP-MS at the LMV). Analysis was performed following procedures outlined in the previous studies (e.g., Rose-Koga et al., 2012). We used a pulse energy of about 3 mJ, a spot diameter between 15 and 33  $\mu\text{m}$ , and a laser pulse frequency of 2–3 Hz, depending in the inclusion size to keep a fluence at sample surface of about  $4 \text{ J/cm}^2$ . The background was measured for 30–40 s before ablation and analysis time was approximately 100 s. Data reduction was performed using the GLITTER software (Van Achterbergh et al., 2001). This technique uses CaO (measured by electron probe) as an internal standard. Reproducibility and accuracy of the analysis were constrained by systematic analysis of two standards BCR2-G and SRM NIST 612 at the beginning, in the middle and at the end of the session. Typical errors on the samples ( $1\sigma$  error of mean) are less than 10% for all trace elements analysed.





**Figure 2.** (a) Raman spectrum of melt inclusion shrinkage bubble (sample Fog 43-8-1). The bubble is made of pure CO<sub>2</sub>, as revealed by the two strong peaks at 1284 and 1387 cm<sup>-1</sup>, identified as Fermi

diad (Kawakami et al., 2003). The separation between the two Raman bands of CO<sub>2</sub> is a reliable densimeter for the CO<sub>2</sub> fluid. **(b)** Raman spectra collected from the shrinkage bubble of Fog 43-8-1 (grey spectrum) and Fog 43-6-2 (orange and light blue spectra, acquired during tomographic analysis) MI samples. In the bubble, minerals such as Fe-magnesite and nahcolite were identified, as evidenced by peaks at 318, 737, and 1092 cm<sup>-1</sup> for Fe-magnesite, (Mg,Fe)CO<sub>3</sub> (Boulard et al., 2012) and the peaks at 685, 1045, and 1266 cm<sup>-1</sup> for nahcolite, NaHCO<sub>3</sub> (Fall et al., 2007). Both spectra also exhibit the characteristic olivine peaks at 822 e 854 cm<sup>-1</sup>.


## 4. Results

### 4.1. Whole-rock and mineral chemistry

The studied tephra samples are nepheline-normative (Ne= 17.6–21.2 wt.%) basanites (SiO<sub>2</sub> = 41.73–43.78 wt.%; K<sub>2</sub>O + Na<sub>2</sub>O = 6.69–7.68 wt.%) (table 1 of Supplementary Data S1), with a common mineral assemblage represented by clinopyroxene, olivine, plagioclase, and Fe-Ti oxides. These tephra samples are part of the same suite analysed (for major elements and fluid inclusions) by Lo Forte et al. (2023). The olivine grains have euhedral shapes and exhibit a narrow range of compositions (forsterite mol% Fo<sub>80-85</sub>) (table S3 of Supplementary Material S1). The olivine phenocrysts of sample SJ (Fog 43) are slightly more forsteritic (Fo<sub>83-85</sub>) than the 1951 (Fog 8) and 2014/15 (Fog 11) olivines (Fo<sub>80-84</sub>). The same SJ olivines exhibit higher CaO (0.34–0.40 wt%) and NiO (0.13–0.15 wt%) contents than the 1951 and 2014/2015 olivines (CaO, 0.31–0.39 wt.%; NiO, 0.03–0.15 wt.%). Mineral chemistry results are detailed in table S3 of Supplementary Material S1.

### 4.2. Melt inclusions: size and morphology.

Olivines contain abundant MI, but only 13 % of them were large enough (> 25 μm) for trace element analyses. All studied inclusions are perfectly glassy, and elliptical in shape (Figure 3), with the long axis (a) ranging between 17.0 μm and 94.0 μm (mean, 49.6 μm) and the short axis (b) between 11.3 μm and 83.1 μm (mean, 40.7 μm). Different criteria are used in the literature to determine the size of the third semi-axis c (e.g., axis a > b =c, Hauri et al., 2018; Shaw et al., 2008, 2010; Wanless et al., 2014, 2015 ; geometric mean, Ni et al., 2017); direct measurements can be used to test different assumptions (Mironov et al., 2020). Here we use the criteria of Moore et al. (2015) that assumes axis (c) equals to the arithmetic mean of axes (a) and (b) of the ellipsoid. The semi-axis (c) is therefore estimated at between 14.2 μm and 88.5 μm (mean, 37.8 μm). Shrinkage bubbles were observed inside all the MI (table S7 of Supplementary Material S1). However, in 31 % of the selected MI the bubbles were lost during the polishing process as the MI surface was exposed for analysis. The MI in which bubbles were lost are not considered further (except samples Fog 43-3-1 for which we report glass CO<sub>2</sub> contents only). There is no textural evidence of MI decrepitation. Roughly half of the shrinkage bubbles contain mineral phases such as Fe-magnesite, nahcolite, and sulphates (Figure 2b).




**Figure 3.** Measurements of the major (a) and minor (b) axes of the MI (Fog 43-6-3) for the determination of the melt volume and measurement of the diameter (d) of the shrinkage bubble for the calculation of its volume.

#### 4.3. Melt inclusion major element compositions and post-entrapment correction (PEC)

Melt inclusion glasses are foiditic to basanitic and compositionally overlap with bulk rocks (Fig. 4a). They typically contain 39.89-45.95 wt.%  $\text{SiO}_2$ , 3.15-5.85 wt.%  $\text{MgO}$ , 3.13-5.32 wt.%  $\text{Na}_2\text{O}$ , and 8.07-10.74 wt.%  $\text{FeO}_t$  (Fig. 4). A variety of processes may alter the melt composition after entrapment (see review in Wallace et al., 2021), including: (i) post-entrapment crystallization (PEC) of olivine at the MI-host mineral interface during slow cooling (Kent et al., 2008); (ii) volatile loss from the melt to the shrinkage bubbles (Wallace et al., 2015) or due to inclusion decrepitation (e.g., Maclennan, 2017); (iii) hydrogen ('water') diffusive loss (Hauri, 2002; Massare et al. 2002; Portnyagin et al. 2008, 2019; Chen et al. 2011; Gaetani et al. 2012); (iv) iron, magnesium diffusive exchange between the inclusion and its host mineral (e.g., Danyushevsky et al., 2000; Cottrell et al., 2002; Gaetani and Watson, 2000, 2002). We correct the composition of olivine-hosted MI for PEC using Melt inclusion modification corrections (MIMiC) (Rasmussen et al., 2020). The code employs the ( $\text{KD}^{\text{Fe-Mg}}$ ) model of Toplis (2005) and the olivine-melt thermometer of Putirka et al. (2007) to calculate the Fe-Mg olivine-melt partitioning (at redox conditions estimated by Moussallam et al., 2019) and to estimate magma temperature, respectively. Melt inclusion compositions, both pre- and post- MIMiC correction, are provided in tables S5 and S6 of Supplementary Material S1. PEC correction requires the addition of 0.03 to 9.28 % olivine to restore melt inclusion compositions to equilibrium with their hosts. The average PEC correction is 3.85 % for São Jorge (SJ) MI and 2.17 % for the two recent eruptions. Following PEC correction, melt inclusion compositions range 39.87-45.74 wt.%  $\text{SiO}_2$ , 4.00-7.36 wt.%  $\text{MgO}$ , 3.02-5.20 wt.%  $\text{Na}_2\text{O}$ , and 8.36-11.57 wt.%  $\text{FeO}_t$ . On the total alkali-silica diagram (Figure 4a), the PEC-corrected MI span the foidite to basanite compositional fields, with the most  $\text{SiO}_2$ -rich MI (41.74-45.74 wt.%) overlapping the range of the whole-rocks (compositions given in Lo Forte et al., 2023) and of bulk rocks from the literature (Hildner et al., 2011, 2012; Mata et al., 2017). These MI have  $\text{MgO}$  contents varying from 4.00 to 5.76 wt.%,  $\text{TiO}_2$  from 2.44 to 3.06 wt.%, and  $\text{P}_2\text{O}_5$  from 0.84 to 1.36 wt.%. The  $\text{CaO}/\text{Al}_2\text{O}_3$  ratios vary between 0.88 and 0.97 for SJ MI, and between 0.44 and 0.84 for the 1951 and 2014/15 MI (Figure 4b).


We model fractional crystallisation (FC) (Figures 4b-d) to determine the extent of crystallisation at the time of entrapment, and to test if the melt follows the liquid line of descent (LLD) prior to be entrapped, or if instead points to some Fe-Mg diffusive exchange during crystallisation (Figure 4b, c). We start our simulations with a hypothetical parental melt composition corresponding to the primitive lava Fogo 16 from Hildner et al. (2012) (MgO = 9.79 wt.%; FeO = 12.03 wt.%). We assume an initial H<sub>2</sub>O content of 2.0 wt.% (see below) and an oxygen fugacity of 1.5 log units above the Fayalite-Magnetite-Quartz buffer ( $\Delta\text{FMQ} +1.5$ ; Moussallam et al., 2019). Starting from this parental melt composition and an initial temperature of 1150 °C, we simulate the cooling-crystallisation trajectory using Petrolog3 (Danyushevsky and Plechov, 2011), using the melt-mineral equilibrium models of Danyushevsky, (2001) for olivine, clinopyroxene, and plagioclase, and Ariskin and Barmina, (1999) for magnetite. Results (Fig. 4b,c,d) show that the modelled LLD accurately reproduces the whole-rock compositional trend. After PEC correction, MI also match more closely the LLD and the bulk rocks trend (Figure 4b,c,d), implying diffusive Fe loss was minor (if any). No correction was therefore applied. All raw and corrected MI compositions are listed in Tables S5 and S6 of Supplementary Material S1.



**Figure 4** Total alkali-silica diagram (a) and major element compositions of melt inclusions (b, c, d) together with whole-rock compositions from Hildner et al. (2011, 2012), Lo Forte et al. (2023) and Mata et al. (2017). The solid lines are liquid lines of descent (LLD), calculated from a primitive parent melt (SiO<sub>2</sub> 42.14 wt.%, MgO 9.79 wt.%, FeO 12.03 wt.%, Al<sub>2</sub>O<sub>3</sub> 12.41 wt.%, CaO 12.96 wt.%) using Petrolog3 (Danyushevsky and Plechov, 2011). The LLD are calculated assuming initial H<sub>2</sub>O content of 2 wt.% and oxygen fugacity of  $\Delta\text{FMQ} + 1.5$  (Moussallam et al., 2019). We used the melt-mineral equilibrium models of Danyushevsky (2001) for olivine, clinopyroxene and plagioclase, and of Ariskin and Barmina (1999) for magnetite. The orange LLD was calculated for FC > 55%, to fit the more evolved MI (d). The grey area represents the range of forsterite component (Fo<sub>80-85</sub>) in the analysed olivines hosting MI (a).

#### 4.4. Trace element composition

Table S5.2 of Supplementary Material S1 reports the concentrations of a selected suite of trace elements in the studied MI. The primitive mantle normalised trace element patterns (Fig. 5) (this study) match those of the bulk-rocks (Hildner et al., 2011, 2012; Mata et al., 2017). No negative Hf anomaly is observed, and the Zr/Hf ratios (averaging at 39) are comparable to the typical primitive mantle value (~36; Palme and O'Neil, 2003). The LILE (Large Ion Lithophile Elements)-enriched composition of Fogo melts is additionally corroborated by elevated Ba/La (15.0-20.3), Rb/Sr (0.05-0.07) and Rb/La (0.98-1.54) ratios (Weaver, 1991).




**Figure 5.** Primitive mantle-normalised (Sun and McDonough, 1989) trace element patterns in MI samples (orange patterns; average values from all samples are used). The trace element patterns are intermediate between the compositional ranges of enriched mantle sources (EM-I and EM-II; data from Willbold and Stracke, 2006) and the HIMU mantle source (Willbold and Stracke, 2006). Note the significant Nb and Ta enrichment relative to the light REE. The red line identifies the compositions of Mid-Oceanic Ridge Basalts (MORB; Sun and McDonough, 1989).


#### 4.5. Volatiles

All the melt inclusion glasses are volatile-rich and contain 1.97 to 2.79 wt.% H<sub>2</sub>O and 9002 to 16044 ppm CO<sub>2</sub> in SJ MI (Fig. 6a). MI glasses from recent (1951-2014/15) eruptions are slightly less CO<sub>2</sub>-rich than the SJ MI glasses and contain 1.20 to 2.92 wt.% H<sub>2</sub>O and 1239 to 9102 ppm CO<sub>2</sub> (Fig. 6; Table S5.1 Supplementary Material S1). CO<sub>2</sub> in MI glasses broadly decreases with increasing degree of melt evolution, e.g., with decreasing CaO/Al<sub>2</sub>O<sub>3</sub> (Fig. 6b) and MgO (Fig. 6d) and increasing FeO<sub>t</sub>/MgO (Fig. 6c). The SJ MI glasses exhibit higher S concentrations (2676-7794 ppm) than the 1951 and 2014/15 MI (1146 to 5509 ppm) (values measured by nano-SIMS, Tables 5 and S5.1 Supplementary Material 1). Chlorine concentrations, measured by nano-SIMS, range from 947 to 1358 ppm in SJ MI and from 956 to 1415 ppm for the recent eruptions. The measured

fluorine concentrations are 1285-2216 ppm in SJ MI glasses and 1617-2249 ppm in the 1951-2014/15 MI glasses. For each element, a general least-squares linear regression was calculated, along with both 95% confidence and prediction intervals. This process allowed us to extract an error associated with the estimation through the 95% confidence interval for each measurement corrected by the determined linear regression. Subsequently, this error was quadratically added to the measurement itself. The absolute and relative errors were most pronounced for very low concentrations outside the range covered by our set of standards. As we extrapolated to higher concentrations, absolute errors increased. However, for measured concentrations (on average, >2000 ppm), the relative error rose less drastically compared to lower concentrations. For further details, please refer to Table S9 in Supplementary Material S1 and Figure S2 in Supplementary Material S2. The PEC-corrected concentrations vary as follows: 1.9-2.7 wt.% (SJ MI) and 1.2-2.8 wt.% (1951 and 2014/15 eruptions) for H<sub>2</sub>O (Fig. 6a); 2621-7718 ppm (SJ) and 1120-5250 ppm (1951-2014/15) for S; 917-1328 ppm (SJ) and 934-1415 ppm (1951-2014/15) for Cl (Fig. 7). As typical of alkaline magmas in general (Webster et al., 2018), the Fogo MI glasses are especially Cl- and F-enriched (up to 1415 and to 2249 ppm) relative to other OIB localities (Fig. 7), being only matched by the El Hierro basanite MI (Taracsák et al., 2019). The Fogo MI have F/Cl ratios in the 1.2 to 2.4 range, i.e., higher than F/Cl values of most arc magmas, exception made for F/Cl reaching 2.5 in the Trans Mexican arc (Vigouroux et al., 2008) and in the Central American arc (Sadofsky et al., 2008; Fig. 7). This F-rich signature is a distinctive characteristic of OIB magmas in general (Kendrick et al., 2014).



**Figure 6.** CO<sub>2</sub> concentrations vs. H<sub>2</sub>O (a), CaO/Al<sub>2</sub>O<sub>3</sub> (b) and FeOt/MgO (c) ratios, and MgO (d). The SJ MI exhibit higher H<sub>2</sub>O and CO<sub>2</sub> contents than those from the more recent eruptions in 1951 and 2014-15. The orange symbols represent data related to the total CO<sub>2</sub> content in the MI (glass + bubble), corresponding to measurements obtained through Raman-based total CO<sub>2</sub> estimates (see section 5.1 for further details).



**Figure 7.** In the F vs. Cl plot, our data are compared to MI of different OIB localities and from volcanoes of different geodynamic contexts, such as Ascension (Webster and Rebert, 2001), Etna (Métrich et al., 1993, 2004), La Soufriere (Heath et al., 1998), St. Helens (Blundy and Cashman, 2005), Paricutin (Rowe et al., 2011), Trans Mexican belt (Vigouroux et al., 2008), Central American belt (Sadofsky et al., 2008), Vanuatu (Métrich et al., 2011), Eifel (Harms and Schmincke, 2000), Aeolian islands (Di Martino et al., 2011), Masaya (Pérez et al., 2020), and Kamchatka volcanoes (Portnyagin et al., 2011). The F/Cl ratios of our MI (1.16-2.41) fall in the field of OIB volcanoes and partially overlap the composition of El Hierro MI.



## 5. Discussion

### 5.1. Reconstructing the total (glass+bubble) CO<sub>2</sub> content in MI

Our results show that the Fogo MI glasses contain up to ~1.6 wt.% CO<sub>2</sub> (Fig. 6), plus ~2 wt.% H<sub>2</sub>O and ~0.48 wt.% S, on average. However, all MI also contain CO<sub>2</sub>-rich bubbles (Fig. 2). There is a positive relationship between total MI volumes and bubbles volumes (Fig. 8a), indicating that all MI contain similarly low bubble volumetric fractions (0.2-8.6 %). We conclude that the contemporaneous entrapment of melt and fluid inclusions is unlikely (Rasmussen et al., 2020), and therefore all fluid bubbles found are shrinkage bubbles. Hence, as recently demonstrated elsewhere (Taracsák et al., 2019), reconstructing the total MI CO<sub>2</sub> budget requires a procedure in which the CO<sub>2</sub> masses contained in (i) shrinkage bubbles and (ii) glass are both fully taken into account.

The CO<sub>2</sub> mass contained in the glass is initially calculated by multiplying glass mass in the MI (glass density × glass volume) by CO<sub>2</sub> concentration in the glass (cfr. 4.5). Glass density is derived using MIMiC (Rasmussen et al., 2020), which uses the relationships of Lesher and Spera (2015), while glass volume is estimated following the procedure of Moore et al. (2015).

Next, we quantify shrinkage bubble CO<sub>2</sub> using three different, independent methodologies: (i) the Raman method (Esposito et al. 2011; Hartley et al. 2014; Moore et al. 2015; Rasmussen et al. 2018); (ii) the Simulated-volume method (Rasmussen et al., 2020); and (iii) the Observed-volume method (e.g., Shaw et al., 2010; Wanless et al., 2014; Hauri et al., 2018; Tucker et al., 2019). Our aim is to use the level of agreement/disagreement of these different methods as a proxy for the level of accuracy/inaccuracy of such bubble CO<sub>2</sub> estimates (Fig. 8).

In the Raman method (Hartley et al., 2014; Moore et al., 2015; Moore and Bodnar, 2019; Rasmussen et al., 2020), the bubble CO<sub>2</sub> mass is derived as the product of bubble CO<sub>2</sub> density × bubble volume. Raman indicates a pure-CO<sub>2</sub> gas phase composition for the shrinkage bubble (Figure 2 a); we hence apply the densimeter by Wang et al. (2011) for the calculation of the bubble CO<sub>2</sub> density.

The other two CO<sub>2</sub> estimates (Simulated and Observed) are derived using the MiMiC code (Rasmussen et al., 2020). The Simulated-volume method relies on estimating bubble volumes and CO<sub>2</sub> contents from modelling the growth of bubbles during the cooling and quenching of the melt inclusions (rather than directly measuring bubble volumes). This bubble growth model assumes that exchange of CO<sub>2</sub> between melt and bubble terminates at a CO<sub>2</sub> closure temperature ( $T_c$ ) for diffusion (generally higher than the glass-transition temperature) followed by further bubble expansion with no CO<sub>2</sub> loss to bubble (Rasmussen et al., 2020).

Finally, the Observed-volume method (e.g., Shaw et al. 2010; Wanless et al. 2014; Hauri et al. 2018; Tucker et al. 2019) uses directly measured bubble volumes and entails determining the CO<sub>2</sub> mass within the vapor bubble by applying the modified Redlich-Kwong (MRK) equation of state. The method relies on the assumption that vapor-melt equilibrium is maintained throughout the entire bubble quenching/growth history. The temperature of vapor-melt equilibrium is determined based on the calculated glass-transition temperature, which however can vary significantly by several hundred degrees Celsius, depending on factors such as melt viscosity and cooling rate (Giordano et al., 2005).

All three CO<sub>2</sub> restoration methods have their caveats and limitations which can potentially lead to overestimation/underestimation of the total MI CO<sub>2</sub>. Use of the Raman restoration method (e.g., Esposito et al., 2011; Moore et al., 2015), in particular, is challenged by difficulties in (i) correctly measuring melt inclusion geometry, and hence glass vs. bubble volumes (Moore et al., 2015) and


(ii) quantifying the relative volumetric fractions occupied by minerals and fluid in the bubble (Schiavi et al., 2020). Determination of melt inclusion geometry (glass and bubble volumes) in principle requires accurate 3D scanning microtomographic analysis, which is rarely available. Our glass/bubble volumes (Table S7 of Supplementary Material S1) are estimated from 2D analysis, and on the assumption that the length of the third semi-axis  $c$  (normal to the polished plane) equals to the arithmetic mean of axes  $a$  and  $b$  (Moore et al., 2015); this assumption should, at least in principle, introduce a relatively minor (5%) error (Wallace et al., 2015). More problematic is to account for the carbon mass sequestered in mineral phases inside the bubble. Recent studies have found that mineral phases such as carbonates, sulphides, sulphates and native sulphur can precipitate inside bubbles during cooling (e.g., Kamenetsky et al., 2002; Hartley et al., 2014; Moore et al., 2015; Esposito et al., 2016; Robidoux et al., 2018; Guzmics et al., 2019; Tucker et al., 2019; Schiavi et al., 2020; Venugopal et al., 2020; Buso et al., 2022; Feignon et al., 2022). Precipitation of these minerals can potentially sequester large fractions of the bubble-carried  $\text{CO}_2$ , hence ignoring their presence can result in significant (up to 40%) underestimation of the total MI  $\text{CO}_2$  (Schiavi et al., 2020). We detected the presence of carbonate phases, such as Fe-magnesite and nahcolite (Supplementary Material S2 Figure S2), and sulphates, in roughly 50 % of the Raman spectra. The proportions of the different phases are highly variable. Raman microtomography revealed significant volumes of carbonates crystallised on the walls of the MI Fog 43-6-2 bubble, amounting to  $\sim 35$  vol% according to an approximate average estimate based on few 2D sections extracted from the 3D map (Supplementary Material S2 Figure S2). This corresponds to  $\sim 25\%$  of the  $\text{CO}_2$  sequestered in the bubble (compared with 9% reported in Table S7 of the Supplementary Material S1). We conclude that the total  $\text{CO}_2$  content can be underestimated (by about 15-20 % for MI Fog 43-6-2) if carbon stored in carbonates inside the bubble is neglected. However, considering the overall positive correlation observed between glass MI  $\text{CO}_2$  and shrinkage bubble density (Fig. 8b), we conclude that precipitation of carbonates is likely to be marginal in all samples except Fog 43-8-1 and 43-6-2 (that coherently exhibit the most intense Raman peaks associated with carbonates).

The second  $\text{CO}_2$  restoration methodology used (Simulated-volume method) overcomes some of the limitations above, as it does not rely on measured bubble volumes, but rather on estimated bubble volumes based on modelling bubble growth during MI cooling and quenching. The detailed sensitivity study of Rasmussen et al. (2020) demonstrates that the amount of cooling experienced by MI, from their entrapment temperature ( $T_e$ ) to their  $\text{CO}_2$  closure temperature ( $T_c$ ), is the key parameter controlling the extent of  $\text{CO}_2$  loss to bubbles. According to Rasmussen et al. (2020), this cooling extent ( $T_e - T_c$ ) depends on MI dimension (measured texturally) and cooling rate, for which we unfortunately have no independent estimate. In such conditions, MIMiC adopts a calculated cooling rate derived by solving the Dodson equation for the magnesium closure temperature ( $T_c(\text{MgO})$ ); this procedure can potentially introduce a maximum error of 50% on the restored  $\text{CO}_2$  concentrations (when the used  $T_c(\text{MgO})$  is in excess of by one order of magnitude).

Finally, the Observed-volume method calculates the  $\text{CO}_2$  mass in the bubble by applying an equation of state and performing a mass balance. In this approach, the observed bubble volume, measured under the microscope, used in the calculation assumes that melt-bubble equilibrium is maintained throughout the entire bubble growth history. However, if equilibrium is not maintained, the calculation based on the observed volume may not be accurate. Additionally, this method assumes the glass transition temperature ( $T_g$ ) as the final melt-bubble equilibrium temperature, but precisely assessing  $T_g$  is complicated by its strong dependence on factors such as cooling rate, melt composition, melt viscosity, and volatile content (Rasmussen et al., 2020).

Ultimately, since all restoration methods have limits and caveats, a multi-method approach is advisable (Rasmussen et al., 2020). The outputs of the three restoration methods used here are compared in Figures 8c and 8d. We find that the Raman spectroscopy measurements and simulated-volume approach yield  $\text{CO}_2$  concentrations that agree within 9.7 % on average (range: 0.4-22.6 %).

The agreement between these two independent methodologies provides confidence in our restored total MI CO<sub>2</sub> values. The total MI CO<sub>2</sub> contents determined with the observed-volume method also positively correlate with those inferred from Raman analysis (Fig. 8d), but typically plot above the 1:1 line, indicating that the observed-volume approach may systematically over-estimate the MI CO<sub>2</sub> budget, as already suggested (Moore et al., 2018; Rasmussen et al., 2020). Rasmussen et al. (2020) also found that the concentrations reconstructed from both Raman measurements and the simulated-volume method are quantitatively consistent with those obtained by re-homogenising MI at high temperature and pressure – this latter procedure being considered as that yielding the most accurate CO<sub>2</sub> reconstructions. Ultimately, while we are well aware of the many existing potential sources of uncertainty, we conclude that our calculations bring nonetheless robust evidences that (i) in some cases of OIB MI, a large fraction (up to ~50 %) of the MI CO<sub>2</sub> is contained in the bubbles (Mironov et al., 2015), and (ii) the total (glass+bubble) CO<sub>2</sub> content in the Fogo parental melt is certainly > 1 wt. %, and potentially as high as ~2 wt. % (the highest value obtained by the Raman and Observed-volume methods). For the sake of simplicity, in the discussion below we refer to the Raman-based bubble (and hence total) CO<sub>2</sub> estimates, but we note that these overlap with those derived from the bubble growth model using MiMIC by a factor of <10%.



**Figure 8.** (a) Plot of bubble volumes as a function of MI volumes; isovolumetric lines corresponding to bubble/MI volume ratios of 0.2 %, 1.0 %, 2.0 %, 5.0 %, and 8.0 % are also shown. (b) The plot shows a positive correlation between the CO<sub>2</sub> calculated in the glass and the density of the shrinkage bubble, except for the two samples (Fog 43-8-1 and 43-6-2) which exhibit intense Raman peaks associated with carbonates, specifically Fe-magnesite and nahcolite crystals on the bubble wall (Fig. 2). As shown in (b), the overlooked presence of carbonates in the bubble leads to an underestimation of the total CO<sub>2</sub> content. Excluding these two outliers, the samples show a good

correlation ( $R^2 = 0.802$ ). Plot of the total  $\text{CO}_2$  concentration in MI, calculated using the Raman and Simulated methods (**c**) and the Raman and Observed methods (**d**). In plot (**c**), there is a strong correlation ( $R^2=0.891$ ), with  $\text{CO}_2$  concentrations from Raman and Simulated-volume methods showing a 1:1.1 relationship. In contrast, in plot (**d**), the total  $\text{CO}_2$  calculated using the Raman and Observed-volume methods is not strongly correlated ( $R^2=0.806$ ), with a ratio of 1:1.6. The dashed lines correspond to a ratio 1:1.

## 5.2. A $\text{CO}_2$ -rich parental melt


Finally, glass  $\text{CO}_2$  mass and shrinkage bubble  $\text{CO}_2$  mass are summed to obtain total  $\text{CO}_2$  mass in MI, which is ultimately converted in total  $\text{CO}_2$  concentration in the MI. All calculations, for each MI analysed, are given in Table 1 and in Table S7 of Supplementary Material S1. Using the Raman method results described above, we find that bubbles contain 281-5546 ppm  $\text{CO}_2$  (for the SJ MI) and 857-2920 ppm  $\text{CO}_2$  (for the 1951 and 2014/15 MI). The total  $\text{CO}_2$  contents thus range from 1309 to 21502 ppm for SJ MI, and from 1084 to 9102 ppm for the 1951 and 2014/15 eruptions (Fig. 6). We thus estimate that 3 to 47 % of the total MI  $\text{CO}_2$  content is contained in the bubbles.

**Table 1.** Summary of the total  $\text{CO}_2$  content (bubble + glass) calculated in the analysed melt inclusions using three different methods (Raman, Simulated, and Observed).

Sample	$\text{CO}_2$ wt% in MI (Raman Method)	$\text{CO}_2$ wt% in MI (Simulated Method)	$\text{CO}_2$ wt% in MI (Observed Method)
Fog 43-3-1	1.25	1.43	/
43-5-1	2.15	1.95	2.75
43-6-1	1.49	1.48	2.31
43-6-2	1.23	1.42	2.15
43-6-3	1.16	1.18	1.74
43-6-4	/	/	/
43-7-1	0.13	/	/
43-8-1	1.01	1.40	/
Fog 8-1-1	0.40	0.54	0.77

8-4-1	0.89	1.03	1.41
8-7-1	0.74	0.92	0.79
8-8-1	0.91	1.16	1.69
8-11-2	0.62	0.47	0.97
8-11-3	/	/	/
8-11-4	0.11	/	/
8-11-5	/	/	/
Fog 11-2-1	0.21	0.22	0.46
11-3-1	0.82	0.99	1.08

Our results show that the basanitic melts from Fogo are exceptionally CO<sub>2</sub>-rich. Their volatile-rich nature is additionally demonstrated by high halogen contents (Fig. 7). The estimated CO<sub>2</sub> content of the parental melt (2.15 wt.%) is up to a factor 10 higher than for other OIB localities (e.g., Erebus in Antarctica, Moussallam et al., 2014; Fernandina and Santiago in the Galapagos, Koleszar et al., 2009; Piton de la Fournaise in Reunion Island, Di Muro et al., 2014; Kilauea in the Hawaii, Sides et al., 2014; Pico in the Azores, Métrich et al., 2014) and Iceland (e.g., Bardabunga, Bali et al., 2018, and Laki, Hartley, 2014) (Fig. 9a). In addition to confirming that bubbles do matter if the CO<sub>2</sub> MI budget is to be reconstructed (Moore et al., 2015; Rasmussen et al., 2020; Buso et al., 2022), our results also corroborate recent work that shows that alkali rich OIB magmas can be especially carbon-rich, as observed at El Hierro (CO<sub>2</sub> up to 4 wt.% ; Longpré et al., 2017; Moussallam et al., 2019; Taracsák et al., 2019) and La Palma (3-5 wt. % ; e.g., Burton et al., 2023) in the Canary Archipelago, and at Piton de La Fournaise in La Reunion Island (3.5 ± 1.4 wt.%, Boudoire et al., 2018) (Fig. 9a) (note that in these studies shrinkage bubble CO<sub>2</sub> has been accounted for with either observations or modelling). The fact that alkali rich OIB volcanoes erupted magmas richer in CO<sub>2</sub> than alkali poor OIBs has been documented at a global scale (Sun and Dasgupta, 2023), although we see no evidence in our MI of the extremely high (9-10 wt.%) CO<sub>2</sub> contents inferred by these authors for the Cape Verde archipelago parental melts. Rather, the ~2 wt. % CO<sub>2</sub> content we estimate for the Fogo parental melt is at the lower range of the measured/inferred (Hudgins et al., 2015; Aiuppa et al., 2021) CO<sub>2</sub>-enriched signatures of alkaline mafic melts from continental rift context (e.g., in the East African rift system; Foley et al., 2012). This link between high CO<sub>2</sub> compositions and melt enrichment in alkalis (and other incompatible elements; Aiuppa et al., 2021) is further explored below (Fig. 10).




**Figure 9.** (a)  $\text{H}_2\text{O}$  and  $\text{CO}_2$  concentration in studied MI and in other OIB occurrences. Isobars were calculated using an average composition of our MI samples at temperature of  $1150\text{ }^\circ\text{C}$  and an oxygen fugacity of  $\Delta\text{FMQ} + 1.5$  (Moussallam et al., 2019). Degassing paths, in a closed system, were calculated using the volatile saturation models by Moretti et al. (2003) and MagmaSat (Ghiorso and Gualda, 2015), with the same input melt composition and temperature and volatile contents of 21502 ppm and 2.0 wt.% for  $\text{CO}_2$  and  $\text{H}_2\text{O}$  respectively. (b) In the  $\text{CO}_2$  content vs. Pressure graph, different saturation models are compared. The solid lines represent the outputs of the different models, instead the dashed lines represent the extrapolations of these models.

### 5.3. Clues from $\text{CO}_2$ vs. trace-element correlations

A potentially useful approach to test the accuracy of our reconstructed MI  $\text{CO}_2$  content is to normalize them to the measured Nb and Ba contents in the same MI glasses (Fig. 10). These trace elements are thought to have similar bulk mineral/melt distribution coefficients to that of carbon (e.g., Rosenthal et al., 2015; Hirschmann, 2018). Hence, the  $\text{CO}_2/\text{Nb}$  and  $\text{CO}_2/\text{Ba}$  ratios should not be fractionated during mantle melting and fractional crystallisation of mafic melts. If a mantle-sourced melt has not degassed  $\text{CO}_2$  during crustal emplacement, it is thus expected to preserve a record of its source mantle  $\text{CO}_2/(\text{Nb}, \text{Ba})$  ratio (e.g., Saal et al., 2002; Michael and Graham, 2015; Shimizu et al., 2016; Le Voyer et al., 2017; Hauri et al., 2018; Tucker et al., 2019). Following these premises, the expected range of mantle  $\text{CO}_2/\text{Nb}$  (Fig. 10a) and  $\text{CO}_2/\text{Ba}$  (Fig. 10b) have been inferred (see grey filled diagonal bands in Figure 10) from analysis of MORB that, owing to their volatile-depleted nature, are thought not to have reached volatile saturation before eruption (undegassed MORB; Michael and Graham, 2015) or entrapped as MI (e.g., Saal et al., 2002; Le

Voyer et al., 2017; Moussallam et al., 2023). OIB magmas, in contrast, owing to their volatile enriched nature, do systematically reach saturation during ascent (Dixon and Stolper, 1995) and, even more significantly, form post-entrapment shrinkage bubbles that can sequester a large fraction of the MI CO<sub>2</sub> upon cooling and decompression (Hartley et al., 2014; Moore et al., 2015; Wallace et al., 2015; Tucker et al., 2019; Schiavi et al., 2020). As noted above, most MI results from the literature refer to glass measurements only, in which no correction for CO<sub>2</sub> in the bubbles has been attempted (with exceptions; Taracsák et al., 2019; Tucker et al., 2019); unsurprisingly, these plot below the mantle field in Figure 10, implying their CO<sub>2</sub>/trace element ratios underestimate the source mantle ratios, owing to extensive CO<sub>2</sub> loss to bubbles, pre- and post-entrapment (e.g., Hartley et al., 2014).

Considering the highest reconstructed total CO<sub>2</sub> contents only, our MI exhibit a relatively narrow range of CO<sub>2</sub>/Nb (277-283) and CO<sub>2</sub>/Ba (26-22) (Fig. 10). This is at the lower limit (CO<sub>2</sub>/Nb), or below (CO<sub>2</sub>/Ba), the interval of representative ratios proposed for the upper mantle, based on the analysis of MORB MI and/or undegassed MORB glasses (Fig. 10). It is important to recall that, in their recent global compilation, Sun and Dasgupta (2023) have proposed that the mantle source of OIB volcanism is characterised by higher CO<sub>2</sub>/Nb (~1850) and CO<sub>2</sub>/Ba (~226) ratios than the depleted mantle source feeding MORB volcanism. Overall, Fig. 10 indicates that, in spite of our applied correction for CO<sub>2</sub> in shrinkage bubbles, our reconstructed total CO<sub>2</sub> concentrations likely still underestimate the real parental melt CO<sub>2</sub> contents, potentially because of early (before MI formation) CO<sub>2</sub> separation from melt into the vapour phase (e.g., Rosenthal et al., 2015; Le Voyer et al., 2017; Hirschmann, 2018; Aiuppa et al., 2021).



**Figure 10.** CO<sub>2</sub>/trace element ratios. (a) In the Nb versus CO<sub>2</sub> plot, most of the literature “degassed” MI and our MI with the CO<sub>2</sub> content measured in the glass only (white diamond with

light blue border) plot below the CO<sub>2</sub>-Nb mantle array (represented by the grey area), as defined by the un-degassed MI and glasses of the MORB (characteristic upper mantle CO<sub>2</sub>/Nb ratios from refs. Saal et al., 2002; Cartigny et al., 2008; Le Voyer et al., 2017; Hirschmann, 2018). Contrarily, our MI with the total CO<sub>2</sub> content (orange symbol, CO<sub>2</sub> glass+CO<sub>2</sub> bubble) fall in the mantle array field. (b) same as (a) but using Ba. Our MI with the total CO<sub>2</sub> content fall within the CO<sub>2</sub>/Ba mantle range, 74-120 (Saal et al., 2002; Cartigny et al., 2008; Le Voyer et al., 2017; Hirschmann, 2018). In contrast, literature “degassed” MI and our MI with CO<sub>2</sub> measured only in the glass plot below the CO<sub>2</sub>/Ba mantle array.

#### 5.4. High CO<sub>2</sub> implies deep magma storage

The important implication of the high CO<sub>2</sub> contents of the studied MI is that magma must be stored deep in the Fogo plumbing system prior to eruption (Figs. 9b, 11). To demonstrate this, we use the inferred H<sub>2</sub>O-CO<sub>2</sub> MI concentrations (Fig. 9a), in combination with available saturation models (Fig. 9b), to estimate the MI entrapment pressures (assuming MI are volatile saturated at the time of their entrapment). However, the conversion of volatile contents into pressures requires selection of one of the several saturation models existing in the literature (e.g., MagmaSat, Ghiorso and Gualda, 2015; Iacono-Marziano et al., 2012; D-Compress, Burgisser et al., 2015; Moretti et al., 2003; VolatileCalc, Newman and Lowenstern, 2002; Jiménez-Mejías et al., 2021; Shishkina et al., 2014), all of which have been calibrated over different compositional ranges and at relatively low pressures, and hence for lower CO<sub>2</sub> concentrations than the concentrations considered here). In addition to being calibrated at relatively low pressures (< 800 MPa, the model of Eguchi and Dasgupta, 2018 being the only exception), these different volatile saturation models predict CO<sub>2</sub> solubilities in Fogo melts that are highly variable, and typically span a factor 2 to 4 (or more) from the least and the most conservative estimate (Fig. 8b). This implies that, for a given MI CO<sub>2</sub> content, a range of MI entrapment pressures can be derived, with the VolatileCalc and Iacono-Marziano et al. (2012) models typically returning the lowest entrapment pressures, and the Jiménez-Mejías et al. (2021) model the highest (Fig. 9b). It is important to remind that some of these models have been calibrated for tholeiitic compositions, and their results are therefore shown for illustrative purposes only. The MagmaSat model outputs some intermediate entrapment pressure conditions in all cases (Fig. 9b).

We focus on the range (2095-21502 ppm) of total CO<sub>2</sub> contents, which converts into a minimum-maximum range of entrapment pressures of 100-290 MPa (217 MPa with MagmaSat) for the most CO<sub>2</sub>-depleted inclusion (Fog 11-2-1; 0.21 wt. %) and to 648-1430 MPa (773 MPa with MagmaSat) for the most CO<sub>2</sub>-rich inclusion (Fog 43-5-1; 2.1 wt. %). The upper range of MI entrapment pressures is higher than the pressures obtained previously at Fogo using clinopyroxene-liquid barometry (300-750 MPa) and fluid inclusion microthermometry (200-740 MPa) (Hildner et al. 2011, 2012; Klügel et al., 2020; Lo Forte et al., 2023). Ultimately, these pressures indicate MI may have been entrapped during early olivine crystallisation at depths of 26 to 35 km, well within the lithospheric mantle (extending from ~12 to ~60 km depth underneath Fogo Island; Liu et al., 2021).


We then use the composition (major elements and volatile contents) of this deeply trapped Fog 43-5-1 MI as the starting condition for model simulations of volatile degassing upon decompression. We run two model simulations of degassing using the saturation codes of Moretti et al. (2003) and MagmaSat (Ghiorso and Gualda, 2015). Both codes have been used previously to model magma degassing in a variety of volcanic systems erupting magmas of different composition (Aiuppa et al., 2007, 2010, 2017, 2022; Edmonds et al., 2010; Marini et al., 2011; Oppenheimer et al., 2011; Pino et al., 2011; Moretti 2013a, 2013b; de Moor et al., 2016; Bamber et al., 2022; Halldórsson et al., 2022; Romero et al., 2023; Sharpe et al., 2022; Boulanger and France, 2023; Liu et al., 2023; Mitchell et al., 2023). With both volatile saturation codes, we model dissolved volatile contents in



melt along a closed-system decompression path, starting from 1200 MPa pressure (final pressure, 0.1 MPa), in isothermal conditions ( $T = 1150\text{ °C}$ ), and at  $\Delta\text{FMQ} +1.5$ . Results of our model calculations, illustrated in Figure 9a, indicate that the modelled melt volatile compositions reproduce well the range of  $\text{H}_2\text{O}$ - $\text{CO}_2$  contents measured in our MI (Fig. 9a), providing confidence in our choice of volatile saturation models and model output. For comparison, we also show in Figure 9 the modelled degassing path calculated for (i) isothermal decompression in open-system conditions (orange curve) and (ii) non-isothermal (temperature range, 1128-985 °C) closed-system degassing with crystallisation taken into account (green curve). The latter model trend was obtained by linking together (a) MI saturation pressures (from  $\text{H}_2\text{O}$ - $\text{CO}_2$  contents) with (b) MI major element compositions, and hence (by matching against the modelled LLD; Fig. 4) melt temperature and extent of crystallisation. The open system run predicts some extreme  $\text{CO}_2$  depletions at low pressure (that are not observed in the MI dataset; see inset in Fig. 9a); while the non-isothermal closed system+crystallisation model trend predicts some larger ( $> 2\text{ wt. \%}$ )  $\text{H}_2\text{O}$  enrichments at low pressure that potentially explain the composition of some more evolved El Hierro MI.

Our results (Fig. 9) suggest magma storage conditions at Fogo to be significantly deeper than previously thought. Previous work, based on fluid inclusion microthermometry (Hildner et al., 2011, 2012; Klügel et al., 2020; Lo Forte et al., 2023) and on clinopyroxene-liquid geothermobarometry (Hildner et al., 2011, 2012; Klügel et al., 2020), have revealed the existence of a complex, vertically stacked magma storage system, with a main zone at  $17\text{--}25 \pm 5\text{ km}$  depth (Fig. 11) overlain, in the  $9\text{--}12 \pm 2\text{ km}$  depth range, by a sequence of levels of temporary magma stagnation, where differentiation eventually takes place prior/during eruption (generating phonotephritic melts, such as those erupted during the 1951, 1995 and 2014-2015 eruptions). Our volatile data, especially those of the  $\text{CO}_2$ -richest – and hence more deeply trapped – MI (2.15 wt.%, Fog 43-5-1), extend the range of magma storage conditions downward. The entrapment (saturation) pressure range, inferred for this MI, is between 648 and 1430 MPa (i.e. about 20 to 45 km depth), depending on the saturation model used (see Fig. 9b). If we concentrate on the models that account for the increase of  $\text{CO}_2$  solubility with melt alkalinity (and silica undersaturation), the inferred MI entrapment pressures are 773 (MagmaSat) and 1020 MPa (Moretti et al., 2003), respectively. With the mantle-crust reference model previously utilised by Lo Forte et al. (2023), these pressures correspond to entrapment depths of  $\sim 27\text{ km}$  (MagmaSat) and  $\sim 36\text{ km}$  (Moretti et al., 2003). The shrinkage bubble in this MI contains a small amount of minerals (cfr. 5.1), implying its Raman-derived  $\text{CO}_2$  bubble density (and hence bubble  $\text{CO}_2$  content) is not severely affected by mineral precipitation. Although it is well possible that the measured  $\text{CO}_2$  content underestimates the real  $\text{CO}_2$  content of the parental melt, because of pre-entrapment  $\text{CO}_2$  loss to bubbles (cfr. 5.2), we interpret such MI saturation pressures as indicative of a deep magma storage at  $31.5 \pm 4.3\text{ km}$  depth, at (or nearby) a main geological boundary bounded by the Elastic effective Thickness ( $T_e$ ; located at  $\sim 30\text{ km}$  depth; Pim et al., 2008).

Ultimately, our data prove that pre-eruptive magma storage/ascent underneath Fogo occurs/starts within/from a deep magma storage zone at  $31.5 \pm 4.3\text{ km}$ , in the lithospheric mantle. This is right above a cluster of deep earthquakes (38-43 km depth; Leva et al. 2019) that may correspond to fracturing events during magma accumulation zone replenishment events with more deeply sourced melts, rising from the convecting mantle source (at  $> 60\text{ km}$ ). Such deep pre-eruptive magma storage conditions are similar to those of other intraplate oceanic islands, e.g., El Hierro (Taracsák et al., 2019), Piton de la Fournaise (Boudoire et al., 2018), La Palma (Dayton et al., 2023), and the Azores islands (e.g., Zanon et al., 2023).



**Figure 11.** Schematic cross-section (not to scale) showing the entire magma plumbing system of Fogo volcano. The model is based on the framework proposed by Lo Forte et al. (2023). Our data provide new constrain on the location of the roots of the volcano at  $31.5 \pm 4.3$  km depth, which feed the upward magma storage zone at 13-25 km depth, as well as the magma stagnation zone located at 9-12 km depth. The blue stars represent the deep earthquakes recorded by Leva et al. (2019), in proximity of the deep magma accumulation zone. The right-hand side boxes present the previous barometry results. Consistent with Liu et al. (2021), the lithosphere-asthenosphere boundary (LAB) is located at 60 km depth. The three isotopic mantle components, which sign the magmatism of Cape Verde, were obtained from prior literature works (e.g., Gerlach et al., 1988; Doucelance et al., 2003; Escrig et al., 2005; Martins et al., 2010; Jackson et al., 2018a).

### 5.5. High carbon in the source mantle, and implications for the global carbon cycle

We use our parental melt CO<sub>2</sub> content inferred above (2.15 wt. %) to estimate the carbon content in the mantle source feeding volcanism at Fogo. To this aim, we use a modal batch-melting equation (Shaw, 1970):

$$C_{pm} = \frac{C_0}{[F + D(1-F)]} \cong C_0/F \quad (1)$$

in which  $C_{pm}$  stands for the CO<sub>2</sub> in mantle-derived parental melts (2.15 wt. % in our case),  $C_0$  is the initial abundance of CO<sub>2</sub> in the source mantle,  $D$  is the CO<sub>2</sub> solid-melt partition coefficient (assumed at 0.00055; Rosenthal et al., 2015), and  $F$  is the melt fraction (0 to 1). This partial melting degree  $F$  in the Fogo mantle source is estimated at 0.06-0.07 by using MI major element composition to derive the final stage mantle-melt equilibration temperature ( $\sim 1597$  °C) and pressure ( $\sim 4.8$  GPa), and hence melting degree, according to the methodology of Sun and Dasgupta (2020, 2023). With these numbers, the 2.15 wt. % parental melt CO<sub>2</sub> content converts into 355-414 ppm C in the Fogo mantle source.

We caution our estimated carbon range should be considered as a conservative lower limit, since our MI potentially entrap an already partially degassed melt (see above). Also, the model is intrinsically simplistic as it assumes a homogeneous (with fixed C content) mantle, while we know the mantle is in fact spatially heterogenous. In spite of these limitations, we conclude our results unambiguously prove the CO<sub>2</sub>-rich nature of alkali-rich mafic melts feeding intraplate volcanism at Cape Verde. Note that, given the very low  $D$  for CO<sub>2</sub> ( $\sim 0$ ; Rosenthal et al., 2015), the use of a fractional melting equation (in place of batch melting) would not lead to any obvious difference.

Our results, combined with recent observations at other OIB localities globally (e.g., Anderson and Poland, 2017; Longprè et al., 2017; Boudoire et al., 2018; Hauri et al., 2018; Miller et al., 2019; Taracsák et al., 2019; Tucker et al., 2019), suggest that a C enriched OIB mantle source is likely to be the rule rather than the exception. Our estimated Fogo mantle content (355-414 ppm C) is well within the OIB mantle source range recently derived by Sun and Dasgupta (2023) (330-400 ppm), but well above the maximal C content of  $164 \pm 55$  ppm inferred by Rosenthal et al. (2015) for the mantle source of intraplate basalts.

This carbon-enriched nature of global OIB, and of their mantle sources, has important consequences for our understanding of the mass/volume of deep carbon reservoirs, and of the deep carbon cycle - the transfer of carbon from the deep reservoir(s) (mantle+core) to the exosphere (ocean+atmosphere). First, our results contribute to the still open debate on the carbon content of the Bulk Silicate Earth, for which available estimates remain divergent, from  $\sim 110$  ppm C (Hirschmann, 2018) to 330-450 ppm C (Marty et al., 2020) or even up to  $\sim 500$  ppm C (Marty, 2012). The high C content of the OIB source mantle manifestly leans toward estimates based upon

Ar (Marty, 2012) or C/N ratio (Marty et al., 2020) terrestrial inventories that require the deep convecting mantle to be C-rich, possibly >10 more enriched than the Depleted Mantle (DMM) that sustains MORB volcanism (~20-56 ppm C; Voyer et al., 2017; Hauri et al., 2019). We need to bear in mind, however, that radiogenic isotopes have clearly proven the presence of a variety of recycled (slab-related) crustal materials in the OIB mantle (e.g., Hofmann and White, 1982; Hofmann et al., 1997; Hoernle et al., 2002; Cabral et al., 2013; Ellam, 2021), and that it is therefore well possible that a substantial fraction of the high C in OIB is in fact derived from a recycled crustal carbon source (Cabral et al., 2014). Carbon isotope results for OIB magmas would be vital to test this hypothesis but are unfortunately available for only a handful set of OIB. Where available, however, results suggest a heavy (crustal) carbon signature in the mantle (Sandoval-Velasquez et al., 2021). This implies that high carbon OIB may in fact reflect local infiltration of C-enriched metasomatic fluids/melts resulting from melting of carbonated lithologies (e.g., carbon-rich sediments and/or carbonated oceanic lithosphere) entrained into the mantle during past subduction events (Dasgupta, 2018). Such a situation may indeed explain the enriched nature of Fogo magmas.

Secondly, our results with those recently obtained elsewhere (Boudoire et al., 2018; Taracsák et al., 2019) are critical to refining estimates of the OIB contribution to deep carbon flux to the atmosphere. Previous work (Dasgupta and Hirschmann, 2010; Hauri et al., 2019) has found carbon outgassing from our planet interior to be primarily controlled by the volumetrically dominant MORB volcanism, and by less voluminous but more C-rich calcalkaline volcanism. However, MORB magmas are CO<sub>2</sub>-poor (typically <<0.1 wt.%; see Fig. 10), while calcalkaline magma CO<sub>2</sub> contents lie somewhere between 0.1 and 1 wt % CO<sub>2</sub> (Plank and Manning, 2019). If the majority of OIB is as C-rich as Fogo, OIB volcanism is expected to contribute disproportionately (relative to other forms of volcanism) to the deep carbon flux to the atmosphere. Previous inventories (Dasgupta and Hirschmann, 2010) have found the carbon flux through OIB volcanism to be highly difficult to estimate, reflecting the often degassed (C-depleted) nature of OIB-carried MI (especially when no correction for shrinkage bubbles CO<sub>2</sub> is applied), and because of their extreme heterogeneity (in terms of depleted vs. enriched compositions). Because of these complexities, Dasgupta and Hirschmann (2010) conservatively derived their OIB carbon flux (1.2-30 Tg/yr) by assuming it to be ~10–50% of the MORB C flux. If for simplicity we assume global OIB to be dominantly alkaline in nature and to contain 2.15 wt. % CO<sub>2</sub> as our Fogo parental melt, then with a cumulative OIB magma extrusive output of 0.3-0.4 km<sup>3</sup>/yr (Crisp, 1984) the global OIB carbon flux would be 16-21 Tg/yr, and > 100 Tg/yr if degassing of intrusive magma (1.5-2 km<sup>3</sup>/yr; Crisp, 1984) is also taken into account.

These initial calculations are subject to large uncertainties. For example, tholeiitic OIB (e.g., Hawaii) or near-ridge hot-spot related OIB (e.g., Iceland) are less CO<sub>2</sub>-poor than alkali rich OIB (Fig. 10), and the fact that all alkali rich OIB are as C-rich as Fogo (or El Hierro; Taracsák et al., 2019) is virtually untested. Ultimately, the volumetric proportions of enriched and non-enriched OIB would need to be constrained with a reasonable level of detail. Thus, our calculations here tentatively indicate that the global OIB carbon flux might be significantly higher than previously thought.

## 6. Conclusions

Olivine-hosted MI from recent ( $\leq 10$  ka) Fogo basanitic tephra samples exhibit some of the highest (glass + bubbles) CO<sub>2</sub> contents (up to 2.15 wt. %) on record. The Fogo MI are also found to additionally contain up to 2.73 wt. % H<sub>2</sub>O, 8205 ppm S, 1459 ppm Cl and 2249 ppm F. Our results are based on the use of three independent methodologies to correct for CO<sub>2</sub> contained in shrinkage bubbles, whose general agreement corroborates our reconstructed total CO<sub>2</sub> concentrations. The inferred MI CO<sub>2</sub>/Nb (277-283) and CO<sub>2</sub>/Ba (26-22) ratios are lower than the characteristic mantle

ratios, suggesting our MI – although exceptionally CO<sub>2</sub>-rich - do still underestimate the real CO<sub>2</sub> contents in primary melts (potentially containing up to 8-9 wt % CO<sub>2</sub>; Sun and Dasgupta, 2023) due to pre-entrapment CO<sub>2</sub> loss to bubbles. This notwithstanding, our results suggest that the basanitic magmas sustaining Fogo activity may originate from a metasomatized, carbon-rich mantle source containing a minimum of 355-414 ppm C, in agreement with recent estimates for global OIB mantle sources. Prior to eruption, these parental basanitic magmas would remain stored in the lithospheric mantle in the ~27 to ~36 km depth range, deeper than previously inferred (Hildner et al., 2011, 2012; Klügel et al., 2020; Lo Forte et al., 2023).

Our results demonstrate the importance of measuring shrinkage bubble CO<sub>2</sub> to correctly reconstructing the architecture of magmatic plumbing systems, and confirm the exceptional carbon-rich signature of alkali rich OIB magmas and their carbon-rich mantle sources. In spite of their relatively low eruptive volumes, OIB eruptions may contribute disproportionately to global volcanic carbon outgassing from the mantle into the Earth's atmosphere. We tentatively estimate the global OIB carbon flux to potentially be as high as 17-23 Tg/yr.

## Acknowledgements

This study was funded by the Deep Carbon Observatory (Alfred P. Sloan Foundation, USA), the Ministero dell'Università e Ricerca (Italy, Grants n. 2017LMNLAW and 2022HA8XCS), and Fundação para a Ciência e Tecnologia, Portugal (MAGAT project, ref. CIRCNA/OCT/0016/2019). LMV and IVAR are acknowledged for providing Francesco Maria Lo Forte access to laboratories. Dr. Jean-Luc Devidal (LMV) is acknowledged for assistance during electron probe analyses. Guilherme Bettencourt (IVAR) helped during sampling fieldwork at Fogo. The authors wish to acknowledge constructive comments from the Editor (Jeffrey G. Catalano), the Associate Editor (Adreas Stracke), M. Portnyagin, and one anonymous reviewer.

## Data Availability

Data are available through Mendeley Data at <https://doi:10.17632/gr2dy9grdn.1>.

## Appendix A. Supplementary Material

The Supplementary Material comprises an excel file labelled as Supplementary Material S1, containing multiple sheets with detailed data information: Whole rock; Sample ubication; Host olivine analysis; Scoria analysis; Uncorrected MIs analysis; Corrected MIs analysis; Total MI CO<sub>2</sub> contents; MI Saturation Pressures. Additionally, there is a .pdf file labelled as Supplementary Material S2, which includes the calibration of nano-SIMS standards (Figure S1) and the Raman tomography of MI Fog 43-6-2 (Figure S2).

## References

- Aiuppa, A., Allard, P., Bernard, B., Lo Forte, F.M., Moretti, R., Hidalgo, S., 2022. Gas Leakage from Shallow Ponding Magma and Trapdoor Faulting at Sierra Negra Volcano (Isabela Island, Galápagos). *Geochem. Geophys. Geosyst.* 23.
- Aiuppa, A., Bertagnini, A., Métrich, N., Moretti, R., Di Muro, A., Liuzzo, M., Tamburello, G., 2010. A model of degassing for Stromboli volcano. *Earth and Planet. Sci. Lett.* 295, 195–204.
- Aiuppa, A., Bitetto, M., Francofonte, V., Velasquez, G., Parra, C.B., Giudice, G., et al., 2017. A CO<sub>2</sub>-gas precursor to the March 2015 Villarrica volcano eruption. *Geochem. Geophys. Geosyst.* 18, 2120–2132.
- Aiuppa, A., Casetta, F., Coltorti, M., Stagno, V., Tamburello, G., 2021. Carbon concentration increases with depth of melting in Earth's upper mantle. *Nat. Geosci.* 14, 697–703.
- Aiuppa, A., Moretti, R., Federico, C., Giudice, G., Gurrieri, S., Liuzzo, M., Papale, P., Shinohara, H., Valenza, M., 2007. Forecasting Etna eruptions by real-time observation of volcanic gas composition. *Geol.* 35, 1115.
- Aiuppa, A., Ronchi, A., Bitetto, M., Rizzo, A.L., Viveiros, F., Allard, P., Frezzotti, M.L., Valenti, V., Zanon, V., 2020. The fumarolic CO<sub>2</sub> output from Pico do Fogo volcano (Cape Verde). *Ital. J. Geosci.* 139, 325–340.
- Anderson, A.T., 1974. Evidence for a Picritic, Volatile-rich Magma beneath Mt. J. *Petrol.* 15, 243–267.
- Anderson, K.R., Poland, M.P., 2017. Abundant carbon in the mantle beneath Hawaii. *Nat. Geosci.* 10: 704–708
- Ariskin, A.A., Barmina, G.S., 1999. An empirical model for the calculation of spinel-melt equilibria in mafic igneous systems at atmospheric pressure: 2. Fe-Ti oxides. *Contrib. Mineral. Petrol.* 134, 251–263.
- Aster, E.M., Wallace, P.J., Moore, L.R., Watkins, J., Gazel, E., Bodnar, R.J., 2016. Reconstructing CO<sub>2</sub> concentrations in basaltic melt inclusions using Raman analysis of vapor bubbles. *J. Volcanol. Geotherm. Res.* 323, 148–162.
- Bali, E., Hartley, M.E., Halldórsson, S.A., Gudfinnsson, G.H., Jakobsson, S., 2018. Melt inclusion constraints on volatile systematics and degassing history of the 2014–2015 Holuhraun eruption, Iceland. *Contrib. Mineral. Petrol.* 173, 9.
- Bamber, E.C., La Spina, G., Arzilli, F., de' Michieli Vitturi, M., Polacci, M., Hartley, M.E., et al., 2022. Basaltic Plinian eruptions at Las Sierras-Masaya volcano driven by cool storage of crystal-rich magmas. *Commun. Earth Environ.* 3, 253.
- Blundy, J., Cashman, K., 2005. Rapid decompression-driven crystallization recorded by melt inclusions from Mount St. Helens volcano. *Geol.* 33, 793.
- Bonadiman, C., Beccaluva, L., Coltorti, M., Siena, F., 2005. Kimberlite-like Metasomatism and 'Garnet Signature' in Spinel-peridotite Xenoliths from Sal, Cape Verde Archipelago: relics of a subcontinental mantle domain within the atlantic oceanic lithosphere? *J. Petrol.* 46, 2465–2493.

- Boudoire G., Padeloup G., Schiavi F., Cluzel N., Rafflin V., Grassa F., et al., 2023. Magma storage and degassing beneath the youngest volcanoes of the Massif Central (France): lessons for the monitoring of a dormant volcanic province. *Chem Geol.*, 634, 121603.
- Boudoire, G., Rizzo, A.L., Di Muro, A., Grassa, F., Liuzzo, M., 2018. Extensive CO<sub>2</sub> degassing in the upper mantle beneath oceanic basaltic volcanoes: First insights from Piton de la Fournaise volcano (La Réunion Island). *Geochim. et Cosmochim. Acta* 235, 376–401.
- Boulard, E., Menguy, N., Auzende, A. L., Benzerara, K., Bureau, H., Antonangeli, D., et al., 2012. Experimental Investigation of the Stability of Fe-Rich Carbonates in the Lower Mantle. *J. Geophys. Res.* 117, B02208.
- Boulanger, M., France, L., 2023. Cumulate Formation and Melt Extraction from Mush-Dominated Magma Reservoirs: The Melt Flush Process Exemplified at Mid-Ocean Ridges. *J. Petrol.* 64, egad005.
- Burgisser, A., Alletti, M., Scaillet, B., 2015. Simulating the behavior of volatiles belonging to the C–O–H–S system in silicate melts under magmatic conditions with the software D-Compress. *Comput. Geosci.* 79, 1–14.
- Burton, M., Aiuppa, A., Allard, P., Asensio-Ramos, M., Confrades, A.P., La Spina, A., et al., 2023. Exceptional eruptive CO<sub>2</sub> emissions from intra-plate alkaline magmatism in the Canary volcanic archipelago. *Commun. Earth Environ.* 4, 467.
- Buso, R., Laporte, D., Schiavi, F., Cluzel, N., Fonquernie, C., 2022. High-pressure homogenization of olivine-hosted CO<sub>2</sub>-rich melt inclusions in a piston cylinder: insight into the volatile content of primary mantle melts. *Eur. J. Mineral.*, 34, 325–349.
- Cabral, R. A., Jackson, M. G., Koga, K. T., Rose-Koga, E. F., Hauri, E. H., M. J. Whitehouse et al., 2014. Volatile cycling of H<sub>2</sub>O, CO<sub>2</sub>, F, and Cl in the HIMU mantle: A new window provided by melt inclusions from oceanic hot spot lavas at Mangaia, Cook Islands, *Geochem. Geophys. Geosyst.*, 15, 4445–4467.
- Cabral, R. A., Jackson, M. G., Rose-Koga, E. F., Koga, K. T., Whitehouse, M. J., Antonelli, M. A., et al., 2013. Anomalous sulphur isotopes in plume lavas reveal deep mantle storage of Archaean crust. *Nature* 496, 490–493
- Cannatelli, C., Doherty, A.L., Esposito, R., Lima, A., De Vivo, B., 2016. Understanding a volcano through a droplet: A melt inclusion approach. *J. of Geochem. Explor.* 171, 4–19.
- Carroll, M.R., and Webster, J.D., 1994. Sulfur, noble gases, and halogens: solubility relations of the less abundant volatile species in magmas. *Rev. Mineral. Geochem.* 30, 231-79.
- Cartigny, P., Pineau, F., Aubaud, C., Javoy, M., 2008. Towards a consistent mantle carbon flux estimate: Insights from volatile systematics (H<sub>2</sub>O/Ce, δD, CO<sub>2</sub>/Nb) in the North Atlantic mantle (14° N and 34° N). *Earth and Planet. Sci. Lett.* 265, 672–685.
- Chen, Y., Provost, A., Schiano, P., and Cluzel, N., 2011. The rate of water loss from olivine-hosted melt inclusions. *Contrib. Mineral. Petrol.*, 162(3), 625–636.
- Cottrell, E., Spiegelman, M., Langmuir, C.H., 2002. Consequences of diffusive reequilibration for the interpretation of melt inclusions: reequilibration of melt inclusions. *Geochem. Geophys. Geosyst.* 3, 1–26.

- Créon, L., Levrèsse, G., Remusat, L., Bureau, H., Carrasco-Nuñez, G. 2018. New method for initial composition determination of crystallized silicate melt inclusions. *Chem. Geol.* 483, 162–173.
- Crisp, J. A. 1984. Rates of magma emplacement and volcanic output, *J. Volcanol. Geotherm. Res.*, 20, 177 – 211.
- Danyushevsky, L.V., 2001. The effect of small amounts of H<sub>2</sub>O on crystallisation of mid-ocean ridge and backarc basin magmas. *J. Volcanol. and Geother. Res.* 110, 265–280.
- Danyushevsky, L.V., Della-Pasqua, F.N., Sokolov, S., 2000. Re-equilibration of melt inclusions trapped by magnesian olivine phenocrysts from subduction-related magmas: petrological implications. *Contrib. Mineral. Petrol.* 138, 68–83.
- Danyushevsky, L.V., McNeill, A.W., Sobolev, A.V., 2002. Experimental and petrological studies of melt inclusions in phenocrysts from mantle-derived magmas: an overview of techniques, advantages and complications. *Chem. Geol.* 183, 5–24.
- Danyushevsky, L.V., Plechov, P., 2011. Petrolog3: Integrated software for modeling crystallization processes: Petrolog3. *Geochem. Geophys. Geosyst.* 12.
- Dasgupta, R., 2018 Volatile-bearing partial melts beneath oceans and continents – where, how much, and of what compositions? *Am. J. Sci.* 318, 141–165.
- Dasgupta, R., Chi, H., Shimizu, N., Buono, A.S., Walker, D., 2013. Carbon solution and partitioning between metallic and silicate melts in a shallow magma ocean: implications for the origin and distribution of terrestrial carbon. *Geochim. Cosmochim. Acta* 102, 191–212.
- Dasgupta, R., Hirschmann, M.M., 2010. The deep carbon cycle and melting in Earth's interior. *Earth and Planet. Sci. Lett.* 298, 1–13.
- Day, S.J., Heleno da Silva, S.I.N., Fonseca, J.F.B.D., 1999. A past giant lateral collapse and present-day flank instability of Fogo, Cape Verde Islands. *J. Volcanol. and Geother. Res.* 94, 191–218.
- de Moor, J.M., Aiuppa, A., Avaró, G., Wehrmann, H., Dunbar, N., Müller, C., et al., 2016. Turmoil at Turrialba Volcano (Costa Rica): Degassing and eruptive processes inferred from high-frequency gas monitoring: Turmoil at Turrialba Volcano. *J. Geophys. Res. Solid Earth* 121, 5761–5775.
- Di Martino, C., Forni, F., Frezzotti, M.L., Palmeri, R., Webster, J.D., Ayuso, R.A., et al., 2011. Formation of cordierite-bearing lavas during anatexis in the lower crust beneath Lipari Island (Aeolian arc, Italy). *Contrib. Mineral. Petrol.* 162, 1011–1030.
- Di Muro, A., Metrich, N., Vergani, D., Rosi, M., Armienti, P., Fougéroux, T., Deloué, E., Arienzo, I., Civetta, L., 2014. The Shallow Plumbing System of Piton de la Fournaise Volcano (La Réunion Island, Indian Ocean) Revealed by the Major 2007 Caldera-Forming Eruption. *J. Petrol.* 55, 1287–1315.
- Dixon J.E. and Stolper E.M., 1995. An Experimental Study of Water and Carbon Dioxide Solubilities in Mid-Ocean Ridge Basaltic Liquids. Part II: Applications to Degassing. *J. Petrol.* 36, 1633–1646



- Doucelance, R., Escrig, S., Moreira, M., Gariépy, C., Kurz, M.D., 2003. Pb-Sr-He isotope and trace element geochemistry of the Cape Verde Archipelago. *Geochim. Cosmochim. Acta* 67, 3717–3733.
- Edmonds, M., Aiuppa, A., Humphreys, M., Moretti, R., Giudice, G., Martin, R.S., et al., 2010. Excess volatiles supplied by mingling of mafic magma at an andesite arc volcano: excess volatiles supplied by underplating. *Geochem. Geophys. Geosyst.* 11.
- Edmonds, M., Liu, E.J., Cashman, K.V., 2022. Open-vent volcanoes fuelled by depth-integrated magma degassing. *Bull. Volcanol.* 84, 28.
- Edmonds M., and Wallace P.J., 2017. Volatiles and Exsolved Vapor in Volcanic Systems. *Elements*, 13, 29–34.
- Eguchi, J. and Dasgupta, R., 2018. A CO<sub>2</sub> solubility model for silicate melts from fluid saturation to graphite or diamond saturation, *Chem. Geol.*, 487, 23–38.
- Ellam, R. M. 2021. Radiogenic isotopes and mantle evolution. In: Alderton, D. and Elias, S. A. (eds.) *Encyclopedia of Geology*. Academic Press: Amsterdam, 330-344
- Escrig, S., Doucelance, R., Moreira, M., Allègre, C.J., 2005. Os isotope systematics in Fogo Island: Evidence for lower continental crust fragments under the Cape Verde Southern Islands. *Chem. Geol.* 219, 93–113.
- Esposito, R., Bodnar, R.J., Danyushevsky, L., De Vivo, B., Fedele, L., Hunter, J., Lima, A., and Shimizu, N. 2011. Volatile evolution of magma associated with the Solchiaro Eruption in the Phlegrean Volcanic District (Italy). *Journal of Petrology.* 52, 2431–2460.
- Esposito, R., Lamadrid, H.M., Redi, D., Steele-MacInnis, M., Bodnar, R.J., Manning, C.E., De Vivo, B., Cannatelli, C., Lima, A., 2016. Detection of liquid H<sub>2</sub>O in vapor bubbles in reheated melt inclusions: Implications for magmatic fluid composition and volatile budgets of magmas? *Am. Mineral.* 101, 1691–1695.
- Fall, A., Bodnar, R.J., Szabó, C., Pál-Molnár, E. 2007. Fluid evolution in the nepheline syenites of the Ditrău alkaline Massif, Transylvania, Romania. *Lithos*, 95, 331-345.
- Feignon J.G., Cluzel N., Schiavi F., Moune S., Roche O., Clavero J., et al., 2022. High CO<sub>2</sub> content in magmas of the explosive andesitic Enco eruption of Mocho-Choshuenco volcano (Chile). *Bull. Volcanol.* 84, 40.
- Foeken, J.P.T., Day, S., Stuart, F.M., 2009. Cosmogenic <sup>3</sup>He exposure dating of the Quaternary basalts from Fogo, Cape Verdes: Implications for rift zone and magmatic reorganisation. *Quaternary Geochronology* 4, 37–49.
- Foley, S.F. 2011. A Reappraisal of Redox Melting in the Earth's Mantle as a Function of Tectonic Setting and Time, *J. Petrol.*, 52, 7-8, 1363–1391
- Foley, S. F., Fischer, T. P. 2017. An essential role for continental rifts and lithosphere in the deep carbon cycle. *Nat. Geosci.* 10, 897–902.
- Foley, S. F., Link, K., Tiberindwa, J. V. & Barifaijo, E. 2012. Patterns and origin of igneous activity around the Tanzanian craton. *J. Afr. Earth Sci.* 62, 1–18.

- Foley, S.F., Yaxley, G.M., Rosenthal, A., Buhre, S., Kiseeva, E.S., Rapp, R.P., Jacob, D.E. 2009. The composition of near-solidus melts of peridotite in the presence of CO<sub>2</sub> and H<sub>2</sub>O between 40 and 60 kbar. *Lithos.* 112S, 274-283.
- French, S., and Romanowicz, B., 2015. Broad plumes rooted at the base of the Earth's mantle beneath major hotspots, *Nature*, 525, 95–99.
- Gaetani, G.A., O'Leary, J.A., Shimizu, N., Bucholz, C.E., and Newville, M., 2012. Rapid reequilibration of H<sub>2</sub>O and oxygen fugacity in olivine-hosted melt inclusions. *Geol.*, 40(10), 915–918.
- Gaetani, G.A., and Watson, E.B., 2000. Open system behavior of olivine-hosted melt inclusions. *Earth and Planet. Sci. Lett.* 183, 27–41.
- Gaetani, G.A., and Watson, E.B., 2002. Modeling the major-element evolution of olivine-hosted melt inclusions. *Chem. Geol.* 183, 25–41.
- Gerlach, D.C., Cliff, R.A., Davies, G.R., Norry, M., Hodgson, N., 1988. Magma sources of the Cape Verdes archipelago: Isotopic and trace element constraints. *Geochim. et Cosmochim. Acta* 52, 2979–2992.
- Ghiorso, M.S., and Gualda, G.A.R., 2015. An H<sub>2</sub>O–CO<sub>2</sub> mixed fluid saturation model compatible with rhyolite-MELTS. *Contrib. Mineral. Petrol.* 169, 53.
- Giordano, D., Nichols, A.R., and Dingwell, D.B. 2005. Glass transition temperatures of natural hydrous melts: a relationship with shear viscosity and implications for the welding process. *J. Volcanol. and Geother. Res.* , 142 (1–2), 105–118.
- Guzmics, T., Berkesi, M., Bodnar, R.J., Fall, A., Bali, E., Milke, R., Vetlényi, E., Szabó, C., 2019. Natrocarbonatites: A hidden product of three-phase immiscibility. *Geol.* 47, 527–530.
- Halldórsson, S.A., Marshall, E.W., Caracciolo, A., Matthews, S., Bali, E., Rasmussen, M.B., et al., 2022. Rapid shifting of a deep magmatic source at Fagradalsfjall volcano, Iceland. *Nature* 609, 529–534.
- Harms, E., and Schmincke, H.U., 2000. Volatile composition of the phonolitic Laacher See magma (12,900 yr BP): implications for syn-eruptive degassing of S, F, Cl and H<sub>2</sub>O. *Contrib. Mineral. Petrol.* 138, 84–98.
- Hartley, M.E., Maclennan, J., Edmonds, M., Thordarson, T., 2014. Reconstructing the deep CO<sub>2</sub> degassing behaviour of large basaltic fissure eruptions. *Earth and Planet. Sci. Lett.* 393, 120–131.
- Hauri, E., 2002. SIMS analysis of volatiles in silicate glasses, 2: isotopes and abundances in Hawaiian melt inclusions. *Chem. Geol.* 183, 115–141.
- Hauri, E.H., Cottrell, E., Kelley, K.A., Tucker, J.M., Shimizu, K., Voyer, M.L., Marske, J., Saal, A.E., 2019. Carbon in the Convecting Mantle, in: Orcutt, B.N., Daniel, I., Dasgupta, R. (Eds.), *Deep Carbon*. Cambridge University Press, pp. 237–275.
- Hauri, E.H., Maclennan, J., McKenzie, D., Gronvold, K., Oskarsson, N., Shimizu, N., 2018. CO<sub>2</sub> content beneath northern Iceland and the variability of mantle carbon. *Geol.* 46, 55–58.

- Heath, E., Macdonald, R., Belkin, H., Hawkesworth, C., Sigurdsson, H., 1998. Magma genesis at Soufriere Volcano, St Vincent, Lesser Antilles Arc. *J. Petrol.* 39, 1721–1764.
- Hekinian, R., and D. Walker (1987), Diversity and spatial zonation of volcanic-rocks from the East Pacific Rise near 21-degrees-N, *Contrib. Mineral. Petrol.*, 96 (3), 265–280
- Hildner, E., Klügel, A., Hansteen, T.H., 2012. Barometry of lavas from the 1951 eruption of Fogo, Cape Verde Islands: Implications for historic and prehistoric magma plumbing systems. *J. Volcanol. and Geother. Res.* 217–218, 73–90.
- Hildner, E., Klügel, A., Hauff, F., 2011. Magma storage and ascent during the 1995 eruption of Fogo, Cape Verde Archipelago. *Contrib. Mineral. Petrol.* 162, 751–772.
- Hirschmann, M.M. 2006. Water, melting, and the deep Earth H<sub>2</sub>O cycle *Annu. Rev. Earth Planet. Sci.* 629-653.
- Hirschmann, M.M., 2018. Comparative deep Earth volatile cycles: The case for C recycling from exosphere/mantle fractionation of major (H<sub>2</sub>O, C, N) volatiles and from H<sub>2</sub>O/Ce, CO<sub>2</sub>/Ba, and CO<sub>2</sub>/Nb exosphere ratios. *Earth and Planet. Sci. Lett.* 502, 262–273.
- Hoernle, K., Tilton, G., Le Bas, M.J., Duggen, S., Garbe-Schönberg, D., 2002. Geochemistry of oceanic carbonatites compared with continental carbonatites: mantle recycling of oceanic crustal carbonate. *Contrib. Mineral. Petrol.* 142, 520–542.
- Hofmann, A. W, 1997. Mantle geochemistry: the message from oceanic volcanism. *Nature* 385, 219–229.
- Hofmann, A.W., White. W.M., 1982. Mantle plumes from ancient oceanic crust *Earth Planet. Sci. Lett.*, 57 (2), 421-436
- Holm, P. M., Wilson, J.R., Christensen B.P., Hansen, L., Hansen, S. L., Hein, K.M., et al., 2006, Sampling the Cape Verde Mantle Plume: Evolution of Melt Compositions on Santo Antão, Cape Verde Islands, *J. Petrol.*, 47, 1, 145–189.
- Hudgins, T.R., Mukasa, S.B., Simon, A.C., Moore, G., Barifajjo, E. 2015. Melt inclusion evidence for CO<sub>2</sub>-rich melts beneath the western branch of the East African Rift: implications for long-term storage of volatiles in the deep lithospheric mantle *Contrib. Mineral. Petrol.*, 169, 1-18.
- Iacono-Marziano, G., Morizet, Y., Le Trong, E., Gaillard, F., 2012. New experimental data and semi-empirical parameterization of H<sub>2</sub>O–CO<sub>2</sub> solubility in mafic melts. *Geochim. Cosmochim. Acta* 97, 1–23.
- Jackson, M.G., Becker, T.W., Konter, J.G., 2018a. Evidence for a deep mantle source for EM and HIMU domains from integrated geochemical and geophysical constraints. *Earth Planet. Sci. Lett.* 484, 154-167
- Jiménez-Mejías, M., Andújar, J., Scaillet, B., Casillas, R., 2021. Experimental determination of H<sub>2</sub>O and CO<sub>2</sub> solubilities of mafic alkaline magmas from Canary Islands. *Comptes Rendus. Géoscience* 353, 289–314.

- Kamenetsky, V.S., Davidson, P., Mernagh, T.P., Crawford, A.J., Gemmell, J.B., Portnyagin, M.V., Shinjo, R., 2002. Fluid bubbles in melt inclusions and pillow-rim glasses: high-temperature precursors to hydrothermal fluids? *Chem. Geol.* 183, 349–364.
- Kawakami, Y., Yamamoto, J., Kagi, H., 2003. Micro-Raman densimeter for CO<sub>2</sub> inclusions in mantle-derived minerals. *Appl. Spectrosc.* 57,11, 1333–1339.
- Kendrick, M.A., Jackson, M.G., Kent, A.J.R., Hauri, E.H., Wallace, P.J., Woodhead, J., 2014. Contrasting behaviours of CO<sub>2</sub>, S, H<sub>2</sub>O and halogens (F, Cl, Br, and I) in enriched-mantle melts from Pitcairn and Society seamounts. *Chem. Geol.* 370, 69–81.
- Kent, A.J.R., 2008. Melt inclusions in basaltic and related volcanic rocks. In: Putirka, K.D., Tepley III, F.J. (Eds.) *Minerals, inclusions and volcanic processes. Reviews in Mineralogy and Petrology* 69, 273–332.
- Klügel, A., Day, S., Schmid, M., Faria, B., 2020. Magma Plumbing During the 2014–2015 Eruption of Fogo (Cape Verde Islands). *Front. Earth Sci.* 8, 157.
- Koleszar, A.M., Saal, A.E., Hauri, E.H., Nagle, A.N., Liang, Y., Kurz, M.D., 2009. The volatile contents of the Galapagos plume; evidence for H<sub>2</sub>O and F open system behavior in melt inclusions. *Earth and Planet. Sci. Lett.* 287, 442–452.
- Le Bas, M.J.L., Maitre, R.W.L., Streckeisen, A., Zanettin, B., 1986. IUGS Subcommittee on the Systematics of Igneous Rocks. A Chemical Classification of Volcanic Rocks Based on the Total Alkali-Silica Diagram. *J. Petrol.* 27,3, 745–750.
- Le Voyer, M., Hauri, E.H., Cottrell, E., Kelley, K.A., Salters, V.J.M., Langmuir, C.H., Hilton, D.R., Barry, P.H., Füre, E., 2019. Carbon Fluxes and Primary Magma CO<sub>2</sub> Contents Along the Global Mid-Ocean Ridge System. *Geochem. Geophys. Geosyst.* 20, 1387–1424.
- Le Voyer, M., Kelley, K.A., Cottrell, E., Hauri, E.H., 2017. Heterogeneity in mantle carbon content from CO<sub>2</sub>-undersaturated basalts. *Nat. Commun.* 8, 14062.
- Leshner, C.E., Spera, F.J., 2015. Thermodynamic and Transport Properties of Silicate Melts and Magma, in: *The Encyclopedia of Volcanoes*. Elsevier, pp. 113–141.
- Liu, B., Zhang, Z., Giuliani, A., Xie, Q., Kong, W., Wang, C., et al., 2023. A Mantle Plume Connection for Alkaline Lamprophyres (Sannaïtes) from the Permian Tarim Large Igneous Province: Petrological, Geochemical and Isotopic Constraints. *J. Petrol.* 64, egad004.
- Liu, X., Zhao, D., 2021. Seismic evidence for a plume-modified oceanic lithosphere–asthenosphere system beneath Cape Verde. *Geophys. J. Int.* 225, 872–886.
- Lo Forte, F.M., Aiuppa, A., Rotolo, S.G., Zanon, V., 2023. Temporal evolution of the Fogo Volcano magma storage system (Cape Verde Archipelago): a fluid inclusions perspective. *J. Volcanol. and Geother. Res.* 433, 107730.
- Longpré, M.-A., Stix, J., Klügel, A., Shimizu, N., 2017. Mantle to surface degassing of carbon- and sulphur-rich alkaline magma at El Hierro, Canary Islands. *Earth and Planet. Sci. Lett.* 460, 268–280.

- Lowenstern, J.B. 1995. Applications of silicate MIs to the study of magmatic volatiles. In: *Magma, Fluids, and Ore Deposits*, 23 (ed. J. F. H. Thompson). Mineralogical Society of Canada, , 71–99.
- Lowenstern, J.B., 2003. Melt inclusions come of age: volatiles, volcanoes, and sorby's legacy. *Develop. Volcanol.* 5, 1–21.
- Lowenstern, J. B., 2015. Bursting the bubble of melt inclusions, *Am. Mineral.*, 100, 672–673.
- MacLennan, J., 2017. Bubble formation and decrepitation control the CO<sub>2</sub> content of olivine-hosted melt inclusions: melt inclusion carbon dioxide. *Geochem. Geophys. Geosyst.* 18, 597–616.
- Marini, L., Moretti, R., Accornero, M., 2011. Sulfur Isotopes in Magmatic-Hydrothermal Systems, Melts, and Magmas. *Reviews in Mineralogy and Geochemistry* 73, 423–492.
- Martínez-Moreno, F.J., Monteiro Santos, F.A., Madeira, J., Pous, J., Bernardo, I., Soares, A., et al., 2018. Investigating collapse structures in oceanic islands using magnetotelluric surveys: The case of Fogo Island in Cape Verde. *J. Volcanol. and Geother. Res.* 357, 152–162.
- Martins, S., Mata, J., Munhá, J., Mendes, M.H., Maerschalk, C., Caldeira, R., et al., 2010. Chemical and mineralogical evidence of the occurrence of mantle metasomatism by carbonate-rich melts in an oceanic environment (Santiago Island, Cape Verde). *Miner. Petrol.* 99, 43–65.
- Marty, B., 2012. The origins and concentrations of water, carbon, nitrogen and noble gases on Earth. *Earth Planet. Sci. Lett.* 313, 56–66.
- Marty, B., Almayrac, M., Barry, P.H., Bekaert, D.V., Broadley, M.W., Byrne, D.J., et al., 2020. An evaluation of the C/N ratio of the mantle from natural CO<sub>2</sub>-rich gas analysis: Geochemical and cosmochemical implications. *Earth Planet. Sci. Lett.* 551, 116574.
- Massare, D., Métrich, N., and Clocchiatti, R. 2002. High-temperature experiments on silicate melt inclusions in olivine at 1 atm: Inference on temperatures of homogenization and H<sub>2</sub>O concentrations. *Chem. Geol.*, 183(1–4), 87–98.
- Mata, J., Martins, S., Mattielli, N., Madeira, J., Faria, B., Ramalho, R.S., et al., 2017. The 2014–15 eruption and the short-term geochemical evolution of the Fogo volcano (Cape Verde): Evidence for small-scale mantle heterogeneity. *Lithos* 288–289, 91–107.
- Melián, G.V., Hernández, P.A., Pérez, N.M., Asensio-Ramos, M., Padrón, E., Alonso, M., et al., 2021. Insights from Fumarole Gas Geochemistry on the Recent Volcanic Unrest of Pico do Fogo, Cape Verde. *Front. Earth Sci.* 9:631190.
- Métrich, N., Allard, P., Aiuppa, A., Bani, P., Bertagnini, A., Shinohara, H., et al., 2011. Magma and Volatile Supply to Post-collapse Volcanism and Block Resurgence in Siwi Caldera (Tanna Island, Vanuatu Arc). *J. of Petrol.* 52, 1077–1105.
- Métrich, N., Allard, P., Spilliaert, N., Andronico, D., Burton, M., 2004. 2001 flank eruption of the alkali- and volatile-rich primitive basalt responsible for Mount Etna's evolution in the last three decades. *Earth Planet. Sci. Lett.* 228, 1–17.
- Métrich, N., Clocchiatti, R., Mosbah, M., Chaussidon, M., 1993. The 1989–1990 activity of Etna magma mingling and ascent of H<sub>2</sub>O, Cl, S rich basaltic magma. Evidence from melt inclusions. *J. Volcanol. and Geother. Res.* 59, 131–144.

- Métrich, N., and Wallace, P.J., 2008. Volatile Abundances in Basaltic Magmas and Their Degassing Paths Tracked by Melt Inclusions. *Reviews in Mineralogy and Geochemistry* 69, 363–402.
- Métrich, N., Zanon, V., Créon, L., Hildenbrand, A., Moreira, M., Marques, F.O., 2014. Is the ‘Azores Hotspot’ a Wetspot? Insights from the Geochemistry of Fluid and Melt Inclusions in Olivine of Pico Basalts. *J. Petrol.* 55, 377–393.
- Michael, P.J., Graham, D.W., 2015. The behavior and concentration of CO<sub>2</sub> in the suboceanic mantle: Inferences from undegassed ocean ridge and ocean island basalts. *Lithos* 236–237, 338–351.
- Miller, W.G.R., Maclennan, J., Shorttle, O., Gaetani, G.A., Le Roux, V., Klein, F., 2019. Estimating the carbon content of the deep mantle with Icelandic melt inclusions. *Earth Planet. Sci. Lett.* 523, 115699.
- Mironov, N.L., Portnyagin, M.V., 2011. H<sub>2</sub>O and CO<sub>2</sub> in parental magmas of Kliuchevskoi volcano inferred from study of melt and fluid inclusions in olivine. *Russian geology and geophysics* 52, 1353-1367.
- Mironov, N., Portnyagin, M., Botcharnikov, R., Gurenko, A., Hoernle, K., Holtz, F., 2015. Quantification of the CO<sub>2</sub> budget and H<sub>2</sub>O–CO<sub>2</sub> systematics in subduction-zone magmas through the experimental hydration of melt inclusions in olivine at high H<sub>2</sub>O pressure. *Earth and Planetary Science Letters* 425, 1-11.
- Mironov, N., Tobelko, D.P., Smirnov, S.Z., Portnyagin, M.V., Krashennnikov, S.P., 2020. Estimation of CO<sub>2</sub> content in the gas phase of melt inclusions using Raman spectroscopy: Case study of inclusions in olivine from the Karymsky volcano (Kamchatka). *Russian Geology and Geophysics* 61, 600-610.
- Mitchell, M.A., Peacock, J.R., Burgess, S.D., 2023. Imaging the magmatic plumbing of the Clear Lake Volcanic Field using 3-D gravity inversions. *J. Volcanol. and Geother. Res.* 435, 107758.
- Moore, L.R., Bodnar, R.J., 2019. A pedagogical approach to estimating the CO<sub>2</sub> budget of magmas. *JGS* 176, 398–407.
- Moore, L.R., Gazel, E., Tuohy, R., Lloyd, A.S., Esposito, R., Steele-MacInnis, M., et al., 2015. Bubbles matter: An assessment of the contribution of vapor bubbles to melt inclusion volatile budgets. *Am. Mineral.* 100, 806–823.
- Moore, L.R., Mironov, N., Portnyagin, M., Gazel, E., Bodnar, R.J. 2018. Volatile contents of primitive bubble-bearing melt inclusions from Klyuchevskoy volcano, Kamchatka: Comparison of volatile contents determined by mass-balance versus experimental homogenization *J. Volcanol. Geotherm. Res.*, 358, 124-131.
- Moretti, R., Arienzo, I., Civetta, L., Orsi, G., Papale, P., 2013a. Multiple magma degassing sources at an explosive volcano. *Earth and Planet. Sci. Lett.* 367, 95–104.
- Moretti, R., Arienzo, I., Orsi, G., Civetta, L., D’Antonio, M., 2013b. The Deep Plumbing System of Ischia: a Physico-chemical Window on the Fluid-saturated and CO<sub>2</sub>-sustained Neapolitan Volcanism (Southern Italy). *J. Petrol.* 54, 951–984.

- Moretti, R., Papale, P., Ottonello, G., 2003. A model for the saturation of C-O-H-S fluids in silicate melts. *SP* 213, 81–101.
- Mourão, C., Moreira, M., Mata, J., Raquin, A., Madeira, J., 2012. Primary and secondary processes constraining the noble gas isotopic signatures of carbonatites and silicate rocks from Brava Island: evidence for a lower mantle origin of the Cape Verde plume. *Contrib. Mineral. Petrol.* 163, 995–1009.
- Moussallam, Y., Georgeais, G., Rose-Koga, E.F., Koga, K.T., Hartley, M.E., Scaillet, B., Oppenheimer, C., Peters, N., 2023. CO<sub>2</sub> -Undersaturated Melt Inclusions From the South West Indian Ridge Record Surprisingly Uniform Redox Conditions. *Geochem Geophys Geosyst* 24, e2023GC011235.
- Moussallam, Y., Longpré, M.-A., McCammon, C., Gomez-Ulla, A., Rose-Koga, E.F., Scaillet, B., Peters, N., Gennaro, E., Paris, R., Oppenheimer, C., 2019. Mantle plumes are oxidised. *Earth and Planet. Sci. Lett.* 527, 115798.
- Moussallam, Y., Oppenheimer, C., Scaillet, B., Gaillard, F., Kyle, P., Peters, N., Hartley, M., Berlo, K., Donovan, A., 2014. Tracking the changing oxidation state of Erebus magmas, from mantle to surface, driven by magma ascent and degassing. *Earth and Planet. Sci. Lett.* 393, 200–209.
- Naumov, V.B., Portnyagin, M.V., Tolstykh, M.L., Yarmolyuk, V.V., 2003. Composition of Magmatic Melts from the Southern Baikal Volcanic Region: A Study of Inclusions in Olivine from Trachybasalts. *Geochemistry International.* 41, 213–223.
- Newman, S., Lowenstern, J.B., 2002. VolatileCalc: a silicate melt–H<sub>2</sub>O–CO<sub>2</sub> solution model written in Visual Basic for excel. *Computers & Geosciences* 28, 597–604.
- Ni, P., Zhang, Y., Guan, Y., 2017. Volatile loss during homogenization of lunar melt inclusions. *Earth and Planet. Sci. Lett.* 478, 214–224.
- Oppenheimer, C., Fischer, T.P., Scaillet, B., 2014. Volcanic Degassing: Process and Impact, in: *Treatise on Geochemistry*. Elsevier, pp. 111–179.
- Oppenheimer, C., Moretti, R., Kyle, P.R., Eschenbacher, A., Lowenstern, J.B., Hervig, R.L., Dunbar, N.W., 2011. Mantle to surface degassing of alkalic magmas at Erebus volcano, Antarctica. *Earth and Planet. Sci. Lett.* 306, 261–271.
- Palme, H., & O'Neill, H. ST C. 2003. Cosmochemical estimates of mantle composition. *The mantle and core*, 1–38.
- Pérez, W., Freundt, A., Kutterolf, S., 2020. The basaltic plinian eruption of the ~6 ka San Antonio Tephra and formation of the Masaya caldera, Nicaragua. *J. Volcanol. and Geother. Res.* 401, 106975.
- Pim, J., Peirce, C., Watts, A.B., Grevemeyer, I., Krabbenhoft, A., 2008. Crustal structure and origin of the Cape Verde rise. *Earth Planet. Sci. Lett.* 272, 422–428.
- Pino, N.A., Moretti, R., Allard, P., Boschi, E., 2011. Seismic precursors of a basaltic paroxysmal explosion track deep gas accumulation and slug upraise. *J. Geophys. Res.* 116, B02312.
- Plank, T., Manning, C.E., 2019. Subducting carbon. *Nature* 574, 343–352.

- Portnyagin, M., Almeev, R., Matveev, S., and Holtz, F. 2008. Experimental evidence for rapid water exchange between melt inclusions in olivine and host magma. *Earth and Planet. Sci. Lett.*, 272(3), 541–552.
- Portnyagin, M., Mironov, N., Botcharnikov, R., Gurenko, A., Almeev, R.R., Luft, C., et al., 2019. Dehydration of melt inclusions in olivine and implications for the origin of silica-undersaturated island-arc melts. *Earth and Planet. Sci. Lett.*, 517, 95–105.
- Portnyagin, M.V., Naumov, V.B., Mironov, N.L., Belousov, I.A., Kononkova, N.N., 2011. Composition and evolution of the melts erupted in 1996 at Karymskoe Lake, Eastern Kamchatka: Evidence from inclusions in minerals. *Geochem. Int.* 49, 1085–1110.
- Putirka, K.D., Perfit, M., Ryerson, F.J., Jackson, M.G., 2007. Ambient and excess mantle temperatures, olivine thermometry, and active vs. passive upwelling. *Chem. Geol.* 241, 177–206.
- Rasmussen, D.J., Plank, T.A., Roman, D.C., Power, J.A., Bodnar, R.J., and Hauri, E.H. 2018. When does eruption run-up begin? Multidisciplinary insight from the 1999 eruption of Shishaldin volcano. *Earth and Planet. Sci. Lett.*, 486, 1–14.
- Rasmussen, D.J., Plank, T.A., Wallace, P.J., Newcombe, M.E., Lowenstern, J.B., 2020. Vapor-bubble growth in olivine-hosted melt inclusions. *Am. Mineral.* 105, 1898–1919.
- Ribeiro O., (1954), *A Ilha do Fogo e as suas erupções*, Memórias da Junta de Investigação Científica do Ultramar, Série Geográfica, I, Lisboa, 327
- Robidoux, P., Frezzotti, M.L., Hauri, E.H., Aiuppa, A., 2018. Shrinkage Bubbles: The C–O–H–S Magmatic Fluid System at San Cristóbal Volcano. *J. Petrol.* 59, 2093–2122.
- Roedder, E., 1979. Origin and significance of magmatic inclusions. *Bull. Mineral.* 102, 467–510.
- Romero, J.E., Morgado, E., Pisello, A., Boschetty, F., Petrelli, M., Cáceres, F., et al., 2023. Pre-eruptive Conditions of the 3 March 2015 Lava Fountain of Villarrica Volcano (Southern Andes). *Bull. Volcanol.* 85,1, 2.
- Rose-Koga, E.F., Bouvier, A.-S., Gaetani, G.A., Wallace, P.J., Allison, C.M., Andrys, J.A., et al., 2021. Silicate melt inclusions in the new millennium: A review of recommended practices for preparation, analysis, and data presentation. *Chem. Geol.* 570, 120145.
- Rose-Koga, E.F., Koga, K.T., Devidal, J.-L., Shimizu, N., Le Voyer, M., Dalou, C., et al., 2020. In-situ measurements of magmatic volatile elements, F, S, and Cl, by electron microprobe, secondary ion mass spectrometry, and heavy ion elastic recoil detection analysis. *Am. Mineral.* 105, 616–626.
- Rose-Koga, E.F., Koga, K.T., Hamada, M., Hélois, T., Whitehouse, M.J., Shimizu, N., 2014. Volatile (F and Cl) concentrations in Iwate olivine-hosted melt inclusions indicating low-temperature subduction. *Earth, Planets and Space* 66, 81.
- Rose-Koga, E.F., Koga, K.T., Moreira, M., Vlastelic, I., Jackson, M.G., Whitehouse, M.J., et al., 2017. Geochemical systematics of Pb isotopes, fluorine, and sulfur in melt inclusions from São Miguel, Azores. *Chem. Geol.* 458, 22–37.



- Rosenthal, A., Hauri, E.H., Hirschmann, M.M., 2015. Experimental determination of C, F, and H partitioning between mantle minerals and carbonated basalt, CO<sub>2</sub>/Ba and CO<sub>2</sub>/Nb systematics of partial melting, and the CO<sub>2</sub> contents of basaltic source regions. *Earth and Planet. Sci. Lett.* 412, 77–87.
- Rowe, M.C., Peate, D.W., Ukstins Peate, I., 2011. An Investigation into the Nature of the Magmatic Plumbing System at Paricutin Volcano, Mexico. *J. Petrol.* 52, 2187–2220.
- Saal, A.E., Hauri, E.H., Langmuir, C.H., Perfit, M.R., 2002. Vapour undersaturation in primitive mid-ocean-ridge basalt and the volatile content of Earth's upper mantle. *Nature* 419, 451–455.
- Sadofsky, S.J., Portnyagin, M., Hoernle, K., 2008. Subduction cycling of volatiles and trace elements through the Central American volcanic arc: evidence from melt inclusions. *Contrib. Mineral. Petrol.* 155:433–456.
- Sandoval-Velasquez, A., Rizzo, A.L., Aiuppa, A., Remigi, S., Padrón, E., Pérez, N.M., Frezzotti, M.L., 2021. Recycled crustal carbon in the depleted mantle source of El Hierro volcano, Canary Islands. *Lithos* 400–401, 106414. <https://doi.org/10.1016/j.lithos.2021.106414>
- Schiano, P., 2003. Primitive mantle magmas recorded as silicate melt inclusions in igneous minerals. *Earth-Sci. Rev.* 63, 121–144.
- Schiavi, F., Bolfan-Casanova, N., Buso, R., Laumonier, M., Laporte, D., Medjoubi, K., Venugopal, S., Gómez-Ulla, A., Cluzel, N., Hardiagon, M., 2020. Quantifying magmatic volatiles by Raman microtomography of glass inclusion-hosted bubbles. *Geochem. Persp. Lett.* 16, 17–24.
- Sharpe, M.S., Barker, S.J., Rooyackers, S.M., Wilson, C.J.N., Chambefort, I., Rowe, M.C., Schipper, C.I., Charlier, B.L.A., 2022. A sulfur and halogen budget for the large magmatic system beneath Taupō volcano. *Contrib. Mineral. Petrol.* 177, 95.
- Shaw, D.M., 1970. Trace element fractionation during anatexis. *Geochimica et Cosmochimica Acta* 34, 237–243.
- Shaw, A.M., Behn, M.D., Humphris, S.E., Sohn, R.A., Gregg, P.M., 2010. Deep pooling of low degree melts and volatile fluxes at the 85°E segment of the Gakkel Ridge: Evidence from olivine-hosted melt inclusions and glasses. *Earth and Planet. Sci. Lett.* 289, 311–322.
- Shaw, A.M., Hauri, E.H., Fischer, T.P., Hilton, D.R., Kelley, K.A., 2008. Hydrogen isotopes in Mariana arc melt inclusions: Implications for subduction dehydration and the deep-Earth water cycle. *Earth and Planet. Sci. Lett.* 275, 138–145.
- Shimizu, K., Alexander, C.M.O'D., Hauri, E.H., Sarafian, A.R., Nittler, L.R., Wang, et al., 2021. Highly volatile element (H, C, F, Cl, S) abundances and H isotopic compositions in chondrules from carbonaceous and ordinary chondrites *Geochim. Cosmochim. Acta*, 301, 230–258.
- Shimizu, K., Saal, A.E., Myers, C.E., Nagle, A.N., Hauri, E.H., Forsyth, D.W., Kamenetsky, V.S., Niu, Y., 2016. Two-component mantle melting-mixing model for the generation of mid-ocean ridge basalts: Implications for the volatile content of the Pacific upper mantle. *Geochim. Cosmochim. Acta* 176, 44–80.

- Shimizu, K., Ushikubo, T., Hamada, M., Itoh, S., Higashi, Y., Takahashi, E., et al., 2017. H<sub>2</sub>O, CO<sub>2</sub>, F, S, Cl, and P<sub>2</sub>O<sub>5</sub> analyses of silicate glasses using SIMS: Report of volatile standard glasses. *Geochem. J.*, 51(4), 299-313.
- Shishkina, T.A., Botcharnikov, R.E., Holtz, F., Almeev, R.R., Jazwa, A.M., Jakubiak, A.A., 2014. Compositional and pressure effects on the solubility of H<sub>2</sub>O and CO<sub>2</sub> in mafic melts. *Chemical Geology* 388, 112-129.
- Sides, I.R., Edmonds, M., MacLennan, J., Swanson, D.A., Houghton, B.F., 2014. Eruption style at Kīlauea Volcano in Hawai‘i linked to primary melt composition. *Nature Geosciences* 7, 464–469.
- Spilliaert, N., Allard, P., Métrich, N., Sobolev, A.V., 2006. Melt inclusion record of the conditions of ascent, degassing, and extrusion of volatile-rich alkali basalt during the powerful 2002 flank eruption of Mount Etna (Italy). *J. Geophys. Res.* 111, B04203.
- Sorby, H.C., 1858. On the microscopical structure of crystals, indicating the origin of minerals and rocks. *Q. J. Geol. Soc.* 14, 453–500.
- Sun, C., Dasgupta, R., 2020. Thermobarometry of CO<sub>2</sub>-rich, silica-undersaturated melts constrains cratonic lithosphere thinning through time in areas of kimberlitic magmatism. *Earth Planet. Sci. Lett.* 550, 116549.
- Sun, C. and Dasgupta, R., 2023. Carbon budget of Earth’s deep mantle constrained by petrogenesis of silica-poor ocean island basalts. *Earth and Planet. Sci. Lett.* 611, 118135.
- Sun, S.S. and McDonough, W.F. 1989. Chemical and Isotopic Systematics of Oceanic Basalts: Implications for Mantle Composition and Processes. In: Saunders, A.D., Norry, M.J., Eds., *Magmatism in the Ocean Basins*, Geological Society, London, Special Publications, 42, 313-345.
- Taracsák, Z., Hartley, M.E., Burgess, R., Edmonds, M., Iddon, F., Longpré, M.-A., 2019. High fluxes of deep volatiles from ocean island volcanoes: Insights from El Hierro, Canary Islands. *Geochim Cosmochim. Acta* 258, 19–36.
- Toplis, M.J., 2005. The thermodynamics of iron and magnesium partitioning between olivine and liquid: criteria for assessing and predicting equilibrium in natural and experimental systems. *Contrib. Mineral. Petrol.* 149, 22–39.
- Torres, P., Silva, L.C., Munhá, J., Caldeira, R., Mata, J., Tassinari, C., 2010. Petrology and geochemistry of lavas from Sal Island: Implication for the variability of the Cape Verde magmatism. *Comunicações Geológicas* 97, 35–62.
- Tucker, J.M., Hauri, E.H., Pietruszka, A.J., Garcia, M.O., Marske, J.P., Trusdell, F.A., 2019. A high carbon content of the Hawaiian mantle from olivine-hosted melt inclusions. *Geochim Cosmochim. Acta* 254, 156–172.
- Van Achterbergh, E., Ryan, C.G., Jackson, S.E., Griffin, W.L., 2001. Data reduction software for LA-ICP-MS. *Laser-Ablation-ICPMS in the earth sciences—principles and applications Miner. Assoc. Can. (short course series)* 29, 239-243.

- Vigouroux, N., Wallace, P.J., Kent, A.J.R., 2008. Volatiles in High-K Magmas from the Western Trans-Mexican Volcanic Belt: Evidence for Fluid Fluxing and Extreme Enrichment of the Mantle Wedge by Subduction Processes. *J. Petrol.* 49, 1589–1618.
- Wallace, P. J. 2005. Volatiles in subduction zone magmas: concentrations and fluxes based on melt inclusions and volcanic gas data. *J. Volcanol. Geotherm. Res.* 140, 217–240.
- Wallace, P.J., Kamenetsky, V.S., Cervantes, P., 2015. Melt inclusion CO<sub>2</sub> contents, pressures of olivine crystallization, and the problem of shrinkage bubbles. *Am. Mineral.* 100, 787–794.
- Wallace, P.J., Plan, T., Bodnar, R.J., Gaetani, G.A., Shea, T., 2021. Olivine-hosted melt inclusions: a microscopic perspective on a complex magmatic world. *Annu. Rev. Earth Planet. Sci.* 49, 465–494.
- Wang, X., Chou, I.-M., Hu, W., Burruss, R.C., Sun, Q., Song, Y., 2011. Raman spectroscopic measurements of CO<sub>2</sub> density: Experimental calibration with high-pressure optical cell (HPOC) and fused silica capillary capsule (FSCC) with application to fluid inclusion observations. *Geochim Cosmochim. Acta* 75, 4080–4093.
- Wanless, V.D., Behn, M.D., Shaw, A.M., Plank, T., 2014. Variations in melting dynamics and mantle compositions along the Eastern Volcanic Zone of the Gakkel Ridge: insights from olivine-hosted melt inclusions. *Contrib. Mineral. Petrol.* 167, 1005.
- Wanless, V.D., Shaw, A.M., Behn, M.D., Soule, S.A., Escartín, J., Hamelin, C., 2015. Magmatic plumbing at Lucky Strike volcano based on olivine-hosted melt inclusion compositions. *Geochem. Geophys. Geosyst.* 16, 126–147.
- Weaver, B.L., 1991. The origin of ocean island basalt end-member compositions: trace element and isotopic constraints. *Earth and Planet. Sci. Lett.* 104, 381–397.
- Webster, J.D., Baker, D.R., Aiuppa, A., 2018. Halogens in mafic and intermediate-silica content magmas. In: Harlov, D.E., Aranovich, L. (Eds.), *The Role of Halogens in Terrestrial and Extraterrestrial Geochemical Processes: Surface, Crust, and Mantle*. Springer International Publishing, Cham, 307–430.
- Webster, J.D., Rebbert, C.R., 2001. The geochemical signature of fluid-saturated magma determined from silicate melt inclusions in Ascension Island granite xenoliths. *Geochim. Cosmochim. Acta* 65, 123–136.
- Wedepohl, K.H., and Hartmann, G., 1994. The composition of the primitive upper earth's mantle. In: Kimberlites, related rocks and mantle xenoliths. In: Meyer, H.O.A. and Leonardos, O.H. (Eds.), *Companhia de Pesquisa de Recursos Minerais*, 1, 486-495. Rio de Janeiro.
- Willbold, M., Stracke, A., 2006. Trace element composition of mantle end-members: Implications for recycling of oceanic and upper and lower continental crust. *Geochem. Geophys. Geosyst.* 7, 2005GC001005.
- Zanon, V., Silva, R., & Goulart, C. (2023). The crust-mantle transition beneath the Azores region (central-north Atlantic Ocean). *Contrib. Mineral. Petrol.* 178 (8), 50.

**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proofs