



**HAL**  
open science

## Success and Failure of Shock Recovery Experiments for Simulating Short Duration Parent Body Heating

E Quirico, H Yabuta, P Beck, L Bonal, A Bardyn, L Nittler, Conel M. O'd. Alexander, Nathalie Carrasco, Cyril Szopa

► **To cite this version:**

E Quirico, H Yabuta, P Beck, L Bonal, A Bardyn, et al.. Success and Failure of Shock Recovery Experiments for Simulating Short Duration Parent Body Heating. 82nd Annual Meeting of The Meteoritical Society 2019, Jul 2019, Sapporo, Japan. pp.LPI Contribution No. 2157, 2019, id.6343. insu-04441583

**HAL Id: insu-04441583**

**<https://insu.hal.science/insu-04441583v1>**

Submitted on 6 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## SUCCESS AND FAILURE OF SCHOCK RECOVERY EXPERIMENTS FOR SIMULATING SHORT DURATION PARENT BODY HEATING

E. Quirico<sup>1</sup>, H. Yabuta<sup>2</sup>, P. Beck<sup>1</sup>, L. Bonal<sup>1</sup>, A. Bardyn<sup>3</sup>, L. Nittler<sup>3</sup>, C.M.O'D. Alexander<sup>3</sup>, N. Carrasco<sup>4</sup>, C. Szopa<sup>4</sup>. <sup>1</sup>UGA/CNRS IPAG Grenoble France, <sup>2</sup>Hiroshima University, Japan. <sup>3</sup>DTM, Carnegie Institution USA. <sup>4</sup>CNRS/INSU LATMOS-IPSL, Guyancourt France

**Introduction:** Short duration heating is a post-accretional process that has effected the chondrite parent bodies to varying degrees, and that might be involved in the formation of Ultra Carbonaceous Antarctic Micro-Meteorites (UCAMMs)[1,2]. Hypervelocity impacts are a plausible source of heating, owing to the fact that the chondrites were excavated from their parent bodies by collisions. Here, we report shock recovery experiments that are dedicated to simulating these short duration heating processes. Three samples were studied: Murchison (CM2), EET 90628 (LL/3.0) and a tholin sample as a presumed analog precursor of some N-rich UCAMMs [3].

**Experiments:** Shock recovery experiments were conducted at Tokyo Institute of Technology (Japan) during four sessions between 2010 and 2016. A 5-50 GPa peak pressure range in the recovery capsule was achieved by combining different flyers with different velocities. The shocked state in the sample was assumed to equilibrate with that of the capsule through multiple reflections at the capsule boundaries. Peak pressures were calculated with the impedance match method, using the Hugoniot curves of iron and of the flyer plate material. Samples were characterized by FTIR and Raman spectroscopy (IPAG Grenoble and LGL Lyon, France) and SIMS (Carnegie Institute, Washington USA). Tholin samples were produced with a cold plasma reactor (LATMOS, France).

### Results and discussion:

*Murchison.* The IOM composition is very sensitive to shocks, even at low peak pressure (5 GPa). Thanks to FTIR analysis, we observe an increase of the CH<sub>2</sub>/CH<sub>3</sub> ratio (estimated from the peak intensity ratio of the antisymmetric stretching modes at 2925 and 2955 cm<sup>-1</sup>), a decrease of the carbonyl C=O group (~1700 cm<sup>-1</sup>), a decrease of the aliphatic (CH, CH<sub>2</sub>, CH<sub>3</sub>) abundance (except at 50 GPa, for which a large increase is observed), and the development of a new narrow band centered at ~ 1650 cm<sup>-1</sup>. At high shock pressure, large chemical variations are observed within the IOM, pointing to strong spatial heterogeneities of pressure and temperature in the capsule. Overall, these variations are consistent with those measured in heated chondrites. A high CH<sub>2</sub>/CH<sub>3</sub> ratio is always associated evidence for thermal events in type 2 and 3 chondrites, as well as the decrease of the aliphatic abundance. Note, however, the exception of the CM QUE 93005 that shows a dramatic increase of aliphatic groups that might be consistent with the 50 GPa experiment. The band at ~ 1650 cm<sup>-1</sup> has never been detected in C1, C2 or type 3 chondrites. This feature is possibly due to the formation of quinones, perhaps as a result of oxidation during the shock event. The gun is maintained under a primary vacuum, possibly favouring oxidation reactions that might not happen in natural conditions. The carbon structure probed by Raman spectroscopy (514 nm) displays moderate variations, which are more significant at 40 and 50 GPa. The data account for stage II heated chondrites, but not for III/IV stages [4]. In this respect, our shock experiments do not properly mimic the highest degrees of the heating process, possibly because the reverberation process to reach the peak pressure in the experiment releases less heat than a single shock process that happens in natural conditions [5]. SIMS measurements report a dramatic effect on the D/H composition of the IOM. At 5 GPa, the D/H ratio is drastically reduced, and at 40 GPa the IOM is isotopically normal. These findings are basically consistent with IOM compositions from heated C2 chondrites [6].

*EET 90628.* The Raman data of EET 90628 IOM display co-variations of the FWHM-D and I<sub>D</sub>/I<sub>G</sub> parameters pointing to a structural grade that is more consistent with a ~ 3.1 petrologic type. At the same time, we observe an increase of FWHM-G and a decrease of ω<sub>G</sub>. These results show that some primitive type 3 chondrites may have been perturbed by shock processes, and are also consistent with the fact that FWHM-G and ω<sub>G</sub> are not reliable tracers of thermal metamorphism.

*Tholin 97:3.* Shock experiments on a 97:3 tholin show major effects on the composition, and the development of a polyaromatic structure at 20 GPa and beyond. However, even at 40 GPa, the degree of structural order remains lower than that measured in N-rich UCAMMs. We may expect that shocks under natural conditions should lead to more mature carbonaceous materials. However, the D/H sensitivity to thermal processing is inconsistent with hypervelocity impact involvement in UCAMM formation (provided that D carriers are similar in Murchison and UCAMMs).

**References:** [1] Tonui E. et al. (2014) *Geochimica Cosmochimica Acta* 126:284–306. [2] Augé B. et al. (2016) *A&A* 592:A99. [3] Bonnet J-Y et al. (2015) *Icarus* 250: 53-63. [4] Quirico E. et al. (2018) *Geochimica Cosmochimica Acta* 126:284–306 241:17-37. [5] Tomeoka K. et al. (1999) *Geochimica Cosmochimica Acta* 63:3683-3703. [6] Alexander C.M.O'D et al. (2010) *Geochimica Cosmochimica Acta* 74:4417-4437